# Generating Function Associated with the Determinant Formula for the Solutions of the Painlevé II Equation

Joshi, Nalini School of Mathematics and Statistics F07, University of Sydney

Kajiwara, Kenji Faculty of Mathematics, Kyushu University

Mazzocco, Marta DPMMS

https://hdl.handle.net/2324/11833

出版情報:Astérisque. 297, pp.67-78, 2004-12. Société mathématique de France バージョン: 権利関係:

## **MHF** Preprint Series

Kyushu University 21st Century COE Program Development of Dynamic Mathematics with High Functionality

# Generating function associated with the determinant formula for the solutions of the Painlevé II equation

# N. Joshi, K. Kajiwara M. Mazzocco

## MHF 2004-14

(Received April 23, 2004)

Faculty of Mathematics Kyushu University Fukuoka, JAPAN

# Generating Function Associated with the Determinant Formula for the Solutions of the Painlevé II Equation

Nalini JOSHI

School of Mathematics and Statistics F07, University of Sydney, Sydney, NSW 2006, Australia

Kenji KAJIWARA

Graduate School of Mathematics, Kyushu University, 6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8512, Japan

Marta MAZZOCCO

DPMMS, Wilberforce Road, Cambridge CB3 0UB, UK

#### Abstract

In this paper we consider a Hankel determinant formula for generic solutions of the Painlevé II equation. We show that the generating functions for the entries of the Hankel determinants are related to the asymptotic solution at infinity of the linear problem of which the Painlevé II equation describes the isomonodromic deformations.

### **1** Introduction

The Painlevé II equation (PII),

$$\frac{d^2u}{dx^2} = 2u^3 - 4xu + 4\left(\alpha + \frac{1}{2}\right),\tag{1}$$

where  $\alpha$  is a parameter, is one of the most important equations in the theory of nonlinear integrable systems. It is well-known that P<sub>II</sub> admits unique rational solution when  $\alpha$  is a half-integer, and one-parameter family of solutions expressible in terms of the solutions of the Airy equation when  $\alpha$  is an integer. Otherwise the solution is non-classical [12, 13, 15].

The rational solutions for  $P_{II}(1)$  are expressed as logarithmic derivative of the ratio of certain special polynomials, which are called the "Yablonski-Vorob'ev polynomials", [16, 17]. Yablonski-Vorob'ev polynomials admit two determinant formulas, namely, Jacobi-Trudi type and Hankel type. The latter is described as follows: For each positive integer N, the unique rational solution for  $\alpha = N + 1/2$  is given by

$$u = \frac{d}{dx} \log \frac{\sigma_{N+1}}{\sigma_N},$$

where  $\sigma_N$  is the Hankel determinant

$$\sigma_N = \begin{vmatrix} a_0 & a_1 & \cdots & a_{N-1} \\ a_1 & a_2 & \cdots & a_N \\ \vdots & \vdots & \ddots & \vdots \\ a_{N-1} & a_N & \cdots & a_{2N-2} \end{vmatrix}$$

with  $a_n = a_n(x)$  being polynomials defined by the recurrence relation

$$a_{0} = x, \qquad a_{1} = 1,$$

$$a_{n+1} = \frac{da_{n}}{dx} + \sum_{k=0}^{n-1} a_{k} a_{n-1-k}.$$
(2)

The Jacobi-Trudi type formula implies that the Yablonski-Vorob'ev polynomials are nothing but the specialization of the Schur functions [9]. Then, what does the Hankel determinant formula mean? In order to answer this question, a generating function for  $a_n$  is constructed in [5]:

**Theorem 1.1** [5] Let  $\theta(x, t)$  be an entire function of two variables defined by

$$\theta(x,t) = \exp\left(2t^3/3\right) \operatorname{Ai}(t^2 - x),\tag{3}$$

where Ai(z) is the Airy function. Then there exists an asymptotic expansion

$$\frac{\partial}{\partial t}\log\theta(x,t) \sim \sum_{n=0}^{\infty} a_n(x) \, (-2t)^{-n},\tag{4}$$

as  $t \to \infty$  in any proper subsector of the sector  $|\arg t| < \pi/2$ .

This result is quite suggestive, because it shows that the Airy functions enter twice in the theory of classical solutions of the  $P_{II}$ :

(i) in the formula [3]

$$u = \frac{d}{dx} \log \operatorname{Ai}\left(2^{1/3}x\right), \qquad \alpha = 0.$$

the one parameter family of classical solutions of  $P_{II}$  for integer values of  $\alpha$  is expressed by Airy functions,

(ii) in formulae (3), (4) the Airy functions generate the entries of determinant formula for the rational solutions.

In this paper we clarify the nature of this phenomenon. First, we reformulate the Hankel determinant formula for generic, namely non-classical, solutions of  $P_{II}$  already found in [10, 11]. We next construct generating functions for the entries of our Hankel determinant formula. We then show that the generating functions are related to the asymptotic solution at infinity of the isomonodromic problem introduced by Jimbo and Miwa [6].

This result explains the appearance of the Airy functions in Theorem 1.1. In fact, for rational solutions of  $P_{II}$ , the asymptotic solution at infinity of the isomonodromic problem is indeed constructed in terms of Airy functions [7, 8, 14].

We expect that the generic solutions of the so-called Painlevé II hierarchy [1, 2, 4] should be expressed by some Hankel determinant formula. Of course the generating functions for the entries of Hankel determinant should be related to the asymptotic solution at infinity of the isomonodromic problem for the Painlevé II hierarchy. We also expect that the similar phenomena can be seen for other Painlevé equations. We shall discuss these generalizations in future publications.

Acknowledgments The authors thank Prof. H. Sakai for informing them of references [7, 8]. They also thank Prof. K. Okamoto for discussions and encouragement. M.M. acknowledges the support from the Engineering and Physical Sciences Research Council Fellowship #GR/S48424/01. K.K. acknowledges the support from the scientist exchange program between Japan Society for the Promotion of Science and Australian Academy of Science #0301002 and the JSPS Grant-in-Aid for Scientific Research (B) #15340057.

### 2 Hankel Determinant Formula and Isomonodromy Problem

#### 2.1 Hankel Determinant Formula

We first prepare the Hankel determinant formula for generic solutions for  $P_{II}$  (1). To show the parameter dependence explicitly, we denote equation (1) as  $P_{II}[\alpha]$ . The formula is based on the fact that the  $\tau$  functions for  $P_{II}$  satisfy the Toda equation,

$$\sigma_n''\sigma_n - (\sigma_n')^2 = \sigma_{n+1}\sigma_{n-1}, \quad n \in \mathbb{Z}, \quad ' = d/dx.$$
(5)

Putting  $\tau_n = \sigma_n / \sigma_0$  so that the  $\tau$  function is normalized as  $\tau_0 = 1$ , equation (5) is rewritten as

$$\tau_{n}^{\prime\prime}\tau_{n} - (\tau_{n}^{\prime})^{2} = \tau_{n+1}\tau_{n-1} - \varphi\psi\tau_{n}^{2}, \quad \tau_{-1} = \psi, \quad \tau_{0} = 1, \quad \tau_{1} = \varphi, \quad n \in \mathbb{Z}.$$
 (6)

Then it is known that  $\tau_n$  can be written in terms of Hankel determinant as follows [11]:

**Proposition 2.1** Let  $\{a_n\}_{n \in \mathbb{N}}$ ,  $\{b_n\}_{n \in \mathbb{N}}$  be the sequences defined recursively as

$$a_n = a'_{n-1} + \psi \sum_{\substack{i+j=n-2\\i,j\ge 0}} a_i a_j, \quad b_n = b'_{n-1} + \varphi \sum_{\substack{i+j=n-2\\i,j\ge 0}} b_i b_j, \quad a_0 = \varphi, \quad b_0 = \psi.$$
(7)

For any  $N \in \mathbb{Z}$ , we define Hankel determinant  $\tau_N$  by

$$\tau_N = \begin{cases} \det(a_{i+j-2})_{i,j \le N} & N > 0, \\ 1, & N = 0, \\ \det(b_{i+j-2})_{i,j \le |N|} & N < 0. \end{cases}$$
(8)

#### Then $\tau_N$ satisfies equation (6).

Since the above formula involves two arbitrary functions  $\varphi$  and  $\psi$ , it can be regarded as the determinant formula for general solution of the Toda equation. Imposing appropriate conditions on  $\varphi$  and  $\psi$ , we obtain determinant formula for the solutions of P<sub>II</sub>:

**Proposition 2.2** Let  $\psi$  and  $\varphi$  be functions in x satisfying

$$\frac{\psi''}{\psi} = \frac{\varphi''}{\varphi} = -2\psi\varphi + 2x,\tag{9}$$

$$\varphi'\psi - \varphi\psi' = 2\alpha,\tag{10}$$

Then we have the following:

- (i)  $u_0 = (\log \varphi)'$  satisfies  $P_{\text{II}}[\alpha]$ .
- (*ii*)  $u_{-1} = -(\log \psi)'$  satisfies  $P_{\text{II}}[\alpha 1]$ .
- (iii)  $u_N = \left(\log \frac{\tau_{N+1}}{\tau_N}\right)'$ , where  $\tau_N$  is defined by equation (8), satisfies  $P_{\text{II}}[\alpha + N]$ .

**Proof.** (i) and (ii) can be directly checked by using the relations (9) and (10). Then (iii) is the reformulation of Theorem 4.2 in [10].

#### 2.2 Riccati Equations for Generating Functions

Consider the generating functions for the entries as the following formal series

$$F_{\infty}(x,t) = \sum_{n=0}^{\infty} a_n(x) t^{-n}, \quad G_{\infty}(x,t) = \sum_{n=0}^{\infty} b_n(x) t^{-n}.$$
 (11)

It follows from the recursion relations (7) that the generating functions formally satisfy the Riccati equations. In fact, multiplying the recursion relations (7) by  $t^{-n}$  and take the summation from n = 0 to  $\infty$ , we have:

**Proposition 2.3** The generating functions  $F_{\infty}(x,t)$  and  $G_{\infty}(x,t)$  formally satisfy the Riccati equations

$$t\frac{\partial F}{\partial x} = -\psi F^2 + t^2 F - t^2 \varphi, \tag{12}$$

$$t\frac{\partial G}{\partial x} = -\varphi G^2 + t^2 G - t^2 \psi, \tag{13}$$

respectively.

Since  $F_{\infty}$  and  $G_{\infty}$  are defined as the formal power series around  $t \sim \infty$ , it is convenient to derive the differential equations with respect to t. In order to do this, the following auxiliary recursion relations are useful.

**Lemma 2.4** Under the condition (9) and (10),  $a_n$  and  $b_n$  ( $n \ge 0$ ) satisfy the following recursion relations,

$$(\psi a_{n+2} - \psi' a_{n+1})' = 2(n+1)\psi a_n, \tag{14}$$

$$(\varphi b_{n+2} - \varphi' b_{n+1})' = 2(n+1)\varphi b_n, \tag{15}$$

respectively.

We omit the details of the proof of Lemma 2.4, because it is proved by straight but tedious induction. Multiplying equations (14) and (15) by  $t^{-n}$  and taking summation over n = 0 to  $\infty$ , we have the following differential equations for  $F_{\infty}$  and  $G_{\infty}$ :

**Lemma 2.5** The generating functions  $F_{\infty}$  and  $G_{\infty}$  formally satisfy the following differential equations,

$$2\psi t \frac{\partial F}{\partial t} = t(\psi' - t\psi) \frac{\partial F}{\partial x} + (\psi'' t - \psi' t^2 + 2\psi)F + t^2(\psi\varphi' + \psi'\varphi), \tag{16}$$

$$2\varphi t \frac{\partial G}{\partial t} = t(\varphi' - t\varphi) \frac{\partial G}{\partial x} + (\varphi'' t - \varphi' t^2 + 2\varphi)G + t^2(\psi\varphi' + \psi'\varphi), \tag{17}$$

respectively.

Eliminating x-derivatives from equations (12), (16), and equations (13), (17), respectively, we obtain the Riccati equations with respect to t:

**Proposition 2.6** The generating functions  $F_{\infty}$  and  $G_{\infty}$  formally satisfy the following Riccati equations,

$$2t\frac{\partial F}{\partial t} = -(\psi' - t\psi)F^2 + \left(\frac{\psi''}{\psi}t + 2 - t^3\right)F + t^2(\varphi' + t\varphi),\tag{18}$$

$$2t\frac{\partial G}{\partial t} = -(\varphi' - t\varphi)G^2 + \left(\frac{\varphi''}{\varphi}t + 2 - t^3\right)G + t^2(\psi' + t\psi),\tag{19}$$

respectively.

#### 2.3 Isomonodromic Problem

The Riccati equations for  $F_{\infty}$  equations (12) and (18) are linearized by standard technique, which yield isomonodromic problem for P<sub>II</sub>. It is easy to derive the following theorem from the Proposition 2.3 and 2.6:

**Theorem 2.7** (i) It is possible to introduce the functions  $Y_1(x, t)$ ,  $Y_2(x, t)$  consistently as

$$F_{\infty}(x,t) = \frac{t}{\psi} \left( \frac{1}{Y_1} \frac{\partial Y_1}{\partial x} + \frac{t}{2} \right) = \frac{2t}{\psi' - t\psi} \left[ \frac{1}{Y_1} \frac{\partial Y_1}{\partial t} + \frac{1}{4} \left( \frac{\psi''}{\psi} - t^2 \right) \right],\tag{20}$$

$$Y_2 = \frac{1}{\psi} \left( \frac{\partial Y_1}{\partial x} + \frac{tY_1}{2} \right).$$
(21)

Then  $Y_1$  and  $Y_2$  satisfy the following linear system for  $Y = \begin{pmatrix} Y_1 \\ Y_2 \end{pmatrix}$ :

$$\frac{\partial}{\partial t}Y = AY, \quad A = \begin{pmatrix} \frac{t^2}{4} - \frac{z}{2} - \frac{x}{2} & -\frac{\psi}{2}(t+u_{-1}) \\ \frac{1}{\psi}\left\{(u_{-1}-t)\frac{z}{2} + \alpha\right\} & -\frac{t^2}{4} + \frac{z}{2} + \frac{x}{2} \end{pmatrix}, \tag{22}$$

$$\frac{\partial}{\partial x}Y = BY, \quad B = \begin{pmatrix} -\frac{t}{2} & \psi \\ \frac{z}{\psi} & \frac{t}{2} \end{pmatrix},$$
(23)

where  $z = -\psi\varphi$ .

#### (ii) Similarly, it is possible to introduce the functions $Z_1(x,t)$ , $Z_2(x,t)$ consistently as

$$G_{\infty}(x,t) = \frac{t}{\varphi} \left( \frac{1}{Z_1} \frac{\partial Z_1}{\partial x} + \frac{t}{2} \right) = \frac{2t}{\varphi' - t\varphi} \left[ \frac{1}{Z_1} \frac{\partial Z_1}{\partial t} + \frac{1}{4} \left( \frac{\varphi''}{\varphi} - t^2 \right) \right],$$
(24)

$$Z_2 = \frac{1}{\varphi} \left( \frac{\partial Y_1}{\partial x} + \frac{tY_1}{2} \right).$$
(25)

Then  $Z_1$  and  $Z_2$  satisfy the following linear system for  $Z = \begin{pmatrix} Z_1 \\ Z_2 \end{pmatrix}$ :

$$\frac{\partial}{\partial t}Z = CZ, \quad C = \begin{pmatrix} \frac{t^2}{4} - \frac{z}{2} - \frac{x}{2} & -\frac{\varphi}{2}(t - u_0) \\ -\frac{1}{\varphi}\left\{(u_0 + t)\frac{z}{2} + \alpha\right\} & -\frac{t^2}{4} + \frac{z}{2} + \frac{x}{2} \end{pmatrix},$$
(26)

$$\frac{\partial}{\partial x}Z = DY, \quad D = \begin{pmatrix} -\frac{t}{2} & \varphi \\ \frac{z}{\varphi} & \frac{t}{2} \end{pmatrix}.$$
(27)

**Remark 2.8** The linear systems (22), (23) and (26), (27) are the isomonodromic problems for  $P_{\text{II}}[\alpha - 1]$  and  $P_{\text{II}}[\alpha]$ , respectively [6]. For example, compatibility condition for equations (22) and (23), namely,

$$\frac{\partial A}{\partial x} - \frac{\partial B}{\partial t} + [A, B] = 0$$

gives

$$\begin{cases} \frac{dz}{dx} = -2u_{-1}z - 2\alpha, \\ \frac{du_{-1}}{dx} = u_{-1}^2 - 2z - 2x, \\ u_{-1} = -\frac{1}{\psi}\frac{d\psi}{dx}, \end{cases}$$
(28)

which yields  $P_{\text{II}}[\alpha - 1]$  for  $u_{-1}$ . This fact also guarantees the consistency of two expressions for  $F_{\infty}$  in terms of  $Y_1$  in equation (20). Similar remark holds true for  $G_{\infty}$  and  $Z_1$ .

**Remark 2.9**  $F_{\infty}$  and  $G_{\infty}$  are also expressed as,

$$F_{\infty} = t \frac{Y_2}{Y_1}, \quad G_{\infty} = t \frac{Z_2}{Z_1},$$
 (29)

respectively. Conversely, it is obvious that for any solution  $Y_1$  and  $Y_2$  for the linear system (22) and (23),  $F = tY_2/Y_1$  satisfies the Riccati equations (12) and (18) (Similar for G).

**Remark 2.10** Theorem 1.1 is recovered by putting  $\psi = 1$ ,  $\varphi = x$ .

**Remark 2.11**  $Y_1$  can be formally expressed in terms of  $a_n$  by using equation (20) as

$$Y_1 = \text{const.} \times \exp\left(\frac{1}{12}t^3 - \frac{x}{2}t\right) t^{-\alpha} \exp\left[\frac{1}{2}\sum_{n=1}^{\infty}\frac{\psi a_{n+1} - \psi' a_n}{n} t^{-n}\right].$$
 (30)

which coincides with known asymptotic behavior of  $Y_1$  around  $t \sim \infty$  [6].

## 3 Solutions of Isomonodromic Problems and Determinant Formula

In the previous section we have shown that the generating functions  $F_{\infty}$  and  $G_{\infty}$  formally satisfy the Riccati equations (12,18) and (13,19), and that their linearization yield isomonodromic problems (22, 23) and (26,27) for P<sub>II</sub>. Now let us consider the converse. We start from the linear system (22) and (23). We have two linearly independent solutions around  $t \sim \infty$ , one of which is related with  $F_{\infty}(x, t)$ . Then, what is another solution? In fact, it is well-known that linear system (22) and (23) admit the formal solutions around  $t \sim \infty$  of the form [6],

$$Y^{(1)} = \begin{pmatrix} Y_1^{(1)} \\ Y_2^{(1)} \end{pmatrix} = \exp\left(\frac{t^3}{12} - \frac{xt}{2}\right) t^{-\alpha} \left[ \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} y_{11}^{(1)} \\ y_{21}^{(1)} \end{pmatrix} t^{-1} + \begin{pmatrix} y_{12}^{(1)} \\ y_{22}^{(1)} \end{pmatrix} t^{-2} + \cdots \right], \quad (31)$$

$$Y^{(2)} = \begin{pmatrix} Y_1^{(2)} \\ Y_2^{(2)} \end{pmatrix} = \exp\left(-\frac{t^3}{12} + \frac{xt}{2}\right) t^{\alpha} \left[\begin{pmatrix} 0 \\ 1 \end{pmatrix} + \begin{pmatrix} y_{11}^{(2)} \\ y_{21}^{(2)} \end{pmatrix} t^{-1} + \begin{pmatrix} y_{12}^{(2)} \\ y_{22}^{(2)} \end{pmatrix} t^{-2} + \cdots \right].$$
(32)

From Remark 2.9 we see that there are two possible power-series solutions for the Riccati equation (18) of the form,

$$Y^{(1)} \to F^{(1)} = t \, \frac{Y_2^{(1)}}{Y_1^{(1)}} = t \, \frac{y_{21}^{(1)} t^{-1} + \dots}{1 + y_{11}^{(1)} t^{-1} + \dots} = c_0 + c_1 t^{-1} + \dots \,, \tag{33}$$

$$Y^{(2)} \to F^{(2)} = t \frac{Y_2^{(2)}}{Y_1^{(2)}} = t \frac{1 + y_{21}^{(2)} t^{-1} + \dots}{y_{11}^{(2)} t^{-1} + \dots} = t^2 (d_0 + d_1 t^{-1} + \dots).$$
(34)

The above two possibilities of power-series solutions for the Riccati equations are verified directly as follows:

**Proposition 3.1** *The Riccati equation (18) admits only the following two kinds of power-series solutions around*  $t \sim \infty$ :

$$F^{(1)} = \sum_{n=0}^{\infty} c_n t^{-n}, \quad F^{(2)} = t^2 \sum_{n=0}^{\infty} d_n t^{-n}.$$
(35)

Proof. We substitute the expression,

$$F = t^{\rho} \sum_{n=0}^{\infty} c_n t^{-n},$$
(36)

for some integer  $\rho$  to be determined, into the Riccati equation (18). We have:

$$\sum_{n=0}^{\infty} 2(\rho-n)c_n t^{\rho+1-n} = \sum_{n=0}^{\infty} \psi' \left(\sum_{k=0}^n c_k c_{n-k}\right) t^{2\rho-2n} - \sum_{n=0}^{\infty} \psi \left(\sum_{k=0}^n c_k c_{n-k}\right) t^{2\rho+1-2n} + \sum_{n=0}^{\infty} \left(\frac{\psi''}{\psi} + 2\right) c_n t^{\rho-n} - \sum_{n=0}^{\infty} c_n t^{\rho+3-n} + t^2(\varphi' + t\varphi)$$

The leading order should be one of  $t^{2\rho+1}$ ,  $t^{\rho+3}$  and  $t^3$ . Investigating the balance of these terms, we have  $\rho = 0$  or  $\rho = 2$ .

We also have the similar result for the solution of the Riccati equation (19):

**Proposition 3.2** The Riccati equation (19) admits only the following two kinds of power-series solutions around  $t \sim \infty$ :

$$G^{(1)} = \sum_{n=0}^{\infty} e_n t^{-n}, \quad G^{(2)} = t^2 \sum_{n=0}^{\infty} f_n t^{-n}.$$
(37)

It is obvious that  $F^{(1)}$  and  $G^{(1)}$  are nothing but our generating functions  $F_{\infty}$  and  $G_{\infty}$ , respectively. Then, what are  $F^{(2)}$  and  $G^{(2)}$ ? In the following, we present two observations regarding this point. First, there are unexpectedly simple relations among those functions:

Proposition 3.3 The following relations holds.

$$F^{(2)}(x,t) = \frac{t^2}{G^{(1)}(x,-t)}, \quad G^{(2)}(x,t) = \frac{t^2}{F^{(1)}(x,-t)}.$$
(38)

**Proof.** Substitute  $F(x,t) = \frac{t^2}{g(x,t)}$  into equation (18). This gives equation (19) for G(x,t) = g(x,-t) by using the relation (9). Choosing  $g(x,t) = G^{(1)}(x,t)$ , F(x,t) must be  $F^{(2)}(x,t)$ , since its leading order is  $t^2$ . We obtain the second equation by the similar argument.

Second,  $F^{(2)}(x,t)$  and  $G^{(2)}(x,t)$  are also interpreted as generating functions of entries of Hankel determinant formula for P<sub>II</sub>. Recall that the determinant formula in Proposition 2.1 is for the  $\tau$  sequence  $\tau_n = \sigma_n/\sigma_0$  so that it is normalized as  $\tau_0 = 1$ . We show that  $F^{(2)}(x,t)$  and  $G^{(2)}(x,t)$  correspond to different normalizations of  $\tau$  sequence:

Proposition 3.4 Let

$$F^{(2)}(x,t) = -\frac{t^2}{\psi^2} \sum_{n=0}^{\infty} d_n \ (-t)^{-n},\tag{39}$$

$$G^{(2)}(x,t) = -\frac{t^2}{\varphi^2} \sum_{n=0}^{\infty} f_n \ (-t)^{-n},$$
(40)

be formal solutions of the Riccati equations (12),(18) and (13), (19), respectively. We put

$$\kappa_{-n-1} = \det(d_{i+j})_{i,j=1,\dots,n} \quad (n>0), \quad \kappa_{-1} = 1,$$
(41)

$$\theta_{n+1} = \det(f_{i+j})_{i,j=1,\dots,n} \quad (n>0), \quad \theta_1 = 1.$$
(42)

Then  $\kappa_n$  and  $\theta_n$  are related to  $\tau_n$  as

$$\kappa_n = \frac{\tau_n}{\psi} = \frac{\tau_n}{\tau_{-1}} \quad (n < 0), \tag{43}$$

$$\theta_n = \frac{\tau_n}{\varphi} = \frac{\tau_n}{\tau_1} \quad (n > 0). \tag{44}$$

To prove Proposition 3.4, we first derive recurrence relations that characterize  $d_n$  and  $f_n$ . By substituting equations (39) and (40) into the Riccati equations (12) and (13), respectively, we easily obtain the following lemma:

**Lemma 3.5** (i)  $d_0$  and  $d_1$  are given by  $d_0 = -\psi$  and  $d_1 = \psi'$ , respectively. For  $n \ge 2$ ,  $d_n$  are characterized by the recursion relation,

$$d_n = d'_{n-1} + \frac{1}{\psi} \sum_{k=2}^{n-2} d_k d_{n-k}, \quad d_2 = \frac{\psi''\psi - (\psi')^2 + \varphi\psi^3}{\psi}.$$
 (45)

(ii)  $f_0$  and  $f_1$  are given by  $d_0 = -\varphi$  and  $d_1 = \varphi'$ , respectively. For  $n \ge 2$ ,  $f_n$  are characterized by the recursion relation,

$$f_n = f'_{n-1} + \frac{1}{\varphi} \sum_{k=2}^{n-2} f_k f_{n-k}, \quad f_2 = \frac{\varphi'' \varphi - (\varphi')^2 + \varphi^3 \psi}{\varphi}.$$
 (46)

Proof of Proposition 3.4. Consider the Toda equations (5) and (6). Let us put

$$\tilde{\tau}_n = \frac{\sigma_n}{\sigma_{-1}} = \frac{\tau_n}{\tau_{-1}} \tag{47}$$

so that  $\tilde{\tau}_{-1} = 1$ . Then it is easy to derive the Toda equation for  $\tilde{\tau}_n$ :

$$\tilde{\tau}_{n}^{\prime\prime}\tilde{\tau}_{n} - (\tilde{\tau}_{n}^{\prime})^{2} = \tilde{\tau}_{n+1}\tilde{\tau}_{n-1} - \frac{\psi^{\prime\prime}\psi - (\psi^{\prime})^{2} + \varphi\psi^{3}}{\psi^{2}}\tilde{\tau}_{n}^{2},$$
(48)

$$\tilde{\tau}_{-2} = \frac{\psi''\psi - (\psi')^2 + \varphi\psi^3}{\psi}, \quad \tilde{\tau}_{-1} = 1, \quad \tilde{\tau}_0 = \frac{1}{\psi}.$$
(49)

We have the determinant formula for  $\tilde{\tau}_n$  as,

$$\tilde{\tau}_n = \begin{cases} \det(\tilde{a}_{i+j-2})_{i,j \le n+1} & n > 0, \\ 1, & n = 0, \\ \det(\tilde{b}_{i+j-2})_{i,j \le |n|-1} & n < 0, \end{cases}$$
(50)

$$\tilde{a}_{n} = \tilde{a}_{n-1}' + \frac{\psi''\psi - (\psi')^{2} + \varphi\psi^{3}}{\psi} \sum_{\substack{i+j=n-2\\i,j>0}} \tilde{a}_{i}\tilde{a}_{j}, \quad \tilde{a}_{0} = \frac{1}{\psi},$$
(51)

$$\tilde{b}_n = \tilde{b}'_{n-1} + \frac{1}{\psi} \sum_{\substack{i+j=n-2\\i,j\ge 0}} \tilde{b}_i \tilde{b}_j, \quad \tilde{b}_0 = \frac{\psi''\psi - (\psi')^2 + \varphi\psi^3}{\psi}.$$
(52)

Now it is obvious from Lemma 3.5 that

$$d_j = \tilde{b}_{j-2} \quad (j \ge 2), \quad \kappa_n = \tilde{\tau}_n \quad (n < 0), \tag{53}$$

which proves equation (41). Equation (42) can be proved in similar manner.

We finally remark that the mysterious relations among the  $\tau$  functions and the solutions of isomonodromic problem in Proposition 3.3 and 3.4 should eventually originate from the symmetry of P<sub>II</sub>, but their meaning is not sufficiently understood yet.

### References

- M. J. Ablowitz and H. Segur. Exact linearization of a Painlevé transcendent. *Phys. Rev. Lett.*, 38:1103–1106, 1977.
- [2] H. Airault. Rational solutions of Painlevé equations. Studies in Applied Mathematics, 61:31-53, 1979.
- [3] N.P. Erugin. On the second transcendent of Painlevé. Dokl. Akad. Nauk BSSR 2:139–142, 1958.
- [4] H. Flaschka and A. C. Newell. Monodromy and Spectrum Preserving Deformations I. Comm. Math. Phys., 76:65–116, 1980.
- [5] K. Iwasaki, K. Kajiwara and T. Nakamura. Generating function associated with the rational solutions of the Painlevé II equation. J. Phys. A: Math. Gen 35:L207–L211, 2002.
- [6] M. Jimbo and T. Miwa, Monodoromy preserving deformation of linear ordinary differential equations with rational coefficients.II. *Physica* 2D:407–448, 1981.
- [7] M. Jimbo. Monodromy problem and the boundary condition for some Painlevé equations. *Publ. RIMS* 18:1137–1161, 1982.
- [8] M. Jimbo. Unpublished work.
- K. Kajiwara and Y. Ohta. Determinant structure of the rational solutions for the Painlevé II equation. J. Math. Phys., 37:4693–4704, 1996.
- [10] K. Kajiwara and T. Masuda. A generalization of the determinant formulae for the solutions of the Painlevé II equation. J. Phys. A: Math. Gen. 32:3763–3778, 1999.
- [11] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada. Determinant formulas for the Toda and discrete Toda equations. *Funkcial. Ekvac.* 44:291–307, 2001.
- [12] Y. Murata. Rational solutions of the second and the fourth Painlevé equations. *Funkcial. Ekvac.*, 28:1–32, 1985.
- [13] M. Noumi and K. Okamoto. Irreducibility of the second and the fourth Painlevé equations. *Funkcial. Ekvac.*, 40:139–163, 1997.
- [14] K. Okamoto. Private communication.
- [15] H. Umemura and H. Watanabe, Solutions of the second and fourth Painlevé equations. I. Nagoya Math. J., 148:151–198, 1997.
- [16] A.P. Vorob'ev. On the rational solutions of the second Painlevé equation. *Differencial'nye Uravnenija* 1:79– 81, 1965.
- [17] A.I. Yablonskii. Vesti Akad. Navuk. BSSR Ser. Fiz. Tkh. Nauk. 3:30-35, 1959.
- [18] W. Wasow. Asymptotic expansions for ordinary differential equations. Pure and Applied Mathematics, John Wiley Sons, Inc., XIV, 1965.

# List of MHF Preprint Series, Kyushu University 21st Century COE Program Development of Dynamic Mathematics with High Functionality

- MHF2003-1 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Yoshitaka WATANABE A numerical method to verify the invertibility of linear elliptic operators with applications to nonlinear problems
- MHF2003-2 Masahisa TABATA & Daisuke TAGAMI Error estimates of finite element methods for nonstationary thermal convection problems with temperature-dependent coefficients
- MHF2003-3 Tomohiro ANDO, Sadanori KONISHI & Seiya IMOTO Adaptive learning machines for nonlinear classification and Bayesian information criteria
- MHF2003-4 Kazuhiro YOKOYAMA On systems of algebraic equations with parametric exponents
- MHF2003-5 Masao ISHIKAWA & Masato WAKAYAMA Applications of Minor Summation Formulas III, Plücker relations, Lattice paths and Pfaffian identities
- MHF2003-6 Atsushi SUZUKI & Masahisa TABATA Finite element matrices in congruent subdomains and their effective use for large-scale computations
- MHF2003-7 Setsuo TANIGUCHI Stochastic oscillatory integrals - asymptotic and exact expressions for quadratic phase functions -
- MHF2003-8 Shoki MIYAMOTO & Atsushi YOSHIKAWA Computable sequences in the Sobolev spaces
- MHF2003-9 Toru FUJII & Takashi YANAGAWA Wavelet based estimate for non-linear and non-stationary auto-regressive model
- MHF2003-10 Atsushi YOSHIKAWA Maple and wave-front tracking — an experiment
- MHF2003-11 Masanobu KANEKO On the local factor of the zeta function of quadratic orders
- MHF2003-12 Hidefumi KAWASAKI Conjugate-set game for a nonlinear programming problem

- MHF2004-1 Koji YONEMOTO & Takashi YANAGAWA Estimating the Lyapunov exponent from chaotic time series with dynamic noise
- MHF2004-2 Rui YAMAGUCHI, Eiko TSUCHIYA & Tomoyuki HIGUCHI State space modeling approach to decompose daily sales of a restaurant into time-dependent multi-factors
- MHF2004-3 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA Cubic pencils and Painlevé Hamiltonians
- MHF2004-4 Atsushi KAWAGUCHI, Koji YONEMOTO & Takashi YANAGAWA Estimating the correlation dimension from a chaotic system with dynamic noise
- MHF2004-5 Atsushi KAWAGUCHI, Kentarou KITAMURA, Koji YONEMOTO, Takashi YANAGAWA & Kiyofumi YUMOTO Detection of auroral breakups using the correlation dimension
- MHF2004-6 Ryo IKOTA, Masayasu MIMURA & Tatsuyuki NAKAKI A methodology for numerical simulations to a singular limit
- MHF2004-7 Ryo IKOTA & Eiji YANAGIDA Stability of stationary interfaces of binary-tree type
- MHF2004-8 Yuko ARAKI, Sadanori KONISHI & Seiya IMOTO Functional discriminant analysis for gene expression data via radial basis expansion
- MHF2004-9 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA Hypergeometric solutions to the q Painlevé equations
- MHF2004-10 Raimundas VIDŪNAS Expressions for values of the gamma function
- MHF2004-11 Raimundas VIDŪNAS Transformations of Gauss hypergeometric functions
- MHF2004-12 Koji NAKAGAWA & Masakazu SUZUKI Mathematical knowledge browser
- MHF2004-13 Ken-ichi MARUNO, Wen-Xiu MA & Masayuki OIKAWA Generalized Casorati determinant and Positon-Negaton-Type solutions of the Toda lattice equation
- MHF2004-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO Generating function associated with the determinant formula for the solutions of the Painlevé II equation