Hypergeometric solutions to the q-Painlevé equation of type \$\${({A}_{1}+{A}_{1}^{prime })}^{left(1¥right)}\$\$

Hamamoto, Taro Faculty of Mathematics, Kyushu University

Kajiwara, Kenji Faculty of Mathematics, Kyushu University

Witte, Nicholas S. Department of Mathematics and Statistics, University of Melbourne

https://hdl.handle.net/2324/11868

出版情報:International Mathematics Research Notices. 2006, pp.84619-, 2006. Hindawi Publishing Corporation バージョン: 権利関係:The article has been accepted for publication in International Mathematics Research Notices (C): 2006 Published by Hindawi Publishing Corporation. All rights reserved.

MHF Preprint Series

Kyushu University 21st Century COE Program Development of Dynamic Mathematics with High Functionality

Hypergeometric Solutions to the q-Painlevé Equation of Type $(A_1 + A'_1)^{(1)}$

T. Hamamoto, K. Kajiwara N. S. Witte

MHF 2006-26

(Received July 28, 2006)

Faculty of Mathematics Kyushu University Fukuoka, JAPAN

Hypergeometric Solutions to the *q*-Painlevé Equation of Type $(A_1 + A'_1)^{(1)}$

Taro Hamamoto¹, Kenji Kajiwara¹ and Nicholas S. Witte²

¹ Graduate School of Mathematics, Kyushu University, 6-10-1 Hakozaki, Fukuoka 812-8581, Japan ² Department of Mathematics and Statistics, University of Melbourne, Victoria 3010, Australia

July 27, 2006

Abstract

A class of classical solutions to the *q*-Painlevé equation of type $(A_1 + A'_1)^{(1)}$ (a *q*-difference analog of the Painlevé II equation) is constructed in a determinantal form with basic hypergeometric function elements. The continuous limit of this *q*-Painlevé equation to the Painlevé II equation and its hypergeometric solutions are discussed. The continuous limit of these hypergeometric solutions to the Airy function is obtained through a uniform asymptotic expansion of their integral representation.

1 Introduction

In this article we consider the following q-difference equation

$$\left(\overline{F}F - 1\right)\left(F\underline{F} - 1\right) = \frac{at^2F}{F + t},\tag{1}$$

where t is the independent variable

$$\overline{t} = qt, \quad \underline{t} = t/q, \quad F = F(t), \quad \overline{F} = F(qt), \quad \underline{F} = F(t/q),$$
(2)

with |q| < 1 and *a* is a parameter. Equation (1) was identified as one of the *q*-difference Painlevé equations by Ramani and Grammaticos[26] with a continuous limit to the Painlevé II equation (P_{II}). In this sense eq.(1) is sometimes regarded as a *q*-analog of P_{II}. Sakai formulated the discrete Painlevé equations as Cremona transformations on a certain family of rational surfaces and developed their classification theory[30]. According to this theory eq.(1) is a discrete dynamical system on the rational surface characterized by the Dynkin diagram of type $A_6^{(1)}$, which posseses the symmetry of the affine Weyl group of type $(A_1 + A'_1)^{(1)}$. Equation (1) may be denoted as $dP(A_6^{(1)})$ by the notation adopted in [20]. Although eq.(1) is the simplest nontrivial *q*-difference Painlevé equation that admits a Bäcklund transformation only a few results are known - its continuous limit[26] and its simplest hypergeometric solution[28, 8, 9]. The first purpose of this article is to construct "higher-order" hypergeometric solutions to eq.(1) explicitly in determinantal form.

The second purpose of this article is to consider the continuous limit in some detail. The limiting procedure works well on the formal level of the defining *q*-difference equation however naïve application of the procedure does not work on the level of their solutions. The application of the continuous limit to the series representation of the basic hypergeometric functions that appear in the solutions does not yield the Airy functions which are the hypergeometric solutions of P_{II} . To obtain the valid limit we follow the procedure used by Prellberg[25] - we construct an appropriate integral representation of the function and derive an asymptotic expansion by applying a generalization of the saddle point method.

Our paper is organized as follows. In section 2 we construct hypergeometric solutions to eq.(1). The simplest solution is obtained in section 2.1, and determinant formula of "higher-order" solutions is presented in section 2.2, whose proof is given in section 2.3. In section 3 we consider the continuous limit as $q \rightarrow 1^-$. The limit on the formal level is discussed in section 3.1. We discuss the limit on the level of hypergeometric functions in section 3.2. Section 4 is devoted to concluding remarks.

2 Hypergeometric Solutions and Their Determinant Formula

2.1 The Simplest Solution

The simplest hypergeometric solution to eq.(1) is obtained by looking for the special case where it reduces to the Riccati equation. Then by linearizing the Riccati equation we obtain a second order linear q-differential equation, which admits basic hypergeometric functions as solutions.

Let us first recall the definition of the basic hypergeometric series[4]

$${}_{r}\varphi_{s}\left(\begin{array}{c}a_{1},\ldots,a_{r}\\b_{1},\ldots,b_{s}\end{array};q,z\right)=\sum_{n=0}^{\infty}\frac{(a_{1},\ldots,a_{r};q)_{n}}{(b_{1},\ldots,b_{s};q)_{n}(q;q)_{n}}\left[(-1)^{n}q^{\binom{n}{2}}\right]^{1+r-s}z^{n},$$
(3)

where

$$(a_1, \dots, a_r; q)_n = \prod_{i=1}^n (a_i; q)_n, \quad (a; q)_n = (1-a)(1-qa)\cdots(1-q^{n-1}a).$$
(4)

Then the simplest hypergeometric solution to eq.(1) is given as follows (see also [8, 9, 28]):

Lemma 2.1. *Eq.(1) admits the following particular solution for a = q:*

$$F = \frac{\overline{\psi}}{\psi},\tag{5}$$

where $\psi(t)$ satisfies the linear q-difference equation

$$\overline{\psi} + t\psi = \underline{\psi}.\tag{6}$$

The general solution of eq.(6) is given by

$$\psi(t) = A_{1}\varphi_{1} \begin{pmatrix} 0 \\ -q \end{pmatrix}; q, -qt + B e^{\pi i \frac{\log q}{\log q}} \varphi_{1} \begin{pmatrix} 0 \\ -q \end{bmatrix}; q, qt$$
(7)

where A and B are arbitrary q-periodic functions.

Proof. It is easy to see that if F is the solution of the Riccati equation

$$\overline{F} = \frac{1}{F} - qt,\tag{8}$$

then F satisfies eq.(1) with a = q. Equation (8) can be linearized via eq.(5) to eq.(6) by putting $F = \phi/\psi$ and equating the numerators and denominators of both sides.

We next substitute $\psi = t^{\rho} \sum_{n=0}^{\infty} a_n t^n$ into eq.(6). Then we have $q^{2\rho} = 1$, which implies $\rho = \frac{m\pi i}{\log q}$ $(m \in \mathbb{Z})$. Furthermore we deduce a recursion relation for a_n from eq.(6)

$$a_n = (-1)^m \frac{q^n}{1 - q^{2n}} a_{n-1}.$$

Accordingly we obtain two fundamental solutions to eq.(6) as

$$\psi_1(t) = \sum_{n=0}^{\infty} \frac{(-1)^n q^{n(n-1)/2}}{(q;q)_n (q;-q)_n} \ (-qt)^n, \quad \psi_2(t) = \mathrm{e}^{\pi i \frac{\log t}{\log q}} \sum_{n=0}^{\infty} \frac{(-1)^n q^{n(n-1)/2}}{(q;q)_n (q;-q)_n} \ (qt)^n,$$

for m = 0, 1 respectively. This completes the proof of Lemma 2.1. \Box

2.2 Bäcklund Transformation and Determinant Formula

Sakai constructed the following transformations for the homogeneous variables *x*, *y*, *z* and the parameters a_0 , a_1 , *b* on the $A_6^{(1)}$ type (*Mul.7*) surface¹:

$$\sigma: (a_1, a_0, b; x: y: z) \mapsto (a_0, a_1, a_1b; z(z+x): bx(z+x): yz)
\sigma_{(15)(24)(60)}: (a_1, a_0, b; x: y: z)
\mapsto (1/a_0, 1/a_1, b; z(x+z)(x+y+z): y((z+bx)(x+z)+yz): bx(x+z)^2)
w_1: (a_1, a_0, b; x: y: z) \mapsto (1/a_1, a_1^2a_0, a_1b; x(y+a_1z): y(y+z): a_1z(y+z))
w_0 = \sigma_{(15)(24)(60)}w_1\sigma_{(15)(24)(60)}.$$
(9)

The action of σ^2

$$\sigma^{2}: (a_{1}, a_{0}, b; x: y: z) \longmapsto (a_{1}, a_{0}, a_{0}a_{1}b; yz^{2}(x+y+z): a_{1}bz^{2}(x+y+z)(x+z): bxyz(x+z)).$$
(10)

gives rise to eq.(1) by putting

$$a = \left(\frac{a_0}{a_1}\right)^{1/2}, \quad t = \left(\frac{b^2}{a_0^3 a_1}\right)^{1/4}, \quad q = (a_0 a_1)^{1/2}, \qquad F = \left(\frac{a_0}{a_1 b^2}\right)^{1/4} \frac{y}{x}.$$
 (11)

Note that these transformations satisfy the fundamental relations,

$$w_0^2 = w_1^2 = 1$$
, $w_0 \sigma^2 = \sigma^2 w_0$, $w_1 \sigma^2 = \sigma^2 w_1$

Then w_0 and w_1 can be regarded as Bäcklund transformations of eq.(1). In particular, the action of $T = \sigma w_1$ is given by

$$T(a) = q^{2}a, \quad T(t) = t, \quad T(F) = \frac{qat\overline{F} + \overline{F}F - 1}{(\overline{F}F - 1)(t\overline{F} + \overline{F}F - 1)}.$$
(12)

Therefore applying *T* to the "seed" solution in Lemma 2.1, we obtain "higher-order" hypergeometric solutions to eq.(1) expressible in terms of a rational function of ψ for $a = q^{2N+1}$ ($N \in \mathbb{Z}$). It is observed that the numerators and denominators of such solutions are factorized and those factors admit the following Casorati determinant formula.

Theorem 2.2. For each $N \in \mathbb{Z}$, we define $\tau_N(t)$ by

$$\tau_{N}(t) = \begin{cases} \begin{pmatrix} \psi(t) & \psi(q^{2}t) & \cdots & \psi(q^{2N-2}t) \\ \psi(q^{-1}t) & \psi(qt) & \cdots & \psi(q^{2N-3}t) \\ \vdots & \vdots & \ddots & \vdots \\ \psi(q^{-N+1}t) & \psi(q^{-N+3}t) & \cdots & \psi(q^{N-1}t) \\ \end{pmatrix} & (N > 0), \\ 1 & (N = 0), \\ \psi(q^{-1}t) & \psi(q^{-3}t) & \cdots & \psi(q^{-2M+1}t) \\ \psi(t) & \psi(q^{-2}t) & \cdots & \psi(q^{-2M+1}t) \\ \vdots & \vdots & \vdots \\ \psi(q^{M-2}t) & \psi(q^{M-4}t) & \cdots & \psi(q^{-M}t) \\ \end{cases} & (M = -N, N < 0). \end{cases}$$

Then

$$F(t) = \begin{cases} \frac{1}{q^{N}} \frac{\tau_{N}(t)\tau_{N+1}(qt)}{\tau_{N}(qt)\tau_{N+1}(t)} & (N \ge 0), \\ -\frac{1}{q^{N+1}} \frac{\tau_{N}(t)\tau_{N+1}(qt)}{\tau_{N}(qt)\tau_{N+1}(t)} & (N < 0), \end{cases}$$
(14)

satisfies eq.(1) with $a = q^{2N+1}$.

¹Actions of these transformations are modified from the original formulae in [30] so that they are subtraction-free. Some typographical errors have been also fixed.

Theorem 2.2 is the direct consequence of the following proposition.

Proposition 2.3. The τ_N satisfy the following bilinear q-difference equations of Hirota type:

(1) $N \ge 0$

$$q^{2N}\tau_{N+1}(t/q)\tau_N(q^2t) - q^N t \tau_{N+1}(t)\tau_N(qt) - \tau_{N+1}(qt)\tau_N(t) = 0,$$
(15)

$$q^{2N}\tau_{N+1}(t/q)\tau_N(qt) - q^{2N}t\,\tau_{N+1}(t)\tau_N(t) - \tau_{N+1}(qt)\tau_N(t/q) = 0.$$
(16)

(2) N < 0

$$q^{2N+2}\tau_{N+1}(t/q)\tau_N(q^2t) + q^{N+1}t\,\tau_{N+1}(t)\tau_N(qt) - \tau_{N+1}(qt)\tau_N(t) = 0,$$
(17)

$$q^{2N+2}\tau_{N+1}(t/q)\tau_N(qt) + q^{2N+1}t\,\tau_{N+1}(t)\tau_N(t) - \tau_{N+1}(qt)\tau_N(t/q) = 0.$$
(18)

In fact one can derive Theorem 2.2 from Proposition 2.3 as follows: for $N \ge 0$ the bilinear equations eqs.(15) and (16) can be rewritten as

$$q^{2N} - \frac{\mu_{N+1}(t/q)}{\mu_N(qt)} \left(q^N t + \frac{\mu_{N+1}(t)}{\mu_N(t)} \right) = 0, \quad q^{2N} - q^{2N} t \, \frac{\mu_{N+1}(t/q)}{\mu_N(t)} - \frac{\nu_N(qt)}{\nu_N(t/q)} = 0, \tag{19}$$

respectively by introducing the variables

$$\nu_N(t) = \frac{\tau_{N+1}(t)}{\tau_N(t)}, \quad \mu_N(t) = \frac{\tau_N(qt)}{\tau_N(t)}.$$
 (20)

Putting

$$F = \frac{1}{q^N} \frac{\nu_N(qt)}{\nu_N(t)} = \frac{1}{q^N} \frac{\tau_{N+1}(qt)\tau_N(t)}{\tau_{N+1}(t)\tau_N(qt)},$$
(21)

and eliminating μ_N and μ_{N+1} from eq.(19) through the use of the identity

$$\frac{\nu_N(qt)}{\nu_N(t)} = \frac{\mu_{N+1}(t)}{\mu_N(t)},$$
(22)

we obtain eq.(1) with $a = q^{2N+1}$. The case of N < 0 can be verified in the same way.

2.3 **Proof of Proposition 2.3**

Our basic idea for proving Proposition 2.3 is to use a determinantal technique. Bilinear *q*-difference equations are derived from the Plücker relations which are quadratic identities among determinants whose columns are shifted. Therefore, we first construct such "difference formulas" that relate "shifted determinants" and τ_N by using the *q*-difference equation of ψ . We then derive bilinear difference equations with the aid of difference formulas from proper Plücker relations. We refer to [7, 10, 12, 13, 21] for applications of this method to hypergeometric solutions of other discrete Painlevé equations.

Let us consider the case of N > 0. We first introduce a notation for the determinants

$$\tau_{N}(t) = \begin{vmatrix} \psi(t) & \psi(q^{2}t) & \cdots & \psi(q^{2N-2}t) \\ \psi(q^{-1}t) & \psi(qt) & \cdots & \psi(q^{2N-3}t) \\ \vdots & \vdots & & \vdots \\ \psi(q^{-N+1}t) & \psi(q^{-N+3}t) & \cdots & \psi(q^{N-1}t) \end{vmatrix} = |\Psi_{0}, \Psi_{2}, \dots, \Psi_{2N-2}|,$$
(23)

where Ψ_k denotes a column vector

$$\Psi_{k} = \begin{pmatrix} \psi(q^{k}t) \\ \psi(q^{k-1}t) \\ \vdots \\ \psi(q^{k-N+1}t) \end{pmatrix}.$$
(24)

Here the height of the column vector is N however we employ the same symbol for determinants with differing heights.

Lemma 2.4. The following formulas hold:

$$|\Psi_0, \Psi_2, \dots, \Psi_{2N-2}| = \tau_N(t), \tag{25}$$

$$|\widehat{\Psi}_{0},\Psi_{1},\Psi_{3},\ldots,\Psi_{2N-3}| = (-1)^{N-1}q^{-(N-1)(N-2)/2}t^{-N+1}\tau_{N}(t),$$
(26)

$$|\Psi_1, \widehat{\Psi}_2, \Psi_3, \dots, \Psi_{2N-3}| = (-1)^{N-2} q^{-(N-1)(N-2)/2} t^{-N+1} \tau_N(t),$$
(27)

where $\widehat{\Psi}_k$ denotes the column vector

$$\widehat{\Psi}_{k} = \begin{pmatrix} \psi(q^{k}t) \\ q\psi(q^{k-1}t) \\ \vdots \\ q^{N-1}\psi(q^{k-N+1}t) \end{pmatrix}.$$
(28)

Proof. Using the linear equation eq.(6) for ψ on the N-th column of the determinant eq.(23) we find

$$\begin{aligned} \tau_N(t) &= \begin{vmatrix} \psi(t) & \psi(q^2t) & \cdots & \psi(q^{2N-4}t) & \psi(q^{2N-4}t) - q^{2N-3}t\psi(q^{2N-3}t) \\ \psi(q^{-1}t) & \psi(qt) & \cdots & \psi(q^{2N-5}t) - q^{2N-4}t\psi(q^{2N-4}t) \\ \vdots & \vdots & \vdots \\ \psi(q^{-N+1}t) & \psi(q^{-N+3}t) & \cdots & \psi(q^{N-3}t) & \psi(q^{N-3}t) - q^{N-2}t\psi(q^{N-2}t) \end{vmatrix}, \\ \\ &= \begin{vmatrix} \psi(t) & \psi(q^{2}t) & \cdots & -q^{2N-3}t\psi(q^{2N-3}t) \\ \psi(q^{-1}t) & \psi(qt) & \cdots & -q^{2N-4}t\psi(q^{2N-4}t) \\ \vdots & \vdots & & \vdots \\ \psi(q^{-N+1}t) & \psi(q^{-N+3}t) & \cdots & -q^{N-2}t\psi(q^{N-2}t) \end{vmatrix}. \end{aligned}$$

Applying the same procedure from the (N - 1)-th column to the second column we have

$$\begin{aligned} \tau_{N}(t) &= \begin{vmatrix} \psi(t) & -qt\psi(qt) & \dots & -q^{2N-3}t\psi(q^{2N-3}t) \\ \psi(q^{-1}t) & -t\psi(t) & \dots & -q^{2N-4}t\psi(q^{2N-4}t) \\ \vdots & \vdots & \vdots \\ \psi(q^{-N+1}) & -q^{-N+2}t\psi(q^{-N+2}t) & \dots & -q^{N-2}t\psi(q^{N-2}t) \end{vmatrix} \\ \\ &= (-t)^{N-1}q^{(N-1)^{2}} \begin{vmatrix} \psi(t) & \psi(qt) & \cdots & \psi(q^{2N-3}t) \\ \psi(q^{-1}t) & q^{-1}\psi(t) & \cdots & q^{-1}\psi(q^{2N-4}t) \\ \vdots & \vdots & \vdots \\ \psi(q^{-N+1}t) & q^{-N+1}\psi(q^{-N+2}t) & \cdots & q^{-N+1}\psi(q^{N-2}t) \end{vmatrix} \\ \\ &= (-t)^{N-1}q^{(N-1)(N-2)/2} \begin{vmatrix} \psi(t) & \psi(qt) & \cdots & \psi(q^{2N-3}t) \\ q\psi(q^{-1}t) & \psi(t) & \cdots & \psi(q^{2N-3}t) \\ q\psi(q^{-1}t) & \psi(t) & \cdots & \psi(q^{2N-4}t) \\ \vdots & \vdots & \vdots \\ q^{N-1}\psi(q^{-N+1}t) & \psi(q^{-N+2}t) & \cdots & \psi(q^{N-2}t) \end{vmatrix} \\ \\ &= (-t)^{N-1}q^{(N-1)(N-2)/2} | \widehat{\Psi}_{0}, \Psi_{1}, \Psi_{3}, \dots, \Psi_{2N-3} |, \end{aligned}$$

which is nothing but eq.(26). At the stage where the above procedure has been employed up to the third column we have $2N_{12} + (2N_{12} + (2$

$$\tau_N(t) = \begin{vmatrix} \psi(t) & \psi(q^2t) & -q^3t\psi(q^3t) & \dots & -q^{2N-3}t\psi(q^{2N-3}t) \\ \psi(q^{-1}t) & \psi(qt) & -q^2t\psi(q^2t) & \dots & -q^{2N-4}t\psi(q^{2N-4}t) \\ \vdots & \vdots & \vdots & \vdots \\ \psi(q^{-N+1}t) & \psi(q^{-N+3}t) & -q^{-N+4}t\psi(q^{-N+4}t) & \dots & -q^{N-2}t\psi(q^{N-2}t) \end{vmatrix}.$$

Using eq.(6) on the first column, we obtain

$$\begin{aligned} \tau_{N}(t) &= \begin{vmatrix} \psi(q^{2}t) + qt\psi(qt) & \psi(q^{2}t) & -q^{3}t\psi(q^{3}t) & \dots & -q^{2N-2}\psi(q^{2N-3}t) \\ \psi(qt) + t\psi(t) & \psi(qt) & -q^{2}t\psi(q^{2}t) & \dots & -q^{2N-3}\psi(q^{2N-4}t) \\ \vdots & \vdots & \vdots & \vdots \\ \psi(q^{-N+3}t) + q^{-N+2}t\psi(q^{-N+2}t) & \psi(q^{-N+3}t) & -q^{-N+4}t\psi(q^{-N+4}t) & \dots & -q^{N-1}\psi(q^{N-2}t) \end{vmatrix} \\ \\ &= (-1)^{N-2}q^{(N-1)(N-2)/2}t^{N-1} \begin{vmatrix} \psi(qt) & \psi(q^{2}t) & \psi(q^{3}t) & \dots & \psi(q^{2N-3}t) \\ \psi(t) & q\psi(qt) & \psi(q^{2}t) & \dots & \psi(q^{2N-4}t) \\ \vdots & \vdots & \vdots & \vdots \\ \psi(q^{-N+2}t) & q^{N-3}\psi(q^{-N+3}t) & \psi(q^{-N+4}t) & \dots & \psi(q^{N-2}t) \end{vmatrix} \\ \\ &= (-1)^{N-2}q^{(N-1)(N-2)/2}t^{N-1} | \Psi_{1}, \widehat{\Psi}_{2}, \Psi_{3}, \dots, \Psi_{2N-3} |, \end{aligned}$$

which is eq. (27). \Box

Now consider the Plücker relation,

$$\begin{aligned} 0 &= \left| \Psi_{-1}, \widehat{\Psi}_{0}, \Psi_{1}, \dots, \Psi_{2N-5} \right| \times |\Psi_{1}, \Psi_{3}, \dots, \Psi_{2N-3}, \phi| \\ &- \left| \widehat{\Psi}_{0}, \Psi_{1}, \Psi_{3}, \dots, \Psi_{2N-5}, \phi \right| \times |\Psi_{-1}, \Psi_{1}, \Psi_{3}, \dots, \Psi_{2N-3}| \\ &+ \left| \Psi_{-1}, \Psi_{1}, \Psi_{3}, \dots, \Psi_{2N-5}, \phi \right| \times \left| \widehat{\Psi}_{0}, \Psi_{1}, \Psi_{3}, \dots, \Psi_{2N-3} \right|, \end{aligned}$$

for an arbitrary column vector ϕ . In particular by choosing ϕ as

$$\phi = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 1 \end{pmatrix},$$

and applying Lemma 2.4 we obtain eq.(15) from the former and eq.(16) from the latter, respectively. The case of N < 0 can be proved in a similar manner. This completes the proof of Proposition 2.3. \Box

3 Continuous Limit to P_{II}

3.1 Continuous Limit on the Formal Level

The continuous limit of "q-P_{II}" to P_{II} involves the "quantum" to classical limit $q \to 1$ but in contrast to the trivial limits usually employed in basic hypergeometric series, i.e. making the substitution $z \mapsto (1 - q)z$ and then setting $q \to 1$, we have a completely different limiting process which is far from trivial and has not been studied much. Let us first recall the formal limits of dP($A_6^{(1)}$) eq.(1).

Proposition 3.1. [26] With the replacements

$$F = ie^{-\delta w}, \quad a = e^{-\frac{\eta}{2}\delta^3}, \quad q = e^{-\frac{\delta^3}{2}}, \quad t = -2ie^{-\frac{s}{2}\delta^2} = -2iq^{\frac{s}{\delta}}, \tag{29}$$

eq.(1) has a limit to P_{II}

$$\frac{d^2w}{ds^2} = 2w^3 + 2sw + \eta,$$
(30)

as $\delta \rightarrow 0$.

Proposition 3.1 can be easily verified by noticing that

$$w(q^{\pm 1}t) = w(s \pm \delta) = w \pm \delta \frac{d}{ds}w + \frac{\delta^2}{2!}\frac{d^2w}{ds^2} + O(\delta^3).$$
(31)

It is well-known that P_{II} eq.(30) admits the hypergeometric solutions for $\eta = 2N + 1$ ($N \in \mathbb{Z}$)[24]:

$$w = -\frac{d}{ds} \log \frac{\kappa_{N+1}}{\kappa_N}, \qquad (32)$$

$$\begin{cases} \left| \begin{array}{cccc} v & \frac{dv}{ds} & \cdots & \left(\frac{d}{ds}\right)^{N-1} v \\ \frac{dv}{ds} & \frac{d^2v}{ds^2} & \cdots & \left(\frac{d}{ds}\right)^N v \\ \vdots & \vdots & \ddots & \vdots \\ \left(\frac{d}{ds}\right)^{N-1} v & \left(\frac{d}{ds}\right)^N v & \cdots & \left(\frac{d}{ds}\right)^{2N-2} v \\ \end{array} \right| \qquad (N > 0), \qquad (33)$$

$$\begin{cases} v & \frac{dv}{ds} & \cdots & \left(\frac{d}{ds}\right)^{M-1} v \\ \frac{dv}{ds} & \frac{d^2v}{ds^2} & \cdots & \left(\frac{d}{ds}\right)^{M-1} v \\ \vdots & \vdots & \ddots & \vdots \\ \left(\frac{d}{ds}\right)^{M-1} v & \left(\frac{d}{ds}\right)^M v & \cdots & \left(\frac{d}{ds}\right)^{2M-2} v \\ \end{array} \right| \qquad (N = -M < 0), \qquad (33)$$
iry equation,

where v satisfies the Airy equation,

$$\frac{d^2v}{ds^2} = -sv. aga{34}$$

The general solution to eq.(34) is given by

$$v(s) = C\operatorname{Ai}(e^{\frac{m}{3}}s) + D\operatorname{Ai}(e^{-\frac{m}{3}}s),$$
(35)

where C, D are arbitrary constants and Ai(s) is the Airy function defined by

$$Ai(s) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{\frac{i}{3}u^3 + isu} \, du.$$
(36)

Proposition 3.2. Under the substitutions eq.(29) and with

$$\psi(t) = e^{\frac{\pi i}{2} \frac{\log t}{\log q}} v(t), \tag{37}$$

the hypergeometric solutions to eq.(1) for $a = q^{2N+1}$ ($N \in \mathbb{Z}$), eqs.(6), (13) and (14) yield, in the limit $\delta \to 0$, the hypergeometric solutions to P_{II} eq.(30) for $\eta = 2N + 1$, eqs. (34), (33) and (32).

Proof. Noticing that

$$\psi(q^{\pm 1}t) = \pm i e^{\frac{\pi i}{2} \frac{\log t}{\log q}} v(s \pm \delta),$$

the linear equation eq.(6) can then be rewritten as

$$v(s+\delta) + v(s-\delta) = 2e^{-\frac{s}{2}\delta^2}v(s),$$
 (38)

which yields eq.(34) in the limit $\delta \to 0$. We next consider the limit of τ_N and the dependent variable transformation. The determinant τ_N (N > 0) can be rewritten as

$$\begin{aligned} \tau_{N} &= e^{\frac{N\pi i}{2} \frac{\log t}{\log q}} \begin{vmatrix} v(t) & i^{2}v(q^{2}t) & \cdots & i^{2N-2}v(q^{2N-2}t) \\ i^{-1}v(q^{-1}t) & iv(qt) & \cdots & i^{2N-3}v(q^{2N-3}t) \\ \vdots & \vdots & \ddots & \vdots \\ i^{-N+1}v(q^{-N+1}t) & i^{-N+3}v(q^{-N+3}t) & \cdots & i^{N-1}v(q^{N-1}t) \end{vmatrix} \\ \\ &= e^{\frac{N\pi i}{2} \frac{\log t}{\log q}} i^{N(N-1)/2} \begin{vmatrix} v(s) & v(s+2\delta) & \cdots & v(s+(2N-2)\delta) \\ v(s-\delta) & v(s+\delta) & \cdots & v(s+(2N-3)\delta) \\ \vdots & \vdots & \vdots \\ v(s-(N-1)\delta) & v(s-(N-3)\delta) & \cdots & v(s+(N-1)\delta) \end{vmatrix} \\ \\ &= e^{\frac{N\pi i}{2} \frac{\log t}{\log q}} i^{N(N-1)/2} \sigma_{N}(s), \end{aligned}$$

so that

$$F = \frac{i}{q^N} \frac{\sigma_N(s)\sigma_{N+1}(s+\delta)}{\sigma_N(s+\delta)\sigma_{N+1}(s)}.$$
(39)

We also note that

$$\sigma_N(s) = (-2\delta^2)^{N(N-1)/2} [\kappa_N + O(\delta)].$$

Therefore we deduce

$$w = -\frac{1}{\delta}\log\frac{F}{i} = -\frac{1}{\delta}\log\left[1 + \delta\left(\frac{d}{ds}\log\frac{\kappa_{N+1}}{\kappa_N}\right) + O(\delta^2)\right] = -\frac{d}{ds}\log\frac{\kappa_{N+1}}{\kappa_N} + O(\delta).$$
(40)

The limit in the case of N < 0 can be verified in a similar manner. \Box

Now let us consider the limit of the solution of the linear equation eq.(6). The two linearly independent powerseries solutions of the Airy equation eq.(34) are given by

$${}_{0}F_{1}\left(\begin{array}{c}-\\\frac{2}{3}\end{array};-\frac{s^{3}}{3^{2}}\right) = 1 - \frac{1}{3!}s^{3} + \frac{1 \cdot 4}{6!}s^{6} - \frac{1 \cdot 4 \cdot 7}{9!}s^{9} + \cdots,$$

$$s_{0}F_{1}\left(\begin{array}{c}-\\\frac{4}{3}\end{array};-\frac{s^{3}}{3^{2}}\right) = s - \frac{2}{4!}s^{4} + \frac{2 \cdot 5}{7!}s^{7} - \frac{2 \cdot 5 \cdot 8}{10!}s^{10} + \cdots.$$
(41)

However as is apparent from the series expansion of hypergeometric functions in eq.(7) the application of the scaling changes of variables in eq.(29) does not yield any meaningful limit as $\delta \to 0$ on a term by term basis. What is required is another representation of these functions and a uniform, possibly asymptotic, expansion with respect to the other parameters as $q \to 1$. This question has been addressed in [25] and for the most part answered there. We discuss the continous limit of the hypergeometric functions $_{1}\varphi_{1}\begin{pmatrix} 0\\ -q \end{pmatrix}$ in the next section.

3.2 Continuous Limit of the Hypergeometric Functions

There are three key ingredients in [25] which are necessary to derive the final formula that we require. The first is a suitable integral representation for the $_1\varphi_1\begin{pmatrix}0\\y\\ \end{bmatrix};q,x\end{pmatrix}$ function and this is the *q*-analog of the Mellin-Barnes inversion integral.

Proposition 3.3. [4, 25] The representation

$${}_{1}\varphi_{1}\begin{pmatrix}0\\y;q,x\end{pmatrix} = \frac{(q;q)_{\infty}}{(y;q)_{\infty}} \int_{\rho-i\infty}^{\rho+i\infty} \frac{dz}{2\pi i} \, z^{-\log x/\log q} \, \frac{(y/z;q)_{\infty}}{(z;q)_{\infty}},\tag{42}$$

is valid for $x, y \in \mathbb{C}$, with $|arg(x)| < \pi$, $y \neq q^{-n}$ $(n \in \mathbb{Z}_{\geq 0})$, $0 < \rho < 1$ and 0 < q < 1.

Remark 3.1. Proposition 3.3 follows by evaluating the integral on the contour described in Fig.1 with the residues at $z = q^{-n}$

$$\operatorname{Res}_{z=q^{-n}}(z;q)_{\infty}^{-1} = (-1)^{n+1} \frac{q^{\binom{n}{2}}}{(q;q)_n (q;q)_{\infty}}, \quad n \in \mathbb{Z}_{\geq 0},$$
(43)

and by deforming the path C appropriately according to Cauchy's Theorem.

The second ingredient is an asymptotic formula for the q-shifted factorial $(t;q)_{\infty}$ as $q \rightarrow 1$ which is uniform with respect to t. Such expansions have only recently been studied and in particular by Meinardus [19], McIntosh [16, 17, 18] and the above cited work [25]. Amongst all the essentially equivalent forms we choose the following statement:

Proposition 3.4. [19, 25, 18] As $q \to 1^-$ the q-shifted factorial $(t; q)_{\infty}$ has an asymptotic expansion

$$\log(t;q)_{\infty} \sim \frac{1}{\log q} \operatorname{Li}_{2}(t) + \frac{1}{2} \log(1-t) + \sum_{n=1}^{\infty} \frac{B_{2n}}{(2n)!} \left(t\frac{d}{dt}\right)^{2n-2} \frac{t}{1-t} (\log q)^{2n-1},$$
(44)



Figure 1: Contour for Proposition 3.3.

for 0 < q < 1 and uniform for t in any compact domain of \mathbb{C} such that $|\arg(1-t)| < \pi$. Here $\operatorname{Li}_2(t)$ is the dilogarithm function defined by

$$\operatorname{Li}_{2}(t) = -\int_{0}^{t} \frac{\log(1-u)}{u} \, du = \sum_{n=1}^{\infty} \frac{t^{n}}{n^{2}},\tag{45}$$

and B_{2n} the even Bernoulli numbers. In the case of t = q we have [22]

$$\log(q;q)_{\infty} = \frac{\pi^2}{6\log q} + \frac{1}{2}\log\frac{2\pi}{-\log q} + O(\log q).$$
(46)

We next apply Proposition 3.4 to the integral representation eq.(42). Noticing that q-shifted factorials in the integral representation are rewritten by putting $q = e^{-\epsilon}$ as

$$\frac{(y/z;q)_{\infty}}{(z;q)_{\infty}}z^{-\log x/\log q} = e^{\frac{1}{\epsilon}\left[-\text{Li}_{2}(y/z) + \text{Li}_{2}(z) + \log x\log z\right]} \times e^{\frac{1}{2}\left[\log\left(1-\frac{y}{z}\right) - \log(1-z)\right]} \times [1+O(\epsilon)]$$

and

$$\frac{(q;q)_{\infty}}{(y;q)_{\infty}} = \mathrm{e}^{\frac{1}{\epsilon} \left[\mathrm{Li}_{2}(y) - \frac{\pi^{2}}{6}\right] + \frac{1}{2}\log\frac{2\pi}{\epsilon} - \frac{1}{2}\log(1-y)} \times [1 + O(\epsilon)],$$

we obtain:

Proposition 3.5. Let $x, y \in \mathbb{C}$ and $q = e^{-\epsilon}$ for $\epsilon > 0$. Then

$${}_{1}\varphi_{1}\begin{pmatrix}0\\y;q,x\end{pmatrix} = \frac{1}{2\pi i} \int_{\rho-i\infty}^{\rho+i\infty} e^{\frac{1}{\epsilon} \left[\log x \log z - \text{Li}_{2}(\frac{y}{z}) + \text{Li}_{2}(z)\right]} e^{\frac{1}{2} \left[\log(1-\frac{y}{z}) - \log(1-z)\right]} dz \times \left[1 + O(\epsilon)\right], \tag{47}$$

where $|\arg(x)| < \pi$, Re $y < \rho$.

Remark 3.2. We take $\log z$ on its principal sheet cut along $(-\infty, 0]$ and $\operatorname{Li}_2(z)$ on its principal sheet cut along $(1, +\infty)$. If $x, y \in (0, 1)$ then for $z \in (y, 1)$ the argument of

$$\log x \log z - \operatorname{Li}_2(\frac{y}{z}) + \operatorname{Li}_2(z) \tag{48}$$

is zero. When $z \in \mathbb{C}$ subject to

$$|\arg z| < \pi$$
, $|\arg(1-z)| < \pi$, $|\arg\left(1-\frac{y}{z}\right)| < \pi$,

that is we exclude the rays $(1, \infty)$, $(-\infty, 0)$ and (0, y), it follows that the argument of eq.(48) lies in the interval $(-\pi, \pi)$. The contour path given in Proposition 3.5 is then just a simple path $z = \rho + it$ (Re $y < \rho < 1$, $t \in (-\infty, \infty)$) satisfying these criteria and the requirement that the endpoints of the contour ensure the existence of the integral. If the contour is deformed then eq.(47) is valid if the contour does not cut across the ray (0, y).

The third ingredient is the application of saddle point method to the Laplace type integral in eq.(47). In this problem two saddle points arise and can coalesce depending on the values of the parameters. Therefore we have to construct an asymptotic approximation that incorporates the contributions from both saddle points uniformly with repsect to their separation. A method for such an asymptotic expansion of this type of integral has been set out by Chester, Friedman and Ursell[2, 32, 33].

To illustrate the method, let us consider the integral

$$I = \frac{1}{2\pi i} \int_C e^{\frac{1}{\epsilon}g(z;d)} f(z) \, dz, \quad \epsilon \to 0$$

where f(z) is analytic with respect to z, and g(z; d) is analytic with respect to z and the parameter d. We assume that there are two saddle points z_1 and z_2 which are determined from g'(z; d) = 0 and that they coalesce when d = 0. The key of this method is to introduce the change of variable $z \rightarrow u$ via the cubic parameterization

$$g(z;d) = \frac{1}{3}u^3 - \alpha u + \beta,$$
 (49)

where α and β are determined as follows. Firstly differentiating eq.(49) we have

$$g'(z;d)\frac{dz}{du} = u^2 - \alpha.$$
(50)

In order for eq.(49) to define a single-valued analytic transformation neither $\frac{dz}{du}$ nor $\frac{du}{dz}$ can vanish in relevant regions. Therefore at the saddle points we have the correspondence

$$z = z_1 \quad \leftrightarrow u = \alpha^{\frac{1}{2}}, \quad z = z_2 \quad \leftrightarrow u = -\alpha^{\frac{1}{2}}, \tag{51}$$

which determines α and β as

$$\frac{4}{3}\alpha^{\frac{2}{3}} = g(z_2; d) - g(z_1; d), \quad 2\beta = g(z_2; d) + g(z_1; d).$$
(52)

The transformation u = u(z; d) defined by eqs.(49) and (52) has three branches. However, it can be shown that there is exactly one branch which has the following properties [2, 33]; (i) $u = u(z; \alpha)$ is expanded into a power series in z with coefficient continuous in d near d = 0, (ii) z_1 and z_2 correspond to $\alpha^{\frac{1}{2}}$ and $-\alpha^{\frac{1}{2}}$ respectively, and (iii) near d = 0 the correspondence $z \leftrightarrow u$ is 1:1.

We next expand f(z) in the form

$$f(z)\frac{dz}{du} = \sum_{m=0}^{\infty} (p_m + q_m u)(u^2 - \alpha)^m,$$
(53)

and define the following integrals

$$F_{m} = \frac{1}{2\pi i} \int_{C'} e^{\frac{1}{\epsilon} \left(\frac{1}{3}u^{3} - \alpha u\right)} (u^{2} - \alpha)^{m} du,$$

$$G_{m} = \frac{1}{2\pi i} \int_{C'} e^{\frac{1}{\epsilon} \left(\frac{1}{3}u^{3} - \alpha u\right)} (u^{2} - \alpha)^{m} u du,$$

(54)

Here C' is the image of C by the transformation given by eqs.(49) and (52). By using recursion relations for F_m and G_m obtained by partial integration, the expansion of I can be written in the form

$$I = e^{\frac{\beta}{\epsilon}} \left[\epsilon^{\frac{1}{3}} V(\alpha \epsilon^{\frac{2}{3}}) \sum_{m=0}^{\infty} a_m \epsilon^m + \epsilon^{\frac{2}{3}} V'(\alpha \epsilon^{\frac{2}{3}}) \sum_{m=0}^{\infty} b_m \epsilon^m \right],$$
(55)

where

$$a_0 = p_0, \quad b_0 = -q_0, \tag{56}$$

and $V(\lambda)$ is the Airy integral

$$V(\lambda) = \frac{1}{2\pi i} \int_{C'} e^{\frac{1}{3}u^3 - \lambda u} du.$$
(57)

The coefficients p_0 and q_0 are determined by putting $z = z_1$, $u = \alpha^{\frac{1}{2}}$ and $z = z_2$, $u = -\alpha^{\frac{1}{2}}$ in eq.(53) as

$$p_{0} = \frac{1}{2} \left[f(z_{1}) \left(\frac{dz}{du} \right)_{z=z_{1}} + f(z_{2}) \left(\frac{dz}{du} \right)_{z=z_{2}} \right],$$

$$q_{0} = \frac{1}{2\alpha^{\frac{1}{2}}} \left[f(z_{1}) \left(\frac{dz}{du} \right)_{z=z_{1}} - f(z_{2}) \left(\frac{dz}{du} \right)_{z=z_{2}} \right].$$
(58)

Prellberg [25] has applied the above expansion to the integral representation eq.(47) to obtain its leading behaviour as $\epsilon \to 0$ for 0 < x, y < 1 as follows:

Proposition 3.6. [25] Let 0 < x, y < 1 and $q = e^{-\epsilon}$ with $\epsilon > 0$. Then as $\epsilon \to 0$ we have

$${}_{1}\varphi_{1}\begin{pmatrix}0\\y;q,x\end{pmatrix} = e^{\frac{1}{\epsilon}[\operatorname{Li}_{2}(y)-\frac{1}{6}\pi^{2}+\frac{1}{2}\log x\log y]}\sqrt{\frac{2\pi}{\epsilon(1-y)}} \times \left[p_{0}\epsilon^{\frac{1}{3}}\operatorname{Ai}(\alpha\epsilon^{-\frac{2}{3}})-q_{0}\epsilon^{\frac{2}{3}}\operatorname{Ai}'(\alpha\epsilon^{-\frac{2}{3}})\right][1+O(\epsilon)],$$
(59)

where the auxiliary variables are

$$\frac{4}{3}\alpha^{3/2} = \log x \log \left(\frac{z_m - \sqrt{d}}{z_m + \sqrt{d}}\right) + 2\mathrm{Li}_2(z_m - \sqrt{d}) - 2\mathrm{Li}_2(z_m + \sqrt{d}),\tag{60}$$

with

$$z_m = \frac{1+y-x}{2}, \qquad d = z_m^2 - y,$$
 (61)

and

$$p_0 = \left(\frac{\alpha}{d}\right)^{1/4} \frac{1 - x - y}{2}, \qquad q_0 = \left(\frac{d}{\alpha}\right)^{1/4}.$$
 (62)

Remark 3.3. Errors in the statement of this result as given in [25] have been fixed, in particular the sign of the q_0 term in eq.(59) and the factor of 2 in the denominator of p_0 in eq.(62).

In our case, the parameters are given by

$$\epsilon = \frac{\delta^3}{2}, \quad q = e^{-\frac{\delta^3}{2}}, \quad y = -e^{-\frac{\delta^3}{2}} = -1 + \frac{\delta^3}{2}, \quad x = \mp qt = \mp 2ie^{-\frac{\delta}{2}\delta^2 + \frac{\delta^3}{2}} = \pm (2i - is\delta^2 - i\delta^3 + \cdots), \tag{63}$$

and unfortunately, they do not match the assumptions of Proposition 3.6. Therefore we have to consider the extension of Proposition 3.6 for our case, taking care of the multi-valuedness of the integrand. We fix the branch of log and fractional power functions as

$$\log z = \ln |z| + i\operatorname{Arg} z, \quad -\pi \le \operatorname{Arg} z < \pi, \qquad z^{\frac{m}{n}} = \exp\left(\frac{m}{n}\log z\right) = |z|^{\frac{m}{n}} e^{\frac{m}{n}i\operatorname{Arg} z}.$$

Note that $\log(XY) = \log X + \log Y$ and $\log(X/Y) = \log X - \log Y$ are valid only mod $2\pi i$.

Substituting eq.(63) into eq.(47) and expanding the integrand with respect to δ , we obtain:

Proposition 3.7. With the substitutions eq.(63) we have as $\delta \rightarrow 0$

$${}_{1}\varphi_{1}\begin{pmatrix}0\\-q\ ;\ q,-qt\end{pmatrix} = \frac{1}{2\pi i} \int_{\rho-i\infty}^{\rho+i\infty} e^{\frac{2}{\delta^{3}}g_{-}(z)} f_{-}(z) \, dz \times \left(\frac{2\pi}{\delta^{3}}\right)^{\frac{1}{2}} e^{\frac{2}{\delta^{3}}\left(-\frac{\pi^{2}}{4} + \frac{\ln 2}{2}\delta^{3}\right)} \times \left[1 + O(\delta^{3})\right], \tag{64}$$

$${}_{1}\varphi_{1}\begin{pmatrix}0\\-q\ ;\ q,qt\end{pmatrix} = \frac{1}{2\pi i} \int_{\rho-i\infty}^{\rho+i\infty} e^{\frac{2}{\delta^{3}}g_{+}(z)} f_{+}(z) \, dz \times \left(\frac{2\pi}{\delta^{3}}\right)^{\frac{1}{2}} e^{\frac{2}{\delta^{3}}\left(-\frac{\pi^{2}}{4} + \frac{\ln 2}{2}\delta^{3}\right)} \times \left[1 + O(\delta^{3})\right],\tag{65}$$

where $0 < \rho < 1$ and

$$g_{-}(z) = \text{Li}_{2}(z) - \text{Li}_{2}\left(-\frac{1}{z}\right) + \log\left(2ie^{-\frac{s\delta^{2}}{2}}\right)\log z,$$
 (66)

$$f_{-}(z) = e^{-\frac{1}{2}\log(1+\frac{1}{z}) - \frac{1}{2}\log(1-z) - \log z},$$
(67)

$$g_{+}(z) = \text{Li}_{2}(z) - \text{Li}_{2}\left(-\frac{1}{z}\right) + \log\left(-2ie^{-\frac{s\delta^{2}}{2}}\right)\log z,$$
 (68)

$$f_{+}(z) = e^{-\frac{1}{2}\log(1+\frac{1}{z}) - \frac{1}{2}\log(1-z) - \log z}.$$
(69)

Let us take the case of x = -qt and apply the saddle point method to eqs.(64), (66) and (67). The saddle points $z_1^{(-)}$ and $z_2^{(-)}$ are determined by

$$g'_{-}(z) = -\frac{\log(1-z)}{z} - \frac{\log\left(1+\frac{1}{z}\right)}{z} + \frac{\log\left(2ie^{-\frac{s\delta^2}{2}}\right)}{z} = 0,$$
(70)

which yields the quadratic equation

$$z^2 + 2ie^{-\frac{s\delta^2}{2}}z - 1 = 0.$$
 (71)

Therefore the saddle points are given by

$$z_1^{(-)} = z_m + D, \quad z_2^{(-)} = z_m - D, \quad z_m = -ie^{-\frac{s\delta^2}{2}}, \quad D^2 = z_m^2 + 1,$$
 (72)

or expanding in terms of δ we obtain

$$z_m = -i + \frac{is}{2}\delta^2 + \cdots, \quad D = s^{\frac{1}{2}}\delta - \frac{s^{\frac{3}{2}}}{4}\delta^3 + \cdots,$$
 (73)

$$z_1^{(-)} = -i + s^{\frac{1}{2}}\delta + \frac{is}{2}\delta^2 + \cdots, \quad z_2^{(-)} = -i - s^{\frac{1}{2}}\delta + \frac{is}{2}\delta^2 + \cdots.$$
(74)

The quantity α is calculated by using eq.(52) as

$$\frac{4}{3}\alpha^{\frac{3}{2}} = g(z_{2}^{(-)}) - g(z_{1}^{(-)})$$

$$= \operatorname{Li}_{2}(z_{m} - D) - \operatorname{Li}_{2}\left(-\frac{1}{z_{m} - D}\right) + \log\left(2ie^{-\frac{s\delta^{2}}{2}}\right)\log(z_{m} - D)$$

$$-\operatorname{Li}_{2}(z_{m} + D) + \operatorname{Li}_{2}\left(-\frac{1}{z_{m} + D}\right) - \log\left(2ie^{-\frac{s\delta^{2}}{2}}\right)\log(z_{m} + D)$$

$$= 2\left[\operatorname{Li}_{2}(z_{m} - D) - \operatorname{Li}_{2}(z_{m} + D)\right] + \log\left(2ie^{-\frac{s\delta^{2}}{2}}\right)\left[\log(z_{m} - D) - \log(z_{m} + D)\right]$$

$$= \frac{2i}{3}s^{\frac{3}{2}}\delta^{3} + O(\delta^{5})$$

where we have used $\frac{1}{z_m \pm D} = -(z_m \mp D)$. Therefore we conclude

$$\alpha = 2^{-\frac{2}{3}} s e^{\frac{\pi i}{3}} \delta^2 + O(\delta^4).$$
(75)

We can derive this for *s* in the sector $-\pi \leq \operatorname{Arg}(s) < \pi/3$ but it actually holds without this restriction. The reason for this is that $_1\varphi_1$ is an analytic function of $t \in \mathbb{C}$ and therefore of $s \in \mathbb{C}$. Consequently the leading term of the expansion of $_1\varphi_1$ as $\delta \to 0$ is analytic with respect to *s* as the remainder terms can be shown to be uniformly bounded in *s* under this limit. Let us next compute p_0 and q_0 according to the formula eq.(58). From the correspondence

$$z = z_1^{(-)} = -i + s^{\frac{1}{2}}\delta + \frac{is}{2}\delta^2 + \dots \quad \longleftrightarrow \quad u = u_1^{(-)} = \alpha^{\frac{1}{2}} = 2^{-\frac{1}{3}}s^{\frac{1}{2}}e^{\frac{\pi i}{6}}\delta + O(\delta^3)$$
$$z = z_2^{(-)} = -i - s^{\frac{1}{2}}\delta + \frac{is}{2}\delta^2 + \dots \quad \longleftrightarrow \quad u = u_2^{(-)} = -\alpha^{\frac{1}{2}} = -2^{-\frac{1}{3}}s^{\frac{1}{2}}e^{\frac{\pi i}{6}}\delta + O(\delta^3)$$

we obtain $(dz/du)_{z=z_1^{(-)}, z_2^{(-)}}$ as

$$\begin{split} \left(\frac{dz}{du}\right)_{z=z_{1}^{(-)}} &= \frac{\left(\frac{dz_{1}}{d\delta}\right)_{\delta\sim0}}{\left(\frac{du_{1}^{(-)}}{d\delta}\right)_{\delta\sim0}} = \frac{s^{\frac{1}{2}} + is\delta + O(\delta^{2})}{2^{-\frac{1}{3}}s^{\frac{1}{2}}e^{\frac{\pi i}{6}} + O(\delta^{2})} = 2^{\frac{1}{3}}e^{-\frac{\pi i}{6}}\left(1 + is^{\frac{1}{2}}\delta + O(\delta^{2})\right),\\ \left(\frac{dz}{du}\right)_{z=z_{2}^{(-)}} &= \frac{\left(\frac{dz_{2}}{d\delta}\right)_{\delta\sim0}}{\left(\frac{du_{2}^{(-)}}{d\delta}\right)_{\delta\sim0}} = \frac{-s^{\frac{1}{2}} + is\delta + O(\delta^{2})}{-2^{-\frac{1}{3}}s^{\frac{1}{2}}e^{\frac{\pi i}{6}} + O(\delta^{2})} = 2^{\frac{1}{3}}e^{-\frac{\pi i}{6}}\left(1 - is^{\frac{1}{2}}\delta + O(\delta^{2})\right). \end{split}$$

Substituting eq.(74) into eq.(67) we have

$$f(z_1^{(-)}) = 2^{-\frac{1}{2}} e^{\frac{\pi i}{4}} \left(1 - is^{\frac{1}{2}} \delta + \cdots \right), \quad f(z_2^{(-)}) = 2^{-\frac{1}{2}} e^{\frac{\pi i}{4}} \left(1 + is^{\frac{1}{2}} \delta + \cdots \right),$$

from which we obtain

$$p_0 = \frac{1}{2} \left[f(z_1^{(-)}) \left(\frac{dz}{du} \right)_{z=z_1^{(-)}} + f(z_2^{(-)}) \left(\frac{dz}{du} \right)_{z=z_2^{(-)}} \right] = 2^{-\frac{1}{6}} e^{\frac{\pi i}{12}} + O(\delta^2),$$
(76)

$$q_0 = \frac{1}{2\alpha^{\frac{1}{2}}} \left[f(z_1^{(-)}) \left(\frac{dz}{du} \right)_{z=z_1^{(-)}} - f(z_2^{(-)}) \left(\frac{dz}{du} \right)_{z=z_2^{(-)}} \right] = 0 + O(\delta).$$
(77)

We compute β by using eq.(52) as

$$2\beta = g(z_{2}^{(-)}) + g(z_{1}^{(-)})$$

$$= \operatorname{Li}_{2}(z_{m} - D) - \operatorname{Li}_{2}\left(-\frac{1}{z_{m} - D}\right) + \log\left(2ie^{-\frac{s\delta^{2}}{2}}\right)\log(z_{m} - D)$$

$$+ \operatorname{Li}_{2}(z_{m} + D) - \operatorname{Li}_{2}\left(-\frac{1}{z_{m} + D}\right) + \log\left(2ie^{-\frac{s\delta^{2}}{2}}\right)\log(z_{m} + D)$$

$$= \log\left(2ie^{-\frac{s\delta^{2}}{2}}\right)\left[\log(z_{m} - D) + \log(z_{m} + D)\right],$$
(78)

which yields

$$\beta = -\frac{\pi i}{2}\ln 2 + \frac{\pi^2}{4} + \frac{\pi i}{4}s\delta^2 + O(\delta^4).$$
⁽⁷⁹⁾

Remark 3.4. The multi-valuedness of the integrand has a critical effect in the calculation of β . One might compute β from eq.(78) as

$$2\beta = \log\left(2ie^{-\frac{s\delta^2}{2}}\right) \left[\log(z_m - D) + \log(z_m + D)\right] = \log\left(2ie^{-\frac{s\delta^2}{2}}\right) \log(z_m^2 - D^2) \\ = \log\left(2ie^{-\frac{s\delta^2}{2}}\right) \log(-1) = -\pi i (\ln 2 + \frac{\pi i}{2} - \frac{s\delta^2}{2}),$$

but the second equality does not hold in general (in this case it is accidentally correct). In fact, the same procedure for the case of x = qt yields wrong result.

Let us finally consider the image of integration path $C : z = \rho + it (-\infty < t < \infty)$ in the *u*-plane. From the identity of dilogarithm[14, 3]

$$\operatorname{Li}_{2}(z) = -\operatorname{Li}_{2}(\frac{1}{z}) - \frac{1}{2}(\log z)^{2} + \pi i \log z + \frac{\pi^{2}}{3},$$
(80)

we see for $t \to \pm \infty$

$$g_{-}(\rho + it) \sim \text{Li}_{2}(\rho + it) = -\text{Li}_{2}(\frac{1}{\rho + it}) - \frac{1}{2}(\log(\rho + it))^{2} + \pi i \log(\rho + it) + \frac{\pi^{2}}{3}$$

$$\sim -\frac{1}{2}(\log|t|)^{2} \pm \frac{\pi i}{2}\log|t|, \quad t \to \pm \infty,$$
(81)

therefore

$$u(\rho + it) \sim (3g_{-}(\rho + it))^{\frac{1}{3}} \sim e^{\pm \frac{\pi i}{3}} \left(\frac{3}{2}(\log |t|)^2\right)^{\frac{1}{3}}, \quad t \to \pm \infty.$$

This gives the integration path C' as $\infty e^{-\frac{\pi i}{3}} \rightarrow O \rightarrow \infty e^{\frac{\pi i}{3}}$, which implies $V(\lambda) = \operatorname{Ai}(\lambda)$. Closer investigation shows that the mapping given by eq.(49) is regular and 1:1 in the domain including C'.

Collecting the above results and performing similar calculations for the case of x = qt, we obtain the following asymptotic expansions from eq.(55):

Proposition 3.8. With the substitutions eq.(29) we have as $\delta \rightarrow 0$

$${}_{1}\varphi_{1}\begin{pmatrix}0\\-q\ ;\ q,-qt\end{pmatrix} = 2\pi^{\frac{1}{2}}\delta^{-\frac{1}{2}}e^{-\frac{\pi i}{\delta^{3}}\ln 2 + \frac{\pi i}{2\delta}s + \frac{\pi i}{12}}\left[\operatorname{Ai}(se^{\frac{\pi i}{3}}) + O(\delta^{2})\right],\tag{82}$$

$${}_{1}\varphi_{1}\begin{pmatrix}0\\-q\ ;\ q,qt\end{pmatrix} = 2\pi^{\frac{1}{2}}\delta^{-\frac{1}{2}}e^{\frac{\pi i}{\delta^{3}}\ln 2 - \frac{\pi i}{2\delta}s - \frac{\pi i}{12}}\left[\operatorname{Ai}(se^{-\frac{\pi i}{3}}) + O(\delta^{2})\right],\tag{83}$$

for s in any compact domain of \mathbb{C} .

Now we are in a position to deduce the limit of the general solution to eq.(6). From eqs.(37) and (7) we note that

$$v(t) = A e^{-\frac{\pi i}{2} \frac{\log t}{\log q}} \varphi_1 \left(\begin{array}{c} 0 \\ -q \end{array}; q, -qt \right) + B e^{\frac{\pi i}{2} \frac{\log t}{\log q}} \varphi_1 \left(\begin{array}{c} 0 \\ -q \end{array}; q, qt \right),$$
(84)

for arbitrary q-periodic functions A, B. Observing that

$$e^{\pm \frac{\pi i}{2} \frac{\log t}{\log q}} = e^{\frac{1}{\delta^3} \left(\mp \pi i \ln 2 \mp \frac{\pi^2}{2} \right) \pm \frac{\pi i}{2\delta} s} \times [1 + O(\delta)],$$
(85)

we find that the *s*-dependence in the exponential pre-factors of eqs. (82) and (83) cancels exactly. Therefore we finally arrive at the desired result:

Theorem 3.9. We have as $\delta \to 0$

$$v(t) = 2\sqrt{\frac{\pi}{\delta}} \left[A \, \mathrm{e}^{\frac{\pi^2}{2\delta^3} + \frac{\pi \mathrm{i}}{12}} \mathrm{Ai}(\mathrm{e}^{\frac{\pi \mathrm{i}}{3}}s) + B \, \mathrm{e}^{-\frac{\pi^2}{2\delta^3} - \frac{\pi \mathrm{i}}{12}} \mathrm{Ai}(\mathrm{e}^{-\frac{\pi \mathrm{i}}{3}}s) \right] \times [1 + O(\delta)],\tag{86}$$

for *s* in any compact domain of \mathbb{C} and constants $A, B \in \mathbb{C}$.

4 Concluding Remarks

In this article we have considered the *q*-Painlevé equation $dP(A_6^{(1)})$, eq.(1), and constructed its classical solutions having a determinantal form with basic hypergeometric function elements. We have also discussed the continuous limit to P_{II}. In particular, we have shown that hypergeometric functions ${}_{1}\varphi_1\begin{pmatrix} 0\\ -q \end{pmatrix}$ actually reduce to the

Airy functions Ai $(e^{\pm \frac{\pi i}{3}})$ by applying a generalization of the saddle point method to their integral representations. We first remark that P_{II} eq.(30) admits rational solutions for $\eta = 2N$ ($N \in \mathbb{Z}$) which can be expressed in terms of

a specialization of 2-core Schur functions[11]. Such solutions are obtained by applying Bäcklund transformations to the simple rational solution that is fixed by the Dynkin diagram automorphism. One would expect similar rational solutions for eq.(1), however Masuda has shown that there is no rational solution fixed by the corresponding Dynkin diagram automorphism [15]. This implies that it is not appropriate to regard eq.(1) simply as a "q-analog of P_{II}".

Secondly, we observe an asymmetry in the structure of determinant formula eq.(13); the shifts in *t* of the entries are different between the horizontal and the vertical directions. It might be natural to regard this structure as originating from the asymmetry of the root lattice. Actually in other cases where this situation arises, such as the $A_5^{(1)}$ surface("*q*-P_{IV}" and "*q*-P_{III}", ($A_2 + A_1$)⁽¹⁾-symmetry)[10, 7], $A_4^{(1)}$ surface("*q*-P_V", $A_4^{(1)}$ -symmetry)[5] or $A_3^{(1)}$ surface("*q*-P_{VI}", $D_5^{(1)}$ -symmetry)[31], the determinant structure is symmetric. Such an asymmetric structure of the

determinant is known for several discrete Painlevé equations - one example is the "standard" discrete Painlevé II equation[27, 13]

$$x_{n+1} + x_{n-1} = \frac{(an+b)x_n + c}{1 - x_n^2},$$
(87)

where a, b, c are parameters. Equation (87) may be regarded as a special case of the "asymmetric" discrete Painlevé II equation²[29, 23, 28]

$$x_{n+1} + x_{n-1} = \frac{(an+b)x_n + c + d(-1)^n}{1 - x_n^2},$$
(88)

when d = 0, which is actually a discrete Painlevé equation associated with the $D_5^{(1)}$ surface and arises as a Bäcklund transformation of the Painlevé V equation (P_V). Therefore the hypergeometric solutions for (88) are expressible in terms of the Whittaker function and are the same as those for P_V[28]. However, the hypergeometric solutions for eq.(87) are quite different; the relevant hypergeometric function is the parabolic cylinder function and the determinant structure has the same asymmetry as that for dP($A_6^{(1)}$) eq.(1). A similar structure is known for the "standard" discrete (*q*-)Painlevé III equation (dP_{III})[27, 12]

$$\frac{x_{n+1}x_{n-1}}{d_1d_2} = \frac{(x_n - c_1q^n)(x_n - c_2q^n)}{(x_n - d_1)(x_n - d_2)},$$
(89)

and the q-Painlevé VI equation $(q-P_{VI})$ [6, 29],

$$y_{n}y_{n+1} = \frac{a_{3}a_{4}(z_{n+1} - b_{1}q^{n})(z_{n+1} - b_{2}q^{n})}{(z_{n+1} - b_{3})(z_{n+1} - b_{4})},$$

$$z_{n}z_{n+1} = \frac{b_{3}b_{4}(y_{n} - a_{1}q^{n})(y_{n} - a_{2}q^{n})}{(y_{n} - a_{3})(y_{n} - a_{4})},$$

$$\frac{b_{1}b_{2}}{b_{3}b_{4}} = q\frac{a_{1}a_{2}}{a_{3}a_{4}},$$
(90)

where a_i , b_i (i = 1, 2, 3, 4), c_j and d_j (j = 1, 2) are parameters. The hypergeometric solutions for eq.(90) are given by the basic hypergeometric series $_2\varphi_1$ [31] while those for eq.(89) are given by Jackson's *q*-Bessel function[12].

Thus the results of our study sheds some light on the "degenerated" equations such as eqs.(87) or (89). They are not just special cases of the original "generic" equations. Our results imply that putting d = 0 in eq.(88) is not just killing the "parity", but causes qualitative change of the root lattice which in turn results in different hypergeometric solutions and an asymmetry of the determinant formula. Therefore they should be studied independently of the "generic" discrete Painlevé equations in the Sakai's classification[30].

Acknowledgement We acknowledge Profs. Takashi Aoki, Takahiro Kawai, Tatsuya Koike and Yoshitsugu Takei for their interests in this work and valuable discussions. We also thank Profs. Tetsu Masuda, Masatoshi Noumi, Yasuhiro Ohta and Yasuhiko Yamada for their continuous encouragement and discussions. NSW acknowledges support from the Australian Research Council.

References

- [1] M. Abramowitz and I. Stegun, Handbook of mathematical functions (Dover, 1972).
- [2] C. Chester, B. Friedman and F. Ursell, An extension of the method of the steepest descents, Proc. Cambridge Philos. Soc. 53(1957) 699-611.
- [3] A. Erdelyj, Higher transcendental functions Vol.1 (McGraw-Hill, New York, 1953).
- [4] G. Gasper and M. Rahman, *Basic hypergeometric series*, second edition, Encyclopedia of Mathematics and Its Applications **96** (Cambridge University Press, Cambridge, 2004).
- [5] T. Hamamoto and K. Kajiwara, *Hypergeometric solutions to the q-P_{II} and q-P_V equations*, Article No. 41, Reports of RIAM Symposium No.16ME-S1 "Physics and Mathematical Structures of Nonlinear Waves" (2005) (in Japanese), available online at http://www.riam.kyushu-u.ac.jp/fluid/meeting/16ME-S1/

²The word "asymmetric" comes from the terminology of the Quispel-Roberts-Thompson mapping which is known as the fundamental family of second-order integrable mappings. It has nothing to do with the asymmetric structure of the determinant formula of their solutions.

- [6] M. Jimbo and H. Sakai, A q-analog of the sixth Painlevé equation, Lett. Math. Phys. 38(1996) 145-154.
- [7] K. Kajiwara and K. Kimura, On a q-Painlevé III equation. I: derivations, symmetry and Riccati type solutions, J. Nonlin. Math. Phys. 10 (2003) 86-102.
- [8] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, *Hypergeometric solutions to the q-Painlevé equations*, Int. Math. Res. Not. **2004**(2004) 2497-2521.
- [9] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, *Construction of hypergeometric solutions to the q-Painlevé equations*, Int. Math. Res. Not. **2005**(2005) 1439-1463.
- [10] K. Kajiwara, M. Noumi and Y. Yamada, A study on the fourth q-Painlevé equation, J. Phys. A: Math. Gen. 34 (2001) 8563-8581.
- [11] K. Kajiwara and Y. Ohta, *Determinant structure of the rational solutions for the Painlevé II equation*, J. Math. Phys. **37**(1996) 4093-4704.
- [12] K. Kajiwara, Y. Ohta, J. Satsuma, Casorati determinant solutions for the discrete Painlevé III equation, J. Math. Phys. 36(1995) 4162–4174.
- [13] K. Kajiwara, Y. Ohta, J. Satsuma, B. Grammaticos and A. Ramani, *Casorati determinant solutions for the discrete Painlevé–II equation*, J.Phys. A: Math. Gen. 27(1994) 915–922.
- [14] A. Kirillov, Dilogarithm identities, Prog. of Theor. Phys. Suppl. 118 (1995) 61-142.
- [15] T. Masuda, private communications.
- [16] R.J. McIntosh, Some asymptotic formulae for q-hypergeometric series, J. London Math. Soc. (2) 51(1995) 120-136.
- [17] R.J. McIntosh, Asymptotic transformation of q-series, Canad. J. Math. 50(1998) 412-425.
- [18] R.J. McIntosh, Some asymptotic formulae for q-shifted factorials, Ramanujan J. 3(1999) 205-214.
- [19] G. Meinardus Über Partitionen mit Differenzenbedingungen, Math. Z. 61(1954) 289-302.
- [20] M. Murata, H. Sakai and J. Yoneda, *Riccati solutions of discrete Painlevé equations with Weyl group symmetry* of type $E_8^{(1)}$, J. Math. Phys. **44**(2003) 1396-1414.
- [21] F. Nijhoff, J. Satsuma, K. Kajiwara, B. Grammaticos and A. Ramani, *A study of the alternate discrete Painlevé II equation*, Inverse Problems **12**(1996) 697–716.
- [22] G.H. Hardy, Ramanujan (Cambridge University Press, 1940).
- [23] Y. Ohta, Self-dual construction of discrete Painlevé equations (in Japanese), Applied mathematics of discrete integrable systems (Kyoto, 1998), RIMS Kokyuroku No. 1098 (1999) 130–137.
- [24] K. Okamoto, Studies on the Painlevé equations. III. second and fourth Painlevé equations, P_{II} and P_{IV}. Math. Ann. 275 (1986) 221–255.
- [25] T. Prellberg, Uniform q-Series asymptotics for staircase polygons, J. Phys. A: Math. Gen. 28(1995) 1289-1304.
- [26] A. Ramani and B. Grammaticos, Discrete Painlevé equations: coalescences, limits and degeneracies, Physica A 228(1996) 160–171.
- [27] A. Ramani, B. Grammaticos and J. Hietarinta, *Discrete Versions of the Painlevé Equations*, Phys. Rev. Lett. 67(1991) 1829-1832.
- [28] A. Ramani, B. Grammaticos, T. Tamizhmani and K.M. Tamizhmani, *Special function solutions of the discrete Painlevé equations*, Comput. Math. Appl. **42**(2001) 603-614.

- [29] A. Ramani, Y. Ohta, B. Grammaticos and J. Satsuma, *Self-duality and Schlesinger chains for the asymmetric d-P*_{II} and *q-P*_{III} equations, Commun. Math. Phys. **192**(1998) 67-76.
- [30] H. Sakai, *Rational Surfaces associated with affine root systems and geometry of the Painlevé equations*, Commun. Math. Phys. **220**(2001) 165-229.
- [31] H. Sakai, *Casorati determinant solutions for the q-difference sixth Painlevé equation*, Nonlinearity **11**(1998) 823-833.
- [32] F. Ursell, *Integrals with large parameter. The continuation of uniformly asymptotic expansions*, Proc. Cambridge Philos. Soc. **61**(1965) 113-128.
- [33] R. Wong, Asymptoric approximation of integrals, Classics in applied mathematics 34 (SIAM, 2001).

List of MHF Preprint Series, Kyushu University 21st Century COE Program Development of Dynamic Mathematics with High Functionality

- MHF2005-1 Hideki KOSAKI Matrix trace inequalities related to uncertainty principle
- MHF2005-2 Masahisa TABATA Discrepancy between theory and real computation on the stability of some finite element schemes
- MHF2005-3 Yuko ARAKI & Sadanori KONISHI Functional regression modeling via regularized basis expansions and model selection
- MHF2005-4 Yuko ARAKI & Sadanori KONISHI Functional discriminant analysis via regularized basis expansions
- MHF2005-5 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA Point configurations, Cremona transformations and the elliptic difference Painlevé equations
- MHF2005-6 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA Construction of hypergeometric solutions to the q Painlevé equations
- MHF2005-7 Hiroki MASUDA Simple estimators for non-linear Markovian trend from sampled data: I. ergodic cases
- MHF2005-8 Hiroki MASUDA & Nakahiro YOSHIDA Edgeworth expansion for a class of Ornstein-Uhlenbeck-based models
- MHF2005-9 Masayuki UCHIDA Approximate martingale estimating functions under small perturbations of dynamical systems
- MHF2005-10 Ryo MATSUZAKI & Masayuki UCHIDA One-step estimators for diffusion processes with small dispersion parameters from discrete observations
- MHF2005-11 Junichi MATSUKUBO, Ryo MATSUZAKI & Masayuki UCHIDA Estimation for a discretely observed small diffusion process with a linear drift
- MHF2005-12 Masayuki UCHIDA & Nakahiro YOSHIDA AIC for ergodic diffusion processes from discrete observations

- MHF2005-13 Hiromichi GOTO & Kenji KAJIWARA Generating function related to the Okamoto polynomials for the Painlevé IV equation
- MHF2005-14 Masato KIMURA & Shin-ichi NAGATA Precise asymptotic behaviour of the first eigenvalue of Sturm-Liouville problems with large drift
- MHF2005-15 Daisuke TAGAMI & Masahisa TABATA Numerical computations of a melting glass convection in the furnace
- MHF2005-16 Raimundas VIDŪNAS Normalized Leonard pairs and Askey-Wilson relations
- MHF2005-17 Raimundas VIDŪNAS Askey-Wilson relations and Leonard pairs
- MHF2005-18 Kenji KAJIWARA & Atsushi MUKAIHIRA Soliton solutions for the non-autonomous discrete-time Toda lattice equation
- MHF2005-19 Yuu HARIYA Construction of Gibbs measures for 1-dimensional continuum fields
- MHF2005-20 Yuu HARIYA Integration by parts formulae for the Wiener measure restricted to subsets in \mathbb{R}^d
- MHF2005-21 Yuu HARIYA A time-change approach to Kotani's extension of Yor's formula
- MHF2005-22 Tadahisa FUNAKI, Yuu HARIYA & Mark YOR Wiener integrals for centered powers of Bessel processes, I
- MHF2005-23 Masahisa TABATA & Satoshi KAIZU Finite element schemes for two-fluids flow problems
- MHF2005-24 Ken-ichi MARUNO & Yasuhiro OHTA Determinant form of dark soliton solutions of the discrete nonlinear Schrödinger equation
- MHF2005-25 Alexander V. KITAEV & Raimundas VIDŪNAS Quadratic transformations of the sixth Painlevé equation
- MHF2005-26 Toru FUJII & Sadanori KONISHI Nonlinear regression modeling via regularized wavelets and smoothing parameter selection
- MHF2005-27 Shuichi INOKUCHI, Kazumasa HONDA, Hyen Yeal LEE, Tatsuro SATO, Yoshihiro MIZOGUCHI & Yasuo KAWAHARA On reversible cellular automata with finite cell array

- MHF2005-28 Toru KOMATSU Cyclic cubic field with explicit Artin symbols
- MHF2005-29 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Kaori NAGATOU A computational approach to constructive a priori and a posteriori error estimates for finite element approximations of bi-harmonic problems
- MHF2005-30 Kaori NAGATOU, Kouji HASHIMOTO & Mitsuhiro T. NAKAO Numerical verification of stationary solutions for Navier-Stokes problems
- MHF2005-31 Hidefumi KAWASAKI A duality theorem for a three-phase partition problem
- MHF2005-32 Hidefumi KAWASAKI A duality theorem based on triangles separating three convex sets
- MHF2005-33 Takeaki FUCHIKAMI & Hidefumi KAWASAKI An explicit formula of the Shapley value for a cooperative game induced from the conjugate point
- MHF2005-34 Hideki MURAKAWA A regularization of a reaction-diffusion system approximation to the two-phase Stefan problem
- MHF2006-1 Masahisa TABATA Numerical simulation of Rayleigh-Taylor problems by an energy-stable finite element scheme
- MHF2006-2 Ken-ichi MARUNO & G R W QUISPEL Construction of integrals of higher-order mappings
- MHF2006-3 Setsuo TANIGUCHI On the Jacobi field approach to stochastic oscillatory integrals with quadratic phase function
- MHF2006-4 Kouji HASHIMOTO, Kaori NAGATOU & Mitsuhiro T. NAKAO A computational approach to constructive a priori error estimate for finite element approximations of bi-harmonic problems in nonconvex polygonal domains
- MHF2006-5 Hidefumi KAWASAKI A duality theory based on triangular cylinders separating three convex sets in \mathbb{R}^n
- MHF2006-6 Raimundas VIDŪNAS Uniform convergence of hypergeometric series
- MHF2006-7 Yuji KODAMA & Ken-ichi MARUNO N-Soliton solutions to the DKP equation and Weyl group actions

- MHF2006-8 Toru KOMATSU Potentially generic polynomial
- MHF2006-9 Toru KOMATSU Generic sextic polynomial related to the subfield problem of a cubic polynomial
- MHF2006-10 Shu TEZUKA & Anargyros PAPAGEORGIOU Exact cubature for a class of functions of maximum effective dimension
- MHF2006-11 Shu TEZUKA On high-discrepancy sequences
- MHF2006-12 Raimundas VIDŪNAS Detecting persistent regimes in the North Atlantic Oscillation time series
- MHF2006-13 Toru KOMATSU Tamely Eisenstein field with prime power discriminant
- MHF2006-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO Generating function associated with the Hankel determinant formula for the solutions of the Painlevé IV equation
- MHF2006-15 Raimundas VIDŪNAS Darboux evaluations of algebraic Gauss hypergeometric functions
- MHF2006-16 Masato KIMURA & Isao WAKANO New mathematical approach to the energy release rate in crack extension
- MHF2006-17 Toru KOMATSU Arithmetic of the splitting field of Alexander polynomial
- MHF2006-18 Hiroki MASUDA Likelihood estimation of stable Lévy processes from discrete data
- MHF2006-19 Hiroshi KAWABI & Michael RÖCKNER Essential self-adjointness of Dirichlet operators on a path space with Gibbs measures via an SPDE approach
- MHF2006-20 Masahisa TABATA Energy stable finite element schemes and their applications to two-fluid flow problems
- MHF2006-21 Yuzuru INAHAMA & Hiroshi KAWABI Asymptotic expansions for the Laplace approximations for Itô functionals of Brownian rough paths
- MHF2006-22 Yoshiyuki KAGEI Resolvent estimates for the linearized compressible Navier-Stokes equation in an infinite layer

MHF2006-23 Yoshiyuki KAGEI

Asymptotic behavior of the semigroup associated with the linearized compressible Navier-Stokes equation in an infinite layer

- MHF2006-24 Akihiro MIKODA, Shuichi INOKUCHI, Yoshihiro MIZOGUCHI & Mitsuhiko FUJIO The number of orbits of box-ball systems
- MHF2006-25 Toru FUJII & Sadanori KONISHI Multi-class Logistic Discrimination via Wavelet-based Functionalization and Model Selection Criteria
- MHF2006-26 Taro HAMAMOTO, Kenji KAJIWARA & Nicholas S. WITTE Hypergeometric Solutions to the q-Painlevé Equation of Type $(A_1 + A'_1)^{(1)}$