Seasonal Fluxes of Silicoflagellates and Actiniscus in the Subarctic Pacific During 1982-1984

Takahashi, Kozo Department of Geology and Geophysics, Woods Hole Oceanographic Institution

https://hdl.handle.net/2324/12437

出版情報:Journal of Marine Research. 45, pp.397-425, 1987. Sears Foundation for Marine Research, Yale University バージョン: 権利関係:

Seasonal fluxes of silicoflagellates and *Actiniscus* in the subarctic Pacific during 1982–1984

by Kozo Takahashi¹

ABSTRACT

A seasonal flux study of silicoflagellates and *Actiniscus* was conducted at subarctic Pacific Station PAPA (50N, 145W; water depth 4200 m) during September, 1982 through August, 1984 using PARFLUX high resolution time series sediment traps deployed at 1000 m and 3800 m depths. The time series sediment trap samples were collected for 4 to 16-day periods depending on the samples; most of the samples were collected for 14 to 16-day periods. From a depth of 3800 m, a total of 47 samples represents a nearly two-year continuous record of the fluxes of silicoflagellate and *Actiniscus* taxa. An additional 12 samples from 1000 m represent a 6-month period, providing synchronized time series samples with the deeper depth which is essential to understand particle sinking processes.

Seven silicoflagellate taxa, several variants of silicoflagellates, and Actiniscus pentasterias (Ehrenberg) group were examined. A total of more than 32×10^3 specimens were identified to species level and counted in this study. Distephanus speculum (Ehrenberg) and Dictyocha mandrai Ling are the dominant taxa, generally contributing >75% of the flux assemblages. Considerable intra-annual and interannual flux variability was observed with the range of total silicoflagellate flux varying from 5×10^3 to 700×10^3 skeletons m⁻²day⁻¹ during the two years. Seasonal flux patterns of D. speculum and its closely related taxa are internally consistent and they conform with the productivity signals shown by diatom, total mass, or opal fluxes. The seasonal flux pattern of D. mandrai exhibits its own unique late fall/early winter signals with most of the cumulative flux concentrated during this season in year 1.

The sinking mechanism of silicoflagellates is large aggregates which sink faster than discrete specimens. Silicoflagellates generally sink faster than marine snow mediated diatom assemblages. The larger influence of fecal material may be responsible for the observed faster sinking speeds than those of diatoms. No significant dissolution of silicoflagellate assemblages occurs in the water column due to innate protection and to accelerated sinking. Most of the silicoflagellates supplied to the sea floor are subsequently dissolved, and only 1% of the total supply is preserved in the surface sediments, with considerably altered assemblages.

1. Introduction

Silicoflagellates are marine chrysophytes which dwell in the euphotic zones of essentially all parts of the world oceans. Their microplankton size skeletons (20–80 μ m) are composed of biogenic opal which can be preserved in the sedimentary fossil record.

1. Department of Geology and Geophysics, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts, 02543, U.S.A.

Journal of Marine Research

[45, 2

The numerical abundance of silicoflagellates in siliceous plankton standing stocks is generally less than that of diatoms and is followed by radiolarians. Due to the small mass of silicoflagellate skeletons, their contribution to SiO_2 flux is no more than a few percent, but numerous skeletons in fluxes make them good tracers for the vertical transport mechanisms of settling particles (Takahashi, 1987a). Some silicoflagellate taxa are considered to be useful as environmental indicators such as for high productivity and oligotrophic waters (Murray and Schrader, 1983; Takahashi, 1987a). The recent rapid progress in using silicoflagellates as a tool to interpret the fossil record yields a promising future for reconstructing paleotemperature variation (e.g., Ciesielski, 1975) and paleoproductivity (e.g., Schrader and Baumgartner, 1983) studies.

In a previous paper (Takahashi, 1987a), silicoflagellate fluxes at meso- and bathypelagic depths were assessed at four different environments ranging from the temperate oligotrophic Atlantic to the tropical eutrophic Pacific in which temperate to tropical species were dealt with. In that paper, not only were detailed vertical processes examined but also indicator species for productivity and oligotrophic environments were proposed. A time-series flux assessment throughout the year was not conducted in the previous study in the low latitudes where relatively small seasonal changes in productivity were expected.

The objectives of this paper are as follows. As the fossil records generally exhibit mixed signals from different seasons with unknown ratios between them, it is difficult to assess summer or winter conditions accurately by studying fossils alone. If we can understand the seasonal variations of supply rates of microfossils to the sea floor, accurate paleoenvironmental assessments will be possible. It is important to study year to year variability of the supply rate in the short term as well as in the long term. For these reasons it is desirable to conduct a high resolution time-series vertical flux study of silicoflagellate assemblages in an area where primary production is seasonally highly variable. Furthermore, it is crucial to understand the processes of fossil preservation in the sedimentary record, in particular, early diagenesis in the water column. Less than a few percent of biogenic opal is generally preserved in surface sediments and the bulk of the supply is dissolved in the vicinity of the sea floor (Takahashi, 1983, 1986), indicating the significance of such a dissolution mechanism to the fossil record. It is likely that all of the preserved fossils go through some level of dissolution process in corrosive water with respect to biogenic opal. Without a better understanding of the dissolution/preservation processes, it is difficult to link the present with the past. Therefore, a part of this study is focused on the effects of dissolution.

Since silicoflagellate flux is a function of organic production in the overlying euphotic zone (Takahashi, 1987a), flux studies on materials collected from the bathypelagic zone will provide productivity signals, a part of which will be subsequently preserved in the fossil record. Therefore, one can expect the results of this study to be of value for interpretive paleoproductivity. Whether temperature signals



Figure 1. A map showing the location of PARFLUX sediment trap Station PAPA with bathymetric contours in meters.

(e.g., Ciesielski, 1975) can be derived from the flux assemblages or not will decide the potential use of silicoflagellates for future paleotemperature studies.

The results of a detailed high resolution time-series vertical flux study of silicoflagellates during 2 years from September, 1982 to August, 1984 at Station PAPA (Fig. 1; 50N, 145W; water depth 4200 m) located in the Alaskan current gyre are presented. Station PAPA was the site of Canadian Weather Station P and numerous hydrographic data were collected there from 1956 to 1981 (Anonymous, 1981; Tabata and Peart, 1985; and unpublished observations by the Institute of Ocean Sciences, Patricia Bay, British Columbia). The study of a continuous 2-year flux record provides an opportunity to learn the extent of intra-annual and interannual variabilities in present-day sedimentation rates. A pair of sediment trap samples collected simultaneously from two depths for 6 months enable us to clarify vertical sinking processes.

The Actiniscus pentasterias group was also studied because of its similar size and abundance to silicoflagellates and is included in this report. Actiniscus pentasterias is a dinoflagellate which secretes a pair of siliceous endoskeletons in the cell (Orr and Conley, 1976). No ebridians occurred in the material studied.

2. Methods

An automated time series PARFLUX Mark 5 sediment trap was deployed at 3800 m for 4 periods of 6 months each for a total of 2 years. Forty-seven samples (September, 1982-August, 1984; PAPA I, II, III, and IV) were collected by bottom tethered mooring array (Table 1). An additional trap was deployed at 1000 m for a 6-month period and its sampling intervals were synchronized with those of the 3800 m trap; 12 samples were collected (PAPA II). The Mark 5 sediment trap has a conical collection funnel made of polypropylene with a surface opening of 1.167 m^2 which is covered with a 94-cell honeycomb baffle. The funnel is connected to a rotating disk located underneath. The disk has 12 screw-on sample containers (250 ml each) and turns according to a preprogrammed schedule set into a command microprocessor (Honjo, 1984; Honjo and Wong, 1985). The containers had sodium azide as a microbial poison. A total of 59 time series sediment trap samples were collected for 4 to 16-day periods; most of the samples were collected for 14 to 16-day periods (Table 1). Dissolution of opal within the sample containers during the deployment and storage in the laboratory was not significant as shown by dissolved silicon measurements of supernatant of the samples (Manganini and Honjo, 1985).

The microslide preparation method used in this work is similar to that employed for a diatom study (Takahashi, 1986). Briefly, a sample is sieved through 1000 μ m and 63 μ m mesh stainless screens. Both 63-1000 μ m and <63 μ m size fractions are split into four aliquots using an Erez-Honjo precision liquid rotary splitter (Honjo, 1978) with an accuracy error of less than 5%. The splitting is repeated until a desirable aliquot size was obtained, requiring several repetitions. Because silicoflagellate fluxes varied greatly with seasons, the aliquot size had to be changed often in consecutive samples. An appropriately split aliquote of 63-1000 μ m was wet sieved through 250 μ m and 125 µm mesh screens. The size-fractioned aliquot of the appropriate sample size (see Takahashi, 1985, for details) is filtered through a 47 mm HA Millipore[®] grid membrane filter with a nominal 0.45 μ m pore size using a rectangular stainless steel filtration funnel with a 19×42 mm opening. The filter is rinsed with distilled water to remove salts and then dried in an oven. After trimming off the excess margins, the dried filter is mounted with Cargile[®] type B immersion oil on a standard glass microslide. The oil-immersed filter is left overnight in a clean box in order to drive air bubbles out of the biogenic particles. A cover glass is placed over the oil with Devcon® 5-minute epoxy adhesive on four corners of the cover glass to secure the sample on the slide.

Silicoflagellate counts were performed with a light transmission microscope with wide-field, $10 \times$ eyepiece lenses and a $20 \times$ S Plan Apochromat objective lens with a total magnification of $200 \times$. Silicoflagellate flux values reported in this paper are the sum of four size fractions actually studied: $<63 \mu$ m, $63-125 \mu$ m, $125-250 \mu$ m, and $250-1000 \mu$ m. Detailed counts and fluxes for each size fraction of silicoflagellate species, as well as detailed methods and statistics, are reported elsewhere (Takahashi,

trap experiment at subarctic Pacific Station PAPA. Dates represent opening and closing times of sample cups placed with a tight seal under the conical PARFLUX Mark 5 traps (surface area = 1.167 m^2). For deployment II there are two synchronized depth samples from 1000 m and 3800 m of each sequenced sample number.

				Duration
Deployment	Sample #	Date open	Date closed	(days)
PAPA I	#1	Sep. 23, 1982	Oct. 8, 1982	15
3800 m	#2	Oct. 8,	Oct. 23	15
	#3	Oct. 23	Nov. 7	15
	#4	Nov. 7	Nov. 22	15
	#5	Nov. 22	Dec. 7	15
	#6	Dec. 7	Dec. 22	15
	#7	Dec. 22, 1982	Jan. 6, 1983	15
	#8	Jan. 6	Jan. 21	15
	#9	Jan. 21	Feb. 5	15
	#10	Feb. 5	Feb. 20	15
	#11	Feb. 20	Mar. 7	15
	#12	Mar. 7	Mar. 18	11
PAPA II	#13	Mar. 27	Apr. 12	16
1000 m & 3800 m	#14	Apr. 12	Apr. 28	16
	#15	Apr. 28	May 14	16
	#16	May 14	May 30	16
	#17	May 30	June 15	16
	#18	June 15	July 1	16
	#19	July 1	July 17	16
	#20	July 17	Aug. 2	16
	#21	Aug. 2	Aug. 18	16
	#22	Aug. 18	Sept. 3	16
	#23	Sept. 3	Sept. 19	16
	#24	Sept. 19	Oct. 5	16
PAPA III	#25	Oct. 14	Oct. 30	16
3800 m	#26	Oct. 30	Nov. 15	16
	#27	Nov. 15	Dec. 1	16
	#28	Dec . 1	Dec. 17	16
	#29	Dec. 17, 1983	Jan. 2, 1984	16
	#30	Jan. 2	Jan. 18	16
	#31	Jan. 18	Feb. 3	16
	#32	Feb. 3	Feb. 19	16
	#33	Feb. 19	Mar. 6	16
	#34	Mar. 6	Mar. 22	16
	#35	Mar. 22	Apr. 7	16
	#36	Apr. 7	Apr. 23	16

Deployment	Sample #	Date open	Date closed	Duration (days)
PAPA IV	#37*	Apr. 23	May 13, 1984	14
3800 m	#38	May 13	May 27	14
	#39	May 27	June 10	14
	#40	June 10	June 24	14
	#41	June 24	July 8	14
	#42	July 8	July 22	14
	#43	July 22	Aug. 5	14
	#44	Aug. 5	Aug. 9	4
	#45	Aug. 9	Aug. 13	4
	#46	Aug. 13	Aug. 17	4
	#47	Aug. 17	Aug. 21	4
	#48	Aug. 21	Aug. 24, 1984	4

Table 1. (Continued)

*No sample/data due to malfunction

1985). Because samples were prepared for both silicoflagellate and radiolarian studies, four different size fractions were examined, but most silicoflagellate and Actiniscus specimens are located in the <63 μ m fraction. Thus, most of the counting effort was spent on the <63 μ m fraction. One slide per size fraction of each sample was studied for census. One to nine traverses of vertical strips (along the minor axis of a microslide) per slide were studied for counts. Each strip represents $15.1 \times 10^6 \ \mu$ m² which is equal to 1/35.9 (or 2.786 $\times 10^{-2}$) of an entire filtered area of the membrane filter on a slide. Raw counts for a sample combining all of the four slides ranged from 74 to 1558 per sample with a mean of 552. Possible major sources of errors in silicoflagellate counts reported here include (i) sample splitting, (ii) imperfect distribution of filtered silicoflagellates, and (iii) counting errors. The second category of the sources of errors is determined to be the largest, based on statistical assessments performed on the representative samples. The first and third categories caused less than 5% of the errors, each. Overall reproducibility for silicoflagellate flux assessment reported here is considered better than 85%.

Diversity indices of silicoflagellate assemblages were computed using the following Shannon and Weaver formula (Pielou, 1969),

$$H' = -\sum_{i=1}^{n} Pi \log Pi \tag{1}$$

where H' is the diveristy index in a "bel" or a "decimal digit" unit (Pielou, 1969), Pi is the proportion of the *i*th species of the total population dealt with, and *n* is the number of species. The nine most abundant counting groups of silicoflagellates, whose seasonal fluxes are illustrated in later sections, were used for the diversity index computations.

The sinking speeds of individual silicoflagellate species were obtained by examining

seasonal flux maxima and minima of 1000 m and 3800 m data. In order to find the best matching lag time between the two depth data sets, a crosscorrelation method was applied (e.g., Davis, 1973). In this method, we first compute a correlation coefficient, r, between data from the two depths without shifting any data (Lag O). This provides a direct data comparison between the two depths. If sinking time between the two depths is less than the sample resolution, a strong correlation coefficient is obtained. Second, the 3800 m data are shifted by one sample to the preceding sample dates (to the left) and a correlation coefficient r is computed (Lag 1). By shifting the 3800 m data further left to earlier sample dates and computing the coefficients, one can get a continuous trend of crosscorrelation.

3. Results

Seven silicoflagellate taxa, their several variants, and the Actiniscus pentasterias group were encountered during the census (Plates 1 and 2; Appendix). More than one species of Actiniscus were encountered; they were lumped together as a group. Detailed taxonomic discussions with illustrations of the taxa used in the paper will be published elsewhere.

The counts were converted into flux and reported as the number of individuals $m^{-2}d^{-1}$ (Appendix). Highly variable fluxes of silicoflagellate and *Actiniscus* taxa were observed, both intra-annually and interannually, during the 1982–1984 period (Figs. 2–4). *Distephanus speculum* and *Dictyocha mandrai* are major contributors to the flux of silicoflagellates throughout the period studied. The combined flux of the two species is generally greater than 75% ($\bar{x} = 83\%$, $\sigma = 8\%$) of the total silicoflagellate flux. During August, 1984, the combined flux decreased to about 67% of total silicoflagellates.

Essentially all silicoflagellates encountered during the census were not associated with fecal pellets. The number of fecal pellets found in the samples was so small that they rarely turned up during the census.

a. Seasonal flux variability in year 1 (September, 1982–September, 1983). The intensity of the flux of silicoflagellates varied significantly from species to species. For example, a maximum flux of Distephanus speculum (Plate 1, Figs. 1, 2; Plate 2, Fig. 4) reached 543×10^3 individuals $m^{-2}d^{-1}$ in October, 1982, at 3,800 m while that of Distephanus quinquangellus was only 5×10^3 individuals $m^{-2}d^{-1}$ (Figs. 2a, 2b). Seasonal flux patterns of silicoflagellate species can be assessed by comparing timing and relative intensity of flux maxima and minima. All species show 3 to 5 major flux maxima at 3,800 m during year 1. The 5 seasonal maxima are: October, December (1982), March, May, and August (1983). A general similarity of the flux patterns of species within a genus is observed in Distephanus and Dictyocha, respectively. The flux of Distephanus presents rather intensive maxima during October, March, and August.



Plate 1. Light micrographs of silicoflagellate species from Station PAPA. All scale bars equal 20 µm. The scale of Figures 2, 13, and 14 are the same as that in Figure 1. Sources of the micrographs are from maximum flux time periods at 3,800 m illustrated in Figures 2-4. (1, 2) Distephanus speculum (Ehrenberg), apical view (1) and side view (2). Radial spine length and ratios of apical/basal ring diameters tend to vary whereas the shapes of the apical ring (mostly circular) and the basal ring (mostly equilateral hexagon) tend to be uniform. (3) Distephanus quinquangellus Bukry and Foster. (4) Distephanus crux ? (Ehrenberg). (5) Distephanus boliviensis (Frenguelli). (6) Distephanus octangulatus Wailes, regular form with one apical window. (7) Distephanus octangulatus Wailes. A form with three apical windows. Forms with two or more apical windows were enumerated separately from the regular form with an apical window. (8) Dictyocha mandrai Ling, regular form. (9) Dictyocha mandrai Ling, form with variable shaped apical rods. (10) Dictyocha messanensis? f. spinosa (Lemmerman). Size and orientation of supporting spines vary considerably. (11) Dictyocha mandrai f. pentagona. (12-14) Actiniscus pentasterias group. (15) An organic aggregate containing many specimens of D. speculum. Compare the size of a discrete specimen of D. boliviensis in the upper left with that of D. speculum. Sample #22 (18 August-3 September, 1983), 3,800 m.



Plate 2. Scanning electron micrographs of silicoflagellate species from Station PAPA. All scale bars equal 10 µm. All specimens were collected from 3,800 m. (1) *Dictyocha mandrai* Ling, apical view, Sample #23 (3 Sept.-19 Sept., 1983). Note wavy ornamentation on skeletal surface which was not observed in low latitude sediment trap material (Takahashi, 1987a). A small specimen on upper right is *D. speculum* for size comparison. (2) *Dictyocha mandrai* Ling, side view, #25 (14 Oct.-30 Oct., 1983). Note that apical spine is relatively short compared to *D. messanensis* ? *spinosa* illustrated in the low latitude study. (3) *Distephanus octangulatus* Wailes, apical view, #2 (8 Oct.-23 Oct. 1982). Note wavy surface ornamentation. (4) *Distephanus speculum* (Ehrenberg), oblique apical view, #25 (14 Oct.-30 Oct., 1983). Note small size and smooth circular shape of apical ring without any spines and compare it to (5). Overall size of this taxon is relatively small compared to *D. mandrai* which is illustrated in (1). (5) *Distephanus boliviensis* (Frenguelli), apical view, #23 (3 Sept.-19 Sept., 1983). Note several spines projecting from apical ring. Shape of its apical ring is a polygon. (6) *Actiniscus pentasterias* (Ehrenberg), apical view, #14 (12 April-28 April, 1983).

APPENDIX

Time-series fluxes (No. $m^{-2}d^{-1}$) of silicoflagellate taxa and *Actiniscus* at 1000 m (for 6 months) and at 3800 m (for 2 years) at the Subarctic Station PAPA during 1982–1984.

					Distephanus
Deployment/Depth	Distephanus	Distephanus	Distephanus	Distephanus	octangulatus
Sample No.	speculum	quinquangellus	boliviensis	octangulatus	(>1 apical window)
	-			-	-
PAPA I 3800 m					<u>^</u>
#1	206710	508	3588	3115	0
#2	542606	5303	16590	19793	306
#3	287841	2188	13230	9721	840
#4	239182	2940	7718	10229	840
#5	200830	1479	10343	7070	44
#6	208040	2984	14980	12644	0
#7	182858	2100	19163	8068	840
#8	95025	420	8024	7560	420
#0	37310	420	3710	5775	304
#10	10162	175	2010	2000	554
#10	19103	1/5	3019	2000	44
#11	8400	0	1094	2888	0
#12	7186	0	858	3873	143
PAPA II 1000 m					
#13	313943	779	5619	24364	0
#14	125430	123	2053	17024	123
#15	62166	125	2755	20540	246
#15	40700	0	5004	20349	1575
#10	40729	0	5094	14421	1373
#17	23404	0	5636	/998	0
#18	13978	0	6276	11419	394
#19	30823	0	21503	8882	0
#20	61481	1316	10866	6003	439
#21	184412	1436	9352	4430	0
#22	180105	2994	7588	2625	0
#23	117800	779	6276	1313	0
#24	104798	1436	6891	1436	Ő
#24	104770	1450	0071	1450	v
PAPA II 3800 m					
#13	276085	2994	7752	23544	779
#14	150409	2092	3897	13700	123
#15	108941	779	5742	12387	1313
#16	115257	656	3568	16735	0
#17	67555	656	5088	17024	656
#19	42024	0.0	J700 4150	7040	799
#10	42034	617	4139	10051	/00
#19	40108	517	489/	10951	911
#20	59357	0	7373	6757	3510
#21	172854	1878	12022	14778	3510
#22	303803	4387	20026	8143	877
#23	97210	369	15135	7957	123
#24	46590	246	1792	3633	439
DIDA III 2000					
PAPA III 3800 m					
#25	41630	1162	3589	2892	164
#26	34413	0	2789	3281	492
#27	23626	164	1641	2338	0
#28	16243	164	2010	1805	0
#29	13659	0	1148	1189	0
#30	11218	0	1528	1641	0
#31	8244	164	1005	523	0
#32	9865	0	872	656	0
#33	6214	ů	1548	118	Ő
#33 #34	0214	174	007	1702	164
#34	9249	174	002	1702	104
#33	16509	390	2789	1/04	0
#36	12141	0	2953	1354	U
PAPA IV 3800 m					
#37	No sample				
#39 #28	22208	, 0	5428	1504	0
#30 #20	23270	0	J430 6000	011	0
#39 #40	5140/	U	0000	544	U
#40	18657	0	3375	1125	U
#41	9141	47	1688	1219	188
#42	10500	188	1875	609	0
#43	1031	0	1734	656	0
#44	2625	0	1969	164	0
#45	1969	0	3117	164	0
#46	656	ñ	2297	328	164
#47	1212	ñ	378	1477	0
11'''''' #AQ	820	0	107	656	ň
#40	020	U	472	0.0	v

APPENDIX (Continued)

	Dictyocha mandrai	Dictyocha	Dictyocha			Actiniscus
Dictyocha	(with variable	mandrai f.	messanensis ?	Distephanus	Total	pentasterias
mandrai	apical rods)	pentagona	spinosa	crux ?	Silicoflagellates	group
10509	219	0	840	1059	226967	37441
94789	1855	1899	0	5915	689055	158708
75355	744	420	700	3404	394486	60813
101159	770	420	1260	2520	367037	99 759
178877	2914	420	2739	3404	408293	73491
280403	551	2275	1260	3824	527380	65958
233039	1059	1680	8864	2144	459813	78978
110408	840	1260	10920	1260	236136	49228
73413	263	350	8094	1094	130419	20125
37231	205	88	4900	219	67725	17588
30844	ő	0	5075	525	48825	13169
27172	0	203	6066	286	45786	11643
21112	Ŭ	205	0000	200	10700	11045
34454	0	0	23093	0	402252	18991
38884	0	0	17678	0	203116	37407
27112	0	0	13905	0	131746	43765
44715	0	0	16242	0	122776	133801
16636	0	Õ	15873	Ō	69548	53207
15554	ŏ	õ	9844	394	57858	52026
17046	Ŏ	122	6991	420	85696	80705
17040	2740	123	0001	1755	142010	105736
4/390	5/40	123	0070	1212	254222	103230
40323	3414	123	1219	1313	234223	77556
31583	1313	636	1559	1148	229371	27030
23134	779	656	3281	0	154018	33306
21739	779	0	3527	0	140606	41919
69319	492	369	13084	123	394541	26251
36997	1025	656	7465	120	216364	20098
44708	123	779	6276	1313	182361	34700
28884	125	,,,,	11526	656	187283	59967
15816	656	656	10278	1212	160400	62061
22740	050	0.0	5499	1515	0/158	40515
33740	102	204	7067	700	115012	47313
44101	123	394	/003	700	113913	4//93
74145	3033	1001	9636	2001	10/412	120575
/5456	10144	0	10//6	2001	303418	144914
52583	5511	2755	5511	1755	405353	/6135
37653	1313	0	6153	1436	16/349	55947
20775	781	0	2755	219	77231	30914
21125	1148	164	1340	335	73550	23552
20303	656	328	205	41	62510	21247
14684	1477	520	369	369	44667	15627
11262	656	0	522	378	33101	16243
12084	820	0	555	520	20558	10275
10001	020	0	650	10	36338	12307
7004	328	0	554	229	20200	12/40
/004	164	0	513	328	1/945	11930
10203	328	10	10	1/4	22118	14489
7824	328	0	338	0	16591	11013
15135	328	0	513	10	28158	15022
6583	21	0	267	21	28343	7404
7465	0	0	697	0	24610	9639
14766	0	0	844	0	46689	33751
21282	0	0	1594	0	61127	19313
18329	0	188	563	47	42283	10313
10828	0	0	609	234	23954	9703
8625	47	0	563	0	22501	10266
5953	94	0	141	0	10219	13969
3445	0	0	492	0	8696	9024
52.50	0	Õ	328	Ō	10828	12633
2625	ñ	ň	497	164	6727	12141
6563	164	ň	328	164	10336	10008
2053	104	ň	0	0	4977	7210
2933	v	v	U	v	7766	1219



Figure 2. Two year time-series fluxes of four species of *Distephanus* at 3800 m. The fluxes at 1000 m are only for six months.

The December maximum is the fourth highest maximum in all *Distephanus* species, with the exception of *Distephanus boliviensis* (Fig. 2c; Plate 1, Fig. 5; Plate 2, Fig. 5) which shows a second highest maximum in December. A flux maximum in *Distephanus octangulatus* (Fig. 2d; Plate 1, Figs. 6, 7; Plate 2, Fig. 3) is observed in May, but is not as well defined in other species of the genus. Overall, the seasonal flux patterns observed in four species of *Distephanus* are similar enough (with a possible exception of *D. octangulatus*; see later discussion of cluster analysis) relative to patterns observed for *Dictyocha* as well as for diatoms, and thus the fluxes of the 4 species can be grouped together.

Dictyocha mandrai (Fig. 3a; Plate 1, Fig. 8; Plate 2, Figs. 1, 2) is numerically a dominant species in the genus (Fig. 3). It shows a well-defined maximum in December but the intensities of the other maxima in year 1 are relatively low. Two variants of *D. mandrai*, a form with variable-shaped apical rods (Fig. 3c) and a form with a pentagonal basal ring (*D. mandrai pentagona*), show high flux values in August. Besides the August peak, *D. mandrai pentagona* shows high flux maxima during October and December. Dictyocha messanensis ? spinosa (Fig. 3b) shows an intense maximum in May which is rather unique compared with other silicoflagellate species observed in year 1 except for *D. octangulatus*.



Figure 3. Two year time-series fluxes of *Dictyocha mandrai*, its closely related forms, and *Distephanus crux*? at 3800 m and 1000 m.

The seasonal flux pattern of *Distephanus crux*? (Fig. 3d) resembles that of the *Actiniscus pentasterias* group (Fig. 4a). The flux patterns of the two taxa are more similar to those of diatoms (Takahashi, 1986) than to the above mentioned silicoflagellate species. The pattern of *Distephanus crux*? has some affinity with that of the remainder of the *Distephanus* taxa (see cluster analysis section). This is of interest and casts doubt about the artificial classification of this taxon as *Distephanus* because of the presence of an apical window, a requirement of the genus. The doubt is based on the similarity of basal ring shape (including variability; from *Dictyocha mandrai* type to *Dictyocha messanensis* type) and size in the studied assemblages; thus one would expect the seasonal flux pattern to be more similar to that of *Dictyocha. Actiniscus* shows seasonal fluxes of up to 160×10^3 individuals m⁻²d⁻¹, which is two orders of magnitude higher than that of *Dictyocha mandrai*.

b. Seasonal flux in year 2 (September, 1983–August, 1984) and its comparison to the year 1 record. The relatively high flux values observed at the beginning of year 2 is part of the August maximum in year 1 which tails off in September/October in year 2. Most silicoflagellate taxa show only one flux maximum in year 2 which occurred in



Figure 4. (a) Two year time-series fluxes the Actiniscus pentasterias group at 3800 m and 1000 m. (b) Two year time-series fluxes of Distephanus and Dictyocha (plotted cumulatively) at 3800 m. (c) Organic carbon, opal, and mass (plotted noncumulatively) at 3800 m. These data are taken from Honjo (1984).

May (Figs. 2-3). The May peak in year 2 is smaller, by approximately a factor of 10, than the major peaks in year 1. A similar trend in decrease of flux peak intensity from year 1 to year 2 is observed in *Actiniscus* (Fig. 4a), opal and mass (Fig. 4c), and diatoms (Takahashi, 1987b).

c. Vertical fluxes at 1000 m and 3800 m trap depths. Species flux observed at the two different trap depths are graphically compared (Figs. 2, 3, 4a). During March through

September, 1983, where two-depth trap data are available, two prominent flux maxima occurred which can be compared with each other. March flux maxima at 3800 m and March flux values at 1000 m of many species tend to show similar levels, especially in *Distephanus*. Exact comparisons of the March maxima between the two depths in terms of lag time is difficult due to lack of the 1000 m data prior to sample No. 13. Comparisons of flux maxima during August, however, are possible. Zero sample lag for August maximum was recognized for *D. quinquangellus* and *D. mandrai* variable form, and one sample lag for *D. speculum*, *D. mandrai*, *D. messanensis* ? *spinosa* and *D. crux* ?, and three sample lags for *D. boliviensis* and *D. octangulatus*. It is noteworthy that many species with lags 0 and 1 are relatively small in size (e.g., *D. speculum*: basal diameter = $18.5 \,\mu$ m, S.D = $2.3 \,\mu$ m, N = 100) and that two species with lag 3 are selectively large (e.g., *D. boliviensis*: basal diameter = $24.6 \,\mu$ m, S.D. = $3.4 \,\mu$ m, N = 4).

4. Discussion

The seasonal fluxes observed for each of the 5 species of *Distephanus* (Figs. 2a-d, 3d) represent unique patterns by themselves. Although details of the patterns are different from one another, the overall configurations are similar so they can be generalized as a group. In the case of year 1, they generally show a pronounced October maximum followed by March and August maxima and an additional December maximum. This type of seasonal flux pattern of *Distephanus* does not exactly conform to patterns shown by other silicoflagellate and diatom species. The closest seasonal flux pattern to that of *Distephanus* is Type 1 of diatoms (Takahashi, 1986). However, in detail most species of *Distephanus* show strong March peaks and generally insignificant flux maxima in May, while the diatoms in Type 1 show insignificant or small March peaks and well-defined May peaks. The relative intensity of October and August peaks in relation to the remaining peaks within a species is similar in both *Distephanus* and the diatom species. Although the intensity is generally not so high, the December peak of *Distephanus* is unique in its occurrence in the early winter season.

There is a definite correlation between the increases of *Distephanus* (and many diatom species) fluxes and the short term decrease of silicate concentrations in surface waters (Reynolds and Thunell, 1985), suggesting that these organisms are productivity indicators. Both of these processes take place within a time scale of 2 weeks or less. The uptake of silicates by these indicators (especially diatoms, a major phytoplankton component) effectively decreases the ambient silicate concentration temporarily until the bloom is completed. It is possible that this bloom is succeeded by nonsiliceous phytoplankton blooms which would explain why nitrate concentration in surface waters does not follow the same trend (Reynolds and Thunell, 1985). Correlation between *Distephanus* (in conjunction with diatoms) blooms and silicate concentrations are observed for time intervals corresponding to the March, May, and August maxima



Figure 5. Time-depth plots of monthly averaged (a) temperature (1956–1981 data); (b) salinity (1956–1981); and (c) daily solar radiation at the sea surface (1960–1967) at Station PAPA. The numbers above each line are water depth in meters. The temperature and salinity data are from Tabata and Peart (1985). The figure of solar radiation is redrawn from Anderson *et al.* (1977).

of the fluxes. An exception is the September-October period when the silicate concentration keeps increasing with time as a result of convective mixing caused by atmospheric cooling.

That 54% of D. mandrai annual flux in year 1 occurred during a 3-month period, from October to January, is an important fact, especially for paleo-oceanographic implications. If specimens of D. mandrai from different seasons are equally preserved, fossil records represent mostly fall/winter signals and few signals from other seasons. The specimens that are supplied to the sea floor in high flux seasons may actually be preserved better than those supplied in low flux seasons due to rapid burial with accompanying large amounts of biogenic opal. The unique timing of this pronounced maximum is considered to be a response to given environmental conditions. Although slightly earlier than that of D. mandrai, a similarly offsetting flux maximum from the October opal flux maximum is shown by the diatom species, Coscinodiscus marginatus (Takahashi, 1986). From fall through early winter, the water temperature in the euphotic layer (maximum is in September) decreases gradually through March or April, depending on depth. During that time, little salinity change is observed; incident solar radiation decreases until December (Fig. 5). Nutrient levels are continuously increasing from August minima to winter maxima (Anderson et al., 1977; Wong and Honjo, 1984). The flux maximum observed in December at 3800 m results from late November production in the euphotic zone (see a later section on sinking speeds). The late November production of D. mandrai corresponds with (1) relative high levels of available nutrients. (2) relatively low but adequate levels of incident solar radiation for this species, and (3) intermediate levels of water temperature ($\sim 8.5^{\circ}$ C at the surface) (Fig. 5), and is perhaps related to biological competition.

The fact that there is no December or August flux maximum in year 2 for any silicoflagellate species leads to questions on the range of year to year variability. The records from year 2 appear to deviate from the other 3 years, as suggested by unpublished mass flux data for years 3 and 4 which show that the flux maxima at 3800 m occurred about October/November, March/May, and August/September (Honjo, pers. comm., 1986). The causes of the large year 1 to year 2 flux variability observed have not yet been determined; this will require long-term time-series flux studies as well as detailed analyses of environmental parameters. The possible effects of the 1982–1983 large El Niño event in the Pacific low to middle latitudes should not be overlooked (Takahashi, 1987b).

The silicoflagellate flux maxima often occurs slightly later than that of diatoms, about 2 weeks, as shown by the November/December and August peaks of year 1 and the May peak of year 2. This suggests that diatoms respond more quickly to favorable environmental conditions than silicoflagellates do. Biological competition may be an important factor in offsetting the flux maxima. Alternatively, a difference in the doubling times of cell populations of diatoms and silicoflagellates may be responsible for the offset in the fluxes. However, in the light of the wide range of diatoms sizes (i.e., 3 orders of magnitude), and in their morphology and physiology, the alternative is an unlikely explanation.

a. Species affinity among silicoflagellate taxa. Seasonal flux patterns and flux values (individuals $m^{-2}day^{-1}$) are 2 major signals in the time-series flux record. Of these, the seasonal flux pattern can be used to clarify taxanomic affinities among silicoflagellate species. Cluster analysis (e.g., Davis, 1973) is a useful technique for this definition. To maximize comparison of the patterns and to estimate the effect of flux values, correlation coefficients were used (rather than using Euclidian distance) for constructing a dendrogram (Fig. 6). Most of the obvious similarities and differences in the patterns can be readily extracted without using a cluster technique. However, some subtle differences and similarities which cannot be detected by visual observation can be numerically expressed by this technique.

D. speculum and D. quinquangellus are the most closely clustered groups among the species studied. The major morphological differences is the shape of the basal ring, whether hexagonal or pentagonal. Some authors attribute these two forms to two varieties of a single species (e.g., Ciesielski, 1975), while others regard them as two distinct species (e.g., Bukry, 1973). As the pentagonal (or very rarely, heptagonal) form is usually rare, it is often lumped together with the dominant hexagonal form. As the seasonal flux patterns for both forms are very similar, they suggest that two taxa could actually represent a single species. However, until conclusive evidence based on further biological information becomes available, it is probably wise to separate these



Figure 6. Dendrogram of Silicoflagellates and *Actiniscus* seasonal flux patterns constructed by the averaging method, using arithmetic averaging of correlation coefficients.

two forms because lumping of split counts can be easily done while the reverse is usually not possible.

It is noteworthy that the seasonal pattern of *Distephanus octangulatus* shows a relatively close affinity with that of *Dictyocha messanensis*? spinosa (r = 0.70) despite different generic assignment, in addition to those of *Distephanus speculum* (r = 0.73) and *Distephanus quinquangellus* (r = 0.70). Although *D. octangulatus* is placed in the genus *Distephanus*, its morphology and size are vastly different from all other species in the genus. The skeletal surface ornamentation observed on *D. octangulatus* also occurs in species assigned to the genus *Dictyocha*, but not in species in *Distephanus* except for the former (Plate 2). This implies that the generic assignment of *D. octangulatus* should be reconsidered.

The unique high August maximum presented by the variety of D. mandrai with variably shaped apical rods (Fig. 3c) caused the extreme position of this counting group in the dendrogram (Fig. 6). I consider this form as part of D. mandrai and not as an independent subspecies. The variably shaped skeletons occur throughout the year, but are more common during the 1983 summer season when other fluxes such as total mass and diatoms are also significantly high. The August maximum of this form of D. mandrai may result from high nutrient concentrations and high temperatures. Such variable morphology in the skeletons of D. mandrai has been observed in coastal waters of the Gulf of San Matias (Frenguelli, 1935) and in Lake Hiruga, Japan, where salinity varies seasonally from 13 to 27% in surface waters (Shitanaka, 1983). Nutrient levels in these coastal environments are apparently higher than those in

Taxon	Mass	Opal	Organic carbon
Distephanus speculum	0.62	0.66	0.44
Distephanus quinquangellus	0.56	0.55	0.51
Distephanus boliviensis	0.66	0.63	0.61
Distephanus octangulatus	0.68	0.66	0.33
Distephanus octangulatus (>1 apical window)	0.77	0.78	0.51
Distephanus crux ?	0.64	0.70	0.35
Dictyocha mandrai	0.38	0.40	0.14
Dictyocha mandrai (variable apical rods)	0.76	0.70	0.73
Dictyocha messanensis ? spinosa	0.55	0.48	0.32
Total Silicoflagellates	0.63	0.66	0.41
Actiniscus pentasterias group	0.90	0.92	0.57

Table 2. Correlation coefficient r between silicoflagellate taxa and mass, opal, and organic carbon fluxes at 3800 m for two year record.

pelagic realms and may be responsible for the variable shapes of the skeletons. The salinity change is very small at Station PAPA (Fig. 5b).

b. Relations with other fluxes and factors. Seasonal flux patterns of silicoflagellates are similar in general but different in detail from those of diatoms (Takahashi, 1986). For example, the March flux maximum of silicoflagellates is as high as that of August, whereas diatom flux shows a rather low maximum in March relative to a high August maximum. This greatly contributes to the level of correlation coefficients between silicoflagellate species and total mass, opal, or organic carbon fluxes (Table 2). All coefficients for total mass and opal are significant at the 99% level and thus silicoflagellates are considered to co-occur with other productivity components. However, the coefficients are not as good as those between diatoms and total mass or opal fluxes (Takahashi, 1986) which are not totally independent of one another. Since silicoflagellates contribute an insignificant amount compared to opal flux (<0.4% using 0.34 ng shell⁻¹ of Takahashi, 1987a), the coefficient of correlation with organic carbon reflects the labile nature of organic carbon which is largely recycled in the upper parts of the water column.

In order to understand factors controlling fluxes of silicoflagellate species it is necessary to reduce a large number of flux patterns into small groups so that we can correlate the patterns with environmental parameters. Results of R-mode factor analysis (Table 3) show in factor 1 high scores of D. speculum and D. quinquangellus and intermediate scores for D. boliviensis, D. octangulatus and D. crux ?. Factor 1 is interpreted as the productivity factor since fluxes of these species show relatively high

	Factors				
Taxon	1	2	3	4	
Distephanus speculum	0.877	0.195	0.157	0.216	
Distephanus quinquangellus	0.856	0.153	0.205	0.230	
Distephanus boliviensis	0.486	0.191	0.300	0.410	
Distephanus octangulatus	0.567	0.761	0.087	0.176	
Distephanus crux?	0.578	0.096	0.187	0.461	
Dictyocha mandrai	0.257	0.204	0.079	0.890	
Dictyocha mandrai (variable apical rods)	0.182	0.159	0.958	0.081	
Dictyocha mandrai pentagona	0.388	0.158	0.120	0.354	
Dictyocha messanensis? spinosa	-0.061	0.962	0.150	0.092	
Variance	2.969	2.282	1.168	1.473	

Table 3. Matrix of varimax rotated factor score.

correlation with total mass, opal fluxes (Table 2) as well as productivity indicator diatom species. Factor 2 is related to high light intensity and *D. messanensis*? spinosa and *D. octangulatus* show high scores. Factors 3 and 4 whose high scores are represented by *D. mandrai* variable form and *D. mandrai*, respectively, show seasonal relationships with summer and fall/winter respectively in year 1 but not in year 2. Acquisition of longer term flux records than two years will be required to clarify these trends.

c. Sinking processes. The time necessary for silicoflagellates in aggregates to sink through the water column from 1000 m to 3800 m is approximately equal to, or shorter than, the sample time interval of 16 days as shown by cross-correlation analysis (Figs. 2–3; Table 4). Cross-correlation seeks the best match of flux maxima and minima

Table 4. Correlation coefficient r between 1000 m and 3800 m data sets for silicoflagellate taxa and *Actiniscus* using the cross-correlation method. Results of sample lag 0, 1, and 2 are listed. The highest r values in the cross-correlation are underlined. Note that the value for total silicoflagellates at lag 0 is the highest. This is due to the effect of amplifying signals and to cancelling noises in silicoflagellate species flux.

	Sample lag			
Taxon	0	1	2	
Distephanus speculum	0.84	0.50	-0.05	
Distephanus quinquangellus	$\overline{0.60}$	0.32	0.12	
Distephanus octangulatus	0.69	0.74	0.73	
Distephanus crux ?	0.68	0.71	0.32	
Dictyocha mandrai	0.57	0.34	-0.41	
Dictyocha messanensis ? spinosa	0.61	0.42	0.36	
Total Silicoflagellates	0.85	0.45	-0.09	
Actiniscus pentasterias group	0.46	0.69	0.36	

417

between the two depths and determines appropriate time lags. The examined silicoflagellate species showed that time lag 0 and lag 1 led the above sinking time. This corresponds to 175 m day⁻¹ or greater sinking speed, and implies accelerated sinking relative to single particle sinking of 1.4 m day^{-1} of silicoflagellates (Takahashi, 1987a). The three most abundant species, D. speculum, D. mandrai, and D. messanensis ? spinosa, show the highest correlation coefficient between 1000 m and 3800 m depths with 0 sample lag (Table 4), suggesting much faster sinking speeds than 175 m day^{-1} . This may be a realistic approximation, because of their smaller size range, silicoflagellates are grazed more efficiently by herbivores than diatoms are. Within silicoflagellate taxa, as noted in the Results section, relatively small sized species tend to sink faster than those of larger size, suggesting a possibility of differential grazing by herbivores. The herbivores are probably excreting readily disintegrated fecal matter (but not pellets) that sinks faster than marine snow. Silicoflagellate transport via biogenic aggregates (Plate 1, Fig. 15) was also documented in the equatorial Atlantic and Pacific (Takahashi, 1987a), which supports the above observation. Perhaps both transport mechanisms via fecal matter and marine snow are responsible for the fluxes observed.

Although it was hypothesized earlier that two copepod species of *Neocalanus* were primary grazers in the subarctic Pacific, the hypothesis has not been vigorously tested yet (Frost *et al.*, 1983). In light of such unsettled grazers and the present observation of insignificant numbers of fecal pellets in the flux, we must consider other grazing candidates such as salps more critically than before.

We note that the Actiniscus pentasterias group, dinoflagellates, shows a maximum coefficient at lag 1 which is sufficiently different from the coefficient at lag 0 while two species of silicoflagellates with maximum coefficient at lag 1 show similar coefficients at lag 0. The A. pentasterias group's sinking mechanism is more like that of diatoms rather than that of silicoflagellates based on results of cross correlations.

Similar values of flux assemblages at 1000 m and 3800 m at sample lag 0 or lag 1 indicate that the effect of dissolution of sinking silicoflagellates is insignificant. The abundance patterns of the assemblage (in percent) at 1000 m are also repeated at 3800 m with a slight time offset (Fig. 7). If significant uniform dissolution and/or species preferential dissolution occurred, the flux values and percentages of soluble and resistant taxa would be different at 3800 m and at 1000 m. That no significant differences are observed suggests that sinking silicoflagellates are unaffected by dissolution.

d. Use of percent signals. Despite considerable change in flux values from year 1 to year 2, the patterns of relative abundance of silicoflagellate species in the flux did not change very much (Fig. 8a). The detailed values of the percentages differ in the 2-year record, but the overall patterns are similar so that the percent signals may be used for a variety of applications, including comparisons with percentages preserved in fossils.





Figure 7. Cumulative percents of silicoflagellate species flux: (a) at 1000 m, May-Sept., 1983; (b) at 3800 m, Sept. 1982-Sept. 1983. Legends for a and b are the same. Shaded areas represent *Distephanus*.

Essentially, *D. speculum* and *D. mandrai* contribute to the largest of the percent values while the remaining species contribute a small percentage of the assemblages. For correlation analysis it is necessary to take a logarithm of fluxes because they tend to vary by orders of magnitude with time whereas percentages vary within a factor of ten. A good correlation of percent *D. speculum* with the logarithm of *D. speculum* flux is seen (r = 0.80) while a fair anti-correlation of percent *D. mandrai* with the logarithm of *D. speculum* flux is evident (r = -0.64). These values are significantly high



Figure 8. (a) Silicoflagellate cumulative percent data for two years. Legends are the same as in Figure 7. (b) Fluxes of *Distephanus speculum* for two years at 3800 m. (c) Plot of silicoflagellate diversity index at 3800 m vs. time. Note that index values are antirelated with both fluxes and % D. speculum.

considering that critical values at 95% and 99% levels with 45 degrees of freedom are 0.288 and 0.372, respectively (Rohlf and Sokal, 1969). The flux of D. speculum in a given interval of time can be predicted once we have a percent value.

$$\log_{10} F_{\rm Ds} = 0.0318 \ P_{\rm Ds} + 2.99 \tag{2}$$

where F_{Ds} and P_{Ds} are flux and proportion of *D. speculum*, respectively.

f. Actiniscus as a productivity indicator. The flux of the Actiniscus pentasterias group is well correlated with the total mass flux (Table 2) and thus can be used to predict it.

$$F_{\rm mass} = 0.00203 \; F_{\rm Ap} + 14.9 \tag{3}$$

where F_{mass} and F_{Ap} are fluxes of total mass and the Actiniscus pentasterias group, respectively. The total mass flux predicted using the Actiniscus pentasterias group flux data with the above regression equation, conforms with the observed total mass flux (Fig. 9). The application of this group in paleoproductivity assessment depends on future studies on downcore preserved samples. Actiniscus is frequently well preserved, often better than silicoflagellates, in many parts of the world (e.g., Dumitrica, 1973; Perch-Nielsen, 1975; Orr and Conley, 1976; Takahashi, unpublished) and deserves attention.

g. Relationship with fossil records in surface sediments. A few percent or less of silicoflagellates that arrive at the sea floor are preserved in the surface sediments of the study area. To compute the percent of skeletons preserved, the following values were used: 10^5 skeletons per gram⁻¹ of dry sediment (Poelchau, 1976), dry density of 0.8 g cm⁻³, sedimentation rate of 1 cm 10^3 yr⁻¹ (see Takahashi, 1986, for details). Use of these values yields 8×10^4 skeletons cm⁻² 10^3 yr⁻¹ that have accumulated in the surface sediments. A mean flux for the 2 year period yields 143,261 skeletons m⁻² day⁻¹ which is equivalent to 5.23×10^6 skeletons cm⁻² 10^3 yr⁻¹. This means that 1.5% of the silicoflagellates in the flux are preserved. Since Poelchau's (1976) values are actually slightly less than 10^5 skeletons g⁻¹ of sediment, approximately 1% of silicoflagellates are preserved. This value is similar to the percentage of diatoms (2%) preserved in the sediments in the same area (Takahashi, 1986). Silicoflagellate dissolution (~ 99% in this case) mostly occurs at the sediment/water interface or in the sediments as shown elsewhere (Takahashi, 1987a); this is also true for other siliceous microfossils (Takahashi, 1983, 1986).

A large discrepancy is observed between species proportion in the flux and in surface sediments (Table 5). This discrepancy may be superficial resulting from different methods used for analysis. In particular, Poelchau's (1976) concentration technique involving settling and decantation may have biased the proportions between species. More likely, however, it reflects the preferential dissolution of susceptible taxa. For

coefficent of 0.46.



Figure 9. Observed (Honjo, 1984) and predicted total mass fluxes at 3,800 m, Station PAPA, 1982–1984. The predicted values were computed using the *Actiniscus pentasterias* group fluxes and a linear regression equation for total mass and *A. pentasterias* group fluxes.

instance, the sum of *D. speculum* and its variants is about 50% in the flux and only 10% occurs in the sediments. *Distephanus speculum* (and *D. quinquangellus*) is one of silicoflagellate taxa of the study area with the smallest and thinnest skeleton. Thus it is likely to be preferentially dissolved relative to other taxa with more robust skeletons. There is a good correlation between the dissolution index (Johnson, 1974) and the thickness of opal skeletons (Takahashi and Hurd, unpublished). *Dictyocha mandrai, D. messanensis* ? *spinosa* and *Distephanus octangulatus* are larger and more robust than *D. speculum* and thus are relatively better preserved. In the case of such preferential dissolution/preservation, for example, 99.8% of *D. speculum* and 98.5% of *D. mandrai* flux to the sea floor would have to be dissolved to acquire the values in the sediment. Alternatively, components of present-day production may be different from that in the surface sediments which represent the past several hundred years or more, causing such a high percent of *D. mandrai* (60%). Detailed down core studies in the future are needed to evaluate all of these three possible explanations.

5. Summary and conclusions

The high resolution time-series flux study of silicoflagellates showed considerable intra-annual and interannual flux variabilities within the studied 2-year period at

Taxon	% in Flux	% in surface sediments
Distephanus speculum	47	10
Distephanus boliviensis	6	—
Distephanus octangulatus	5	15
Distephanus crux ?	1	
Total Distephanus	59	25
Dictyocha mandrai	37	60
D. mandrai (variable apical rods)	1	
Dictyocha messanensis? spinosa	3	15
Total Dictyocha	41	75

Table 5. Comparison of relative abundances in flux and surface sediment assemblages in the study area. The percentages in flux are listed taking mean values of the two year record. The surface sediment data are taken from Poelchau (1976).

pelagic Station PAPA in the northeastern North Pacific. Seasonal flux patterns tend to differ between morphologically distant species groups while the patterns are similar within a morphologically confined group.

Silicoflagellate flux is generally numerically dominated by two major taxa, *Distephanus speculum* and *Dictyocha mandrai*, which contribute on the average 83% $(6.7\% < \sigma < 10.1\%$ at 95% confidence limits) of the total sinking assemblages. *D. speculum* is considered to be a productivity indicator based on its affinity of the seasonal flux pattern to the total mass, opal, and diatom flux patterns. Seasonal patterns show by *Distephanus speculum* and its two other closely related species are similar to the seasonal pattern of diatom flux Type 1, which is a productivity indicator. *D. mandrai* carries its unique late fall/early winter signals. More than half of the cummulative annual flux of this species occurred in a three month period (October, 1982–January, 1983), implying a significance to the fossil record.

The silicoflagellate seasonal flux patterns are generally similar enough to those of total mass, opal, and diatoms so that an assessment of surface water opal or total mass productivity using silicoflagellates is possible. Moreover, the silicoflagellate flux is nearly an independent measure from opal due to its insignificant mass contribution to opal. Although general patterns are similar, as mentioned above, details of silicoflagellate seasonal flux deviate from opal, whose main contributor is diatoms, providing unique signals. Some of the silicoflagellate flux maxima tend to be temporally offset from diatoms by one sample interval, indicating a different response to environmental changes from that of diatoms.

The level of flux changes significantly from year 1 to year 2. Despite such annual flux variations in the 2-year record, the annual patterns of species percents are similar for the two years, suggesting a possible use of percent data for productivity assessment.

Sinking speeds of silicoflagellates are determined to be approximately 175 m day⁻¹ or greater based on cross-correlation analysis of the synchronized 1000 m and 3800 m

samples. Generally, silicoflagellates sink faster than diatoms. Both fecal material and marine snow are likely mechanisms for the observed accelerated sinking speeds. No significant dissolution of silicoflagellate skeletons occurs in the water column due to protection and accelerated sinking via aggregates. Therefore, species composition and absolute flux of each species in the sinking assemblages at 1000 m and 3800 m are similar, with or without the slight temporal offset.

Approximately 1% of silicoflagellates supplied to the sea floor are preserved and the bulk of the supplied skeletons are dissolved in the sediments. The preferential dissolution of D. speculum, a solution susceptible taxon, relative to D. mandrai, a solution resistant taxon, is a probable reason for the preserved species percentages in the surface sediments of the study area.

Acknowledgments. I am grateful to Dr. Susumu Honjo at WHOI for his indispensable leadership in conducting the PARFLUX sediment trap program at Station PAPA with the cooperation of Dr. C. S. Wong at the Institute of Ocean Sciences, Canada. I thank Drs. D. J. DeMaster, M-P. Aubry, and anonymous reviewers for critical reviews of the manuscript and Drs. S. Honjo and D. Lazarus for helpful discussions. Thanks are also due to John Billings for skillful technical assistance, and to Steve Manganini, Vernon Asper, R. Ostermann, A. Spencer, R. Bellegay and many others who participated in numerous cruises to Station PAPA including the captains and crews of C.F.A.V. Endeavor and Parizeau. Special thanks go to Emily Evans for her editorial assistance and typing.

This study was funded by NSF Grants OCE-8502472 and OCE-8608255. The collection of sediment trap samples was supported by NSF Grants OCE-8309024 and OCE-8417106.

This is Woods Hole Oceanographic Institution Contribution No. 6200.

REFERENCES

- Anderson, G. C., R. K. Lam, B. C. Booth and J. M. Glass. 1977. A description and numerical analysis of the factors affecting the processes of production in the Gulf of Alaska. Univ. of Washington Dept. of Oceanography Special Report 76 (Ref. M-77-40). Also as NOAA03-5-022067, Ev. Assess. Alaskan Cont. Shelf, VII, pages 477-798.
- Anonymous. 1981. Oceanographic observations at Ocean Station P. Pacific Marine Science Report Series, volumes 1 to 125. Institute of Ocean Sciences, Patricia Bay, Sidney, B.C., Canada.
- Bukry, D. 1973. Coccolith and silicoflagellate stratigraphy, Deep Sea Drilling Project, Leg 18, Eastern North Pacific, *in* Initial Reports Deep Sea Drilling Project, L.D. Kulm *et al.*, eds., U.S. Government Printing Office, Washington, DC, 18, 817–831.
- Ciesielski, P. F. 1975. Biostratigraphy and paleoecology of Neogene and Oligocene silicoflagellates from cores recovered during Antarctic Leg 28, Deep Sea Drilling Project, *in* Initial Reports Deep Sea Drilling Project, D. E. Hayes *et al.*, eds., U.S. Government Printing Office, Washington, DC, 28, 625-691.
- Davis, J. C. 1973. Statistics and Data Analysis in Geology. John Wiley & Sons, NY, 550 pp.
- Dumitrica, P. 1973. Cenozoic endoskeletal dinoflagellates in southwestern Pacific sediments cored during Leg 21 of the DSDP, *in* Initial Reports Deep Sea Drilling Project, J. E. Andrews *et al.*, eds., U.S. Government Printing Office, Washington, DC, 21, 819–884.
- Frenguelli, J. 1935. Variaciones de Dictyocha fibula en el Golfo de San Matias (Patagonia septentrional): An. Mus. Argentina Cienc. Nat. "Bernadino Rivadavia," v. 38, Protistol. No. 4, 265-281.

- Frost, B. W., M. R. Landry, and R. P. Hassett. 1983. Feeding behavior of large calanoid copepods Neocalanus cristatus and N. plumchrus from the subarctic Pacific Ocean. Deep-Sea Res., 30, 1-13.
- Honjo, S. 1978. Sedimentation of materials in the Sargasso Sea at a 5,367 m deep station. J. Mar. Res., 36, 469-492.
- ----- 1984. Study of ocean fluxes in time and space by bottom-tethered sediment trap arrays: a recommendation. Global Ocean Flux Study Workshop, National Research Council, Washington, DC, 304-324.
- Honjo, S. and C. S. Wong. 1985, abstract. Interannual time series sediment trap experiment at Ocean Station Papa, North Pacific. EOS, 66, 1282.
- Johnson, T. C. 1974. The dissolution of siliceous microfossils in surface sediments of the eastern tropical Pacific. Deep-Sea Res., 21, 851–864.
- Ling. H. Y. 1970. Silicoflagellates from central North Pacific core sediments. Bull. Amer. Paleont., 58, 85-129.
- —— 1977. Late Cenozoic silicoflagellate and ebridians from the eastern North Pacific region. Ist International Congress Pacific Neogene Stratigraphy Proceedings, Tokyo, 205–233.
- Manganini, S. J. and S. Honjo. 1985, abstract. *In-situ* preservation of biogenic material collected by a deep ocean sediment trap. EOS, 66, 1292.
- Murray, D. and H. Schrader. 1983. Distribution of silicoflagellates in plankton and core top samples from the Gulf of California. Mar. Micropaleont., 7, 517–539.
- Orr, W. N. and S. Conley. 1976. Siliceous dinoflagellates in the northeast Pacific rim. Micropaleont., 22, 92-99.
- Perch-Nielsen, K. 1975. Late Cretaceous to Pleistocene Archaeomonads, Ebridians, Endoskeletal Dinoflagellates, and other Siliceous Microfossils from the Subantarctic Southwest Pacific, DSDP, Leg 29, *in* Initial Reports Deep Sea Drilling Project, Vol. 29:873–907, J. P. Kennett *et al.*, eds., Washington, DC (U.S. Government Printing Office).
- Pielou, E. C. 1969. An Introduction to Mathematical Ecology. Wiley-Interscience, NY 286 pp.
- Poelchau, H. S. 1976. Distribution of Holocene silicoflagellates in North Pacific sediments. Micropaleont., 22, 164–193.
- Reynolds, L. and R. C. Thunell. 1985. Seasonal succession of planktonic foraminifera in the subpolar North Pacific. J. Foraminiferal Res., 15, 282–301.
- Rohlf, F. J. and R. R. Sokal. 1969. Statistical Tables. Freeman and Company, San Francisco, 253 pp.
- Schrader, H-J. and T. Baumgartner. 1983. Decadal variation of upwelling in the central Gulf of California, *in* Coastal Upwelling, Part B., J. Thiede, and E. Suess, eds., Plenum Publishing Co., 247–276.
- Shitanaka, M. 1983. Silicoflagellate remains in the sediments of Lake Hiruga, Fukui, Japan.
- Mizunami fossil Museum Research Report, v. 10, 171-180 (in Japanese).
- Tabata, S. and J. L. Peart. 1985. Statistics of oceanographic data based on hydrographic/STD casts made at Ocean Station P during August 1956 through June 1981. Canadian Data Report of Hydrography and Ocean Sciences, No. 31.
- Takahashi, K. 1983. Radiolarian biocoenosis-thanatocoenosis relationship in pelagic oceans. Amer. Assoc. Petrol. Geol. Bull., 67, 556-557.

— 1985. Two year time-series fluxes of silicoflagellates and *Actiniscus*: size fractioned results from subarctic Pacific Station PAPA, 1982–1984. Woods Hole Oceanographic Institution Technical Report, WHOI-85-41.

----- 1986. Seasonal fluxes of pelagic diatoms in the subarctic Pacific, 1982-1983. Deep-Sea Res., 33, 1225-1251.

trap stations, in Ocean Biocoenosis, S. Honjo, ed., Micropaleontology Press, NY.

Wong, C. S. and S. Honjo. 1984, abstract. Material flux at weather station PAPA, high frequency time series observations through production cycles. EOS, 65, 225.

^{----- 1987}b. Response of subarctic Pacific diatom fluxes to 1982-1983 El Niño. J. Geophys. Res., (submitted).