九州大学学術情報リポジトリ Kyushu University Institutional Repository

# CERTAIN OPERATORS OF RIEMANN-LIOUVILLE TYPE AND STATIONARY PROCESSES

Seguchi, Tsunetami College of General Education, Kyushu University

https://doi.org/10.15017/1448939

出版情報:九州大学教養部数学雑誌. 7 (2), pp.23-35, 1970-09. College of General Education, Kyushu University バージョン: 権利関係: Math. Rep. VII-2, 1970.

# CERTAIN OPERATORS OF RIEMANN-LIOUVILLE TYPE AND STATIONARY PROCESSES

By

Tsunetami SEGUCHI [1969. 8. 15.]

#### 1. Introduction.

T. Hida [1] has proposed the problem that "what properties has the stationary process  $x(t) \equiv x(t,\omega)$ , which satisfies the formal equation (1.1)  $D^{\alpha}x(t) = \dot{\xi}(t)$ , where  $D \equiv d/dt$ ,  $0 < \alpha < 1$  and  $\dot{\xi}(t)$  is a white noise". K. Yosida [2] had interpreted and studied the operators such as  $D^{\alpha}$ , which proposed by A. V. Balakrishnan [3], as the fractional powers of infinitesimal generators of the semi-groups of bounded linear operators  $g_t$  (t > 0) on a Banach space X into X.

In this note, we shall interprete and study the operators such as  $D^{\alpha}$ ,  $(D+\beta I)^{\alpha}$ ,  $(D+\beta I)^{-\alpha}$ , as the special cases of shift-commutative linear operators which are defined by T. Seguchi [4] and [5]. And then we shall refer to the properties of stationary processes which satisfy the stochastic functional equations concerning to the operators such as above S-C.L.Op.s of Riemann-Liouville type.

# 2. Preliminaries.

DEFINITION 2.1. Let F be the all complex- (or real-) valued functions defined on  $\mathbb{R}^{10}$ , and let  $(\Lambda, D, \{\tau_t\}_{t \in \mathbb{R}})$  be such that:

(1)  $\{\tau_t\}_{t\in R}$  is the shift operator defined on F; i.e. for any  $f\in F$  and for any  $t\in R$ 

$$\tau_t f(\cdot) \equiv f(\cdot + t).$$

1)  $R \equiv (-\infty, \infty)$ : "the set of all real numbers".

(2) D is a non-trivial complex (or real) linear sub-space of F, i.e.  $D \neq (0)$ , and satisfies the condition

(L 1) τ<sub>t</sub>D⊂D, for any t∈ R.
(3) Λ is a linear operator defined on D and satisfies the condition
(L 2) ΛD⊂F ≡ 𝒴(τ<sub>t</sub>)<sup>2</sup>).
(4) Between the linear operator Λ and the shift τ<sub>t</sub>, the condition
(L 3) for any f∈D and for any t∈ R

$$\tau_t(\Lambda f) = \Lambda(\tau_t f)$$

holds.

Then we shall call the linear operator  $\Lambda$  is the *shift-commutative linear* operator (S-C.L.Op.) defined on D w.r.t.  $\tau_i$ . The notation  $\widetilde{\Lambda}$  means the class of all S-C.L.Op.s defined above and  $\widetilde{\mathscr{D}}(\Lambda)$  means the domain D of  $\Lambda \in \widetilde{\Lambda}$ .

Let  $\Lambda \in \widetilde{\Lambda}$ , and  $\Lambda f(t) \equiv (\Lambda f)(t)$  be the value of  $\Lambda f$  at t, then the following relations hold; there exist two functions such that for any  $\lambda \in \mathscr{A}(\Lambda) \equiv \{\lambda; e^{i\lambda} \in \widetilde{\mathscr{D}}(\Lambda), \lambda \in R\}$  and  $z \in \mathscr{B}(\Lambda) \equiv \{z; e^{z} \in \widetilde{\mathscr{D}}(\Lambda), z \in C\}^{3}$ ,

(2.1)  $\Lambda e^{i\lambda t} \equiv e^{i\lambda t} G_A(\lambda), \quad \Lambda e^{zt} \equiv e^{zt} C_A(z), \text{ for every } t \in R.$ 

DEFINITION 2.2. The above two functions  $G_{\Lambda}(\lambda)$  and  $C_{\Lambda}(z)$  are called the generating function (g. f.) and the characterizing function (c. f.) of S-C. L. Op.  $\Lambda \in \widetilde{\Lambda}$ , respectively.

Let us use the following notations:

% "#" the space of all rapidly decreasing functions in the sense of L.
Schwartz [6]".

$$M = \left\{ \mu; \int_{-\infty}^{\infty} |\varphi(\lambda)|^2 \mu(d\lambda) < \infty, \text{ for every } \varphi \in \mathscr{S}, \text{ non-negative measures} \right\}$$
$$M_k = \left\{ \mu; \int_{-\infty}^{\infty} \mu(d\lambda) / (1+\lambda^2)^k < \infty, \ \mu \in \mathbf{M} \right\}.$$
$$L^2(\mu; \mathscr{S}) = \left\{ f; \int_{-\infty}^{\infty} |f(\lambda)\varphi(\lambda)|^2 \mu(d\lambda) > \infty, \text{ for every } \varphi \in \mathscr{S} \right\}.$$
$$L^2(\mu; k) = \left\{ f; \int_{-\infty}^{\infty} |f(\lambda)|^2 / (1+\lambda^2)^k \mu(d\lambda) < \infty \right\}.$$
$$\mathcal{Q}(\mathfrak{B}, \mathbf{P}) = \text{``the basic probability space''.}$$
$$L^2(\mathcal{Q}) = \left\{ x; \ E[|x(\omega)|^2] < \infty \right\}.$$
$$\mathfrak{E} = \text{``the totality of (weakly) stationary distributions''.}$$

<sup>2)</sup>  $\mathscr{D}(*)$  means the *domain* of a operator "\*".

<sup>3)</sup> C means "the set of all complex numbers".

(2.2) 
$$[\beta/(D-\alpha I)]f(t) \equiv -\beta \int_{0}^{0} e^{\alpha u} \tau_{-u} f(t) du,$$

where  $\alpha, \beta \in C, \neq 0$  and

$$\int_{0}^{0} e^{\alpha u} \tau_{-u} f(t) du \equiv -\int_{0}^{t} e^{\alpha (t-s)} f(s) ds \equiv -e^{\alpha t} F_{\alpha}(t)$$

$$= \begin{cases} \int_{-\infty}^{0} e^{\alpha u} \tau_{-u} f(t) du = \int_{t}^{\infty} e^{\alpha (t-s)} f(s) ds, & \text{if } \Re t \alpha > 0, \\ -\int_{0}^{\infty} e^{\alpha u} \tau_{-u} f(t) du = \int_{-\infty}^{t} e^{\alpha (t-s)} f(s) ds, & \text{if } \Re t \alpha < 0, \\ -e^{\alpha t} F_{\alpha}(t), & \text{if } \Re t \alpha = 0. \end{cases}$$

(2.3) 
$$(1/[(D-\alpha I)(D-\alpha I)])f(t) \equiv -(1/b) \int_{0}^{0} e^{au} \sin bu \tau_{-u} f(t) du,$$

where  $\alpha \equiv a + ib$  (a,  $b \in R$ ,  $\neq 0$ ).

DEFINITION 2.3. Let  $T_t(t \in R)$  be the *shift operator* on  $\mathfrak{S}$ , such that for any  $x \in \mathfrak{S}$ , for every  $\varphi \in \mathscr{S}$  and for every  $t \in R$  the relation

$$T_t x(\varphi) \equiv x(\tau_{-t} \varphi)$$

holds, where  $\tau_t$  is the shift defined on F.

Let K(t) be a right continuous step-function defined on R, and has finite jumping points, i.e.

4)  $L^{2}$ {\*} means the  $L^{2}$ -space generated by "\*".

T. SEGUCHI

$$K(t) \equiv \sum_{t_n \leq t} k_n, \quad \sum_n |k_n| < \infty, \quad k_n \in R \text{ or } C, \quad n = 1, 2, \cdots.$$

And for any  $x \in \mathfrak{S}$  and for every  $\varphi \in \mathcal{S}$ , let us put

(2.4) 
$$\Lambda x(\varphi) \equiv \sum_{n} T_{t_n} x(\varphi) k_n = \sum_{n} x(\tau_{-t_n} \varphi) k_n = \int_{-\infty}^{\infty} x(\tau_{-t} \varphi) dK(t).$$

The all these operators  $\Lambda$  such as above will be called the *shift-commuta*tive linear operators on  $\mathfrak{S}$ , and are denoted class  $\Lambda_s$ .

PROPOSITION 2.1 (Proposition 3.1.1).<sup>5)</sup> If the above operator  $\Lambda$  on  $\mathfrak{S}$  defined by (2.4) is corresponding to the S-C.L.Op.  $\Lambda \in \widetilde{\Lambda}$  such that, for any  $f \in F$ 

(2.5) 
$$\Lambda f(\cdot) \equiv \sum_{n} \tau_{tn} f(\cdot) k_n = \sum_{n} f(\cdot + t_n) k_n = \int_{-\infty}^{\infty} f(\cdot + s) dK(s) = \int_{-\infty}^{\infty} \tau^s f(\cdot) dK(s).$$

Then we have the following results:

(1°)  $G_A$  and  $C_A$  of  $A \in A$  defined above are given by

$$G_{A}(\lambda) = \sum_{n} e^{i\lambda t_{n}} k_{n} = \int_{-\infty}^{\infty} e^{i\lambda t} dK(t),$$
$$C_{A}(z) = \sum_{n} e^{zt_{n}} k_{n} = \int_{-\infty}^{\delta} e^{zt} dK(t).$$

(2°) Especially, if  $x \in \mathfrak{S}_0$ , then for the corresponding usual stationary process  $x \in \mathfrak{S}^\circ$ ,  $\Lambda x(t)$  is given by

$$\Lambda x(t) \equiv \sum_{n} \tau_{t_n} x(t) k_n = \sum_{n} x(t+t_n) k_n = \int_{-\infty}^{\infty} x(t+s) dK(s) = \int_{-\infty}^{\infty} \tau_s x(t) dK(s).$$

DEFINITION 2.4. If  $x \in \mathfrak{S}$  has the orthogonal random measure and the spectral measure dZ and  $\mu$  respectively, and if S-C. L. Op.  $\Lambda \in \widetilde{\Lambda}$  has the g.f.  $G_A \in L^2(\mu; \mathscr{S})$ , then the corresponding S-C. L. Op.  $\Lambda$  in  $\mathfrak{S}$  to  $\Lambda \in \widetilde{\Lambda}$  is defined by, for every  $\varphi \in \mathscr{S}$ 

(2.6) 
$$\Lambda x(\varphi) = \int_{-\infty}^{\infty} (\mathscr{F}\varphi)(\lambda) G_{\Lambda}(\lambda) dZ(\lambda).$$

The domain of  $\Lambda$  in  $\mathfrak{S}$  is denoted by  $\mathscr{D}(\Lambda) \equiv \{x; G_{\Lambda} \in L^{2}(\mu_{x}; \mathscr{S})\}$ , The class of all operators  $\Lambda$  in  $\mathfrak{S}$  defined above and whose domain in  $\mathfrak{S}$  is not empty, is called the *shift-commutative linear operators* (S-C.L.Op.s) in  $\mathfrak{S}$  and denoted by  $\Lambda$ .

REMARK. Here we are treating the weakly stationary processes, i. e. stationary process x is regarded as a point of the closed linear sub-manifold of the Hilbert space  $L^2(\mathcal{Q})$ . Hence, in its representation (2.6), it

.26

<sup>5)</sup> The inside of "()" shows the the number of result in T. Seguchi [4].

27

will be in question that the difference of measure zero. Here we wish to point out that the above definition of  $\Lambda x$  give the answer to this question. That is, as the g.f.  $G_A$  of S-C. L. Op.  $\Lambda$  is uniquely determined by the operator  $\Lambda$ , hence the representation (2.6) is unique, and even if  $G_A \simeq G_r$  $(\mu_x)$  (in this case  $\Lambda x(\varphi) = \int_{-\infty}^{\infty} (\mathscr{F}\varphi)(\lambda) G_A(\lambda) dZ_x(\lambda) = \int_{-\infty}^{\infty} (\mathscr{F}\varphi)(\lambda) G_r(\lambda) dZ_x(\lambda)$  $= \Gamma x(\varphi)$  is considered as the same point in  $L^2(\mathcal{Q})$ ), we must consider that  $\Lambda x \neq \Gamma x$ .

PROPOSITION 2.2. (1°) If  $x \in \mathfrak{S}$  has an orthogonal random measure  $dZ_x$  and a spectral measure  $\mu_x$ , then for any  $\Lambda \in \Lambda$  satisfying  $G_{\Lambda} \in L^2(\mu_x; \mathscr{S})$ , expressions

$$\Lambda x(\varphi) = \int_{-\infty}^{\infty} (\mathscr{F}\varphi)(\lambda) G_{\lambda}(\lambda) dZ_{\lambda}(\lambda),$$

(2.7) 
$$r_{\Lambda x}(\varphi) = \int_{-\infty}^{\infty} (\mathscr{F}\varphi)(\lambda) |G_{\Lambda}(\lambda)|^{2} \mu_{x}(d\lambda),$$

hold, for every  $\varphi \in \mathscr{S}$ , where  $r_{Ax}(\varphi)$  is the covariance distribution of Ax.

(2°) Especially, if  $x \in \mathfrak{S}_0$  and  $G_A \in L^2(\mu_x; 0)$ , then for the corresponding usual stationary process  $x \in \mathfrak{S}^\circ$  to  $x \in \mathfrak{S}_0$ , the following expressions hold:

(2.8) 
$$\Lambda x(t) = \int_{-\infty}^{\infty} e^{i\lambda t} G_{\Lambda}(\lambda) dZ_{\chi}(\lambda),$$

(2.9) 
$$r_{Ax}(t) = \int_{-\infty}^{\infty} e^{i\lambda t} |G_A(\lambda)|^2 \mu_x(d\lambda),$$

where  $dZ_x$  and  $\mu_x$  are those of  $x \in \mathfrak{S}_0$ .

**PROPOSITION 2.3** (Theorem 3.4.4). Let  $\Lambda$ ,  $\Gamma$ ,  $\Delta \in \Lambda$  (or  $\widetilde{\Lambda}$ ), then we have

- (1°) if  $\Delta = a \Lambda + b \Gamma$  then  $G_{\Delta} = a G_{\Lambda} + b G_{\Gamma}$  and  $C_{\Delta} = a C_{\Lambda} + b C_{\Gamma}$ ,
- (2°) if  $\Delta = \Lambda \Gamma$  then  $G_{\Delta} = G_{\Lambda}G_{\Gamma} = G_{\Gamma}G_{\Lambda}$  and  $C_{\Delta} = C_{\Lambda}C_{\Gamma} = C_{\Gamma}C_{\Lambda}$ ,

(3°) if there exists the inverse operator  $\Lambda^{-1} \in \Lambda$  (or  $\widetilde{\Lambda}$ ), then  $G_{\Lambda^{-1}} = 1/G_{\Lambda}$  and  $C_{\Lambda^{-1}} = 1/C_{\Lambda}$ , where  $\Lambda^{-1}$  is an operator such that  $\Lambda \Lambda^{-1} = I$  (identity operator).

PROPOSITION 2.4. (1°) If  $\Lambda \in \widetilde{\Lambda}$  is given by

$$\Lambda f(\cdot) \equiv \int_a^b \tau_{-s} f(\cdot) dK(s),$$

then  $\Lambda \in \Lambda$  corresponding to the above  $\Lambda \in \widetilde{\Lambda}$  can be expressed by

$$\Lambda x(\varphi) \equiv \int_a^b T_{-s} x(\varphi) dK(s) = \int_a^b x(\tau_s \varphi) dK(s),$$

T. Seguchi

where  $K \in BV[a, b]^{(i)}(-\infty \le a \le b \le +\infty), x \in \mathcal{D}(\Lambda).$ 

(2°) Especially, when  $x \in \mathfrak{S}^{\circ}$ , then

$$\Lambda x(t) \equiv \int_a^b \tau_{-s} x(t) dK(s) = \int_a^b x(t-s) dK(s).$$

**PROPOSITION 2.5** (Theorem 4.2.3). (1°) Let x and  $y \in \mathfrak{S}$  satisfy the equation

 $\Lambda y = x$ ,

where  $\Lambda \in \Lambda_{-}$ . Then for every  $t \in R$ 

$$L^2(Y; t) \supset L^2(X; t)$$

holds.

(2°) Furthermore, if there exists the inverse operator  $\Lambda^{-1} \in \Lambda_{-}$ , then

 $L^{2}(Y; t) = L^{2}(X; t)$ 

holds, for every  $t \in \mathbf{R}$ .

PROPOSITION 2.6 (Theorem 5.2.1). Let  $x \in \mathfrak{S}_0$ ,  $\dot{\xi}$  be a differentiation in the sense of random distribution of a homogeneous orthogonal random measure  $d\xi$  and  $\Lambda \in \Lambda$ . Then

(1°) the relation

$$(2.10) \qquad \qquad \Lambda x = \dot{\xi}$$

holds if and only if the conditions (G 1)  $N_A = \phi$  and (G 4°)  $1/G_A \in L^2(m; 0)^{\gamma}$  are satisfied.

(2°) any  $x \in \mathfrak{S}_0$  satisfying (2.10) is given by uniquely

$$\mathbf{x}(\varphi) = \boldsymbol{\Lambda}^{-1} \dot{\boldsymbol{\xi}}(\varphi) = \int_{-\infty}^{\infty} (\mathscr{F}\varphi)(\lambda) d\tilde{\boldsymbol{\xi}}(\lambda) / G_{\boldsymbol{\lambda}}(\lambda),$$

where  $d\tilde{\xi}$  is the homogeneous orthogonal random measure of  $\dot{\xi} \in \mathfrak{S}_1$ .

(3°)  $x \in \mathfrak{S}^{\circ}$ , corresponding to  $x \in \mathfrak{S}_{0}$ , is represented by

$$x(t) = \int_{-\infty}^{\infty} e^{i\lambda t} d\tilde{\xi}(\lambda) / G_{\Lambda}(\lambda),$$
$$r_{x}(t) = \int_{-\infty}^{\infty} e^{i\lambda t} / |G_{\Lambda}(\lambda)|^{2} d\lambda.$$

<sup>6)</sup> BV[a, b] means "the all real- or complex-valued functions of bounded variations in [a, b]."

<sup>7)</sup> *m* means "the usual Lebesgue measure on R", i. e.  $m(d\lambda) \equiv d\lambda$ .

The above all notations, notions, definitions and propositions are in T. Seguchi [4], or can be easily proved.

# 3. S-C.L.Op.s of Riemann-Liouville type belonging to $\widetilde{\Lambda}$ .

#### [1°] S-C.L.Op. $(D+\beta I)^{-\alpha}$ .

We shall first consider the case that the operator  $\Lambda$  defined by the following form: For  $\alpha > 0$ ,  $\beta > 0$  ( $\alpha$ ,  $\beta \in \mathbb{R}$ )

(3.1) 
$$\Lambda f(t) \equiv 1/\Gamma(\alpha) \int_{-\infty}^{t} (t-s)^{\alpha-1} e^{-\beta(t-s)} f(s) ds$$
$$= 1/\Gamma(\alpha) \int_{0}^{\infty} u^{\alpha-1} e^{-\beta u} \tau_{-u} f(t) du.$$

The following formula holds, for any  $\alpha > 0$  ( $\alpha \in \mathbb{R}$ ), and  $z \in C$  such as  $\Re e z > 0$ 

(3.2) 
$$\int_0^\infty e^{-zt} t^{\alpha-1} dt = \Gamma(\alpha)/z^{\alpha}.$$

If we put  $f(t) = e^{i\lambda t} (\lambda \in \mathbb{R})$ , we have from (3.1)

$$\Lambda e^{i\lambda t} = 1/\Gamma(\alpha) \int_{-\infty}^{t} (t-s)^{\alpha-1} e^{-\beta(t-s)} e^{i\lambda s} ds$$
$$= e^{i\lambda t}/\Gamma(\alpha) \int_{0}^{\infty} u^{\alpha-1} e^{-(\beta+i\lambda)u} du$$
$$= e^{i\lambda t}/(\beta+i\lambda).$$

Hence, from the above formula, we can see that the operator  $\Lambda$  belongs to the class  $\widetilde{\Lambda}$  of S-C.L.Op.s and the g.f. of  $\Lambda$  is  $G_{\Lambda}(\lambda) = (i\lambda + \beta)^{-\alpha}$ .

If  $\alpha=1$  then we can easily assure the operator  $\Lambda$  defined by (3.1) is the operator  $(D+\beta I)^{-1}$ , which is given by (2.2). Hence, we shall denote the operator  $\Lambda$  defined by (3.1)  $\Lambda \equiv (D+\beta I)^{-\alpha}$ .

### [2°] S-C.L.Op. $(D+\beta I)^{\alpha}$ .

Next, we shall consider the following case;  $0 < \alpha < 1$ ,  $\beta > 0$  ( $\alpha$ ,  $\beta \in R$ ). Using the formula (3.2) and

(3.3) 
$$\Gamma(1-\alpha) = -\alpha \Gamma(-\alpha), \qquad 0 < \alpha < 1,$$

we have

T. SEGUCHI

(3.4)  $\int_{0}^{\infty} u^{-\alpha-1} (1-e^{-(\beta+i\lambda)u}) du$  $= -1/\alpha [u^{-\alpha} (1-e^{-(\beta+i\lambda)u})] \int_{0}^{\infty} -[(\beta+i\lambda)/\alpha] \int_{0}^{\infty} u^{-\alpha} e^{-(\beta+i\lambda)u} du$  $= -[(\beta+i\lambda)/\alpha] \int_{0}^{\infty} u^{(1-\alpha)-1} e^{-(\beta+i\lambda)u} du$  $= -[(\beta+i\lambda)/\alpha] [\Gamma(1-\alpha)]/(\beta+i\lambda)^{1-\alpha}$  $= -[\Gamma(1-\alpha)/\alpha] (\beta+i\lambda)^{\alpha}.$ 

On the other hand, if we put u=t-s, from (3.4) the following formula holds:

(3.5) 
$$\int_{0}^{\infty} u^{-\alpha-1} (1-e^{-(\beta+i\lambda)u}) du$$
$$= \int_{-\infty}^{t} (t-s)^{-\alpha-1} (1-e^{-(\beta+i\lambda)(t-s)}) ds$$
$$= e^{-i\lambda t} \int_{-\infty}^{t} (t-s)^{-\alpha-1} (e^{i\lambda t}-e^{-\beta(t-s)}e^{i\lambda s}) ds$$

Hence, from (3.4) and (3.5), if we consider the operator  $\Lambda$  such that

(3.6) 
$$\Lambda f(t) \equiv 1/\Gamma(-\alpha) \int_{-\infty}^{t} (t-s)^{-\alpha-1} [f(t) - e^{-\beta(t-s)}f(s)] ds$$
$$= 1/\Gamma(-\alpha) \int_{0}^{\infty} u^{-\alpha-1} (I - e^{-\beta u} \tau_{-u}) f(t) du,$$

then, putting  $f(t) \equiv e^{i\lambda t}$ , we get

(3.7) 
$$\Lambda e^{i\lambda t} = 1/\Gamma(-\alpha) \int_{-\infty}^{t} (t-s)^{-\alpha-1} (e^{i\lambda t} - e^{-\beta(t-s)} e^{i\lambda s}) ds$$
$$= e^{i\lambda t}/\Gamma(-\alpha) \int_{-\infty}^{t} (t-s)^{-\alpha-1} (1 - e^{-(\beta+i\lambda)(t-s)}) ds$$
$$= e^{i\lambda t} (\beta+i\lambda)^{\alpha}.$$

The formula (3.7) shows that the operator  $\Lambda$  defined by (3.6) is belonging to the class  $\widetilde{\Lambda}$  and its g.f. is  $(i\lambda + \beta)^{\alpha}$ . In accordance with [1°], we shall denote the above operator  $\Lambda \equiv (D + \beta I)^{\alpha}$ .

[3°] S-C.L.Op.  $D^{\alpha}$ .

If f(t) be the function such that the right hand side of formula (3.6) is meaningful, and let  $0 < \alpha < 1$ , then we have

30

Certain operators of Riemann-Liouville type and stationary processes 31

(3.8) 
$$\lim_{d \to 0} 1/\Gamma(-\alpha) \int_{-\infty}^{t} (t-s)^{-\alpha-1} [f(t) - e^{-\theta(t-s)} f(s)] ds$$
$$= 1/\Gamma(-\alpha) \int_{-\infty}^{t} (t-s)^{-\alpha-1} [f(t) - f(s)] ds$$
$$= 1/\Gamma(-\alpha) \int_{0}^{\infty} u^{-\alpha-1} (I - \tau_{-u}) f(t) du.$$

Let the operator, which is defined by the right hand side of (3.8), be  $D^{\alpha}$ , then from (3.7) we can easily see that

$$D^{\alpha}e^{i\lambda t}=e^{i\lambda t}(i\lambda)^{\alpha}.$$

Hence we see that  $D^{\alpha} \in \widetilde{A}$ ,  $G_{D^{\alpha}}(\lambda) = (i\lambda)^{\alpha}$ ,  $C_{D^{\alpha}}(z) = z^{\alpha}$ . We shall call the operator  $D^{\alpha}$  the differential operator of order  $\alpha$ .

[4°] For general  $\alpha \in R$ , there exists an integer *n* such that  $\alpha \equiv n + \alpha'$ ( $0 < \alpha' < 1$ ). Then above three operators are given by the followings:

- (1)  $(D+\beta I)^{-\alpha}f(t) \equiv (D+\beta I)^{-n}(D+\beta I)^{-\alpha'}f(t),$
- (2)  $(D+\beta I)^{\alpha}f(t) \equiv (D+\beta I)^{n}(D+\beta I)^{\alpha'}f(t),$

(3) 
$$D^{\alpha}f(t) \equiv D^n D^{\alpha'}f(t),$$

where if n > 0,  $(D+\beta I)^{-n}$ ,  $(D+\beta I)^n$  and  $D^n$  are given by the definition of differential operator D and the integral operators (2.2) and (2.3) and Proposition 2.3, and if n > 0, the operator  $D^{-n}$  is defined as the following; let F(t) be a continuous primitive of f(t) at t, satisfying F(0) = 0 and let  $D^{-1}f(t) \equiv F(t)$ , then from Proposition 2.3 we can see that  $D^{-n} \equiv D^{-1}D^{-1}\cdots D^{-1}$ .

[5°] From Proposition 2.3 and [1°]—[4°] in this section, we can easily see that for any  $\alpha \in \mathbb{R}$ 

(1) 
$$[(D+\beta I)^{\alpha}]^{-1}=(D+\beta I)^{-\alpha},$$

- (2)  $[(D+\beta I)^{-\alpha}]^{-1}=(D+\beta I)^{\alpha},$
- (3)  $(D^{\alpha})^{-1} = D^{-\alpha}.$

#### 4. S-C.L.Op.s of Riemann-Liouville type belonging to $\Lambda$ .

From the preceding Section 3, Definition 2.4 and Propositions 2.2 and 2.4, we can immediately obtain the followings:

[1°] For any 
$$x \in \mathfrak{S}$$
 such as  $(i\lambda + \beta)^{-\alpha} \in L^2(\mu_x; \mathscr{S}), \alpha > 0, \beta > 0 \ (\alpha, \beta \in \mathbb{R})$ 

T. SEGUCHI

(4.1) 
$$(D+\beta I)^{-\alpha} x(\varphi) \equiv \int_{-\infty}^{\infty} (\mathscr{F}\varphi)(\lambda) (i\lambda+\beta)^{-\alpha} dZ_x(\lambda)$$
$$= 1/\Gamma(\alpha) \int_{0}^{\infty} u^{\alpha-1} e^{-\beta u} T_{-u} x(\varphi) du$$
$$= 1/\Gamma(\alpha) \int_{0}^{\infty} u^{\alpha-1} e^{-\beta u} x(\tau_u \varphi) du.$$

[2°] For any  $x \in \mathfrak{S}$  such as  $(i\lambda + \beta)^{\alpha} \in L^{2}(\mu_{x}; \mathscr{S}), 0 < \alpha < 1, \beta > 0 \ (\alpha, \beta \in R)$ 

(4.2) 
$$(D+\beta I)^{\alpha} x(\varphi) = \int_{-\infty}^{\infty} (\mathscr{F}\varphi)(\lambda) (i\lambda+\beta)^{\alpha} dZ_{x}(\lambda)$$
$$= 1/\Gamma(-\alpha) \int_{0}^{\infty} u^{-\alpha-1} (I-e^{-\beta u}T_{-u}) x(\varphi) du$$
$$= 1/\Gamma(-\alpha) \int_{0}^{\infty} u^{-\alpha-1} [x(\varphi)-e^{-\beta u} x(\tau_{u}\varphi)] du$$

[3°] For any  $x \in \mathfrak{S}$  such as  $(i\lambda)^{\alpha} \in L^{2}(\mu_{x}; \mathscr{S}), 0 < \alpha < 1 \ (\alpha \in \mathbb{R})$ 

(4.3) 
$$D^{\alpha}x(\varphi) \equiv \int_{-\infty}^{\infty} (\mathscr{F}\varphi)(\lambda)(i\lambda)^{\alpha}dZ_{x}(\lambda)$$
$$= 1/\Gamma(-\alpha)\int_{0}^{\infty} u^{-\alpha-1}(I-T_{-u})x(\varphi)du$$
$$= 1/\Gamma(-\alpha)\int_{0}^{\infty} u^{-\alpha-1}[x(\varphi)-x(\tau_{u}\varphi)]du.$$

[4°] The results of preceding Section 3, [4°] and [5°] are all obtained for the S-C.L.Op.s in  $\mathfrak{S}$  except the case  $D^{-\alpha}$  ( $\alpha > 0$ ). And for  $x \in \mathfrak{S}$ , from Proposition 2.2 the results are trivial, furthermore, for the covariance functions or covariance distributions the results are trivial from Proposition 2.2 and Definition 2.4, hence we shall omit to show the results. As for the operator  $D^{\alpha}$  ( $\alpha > 0$ ), from Proposition 2.3, the following relation is not satisfied;

$$(D^{\alpha})^{-1}=D^{-\alpha},$$

(This relation does not hold in the case of  $D^{\alpha} \in A$ , but in the Section 3, we defined formally that  $D^{-1}f(t) \equiv F(t)$ . In this case, from Proposition 2.3 the definition of  $(D^{\alpha})^{-1}$  can not possible, as  $G_{D^{\alpha}}(\lambda)$  has a zero on pure imaginary axis, i.e.  $G_{D^{\alpha}}(0)=0$  hence the relation  $(D^{\alpha})^{-1}=D^{-\alpha}$  does not hold from Proposition 2.3.)

32

Certain operators of Riemann-Liouville type and stationary processes

# 5. Applications to the functional equations.

Let us consider the following stochastic functional equations:

(5.1) 
$$(D+\beta I)^{-\alpha}y=x, \quad \alpha>0, \ \beta>0 \ (\alpha, \ \beta\in R),$$

$$(5.2) (D+\beta I)^{\alpha}y = x, 0 < \alpha < 1, \ \beta > 0 \ (\alpha, \ \beta \in R),$$

$$(5.3) D^{\alpha}y = x, 0 < \alpha < 1, \ (\alpha \in \mathbf{R}),$$

where x and  $y \in \mathfrak{S}$ .

The above equations are meanigful if and only if the following relations

$$(5.1)' \qquad (i\lambda+\beta)^{-\alpha}dZ_{\nu}(\lambda) = dZ_{\kappa}(\lambda).$$

$$(5.2)' \qquad (i\lambda + \beta)^{\alpha} dZ_{y}(\lambda) = dZ_{x}(\lambda),$$

$$(5.3)' \qquad (i\lambda)^{\alpha} dZ_{y} = dZ_{x}(\lambda)$$

hold for any  $\lambda \in R$ . Then from the preceding Section 4, [4°], we can see for the equations (5.1) and (5.2) the following relations hold:

(5.1)" 
$$y = ((D+\beta I)^{-\alpha})^{-1}x = (D+\beta I)^{\alpha}x,$$

(5.2)'' 
$$y = ((D+\beta I)^{\alpha})^{-1}x = (D+\beta I)^{-\alpha}x.$$

Furthermore, from the definitions of above operators  $(D+\beta I)^{\alpha}$  and  $(D+\beta I)^{\alpha}$  $\beta I$ )<sup>- $\alpha$ </sup> (Section 4, [1°] and [2°]) and Proposition 2.5, we can obtain the following result:

(5.4) 
$$(D+\beta I)^{-\alpha} = ((D+\beta I)^{\alpha})^{-1} \in \Lambda_{-},$$

(5.5) 
$$(D+\beta I)^{\alpha} = ((D+\beta I)^{-\alpha})^{-1} \in A_{-}$$

$$(5.6) L2(Y; t) = L2(X; t), for every t \in R.$$

The other results are obtained from Section 4 in T. Seguchi [4].

# 6. Applications to the representations of certain stationary processes.

Now let us consider the following stochastic functional equations

(6.1) 
$$(D+\beta I)^{-\alpha}x=\xi, \quad \alpha>0, \ \beta>0 \ (\alpha, \ \beta\in R),$$

$$(6.2) (D+\beta I)^{\alpha} x = \xi, 0 < \alpha < 1, \ \beta > 0 \ (\alpha, \ \beta \in R),$$

 $D^{\alpha}x = \dot{\xi}, \qquad 0 < \alpha < 1 \ (\alpha \in R),$ (6.3)

where  $x \in \mathfrak{S}_0$ ,  $\dot{\xi} \in \mathfrak{S}_1$  which is the differentiation in the sense of random distribution of the homogeneous orthogonal random measure  $d\xi$ .

33

T. Seguchi

From Proposition 2.6 the equations (6.1), (6.2) and (6.3) are meaningful if and only if the conditions

(6.4) 
$$\frac{1}{(i\lambda+\beta)^{\alpha'} \in L^2(m; 0), \quad N_{\lambda}=\phi,}{(\beta\geq 0, \ \alpha' \in R, \ \lambda \in R)}$$

are satisfied, where  $\Lambda$  is  $(D+\beta I)^{-\alpha}$  in (6.1),  $(D+\beta I)^{\alpha}$  in (6.2),  $D^{\alpha}$  in (6.3) and  $\alpha' = \alpha \operatorname{or} - \alpha$ ,  $\alpha > 0$ .

But the last condition of (6.4) is not satisfied in the case of  $D^{\alpha}$ , hence the equation (6.3) is not meaningful.

For (6.2), if  $1/2 < \alpha < 1$  the conditions of (6.4) are all satisfied and otherwise the first condition of (6.4) is not satisfied. Hence the equation (6.2) is meaningful if and only if the case that  $1/2 < \alpha < 1$ .

For (6.1), the first condition of (6.4) is not satisfied, hence the equation (6.1) is not meaningful.

From the above discussions, finally, among the above equations, only the equation (6.2) is meaningful in the case  $1/2 < \alpha < 1$ ,  $\beta > 0$  ( $\alpha$ ,  $\beta \in R$ ). In this case, we can obtain, from Section 3, [5°] and Propositions 2.2, 2.5, 2.6, the following results: (1) There exists unique solution of equation (6.2) and is represented by

(6.5) 
$$x(\varphi) = (D + \beta I)^{-\alpha} \dot{\xi}(\varphi)$$
$$= 1/\Gamma(-\alpha) \int_{0}^{\infty} u^{-\alpha-1} (I - e^{-\beta u} T_{-u}) \dot{\xi}(\varphi) du$$
$$= \int_{-\infty}^{\infty} (\mathscr{F}\varphi)(\lambda) (i\lambda + \beta)^{-\alpha} d\tilde{\xi}(\lambda),$$

where  $d\tilde{\xi}(\lambda)$  is the homogeneous orthogonal random measure corresponding to the white noise  $\dot{\xi}$ , and the covariance distribution of x is given by

(6.6) 
$$r_{x}(\varphi) = \int_{-\infty}^{\infty} (\mathscr{F}\varphi)(\lambda) |i\lambda + \beta|^{-2\alpha} d\lambda.$$

(2) Let  $x \in \mathfrak{S}^{\circ}$  be the stationary process corresponding to the solution  $x \in \mathfrak{S}_{0}$  of the equation (6.2), then we have, from Proposition 2.2 and Definition 2.4

(6.7) 
$$x(t) = 1/\Gamma(-\alpha) \int_0^\infty u^{-\alpha-1} (I - e^{-\beta u} \tau_{-u}) d\xi(u)$$
$$= \int_{-\infty}^\infty e^{i\lambda t} (i\lambda + \beta)^{-\alpha} d\tilde{\xi}(\lambda),$$

35

(6.8) 
$$r_{x}(t) = \int_{-\infty}^{\infty} e^{i\lambda t} |i\lambda + \beta|^{-2\alpha} d\lambda,$$

where  $d\tilde{\xi}$  is that of relation (6.5). (3) Furthermore, in this case, the Paley-Wiener's condition is satisfied in addition to the conditions (6.4), by the same way as Proposition 2.6, we can see that the canonical representation of  $x \in \mathbb{S}^{\circ}$  satisfying the equation (6.2) is the first term of the expression (6.7).

#### References.

- [1] HIDA, T., Representations of Gaussian processes and their applications (in Japanese), Seminar on Prob., Vol. 7, 1961.
- [2] YOSIDA, K., Fractional powers of infinitesimal generators and the analyticity of the semi-groups generated by them, Proc. Japan Acad., 36 (1960), 86-89.
- [3] BALAKIRISHNAN, A. V., Representation of abstract Riesz potentials of the elliptic type, Bull. Amer. Math. Soc., 64 (1958), 288-289.
- [4] SEGUCHI, T., Shift-commutative linear operators and stationary processes, Mem. Fac. Sci. Kyushu Univ., Ser. A, 23 (1969), 106-155.
- [5] , An application of shift-commutative linear operators to linear automatic control systems with randomness — A theoretical formulation—, Mem. Fac. Sci. Kyushu Univ., Ser. A, 24 (1970), 120-145.
- [6] SCHWARTZ, L., Theorie des distributions I, Hermann et Cie, Paris, 1950.