# An explicit formula for the discrete power function associated with circle patterns of Schramm type

Ando, Hisashi Graduate School of Mathematics, Kyushu University

Hay, Mike

Kajiwara, Kenji Faculty of Mathematics, Kyushu University

Masuda, Tetsu 3Department of Physics and Mathematics, Aoyama Gakuin University

https://hdl.handle.net/2324/19573

出版情報:MI Preprint Series. 2011-11, 2011-05-10. 九州大学大学院数理学研究院 バージョン: 権利関係:

## **MI Preprint Series**

Kyushu University The Global COE Program Math-for-Industry Education & Research Hub

## An explicit formula for the discrete power function associated with circle patterns of Schramm type

## Hisashi Ando, Mike Hay, Kenji Kajiwara & Tetsu Masuda

MI 2011-11

(Received May 10, 2011)

Faculty of Mathematics Kyushu University Fukuoka, JAPAN

# An Explicit Formula for the Discrete Power Function Associated with Circle Patterns of Schramm Type

Hisashi Ando<sup>1</sup>, Mike Hay<sup>2</sup>, Kenji Kajiwara<sup>2</sup> and Tetsu Masuda<sup>3</sup>,

<sup>1</sup>Graduate School of Mathematics, Kyushu University, 744 Motooka, Fukuoka 819-0395, Japan

<sup>2</sup>Institute of Mathematics for Industry, Kyushu University, 744 Motooka, Fukuoka 819-0395, Japan

<sup>3</sup>Department of Physics and Mathematics, Aoyama Gakuin University, Sagamihara, Kanagawa 229-8558, Japan

May 10, 2011

#### Abstract

We present an explicit formula for the discrete power function introduced by Bobenko, which is expressed in terms of the hypergeometric  $\tau$  functions for the sixth Painlevé equation. The original definition of the discrete power function imposes strict conditions on the domain and the value of the exponent. However, we show that one can extend the value of the exponent to arbitrary complex numbers except even integers and the domain to a discrete analogue of the Riemann surface.

## **1** Introduction

The theory of discrete analytic functions has been developed in recent years based on the theory of circle packings or circle patterns, which was initiated by Thurston's idea of using circle packings as an approximation of the Riemann mapping [17]. So far many important properties have been established for discrete analytic functions, such as the discrete maximum principle and Schwarz's lemma [5], the discrete uniformization theorem [14], and so forth. For a comprehensive introduction to the theory of discrete analytic functions, we refer to [16].

It is known that certain circle patterns with fixed regular combinatorics admit rich structure. For example, it has been pointed out that the circle patterns with square grid combinatorics introduced by Schramm [15] and the hexagonal circle patterns [4, 7, 8] are related to integrable systems. Some explicit examples of discrete analogues of analytic functions have been presented which are associated with Schramm's patterns:  $\exp(z)$ ,  $\operatorname{erf}(z)$ , Airy function [15],  $z^{\gamma}$ ,  $\log(z)$  [3]. Also, discrete analogues of  $z^{\gamma}$  and  $\log(z)$  associated with hexagonal circle patterns are discussed in [4, 7, 8].

Among those examples, it is remarkable that the discrete analogue of the power function  $z^{\gamma}$  associated with the circle patterns of Schramm type has a close relationship with the sixth Painlevé equation (P<sub>VI</sub>) [6], and this fact has been used to establish the immersion property [3] and embeddedness [1] of the discrete power function. It is desirable to construct a representation formula for the discrete power function in terms of the Painlevé transcendents as was mentioned in [6]. The discrete power function can be formulated as a solution to a system of difference equations on the square lattice  $(n, m) \in \mathbb{Z}^2$  with a certain initial condition. A correspondence between the dependent variable of this system and the Painlevé transcendents can be found in [13], but the formula seems somewhat indirect. Agafonov has constructed an explicit representation formula in terms of the Gauss hypergeometric function [2], however, this formula is valid only on some special points on  $\mathbb{Z}^2$ . In this paper, generalizing Agafonov's result, we aim to establish an explicit representation of P<sub>VI</sub> which is valid on  $\mathbb{Z}^2_+ = \{(n,m) \in \mathbb{Z}^2 \mid n, m \ge 0\}$  and for  $\gamma \in \mathbb{C} \setminus \mathbb{Z}$ . Based on this formula, we generalize the domain of the discrete power function to a discrete analogue of the Riemann surface.

This paper is organized as follows. In section 2, we give a brief review of the definition of the discrete power function and its relation to  $P_{VI}$ . The main result and its proof are given in section 3. We discuss the extension of the domain of the discrete power function in section 4. Section 5 is devoted to concluding remarks.

## **2** Discrete power function

#### 2.1 Definition of the discrete power function

For maps, a discrete analogue of conformality has been proposed by Bobenko and Pinkall in the framework of discrete differential geometry [9].

**Definition 2.1** A map  $f : \mathbb{Z}^2 \to \mathbb{C}$ ;  $(n,m) \mapsto f_{n,m}$  is called discrete conformal if the cross-ratio with respect to every elementary quadrilateral is equal to -1:

$$\frac{(f_{n,m} - f_{n+1,m})(f_{n+1,m+1} - f_{n,m+1})}{(f_{n+1,m} - f_{n+1,m+1})(f_{n,m+1} - f_{n,m})} = -1.$$
(2.1)

The condition (2.1) is a discrete analogue of the Cauchy-Riemann relation. Actually, a smooth map  $f: D \subset \mathbb{C} \to \mathbb{C}$  is conformal if and only if it satisfies

$$\lim_{\epsilon \to 0} \frac{(f(x,y) - f(x+\epsilon,y))(f(x+\epsilon,y+\epsilon) - f(x,y+\epsilon))}{(f(x+\epsilon,y) - f(x+\epsilon,y+\epsilon))(f(x,y+\epsilon) - f(x,y))} = -1$$
(2.2)

for all  $(x, y) \in D$ . However, using Definition 2.1 alone, one cannot exclude maps whose behavior is far from that of usual holomorphic maps. Because of this, an additional condition for a discrete conformal map has been considered [1, 3, 6, 10].

**Definition 2.2** A discrete conformal map  $f_{n,m}$  is called embedded if inner parts of different elementary quadrilaterals  $(f_{n,m}, f_{n+1,m}, f_{n+1,m+1}, f_{n,m+1})$  do not intersect. An example of an embedded map is presented in Figure 1. This condition seems to require that  $f = f_{n,m}$  is a univalent function in the continuous limit, and is too strict to capture a wide class of discrete holomorphic functions. In fact, a relaxed requirement has been considered as follows [1,3].

**Definition 2.3** A discrete conformal map  $f_{n,m}$  is called immersed, or an immersion, if inner parts of adjacent elementary quadrilaterals  $(f_{n,m}, f_{n+1,m}, f_{n+1,m+1}, f_{n,m+1})$  are disjoint.

See Figure 2 for an example of an immersed map.

Let us give the definition of the discrete power function proposed by Bobenko [3, 6, 10].

**Definition 2.4** Let  $f : \mathbb{Z}^2_+ \to \mathbb{C}$ ;  $(n, m) \mapsto f_{n,m}$  be a discrete conformal map. If  $f_{n,m}$  is the solution to the difference equation

$$\gamma f_{n,m} = 2n \frac{(f_{n+1,m} - f_{n,m})(f_{n,m} - f_{n-1,m})}{f_{n+1,m} - f_{n-1,m}} + 2m \frac{(f_{n,m+1} - f_{n,m})(f_{n,m} - f_{n,m-1})}{f_{n,m+1} - f_{n,m-1}}$$
(2.3)

with the initial conditions

$$f_{0,0} = 0, \quad f_{1,0} = 1, \quad f_{0,1} = e^{\gamma \pi i/2}$$
 (2.4)

for  $0 < \gamma < 2$ , then we call f a discrete power function.

The difference equation (2.3) is a discrete analogue of the differential equation  $\gamma f = z \frac{\partial f}{\partial z}$  for the power function  $f(z) = z^{\gamma}$ , which means that the parameter  $\gamma$  corresponds to the exponent of the discrete power function.

It is easy to get the explicit formula of the discrete power function for m = 0 (or n = 0). When m = 0, (2.3) is reduced to a three-term recurrence relation. Solving it with the initial condition  $f_{0,0} = 0$ ,  $f_{1,0} = 1$ , we have

$$f_{n,0} = \begin{cases} \frac{2l}{2l+\gamma} \prod_{k=1}^{l} \frac{2k+\gamma}{2k-\gamma} & (n=2l), \\ \prod_{k=1}^{l} \frac{2k+\gamma}{2k-\gamma} & (n=2l+1), \end{cases}$$
(2.5)

for  $n \in \mathbb{Z}_+$ . When m = 1 (or n = 1), Agafonov has shown that the discrete power function can be expressed in terms of the hypergeometric function [2]. One of the aims of this paper is to give an explicit formula for the discrete power function  $f_{n,m}$  for arbitrary  $(n,m) \in \mathbb{Z}_+^2$ .

In Definition 2.4, the domain of the discrete power function is restricted to the "first quadrant"  $\mathbb{Z}^2_+$ , and the exponent  $\gamma$  to the interval  $0 < \gamma < 2$ . Under this condition, it has been shown that the discrete power function is embedded [1]. For our purpose, we do not have to persist with such a restriction. In fact, the explicit formula we will give is applicable to the case  $\gamma \in \mathbb{C} \setminus 2\mathbb{Z}$ . Regarding the domain, one can extend it to a discrete analogue of the Riemann surface.



Figure 1: An example of the embedded discrete conformal map.



Figure 2: An example of the discrete conformal map that is not embedded but immersed.

### **2.2** Relationship to $P_{\rm VI}$

In order to construct an explicit formula for the discrete power function  $f_{n,m}$ , we will move to a more general setting. The cross-ratio condition (2.1) can be regarded as a special case of the discrete Schwarzian KdV equation

$$\frac{(f_{n,m} - f_{n+1,m})(f_{n+1,m+1} - f_{n,m+1})}{(f_{n+1,m} - f_{n+1,m+1})(f_{n,m+1} - f_{n,m})} = \frac{p_n}{q_m},$$
(2.6)

where  $p_n$  and  $q_m$  are arbitrary functions in the indicated variables. Some of the authors have constructed various special solutions to the above equation [11]. In particular, they have shown that an autonomous case

$$\frac{(f_{n,m} - f_{n+1,m})(f_{n+1,m+1} - f_{n,m+1})}{(f_{n+1,m} - f_{n+1,m+1})(f_{n,m+1} - f_{n,m})} = \frac{1}{t},$$
(2.7)

where *t* is independent of *n* and *m*, can be regarded as a part of the Bäcklund transformations of  $P_{VI}$ , and given special solutions to (2.7) in terms of the  $\tau$  functions of  $P_{VI}$ .

We here give a brief account of the derivation of  $P_{VI}$  according to [13]. The derivation is achieved by imposing a certain similarity condition on the discrete Schwarzian KdV equation (2.7) and the difference equation (2.3) simultaneously. The discrete Schwarzian KdV equation (2.7) is automatically satisfied if there exists a function  $v_{n,m}$  satisfying

$$f_{n,m} - f_{n+1,m} = t^{-1/2} v_{n,m} v_{n+1,m}, \quad f_{n,m} - f_{n,m+1} = v_{n,m} v_{n,m+1}.$$
(2.8)

By eliminating the variable  $f_{n,m}$ , we get for  $v_{n,m}$  the following equation

$$t^{1/2}v_{n,m}v_{n,m+1} + v_{n,m+1}v_{n+1,m+1} = v_{n,m}v_{n+1,m} + t^{1/2}v_{n+1,m}v_{n+1,m+1},$$
(2.9)

which is equivalent to the lattice modified KdV equation. It can be shown that the difference equation (2.3) is reduced to

$$n\frac{v_{n+1,m} - v_{n-1,m}}{v_{n+1,m} + v_{n-1,m}} + m\frac{v_{n,m+1} - v_{n,m-1}}{v_{n,m+1} + v_{n,m-1}} = \mu - (-1)^{m+n}\lambda$$
(2.10)

with  $\gamma = 1 + 2\mu$ , where  $\lambda \in \mathbb{C}$  is an integration constant. In the following we take  $\lambda = \mu$  so that (2.10) is consistent when n = m = 0 and  $v_{1,0} + v_{-1,0} \neq 0 \neq v_{0,1} + v_{0,-1}$ .

Assume that the dependence of the variable  $v_{n,m} = v_{n,m}(t)$  on the deformation parameter *t* is given by

$$-2t\frac{d}{dt}\log v_{n,m} = n\frac{v_{n+1,m} - v_{n-1,m}}{v_{n+1,m} + v_{n-1,m}} + \chi_{n+m},$$
(2.11)

where  $\chi_{n+m} = \chi_{n+m}(t)$  is an arbitrary function satisfying  $\chi_{n+m+2} = \chi_{n+m}$ . Then we have the following Proposition.

**Proposition 2.5** Let  $q = q_{n,m} = q_{n,m}(t)$  be the function defined by  $q_{n,m} = t^{1/2} \frac{v_{n+1,m}}{v_{n,m+1}}$ . Then q satisfies  $P_{\text{VI}}$ 

$$\frac{d^2q}{dt^2} = \frac{1}{2} \left( \frac{1}{q} + \frac{1}{q-1} + \frac{1}{q-t} \right) \left( \frac{dq}{dt} \right)^2 - \left( \frac{1}{t} + \frac{1}{t-1} + \frac{1}{q-t} \right) \frac{dq}{dt} + \frac{q(q-1)(q-t)}{2t^2(t-1)^2} \left[ \kappa_{\infty}^2 - \kappa_0^2 \frac{t}{q^2} + \kappa_1^2 \frac{t-1}{(q-1)^2} + (1-\theta^2) \frac{t(t-1)}{(q-t)^2} \right],$$
(2.12)

with

$$\kappa_{\infty}^{2} = \frac{1}{4}(\mu - \nu + m - n)^{2}, \qquad \kappa_{0}^{2} = \frac{1}{4}(\mu - \nu - m + n)^{2},$$
  

$$\kappa_{1}^{2} = \frac{1}{4}(\mu + \nu - m - n - 1)^{2}, \quad \theta^{2} = \frac{1}{4}(\mu + \nu + m + n + 1)^{2},$$
(2.13)

where we denote  $v = (-1)^{m+n} \mu$ .

In general,  $P_{VI}$  contains four complex parameters denoted by  $\kappa_{\infty}$ ,  $\kappa_0$ ,  $\kappa_1$  and  $\theta$ . Since  $n, m \in \mathbb{Z}_+$ , a special case of  $P_{VI}$  appears in the above proposition, which corresponds to the case where  $P_{VI}$  admits special solutions expressible in terms of the hypergeometric function. In fact, the special solutions to  $P_{VI}$  of hypergeometric type are given as follows:

**Proposition 2.6** [12] Define the function  $\tau_{n'}(a, b, c; t)$  ( $c \notin \mathbb{Z}, n' \in \mathbb{Z}_+$ ) by

$$\tau_{n'}(a,b,c;t) = \begin{cases} \det\left(\varphi(a+i-1,b+j-1,c;t)\right)_{1 \le i,j \le n'} & (n' > 0), \\ 1 & (n' = 0), \end{cases}$$
(2.14)

with

$$\varphi(a, b, c; t) = c_0 \frac{\Gamma(a)\Gamma(b)}{\Gamma(c)} F(a, b, c; t) + c_1 \frac{\Gamma(a - c + 1)\Gamma(b - c + 1)}{\Gamma(2 - c)} t^{1 - c} F(a - c + 1, b - c + 1, 2 - c; t).$$
(2.15)

Here, F(a, b, c; t) is the Gauss hypergeometric function,  $\Gamma(x)$  is the Gamma function, and  $c_0$  and  $c_1$  are arbitrary constants. Then

$$q = \frac{\tau_{n'}^{0,-1,0}\tau_{n'+1}^{-1,-1,-1}}{\tau_{n'}^{-1,-1,-1}\tau_{n'+1}^{0,-1,0}}$$
(2.16)

with  $\tau_{n'}^{k,l,m} = \tau_{n'}(a+k+1,b+l+2,c+m+1;t)$  gives a family of hypergeometric solutions to  $P_{\text{VI}}$  with the parameters

$$\kappa_{\infty} = a + n', \quad \kappa_0 = b - c + 1 + n', \quad \kappa_1 = c - a, \quad \theta = -b.$$
(2.17)

We call  $\tau_{n'}(a, b, c; t)$  or  $\tau_{n'}^{k,l,m}$  the hypergeometric  $\tau$  function of P<sub>VI</sub>.

## 3 Main Results

#### **3.1** Explicit formulae for $f_{n,m}$ and $v_{n,m}$

We present the solution to the simultaneous system of the discrete Schwarzian KdV equation (2.7) and the difference equation (2.3) under the initial conditions

$$f_{0,0} = 0, \quad f_{1,0} = c_0, \quad f_{0,1} = c_1 t^r,$$
 (3.1)

where  $\gamma = 2r$ , and  $c_0$  and  $c_1$  are arbitrary constants. We set  $c_0 = c_1 = 1$  and  $t = e^{\pi i}(=-1)$  to obtain the explicit formula for the original discrete power function. Note that  $\tau_{n'}(b, a, c; t) = \tau_{n'}(a, b, c; t)$ by the definition. Moreover, we interpret F(k, b, c; t) for  $k \in \mathbb{Z}_{>0}$  as F(k, b, c; t) = 0 and  $\Gamma(-k)$  for  $k \in \mathbb{Z}_{\geq 0}$  as  $\Gamma(-k) = \frac{(-1)^k}{k!}$ .

**Theorem 3.1** For  $(n,m) \in \mathbb{Z}^2_+$ , the function  $f_{n,m} = f_{n,m}(t)$  can be expressed as follows.

(1) Case where  $n \le m$  (or n' = n). When n + m is even, we have

$$f_{n,m} = c_1 t^{r-n} N \frac{(r+1)_{N-1}}{(-r+1)_N} \frac{\tau_n(-N, -r-N+1, -r; t)}{\tau_n(-N+1, -r-N+2, -r+2; t)},$$
(3.2)

where  $N = \frac{n+m}{2}$  and  $(u)_j = u(u+1)\cdots(u+j-1)$  is the Pochhammer symbol. When n+m is odd, we have

$$f_{n,m} = c_1 t^{r-n} \frac{(r+1)_{N-1}}{(-r+1)_{N-1}} \frac{\tau_n(-N+1,-r-N+1,-r;t)}{\tau_n(-N+2,-r-N+2,-r+2;t)},$$
(3.3)

where  $N = \frac{n+m+1}{2}$ .

(2) Case where  $n \ge m$  (or n' = m). When n + m is even, we have

$$f_{n,m} = c_0 N \frac{(r+1)_{N-1}}{(-r+1)_N} \frac{\tau_m(-N+2, -r-N+1, -r+2; t)}{\tau_m(-N+1, -r-N+2, -r+2; t)},$$
(3.4)

where  $N = \frac{n+m}{2}$ . When n + m is odd, we have

$$f_{n,m} = c_0 \frac{(r+1)_{N-1}}{(-r+1)_{N-1}} \frac{\tau_m(-N+2, -r-N+1, -r+1; t)}{\tau_m(-N+1, -r-N+2, -r+1; t)},$$
(3.5)

where  $N = \frac{n+m+1}{2}$ .

**Proposition 3.2** For  $(n,m) \in \mathbb{Z}^2_+$ , the function  $v_{n,m} = v_{n,m}(t)$  can be expressed as follows.

(1) Case where  $n \le m$  (or n' = n). When n + m is even, we have

$$v_{n,m} = t^{-\frac{n}{2}} \frac{(r)_N}{(-r+1)_N} \frac{\tau_n(-N+1,-r-N+1,-r+1;t)}{\tau_n(-N+1,-r-N+2,-r+2;t)},$$
(3.6)



Figure 3: The discrete power function Figure 4: The ordinary power function with  $\gamma = 1 + i$ .

where  $N = \frac{n+m}{2}$ . When n + m is odd, we have

$$v_{n,m} = -c_1 t^{r-\frac{n}{2}} \frac{\tau_n(-N+1, -r-N+2, -r+1; t)}{\tau_n(-N+2, -r-N+2, -r+2; t)},$$
(3.7)

where  $N = \frac{n+m+1}{2}$ .

(2) Case where  $n \ge m$  (or n' = m). When n + m is even, we have

$$v_{n,m} = t^{-\frac{m}{2}} \frac{(r)_N}{(-r+1)_N} \frac{\tau_m(-N+1, -r-N+1, -r+1; t)}{\tau_m(-N+1, -r-N+2, -r+2; t)},$$
(3.8)

where  $N = \frac{n+m}{2}$ . When n + m is odd, we have

$$v_{n,m} = -c_0 t^{\frac{m+1}{2}} \frac{\tau_m(-N+2, -r-N+2, -r+2; t)}{\tau_m(-N+1, -r-N+2, -r+1; t)},$$
(3.9)

where 
$$N = \frac{n+m+1}{2}$$
.

Note that these expressions are applicable to the case where  $r \in \mathbb{C}\setminus\mathbb{Z}$ . A typical example of the discrete power function and its continuous counterpart are illustrated in Figure 3 and Figure 4, respectively. Figure 5 shows an example of the case suggesting multivalency of the map. The proof of the above theorem and proposition is given in the next subsection.

#### Remark 3.3

- (1) When m = 1 (or n = 1), the above results correspond to the case where  $P_{VI}$  is reduced to a Riccati equation and solved by the hypergeometric function. This case recovers the result obtained by Agafonov [2].
- (2) Agafonov also has shown that the generalized discrete power function  $f_{n,m}$ , under the setting of  $c_0 = c_1 = 1$ ,  $t = e^{2i\alpha}$  ( $0 < \alpha < \pi$ ) and 0 < r < 1, is embedded [2].



Figure 5: The discrete power function with  $\gamma = 0.25 + 3.35i$ .

**Remark 3.4** As we mention above, some special solutions to (2.7) in terms of the  $\tau$  functions of  $P_{\text{VI}}$  have been presented [11]. It is easy to show that these solutions also satisfy a difference equation which is a deformation of (2.3) in the sense that the coefficients n and m of (2.3) are replaced by arbitrary complex numbers. For instance, a class of solutions presented in Theorem 6 of [11] satisfies

$$(\alpha_0 + \alpha_2 + \alpha_4)f_{n,m} = (n - \alpha_2)\frac{(f_{n+1,m} - f_{n,m})(f_{n,m} - f_{n-1,m})}{f_{n+1,m} - f_{n-1,m}} - (\alpha_1 + \alpha_2 + \alpha_4 - m)\frac{(f_{n,m+1} - f_{n,m})(f_{n,m} - f_{n,m-1})}{f_{n,m+1} - f_{n,m-1}},$$
(3.10)

where  $\alpha_i$  are parameters of  $P_{VI}$  introduced in Appendix A. Setting the parameters as  $(\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4) = (r, 0, 0, -r+1, 0)$ , we see that the above equation is reduced to (2.3) and that the solutions are given by the hypergeometric  $\tau$  functions under the initial conditions (3.1).

#### **3.2 Proof of main results**

In this subsection, we give the proof of Theorem 3.1 and Proposition 3.2. One can easily verify that  $f_{n,m}$  satisfies the initial condition (3.1) by noticing  $\tau_0(a, b, c; t) = 1$ . We then show that  $f_{n,m}$  and  $v_{n,m}$  given in Theorem 3.1 and Proposition 3.2 satisfy the relation (2.8), the difference equation (2.3), the compatibility condition (2.9) and the similarity condition (2.11) by means of the various bilinear relations for the hypergeometric  $\tau$  function. Note in advance that we use the bilinear relations by specializing the parameters a, b and c as

$$a = -N, \quad b = -r - N, \quad c = -r + 1, \quad N = \frac{n+m}{2},$$
 (3.11)

when n + m is even, or

$$a = -r - N + 1, \quad b = -N, \quad c = -r + 1, \quad N = \frac{n + m + 1}{2},$$
 (3.12)

when n + m is odd.

We first verify the relation (2.8). Note that we have the following bilinear relations

$$(c-1)\tau_n^{0,-1,-1}\tau_{n+1}^{-1,-1,-1} = (c-b-1)t\tau_{n+1}^{0,-1,0}\tau_n^{-1,-1,-2} + b\tau_n^{0,0,0}\tau_{n+1}^{-1,-2,-2},$$
  

$$(c-1)\tau_n^{-1,-1,-1}\tau_n^{0,-1,-1} = (c-b-1)\tau_n^{0,-1,0}\tau_n^{-1,-1,-2} + b\tau_n^{0,0,0}\tau_n^{-1,-2,-2},$$
(3.13)

$$(a-b)\tau_m^{0,-1,-1}\tau_m^{0,-1,0} = a\tau_m^{-1,-1,-1}\tau_m^{1,-1,0} - b\tau_m^{0,0,0}\tau_m^{0,-2,-1},$$
  

$$(a-b)t\tau_{m+1}^{0,-1,0}\tau_m^{0,-1,-1} = a\tau_{m+1}^{-1,-1,-1}\tau_m^{1,-1,0} - b\tau_m^{0,0,0}\tau_{m+1}^{0,-2,-1},$$
(3.14)

$$(b-a+1)\tau_m^{0,0,0}\tau_m^{-1,-1,-1} = (b-c+1)\tau_m^{0,-1,0}\tau_m^{-1,0,-1} + (c-a)\tau_m^{0,-1,-1}\tau_m^{-1,0,0},$$
  

$$(b-a+1)\tau_{m+1}^{-1,-1,-1}\tau_m^{0,0,0} = (b-c+1)\tau_{m+1}^{0,-1,0}\tau_m^{-1,0,-1} + (c-a)\tau_m^{0,-1,-1}\tau_{m+1}^{-1,0,0},$$
(3.15)

for the hypergeometric  $\tau$  functions, the derivation of which is discussed in Appendix A. Let us consider the case where n' = n. When n + m is even, the relation (2.8) is reduced to

$$-r\tau_n^{[1,1,1]}\tau_{n+1}^{[0,1,1]} = Nt\tau_{n+1}^{[1,1,2]}\tau_n^{[0,1,0]} - (r+N)\tau_n^{[1,2,2]}\tau_{n+1}^{[0,0,0]},$$
  
$$-r\tau_n^{[0,1,1]}\tau_n^{[1,1,1]} = N\tau_n^{[1,1,2]}\tau_n^{[0,1,0]} - (r+N)\tau_n^{[1,2,2]}\tau_n^{[0,0,0]},$$
  
(3.16)

where we denote

$$\tau_{n'}^{[i_1,i_2,i_3]} = \tau_{n'}(-N+i_1,-r-N+i_2,-r+i_3;t),$$
(3.17)

for simplicity. We see that the relations (3.16) can be obtained from (3.13) with the parameters specialized as (3.11). In fact, the hypergeometric  $\tau$  functions can be rewritten as

$$\tau_n^{0,-1,-1} = \tau_n(a+1,b+1,c) = \tau_n(-N+1,-r-N+1,-r+1) = \tau_n^{[1,1,1]},$$
(3.18)

for instance. When n + m is odd, (2.8) yields

$$-r\tau_n^{[1,2,1]}\tau_{n+1}^{[1,1,1]} = (-r+N)t\tau_{n+1}^{[1,2,2]}\tau_n^{[1,1,0]} - N\tau_n^{[2,2,2]}\tau_{n+1}^{[0,1,0]},$$
  
$$-r\tau_n^{[1,1,1]}\tau_n^{[1,2,1]} = (-r+N)\tau_n^{[1,2,2]}\tau_n^{[1,1,0]} - N\tau_n^{[2,2,2]}\tau_n^{[0,1,0]},$$
  
(3.19)

which is also obtained from (3.13) by specializing the parameters as (3.12). Note that the hypergeometric  $\tau$  functions can be rewritten as

$$\tau_n^{0,-1,-1} = \tau_n(a+1,b+1,c) = \tau_n(-r-N+2,-N+1,-r+1) = \tau_n(-N+1,-r-N+2,-r+1) = \tau_n^{[1,2,1]},$$
(3.20)

this time. In the case where n' = m, one can similarly verify the relation (2.8) by using the bilinear relations (3.14) and (3.15).

Next, we prove that (2.3) is satisfied, which is rewritten by using (2.8) as

$$-r\frac{f_{n,m}}{v_{n,m}} = \frac{nt^{-\frac{1}{2}}}{v_{n+1,m}^{-1} + v_{n-1,m}^{-1}} + \frac{m}{v_{n,m+1}^{-1} + v_{n,m-1}^{-1}}.$$
(3.21)

We use the bilinear relations

$$n'\tau_{n'}^{0,0,0}\tau_{n'}^{0,-1,-1} = (b-c+1)\tau_{n'+1}^{0,-1,0}\tau_{n'-1}^{0,0,-1} + at^{-1}\tau_{n'+1}^{-1,-1,-1}\tau_{n'-1}^{1,0,0},$$

$$(a+b-c+n'+1)\tau_{n'}^{0,0,0}\tau_{n'}^{0,-1,-1} = a\tau_{n'}^{-1,-1,-1}\tau_{n'}^{1,0,0} + (b-c+1)\tau_{n'}^{0,-1,0}\tau_{n'}^{0,0,-1},$$
(3.22)

and

$$\begin{aligned} \tau_n^{0,0,0} \tau_n^{-1,-1,-2} &= -t^{-1} \tau_{n+1}^{-1,-1,-1} \tau_{n-1}^{0,0,-1} + \tau_n^{-1,-1,-1} \tau_n^{0,0,-1}, \\ \tau_m^{0,0,0} \tau_m^{1,-1,0} &= \tau_m^{0,-1,0} \tau_m^{1,0,0} - \tau_{m+1}^{0,-1,0} \tau_{m-1}^{1,0,0}, \\ \tau_m^{0,-1,-1} \tau_m^{-1,0,-1} &= -\tau_{m+1}^{-1,-1,-1} \tau_{m-1}^{0,0,-1} + \tau_m^{-1,-1,-1} \tau_m^{0,0,-1}, \end{aligned}$$
(3.23)

for the proof. Their derivation is also shown in Appendix A. Let us consider the case where n' = n. When n + m is even, we have

$$-n\tau_n^{[1,2,2]}\tau_n^{[1,1,1]} = N\tau_{n+1}^{[1,1,2]}\tau_{n-1}^{[1,2,1]} + Nt^{-1}\tau_{n+1}^{[0,1,1]}\tau_{n-1}^{[2,2,2]},$$
  

$$m\tau_n^{[1,2,2]}\tau_n^{[1,1,1]} = N\tau_n^{[0,1,1]}\tau_n^{[2,2,2]} + N\tau_n^{[1,1,2]}\tau_n^{[1,2,1]},$$
(3.24)

from the bilinear relations (3.22) by specializing the parameters a, b and c as given in (3.11). These lead us to

$$v_{n+1,m}^{-1} + v_{n-1,m}^{-1} = c_1^{-1} t^{-r+\frac{n+1}{2}} \frac{n}{N} \frac{\tau_n^{[1,2,2]} \tau_n^{[1,1,1]}}{\tau_{n+1}^{[0,1,1]} \tau_{n-1}^{[1,2,1]}},$$

$$v_{n,m+1}^{-1} + v_{n,m-1}^{-1} = -c_1^{-1} t^{-r+\frac{n}{2}} \frac{m}{N} \frac{\tau_n^{[1,2,2]} \tau_n^{[1,1,1]}}{\tau_n^{[0,1,1]} \tau_n^{[1,2,1]}}.$$
(3.25)

By using

$$\tau_n^{[1,2,2]}\tau_n^{[0,1,0]} = -t^{-1}\tau_{n+1}^{[0,1,1]}\tau_{n-1}^{[1,2,1]} + \tau_n^{[0,1,1]}\tau_n^{[1,2,1]},$$
(3.26)

which is obtained from the first relation in (3.23), one can verify (3.21). When n + m is odd, we have the bilinear relations

$$-n\tau_n^{[2,2,2]}\tau_n^{[1,2,1]} = (-r+N)\tau_{n+1}^{[1,2,2]}\tau_{n-1}^{[2,2,1]} + (r+N-1)t^{-1}\tau_{n+1}^{[1,1,1]}\tau_{n-1}^{[2,3,2]},$$
  

$$m\tau_n^{[2,2,2]}\tau_n^{[1,2,1]} = (r+N-1)\tau_n^{[1,1,1]}\tau_n^{[2,3,2]} + (-r+N)\tau_n^{[1,2,2]}\tau_n^{[2,2,1]},$$
(3.27)

from (3.22) with (3.12), and

$$\tau_n^{[2,2,2]} \tau_n^{[1,1,0]} = -t^{-1} \tau_{n+1}^{[1,1,1]} \tau_{n-1}^{[2,2,1]} + \tau_n^{[1,1,1]} \tau_n^{[2,2,1]},$$
(3.28)

from the first relation in (3.23). These lead us to (3.21). We next consider the case where n' = m. When n + m is even, we get the bilinear relations

$$-m\tau_m^{[1,2,2]}\tau_m^{[1,1,1]} = N\tau_{m+1}^{[1,1,2]}\tau_{m-1}^{[1,2,1]} + Nt^{-1}\tau_{m+1}^{[0,1,1]}\tau_{m-1}^{[2,2,2]},$$
  

$$n\tau_m^{[1,2,2]}\tau_m^{[1,1,1]} = N\tau_m^{[0,1,1]}\tau_m^{[2,2,2]} + N\tau_m^{[1,1,2]}\tau_m^{[1,2,1]},$$
(3.29)

and

$$\tau_m^{[1,2,2]} \tau_m^{[2,1,2]} = \tau_m^{[1,1,2]} \tau_m^{[2,2,2]} - \tau_{m+1}^{[1,1,2]} \tau_{m-1}^{[2,2,2]}, \qquad (3.30)$$

from (3.22) and the second relation in (3.23), respectively. By using these relations, one can show (3.21) in a similar way to the case where n' = n. When n + m is odd, we use the bilinear relations

$$-m\tau_m^{[2,2,2]}\tau_m^{[1,2,1]} = (-r+N)\tau_{m+1}^{[1,2,2]}\tau_{m-1}^{[2,2,1]} + (r+N-1)t^{-1}\tau_{m+1}^{[1,1,1]}\tau_{m-1}^{[2,3,2]},$$
  

$$n\tau_m^{[2,2,2]}\tau_m^{[1,2,1]} = (r+N-1)\tau_m^{[1,1,1]}\tau_m^{[2,3,2]} + (-r+N)\tau_m^{[1,2,2]}\tau_m^{[2,2,1]},$$
(3.31)

$$\tau_m^{[1,2,1]} \tau_m^{[2,1,1]} = -\tau_{m+1}^{[1,1,1]} \tau_{m-1}^{[2,2,1]} + \tau_m^{[1,1,1]} \tau_m^{[2,2,1]},$$
(3.32)

which are obtained from (3.22) and the third relation in (3.23), respectively, to show (3.21).

We next give the verification of the compatibility condition (2.9) by using the bilinear relations

$$(c-a)\tau_{n'}^{0,-1,-1}\tau_{n'+1}^{-1,-1,0} - b\tau_{n'}^{0,0,0}\tau_{n'+1}^{-1,-2,-1} = (t-1)\tau_{n'}^{-1,-1,-1}\tau_{n'+1}^{0,-1,0},$$
  

$$(c-a)t\tau_{n'}^{0,-1,-1}\tau_{n'+1}^{-1,-1,0} - b\tau_{n'}^{0,0,0}\tau_{n'+1}^{-1,-2,-1} = (t-1)\tau_{n'}^{0,-1,0}\tau_{n'+1}^{-1,-1,-1}.$$
(3.33)

The derivation of these is discussed in Appendix A. We first consider the case where n' = n. When n + m is even, we get

$$(-r+N+1)\tau_n^{[1,1,1]}\tau_{n+1}^{[0,1,2]} + (r+N)\tau_n^{[1,2,2]}\tau_{n+1}^{[0,0,1]} = (t-1)\tau_n^{[0,1,1]}\tau_{n+1}^{[1,1,2]},$$
  

$$(-r+N+1)t\tau_n^{[1,1,1]}\tau_{n+1}^{[0,1,2]} + (r+N)\tau_n^{[1,2,2]}\tau_{n+1}^{[0,0,1]} = (t-1)\tau_n^{[1,1,2]}\tau_{n+1}^{[0,1,1]},$$
(3.34)

from the bilinear relations (3.33). Then we have

$$t^{\frac{1}{2}}v_{n,m} + v_{n+1,m+1} = t^{-\frac{n+1}{2}}(t-1)\frac{(r)_N}{(-r+1)_{N+1}}\frac{\tau_n^{[1,1,2]}\tau_{n+1}^{[0,1,1]}}{\tau_n^{[1,2,2]}\tau_{n+1}^{[0,1,2]}},$$

$$v_{n,m} + t^{\frac{1}{2}}v_{n+1,m+1} = t^{-\frac{n}{2}}(t-1)\frac{(r)_N}{(-r+1)_{N+1}}\frac{\tau_n^{[0,1,1]}\tau_{n+1}^{[1,2,2]}}{\tau_n^{[1,2,2]}\tau_{n+1}^{[0,1,2]}},$$
(3.35)

from which we arrive at the compatibility condition (2.9). When n + m is odd, we have

$$N\tau_{n}^{[1,2,1]}\tau_{n+1}^{[1,1,2]} + N\tau_{n}^{[2,2,2]}\tau_{n+1}^{[0,1,1]} = (t-1)\tau_{n}^{[1,1,1]}\tau_{n+1}^{[1,2,2]},$$

$$Nt\tau_{n}^{[1,2,1]}\tau_{n+1}^{[1,1,2]} + N\tau_{n}^{[2,2,2]}\tau_{n+1}^{[0,1,1]} = (t-1)\tau_{n}^{[1,2,2]}\tau_{n+1}^{[1,1,1]},$$
(3.36)

from (3.33). Calculating  $t^{\frac{1}{2}}v_{n,m} + v_{n+1,m+1}$  and  $v_{n,m} + t^{\frac{1}{2}}v_{n+1,m+1}$  by means of these relations, we see that we have (2.9). In the case where n' = m, one can verify the compatibility condition (2.9) in a similar manner.

Let us finally verify the similarity condition (2.11), which can be written as

$$\frac{n}{2} - \frac{1}{2}\chi_{n+m} - t\frac{d}{dt}\log v_{n,m} = \frac{nv_{n+1,m}}{v_{n+1,m} + v_{n-1,m}}.$$
(3.37)

Here, we take the factor  $\chi_{n+m}$  as  $\chi_{n+m} = r[(-1)^{n+m} - 1]$ . The relevant bilinear relations for the hypergeometric  $\tau$  function are

$$(D+n)\tau_n^{0,0,0} \cdot \tau_n^{0,-1,-1} = at^{-1}\tau_{n+1}^{-1,-1,-1}\tau_{n-1}^{1,0,0},$$

$$(D+b-c+1)\tau_m^{0,-1,-1} \cdot \tau_m^{0,0,0} = (b-c+1)\tau_m^{0,-1,0}\tau_m^{0,0,-1},$$

$$(D+a+m)\tau_m^{0,0,0} \cdot \tau_m^{0,-1,-1} = a\tau_m^{-1,-1,-1}\tau_m^{1,0,0}.$$
(3.38)

The derivation of these is obtained in Appendix A. We first consider the case where n' = n. When n + m is even, it is easy to see that we have

$$n\frac{v_{n+1,m}}{v_{n+1,m}+v_{n-1,m}} = -Nt^{-1}\frac{\tau_{n+1}^{[0,1,1]}\tau_{n-1}^{[2,2,2]}}{\tau_n^{[1,2,2]}\tau_n^{[1,1,1]}},$$
(3.39)

from the bilinear relation (3.24). We get

$$(D+n)\tau_n^{[1,2,2]} \cdot \tau_n^{[1,1,1]} = -Nt^{-1}\tau_{n+1}^{[0,1,1]}\tau_{n-1}^{[2,2,2]},$$
(3.40)

from the first relation in (3.38) with (3.11). From this we can obtain the similarity condition (3.37) as follows. When n + m is odd, we have

$$(D+n)\tau_n^{[2,2,2]} \cdot \tau_n^{[1,2,1]} = -t^{-1}(r+N-1)\tau_{n+1}^{[1,1,1]}\tau_{n-1}^{[2,3,2]}, \tag{3.41}$$

from the first relation in (3.38). This relation together with the first relation in (3.27) leads us to (3.37). Next, we discuss the case where n' = m. When n + m is even, we have

$$(D+N)\tau_m^{[1,2,2]} \cdot \tau_m^{[1,1,1]} = N\tau_m^{[1,1,2]}\tau_m^{[1,2,1]}, \qquad (3.42)$$

from the second relation in (3.38). Then we arrive at (3.37) by virtue of the second relation in (3.29). When n + m is odd, we get

$$(D+r+\frac{n-m-1}{2})\tau_m^{[1,2,1]}\cdot\tau_m^{[2,2,2]} = (r+N-1)\tau_m^{[1,1,1]}\tau_m^{[2,3,2]},$$
(3.43)

from the third relation in (3.38). Then we derive the similarity condition (3.37) by using the second relation in (3.31). This completes the proof of Theorem 3.1 and Proposition 3.2.

## 4 Extension of the domain

First, we extend the domain of the discrete power function to  $\mathbb{Z}^2$ . To determine the values of  $f_{n,m}$  in the second, third and fourth quadrants, we have to give the values of  $f_{-1,0}$  and  $f_{0,-1}$  as the initial conditions. Set the initial conditions as

$$f_{-1,0} = c_2 t^{2r}, \quad f_{0,-1} = c_3 t^{3r}, \tag{4.1}$$

where  $c_2$  and  $c_3$  are arbitrary constants. This is natural because these conditions reduce to

$$f_{1,0} = 1, \quad f_{0,1} = e^{\pi i r}, \quad f_{-1,0} = e^{2\pi i r}, \quad f_{0,-1} = e^{3\pi i r}$$
 (4.2)

at the original setting. Due to the symmetry of equations (2.7) and (2.3), we immediately obtain the explicit formula of  $f_{n,m}$  in the second and third quadrant.

**Corollary 4.1** Under the initial conditions  $f_{0,1} = c_1 t^r$  and (4.1), we have

$$f_{-n,m} = f_{n,m}|_{c_0 \mapsto c_2 t^{2r}}, \quad f_{-n,-m} = f_{n,m}|_{c_0 \mapsto c_2 t^{2r}, c_1 \mapsto c_3 t^{2r}}, \tag{4.3}$$

for  $n, m \in \mathbb{Z}_+$ .

Next, let us discuss the explicit formula in the fourth quadrant. Naively, we use the initial conditions  $f_{0,-1} = c_3 t^{3r}$  and  $f_{1,0} = c_0$  to get the formula  $f_{n,-m} = f_{n,m}|_{c_1 \mapsto c_3 t^{2r}}$ . However, this setting makes the discrete power function  $f_{n,m}$  become a single-valued function on  $\mathbb{Z}^2$ . In order to allow  $f_{n,m}$  to be multi-valued on  $\mathbb{Z}^2$ , we introduce a discrete analogue of the Riemann surface by the following procedure. Prepare an infinite number of  $\mathbb{Z}^2$ -planes, cut the positive part of the "real axis" of each  $\mathbb{Z}^2$ -plane and glue them in a similar way to the continuous case. The next step is to write the initial conditions (3.1) and (4.1) in polar form as

$$f(1,\pi k/2) = c_k t^{kr} \quad (k = 0, 1, 2, 3), \tag{4.4}$$



Figure 6: The discrete power function with  $\gamma = 5/2$  whose domain is  $\mathbb{Z}^2$ .



Figure 7: The discrete power function with  $\gamma = 5/2$  whose domain is the discrete Riemann surface.

where the first component, 1, denotes the absolute value of n + im and the second component,  $\pi k/2$ , is the argument. We must generalize the above initial conditions to those for arbitrary  $k \in \mathbb{Z}$  so that we obtain the explicit expression of  $f_{n,m}$  for each quadrant of each  $\mathbb{Z}^2$ -plane. Let us illustrate a typical case. When  $\frac{3}{2}\pi \leq \arg(n + im) \leq 2\pi$ , we solve the equations (2.7) and (2.3) under the initial conditions

$$f(1, 3\pi/2) = c_3 t^{3r}, \quad f(1, 2\pi) = c_4 t^{4r},$$
(4.5)

to obtain the formula

$$f_{-n,-m} = f_{n,m}|_{c_0 \mapsto c_4 t^{4r}, c_1 \mapsto c_3 t^{2r}} \quad (n, m \in \mathbb{Z}_+).$$
(4.6)

We present the discrete power function with  $\gamma = 5/2$  whose domain is  $\mathbb{Z}^2$  and the discrete Riemann surface in Figure 6 and 7, respectively. Note that the necessary and sufficient condition for the discrete power function to reduce to a single-valued function on  $\mathbb{Z}^2$  is  $(c_k = c_{k+4} \text{ and}) e^{4\pi i r} = 1$ , which means that the exponent  $\gamma$  is an integer.

## 5 Concluding remarks

The discrete logarithmic function and cases where  $\gamma \in 2\mathbb{Z}$  were excluded from the considerations in the previous sections. From the viewpoint of the theory of hypergeometric functions, these cases lead to integer differences in the characteristic exponents. Thus we need a different treatment for precise description of these cases. However, they may be obtained by some limiting procedures in principle. In fact, Agafonov has examined the case where  $\gamma = 2$  and  $\gamma = 0$  by using a limiting procedure [1,2], the former is the discrete power function  $Z^2$  and latter is the discrete logarithmic function. In general, one may obtain a description of these cases by introducing the functions  $\tilde{f}_{n,m}$ and  $\hat{f}_{n,m}$  as

$$\widetilde{f}_{n,m} := \begin{cases} \lim_{r \to j} \frac{1}{j} \frac{(-r+1)_j}{(r+1)_{j-1}} f_{n,m}, & \text{for } \gamma = 2j \in 2\mathbb{Z}_{>0} \\ \lim_{r \to -j} \frac{(-r+1)_j}{(r+1)_j} f_{n,m}, & \text{for } \gamma = -2j \in 2\mathbb{Z}_{<0} \end{cases}$$
(5.1)

and

$$\widehat{f}_{n,m} = \lim_{r \to 0} \frac{f_{n,m} - 1}{r},\tag{5.2}$$

respectively. The function  $\tilde{f}_{n,m}$  might coincide with the counterpart defined in section 6 of [3].

Moreover, it has been shown that the discrete power function and logarithmic function associated with hexagonal patterns are also described by some discrete Painlevé equations [4]. It may be an interesting problem to construct the explicit formula for them.

## Acknowledgements

The authors would like to express our sincere thanks to Professor Masaaki Yoshida for valuable suggestions and discussions. This work was partially supported by JSPS Grant-in-Aid for Scientific Research No. 21740126, 21656027 and 23340037, and by the Global COE Program Education and Research Hub for Mathematics-for-Industry from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

## A Bäcklund transformations of the sixth Painlevé equation

As a preparation, we give a brief review of the Bäcklund transformations and some of the bilinear equations for the  $\tau$  functions [12]. It is well-known that P<sub>VI</sub> (2.12) is equivalent to the Hamilton system

$$q' = \frac{\partial H}{\partial p}, \quad p' = -\frac{\partial H}{\partial q}, \quad ' = t(t-1)\frac{d}{dt},$$
 (A.1)

whose Hamiltonian is given by

$$H = f_0 f_3 f_4 f_2^2 - [\alpha_4 f_0 f_3 + \alpha_3 f_0 f_4 + (\alpha_0 - 1) f_3 f_4] f_2 + \alpha_2 (\alpha_1 + \alpha_2) f_0.$$
(A.2)

Here  $f_i$  and  $\alpha_i$  are defined by

$$f_0 = q - t, \quad f_3 = q - 1, \quad f_4 = q, \quad f_2 = p,$$
 (A.3)

and

$$\alpha_0 = \theta, \quad \alpha_1 = \kappa_{\infty}, \quad \alpha_3 = \kappa_1, \quad \alpha_4 = \kappa_0$$
 (A.4)

with  $\alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4 = 1$ . The Bäcklund transformations of P<sub>VI</sub> are described by

$$s_i(\alpha_j) = \alpha_j - a_{ij}\alpha_i$$
 (*i*, *j* = 0, 1, 2, 3, 4), (A.5)

$$s_2(f_i) = f_i + \frac{\alpha_2}{f_2}, \quad s_i(f_2) = f_2 - \frac{\alpha_i}{f_i} \quad (i = 0, 3, 4),$$
 (A.6)

$$s_{5}: \alpha_{0} \leftrightarrow \alpha_{1}, \quad \alpha_{3} \leftrightarrow \alpha_{4}, \quad f_{2} \mapsto -\frac{f_{0}(f_{2}f_{0} + \alpha_{2})}{t(t-1)}, \quad f_{4} \mapsto t\frac{f_{3}}{f_{0}},$$

$$s_{6}: \alpha_{0} \leftrightarrow \alpha_{3}, \quad \alpha_{1} \leftrightarrow \alpha_{4}, \quad f_{2} \mapsto -\frac{f_{4}(f_{4}f_{2} + \alpha_{2})}{t}, \quad f_{4} \mapsto \frac{t}{f_{4}},$$

$$s_{7}: \alpha_{0} \leftrightarrow \alpha_{4}, \quad \alpha_{1} \leftrightarrow \alpha_{3}, \quad f_{2} \mapsto \frac{f_{3}(f_{3}f_{2} + \alpha_{2})}{t-1}, \quad f_{4} \mapsto \frac{f_{0}}{f_{3}},$$
(A.7)

where  $A = (a_{ij})_{i,j=0}^4$  is the Cartan matrix of type  $D_4^{(1)}$ . Then the group of birational transformations  $\langle s_0, \ldots, s_7 \rangle$  generate the extended affine Weyl group  $\widetilde{W}(D_4^{(1)})$ . In fact, these generators satisfy the fundamental relations

$$s_i^2 = 1$$
  $(i = 0, ..., 7), \quad s_i s_2 s_i = s_2 s_i s_2$   $(i = 0, 1, 3, 4),$  (A.8)

and

$$s_{5}s_{\{0,1,2,3,4\}} = s_{\{1,0,2,4,3\}}s_{5}, \quad s_{6}s_{\{0,1,2,3,4\}} = s_{\{3,4,2,0,1\}}s_{6}, \quad s_{7}s_{\{0,1,2,3,4\}} = s_{\{4,3,2,1,0\}}s_{7}, \\ s_{5}s_{6} = s_{6}s_{5}, \quad s_{5}s_{7} = s_{7}s_{5}, \quad s_{6}s_{7} = s_{7}s_{6}.$$
(A.9)

We add a correction term to the Hamiltonian H as follows,

$$H_0 = H + \frac{t}{4} \left[ 1 + 4\alpha_1 \alpha_2 + 4\alpha_2^2 - (\alpha_3 + \alpha_4)^2 \right] + \frac{1}{4} \left[ (\alpha_1 + \alpha_4)^2 + (\alpha_3 + \alpha_4)^2 + 4\alpha_2 \alpha_4 \right].$$
(A.10)

This modification gives a simpler behavior of the Hamiltonian with respect to the Bäcklund transformations. From the corrected Hamiltonian, we introduce a family of Hamiltonians  $h_i$  (i = 0, 1, 2, 3, 4) as

$$h_0 = H_0 + \frac{t}{4}, \quad h_1 = s_5(H_0) - \frac{t-1}{4}, \quad h_3 = s_6(H_0) + \frac{1}{4}, \quad h_4 = s_7(H_0), \quad h_2 = h_1 + s_1(h_1).$$
 (A.11)

Next, we also introduce  $\tau$  functions  $\tau_i$  (i = 0, 1, 2, 3, 4) by  $h_i = (\log \tau_i)'$ . Imposing the condition that the action of the  $s_i$ 's on the  $\tau$  functions also commute with the derivation ', one can lift the Bäcklund transformations to the  $\tau$  functions. The action of  $\widetilde{W}(D_4^{(1)})$  is given by

$$s_0(\tau_0) = f_0 \frac{\tau_2}{\tau_0}, \quad s_1(\tau_1) = \frac{\tau_2}{\tau_1}, \quad s_2(\tau_2) = \frac{f_2}{\sqrt{t}} \frac{\tau_0 \tau_1 \tau_3 \tau_4}{\tau_2}, \quad s_3(\tau_3) = f_3 \frac{\tau_2}{\tau_3}, \quad s_4(\tau_4) = f_4 \frac{\tau_2}{\tau_4}, \quad (A.12)$$

and

$$s_{5}: \tau_{0} \mapsto [t(t-1)]^{\frac{1}{4}}\tau_{1}, \quad \tau_{1} \mapsto [t(t-1)]^{-\frac{1}{4}}\tau_{0}, \tau_{3} \mapsto t^{-\frac{1}{4}}(t-1)^{\frac{1}{4}}\tau_{4}, \quad \tau_{4} \mapsto t^{\frac{1}{4}}(t-1)^{-\frac{1}{4}}\tau_{3}, \quad \tau_{2} \mapsto [t(t-1)]^{-\frac{1}{2}}f_{0}\tau_{2},$$
(A.13)

$$s_{6}: \tau_{0} \mapsto it^{\frac{1}{4}}\tau_{3}, \quad \tau_{3} \mapsto -it^{-\frac{1}{4}}\tau_{0}, \quad \tau_{1} \mapsto t^{-\frac{1}{4}}\tau_{4}, \quad \tau_{4} \mapsto t^{\frac{1}{4}}\tau_{1}, \quad \tau_{2} \mapsto t^{-\frac{1}{2}}f_{4}\tau_{2}, \tag{A.14}$$

$$s_{7}: \quad \tau_{0} \mapsto (-1)^{-\frac{3}{4}}(t-1)^{\frac{1}{4}}\tau_{4}, \quad \tau_{4} \mapsto (-1)^{\frac{3}{4}}(t-1)^{-\frac{1}{4}}\tau_{0}, \\ \tau_{1} \mapsto (-1)^{\frac{3}{4}}(t-1)^{-\frac{1}{4}}\tau_{3}, \quad \tau_{3} \mapsto (-1)^{-\frac{3}{4}}(t-1)^{\frac{1}{4}}\tau_{1}, \\ \tau_{2} \mapsto -i(t-1)^{-\frac{1}{2}}f_{3}\tau_{2}.$$
(A.15)

We note that some of the fundamental relations are modified

$$s_i s_2(\tau_2) = -s_2 s_i(\tau_2)$$
 (*i* = 5, 6, 7), (A.16)

$$s_{5}s_{6}\tau_{\{0,1,2,3,4\}} = \{i, -i, -1, -i, i\}s_{6}s_{5}\tau_{\{0,1,2,3,4\}},$$

$$s_{5}s_{7}\tau_{\{0,1,2,3,4\}} = \{i, -i, -1, i, -i\}s_{7}s_{5}\tau_{\{0,1,2,3,4\}},$$

$$s_{6}s_{7}\tau_{\{0,1,2,3,4\}} = \{-i, -i, -1, i, i\}s_{7}s_{6}\tau_{\{0,1,2,3,4\}}.$$
(A.17)

Let us introduce the translation operators

$$\begin{aligned} \widehat{T}_{13} &= s_1 s_2 s_0 s_4 s_2 s_1 s_7, \quad \widehat{T}_{40} &= s_4 s_2 s_1 s_3 s_2 s_4 s_7, \\ \widehat{T}_{34} &= s_3 s_2 s_0 s_1 s_2 s_3 s_5, \quad T_{14} &= s_1 s_4 s_2 s_0 s_3 s_2 s_6, \end{aligned}$$
(A.18)

whose action on the parameters  $\vec{\alpha} = (\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4)$  is given by

$$T_{13}(\vec{\alpha}) = \vec{\alpha} + (0, 1, 0, -1, 0),$$
  

$$\widehat{T}_{40}(\vec{\alpha}) = \vec{\alpha} + (-1, 0, 0, 0, 1),$$
  

$$\widehat{T}_{34}(\vec{\alpha}) = \vec{\alpha} + (0, 0, 0, 1, -1),$$
  

$$T_{14}(\vec{\alpha}) = \vec{\alpha} + (0, 1, -1, 0, 1).$$
  
(A.19)

We denote  $\tau_{k,l,m,n'} = T_{14}^{n'} \widehat{T}_{34}^m \widehat{T}_{40}^l \widehat{T}_{13}^k (\tau_0) (k, l, m, n' \in \mathbb{Z})$ . By using this notation, we have

$$\begin{aligned} \tau_{0,0,0,0} &= \tau_{0}, & \tau_{-1,-1,-1,0} &= [t(t-1)]^{\frac{1}{4}} \tau_{1}, \\ \tau_{0,-1,-1,0} &= (-1)^{-\frac{3}{4}} t^{\frac{1}{4}} \tau_{3}, & \tau_{0,-1,0,0} &= (-1)^{-\frac{3}{4}} (t-1)^{\frac{1}{4}} \tau_{4}, \\ \tau_{-1,-2,-1,1} &= (-1)^{-\frac{1}{4}} s_{0}(\tau_{0}), & \tau_{0,-1,0,1} &= (-1)^{-\frac{3}{4}} [t(t-1)]^{\frac{1}{4}} s_{1}(\tau_{1}), \\ \tau_{-1,-1,0,1} &= -it^{\frac{1}{4}} s_{3}(\tau_{3}), & \tau_{-1,-1,-1,1} &= (t-1)^{\frac{1}{4}} s_{4}(\tau_{4}), \end{aligned}$$
(A.20)

for instance. When the parameters  $\vec{\alpha}$  take the values

$$(\alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4) = (-b, a + n', -n', c - a, b - c + 1 + n'),$$
(A.21)

the function  $\tau_{k,l,m,n'}$  relates to the hypergeometric  $\tau$  function  $\tau_{n'}^{k,l,m}$  introduced in Proposition 2.6 by [12]

$$\tau_{k,l,m,n'} = \omega_{k,l,m,n'} \tau_{n'}^{k,l,m} t^{-(\hat{a}+\hat{b}-\hat{c}+2n')^2/4 - (\hat{a}-\hat{b}-n')^2/4 + n'(\hat{b}+n') - n'(n'-1)/2} (t-1)^{(\hat{a}+\hat{b}-\hat{c}+2n')^2/4 + 1/2},$$
(A.22)

where we denote  $\hat{a} = a + k$ ,  $\hat{b} = b + l + 1$  and  $\hat{c} = c + m$ , and the constants  $\omega_{k,l,m,n'} = \omega_{k,l,m,n'}(a, b, c)$  are determined by the recurrence relations

$$\begin{aligned}
\omega_{k+1,l,m,i}\omega_{k-1,l,m,i} &= i\hat{a}(\hat{c} - \hat{a})\omega_{k,l,m,i}^{2}, \\
\omega_{k,l+1,m,i}\omega_{k,l-1,m,i} &= -i\hat{b}(\hat{c} - \hat{b})\omega_{k,l,m,i}^{2}, \\
\omega_{k,l,m+1,i}\omega_{k,l,m-1,i} &= (\hat{c} - \hat{a})(\hat{c} - \hat{b})\omega_{k,l,m,i}^{2}
\end{aligned}$$
(A.23)

and

$$\omega_{k,l,m,n'+1}\omega_{k,l,m,n'-1} = -\omega_{k,l,m,n'}^2$$
(A.24)

with initial conditions

$$\begin{aligned}
\omega_{-1,-2,-1,1} &= (-1)^{-1/4}b, & \omega_{0,-2,-1,1} &= b, \\
\omega_{-1,-1,-1,1} &= 1, & \omega_{0,-1,-1,1} &= (-1)^{-1/4}, \\
\omega_{-1,0,0,1} &= -(-1)^{-3/4}(c-a), & \omega_{0,0,0,1} &= -i, \\
\omega_{-1,-1,0,1} &= -i(c-a), & \omega_{0,-1,0,1} &= (-1)^{-3/4},
\end{aligned}$$
(A.25)

$$\begin{split} \omega_{-1,-2,-1,0} &= (-1)^{-3/4} b, & \omega_{0,-2,-1,0} &= -b, \\ \omega_{-1,-1,-1,0} &= 1, & \omega_{0,-1,-1,0} &= (-1)^{-3/4}, \\ \omega_{-1,0,0,0} &= (-1)^{-3/4} (c-a), & \omega_{0,0,0,0} &= 1, \\ \omega_{-1,-1,0,0} &= c-a, & \omega_{0,-1,0,0} &= (-1)^{-3/4}. \end{split}$$
(A.26)

From the above formulation, one can obtain the bilinear equations for the  $\tau$  functions. For examle, let us express the Bäcklund transformations  $s_2(f_i) = f_i + \frac{\alpha_2}{f_2}$  (i = 0, 3, 4) in terms of the  $\tau$ functions  $\tau_i$  (j = 0, 1, 3, 4). We have by using (A.12)

$$\alpha_{2}t^{-\frac{1}{2}}\tau_{3}\tau_{4} - s_{1}(\tau_{1})s_{2}s_{0}(\tau_{0}) + s_{0}(\tau_{0})s_{2}s_{1}(\tau_{1}) = 0,$$
  

$$\alpha_{2}t^{-\frac{1}{2}}\tau_{0}\tau_{4} - s_{1}(\tau_{1})s_{2}s_{3}(\tau_{3}) + s_{3}(\tau_{3})s_{2}s_{1}(\tau_{1}) = 0,$$
  

$$\alpha_{2}t^{-\frac{1}{2}}\tau_{0}\tau_{3} - s_{1}(\tau_{1})s_{2}s_{4}(\tau_{4}) + s_{4}(\tau_{4})s_{2}s_{1}(\tau_{1}) = 0.$$
  
(A.27)

Applying the affine Weyl group  $\widetilde{W}(D_4^{(1)})$  on these equations, we obtain

$$(\alpha_0 + \alpha_2 + \alpha_4) t^{-\frac{1}{2}} \tau_3 s_4(\tau_4) - s_1(\tau_1) s_4 s_2 s_0(\tau_0) + \tau_0 s_0 s_4 s_2 s_1(\tau_1) = 0,$$
  

$$(\alpha_0 + \alpha_2 + \alpha_4) t^{\frac{1}{2}} \tau_1 \tau_3 - \tau_4 s_4 s_2 s_0(\tau_0) + \tau_0 s_0 s_2 s_4(\tau_4) = 0,$$
(A.28)

$$(\alpha_{0} + \alpha_{1} + \alpha_{2}) t^{-\frac{1}{2}} \tau_{3} \tau_{4} - \tau_{1} s_{1} s_{2} s_{0}(\tau_{0}) + \tau_{0} s_{0} s_{2} s_{1}(\tau_{1}) = 0,$$

$$(\alpha_{0} + \alpha_{1} + \alpha_{2}) t^{\frac{1}{2}} s_{1}(\tau_{1}) \tau_{3} - s_{4}(\tau_{4}) s_{1} s_{2} s_{0}(\tau_{0}) + \tau_{0} s_{0} s_{1} s_{2} s_{4}(\tau_{4}) = 0,$$

$$(\alpha_{2} + \alpha_{3} + \alpha_{4}) t^{-\frac{1}{2}} \tau_{0} \tau_{1} - \tau_{4} s_{4} s_{2} s_{3}(\tau_{3}) + \tau_{3} s_{3} s_{2} s_{4}(\tau_{4}) = 0,$$
(A.29)
$$(\alpha_{2} + \alpha_{3} + \alpha_{4}) t^{-\frac{1}{2}} \tau_{0} \tau_{1} - \tau_{4} s_{4} s_{2} s_{3}(\tau_{3}) + \tau_{3} s_{3} s_{2} s_{4}(\tau_{4}) = 0,$$
(A.29)

$$\begin{aligned} \alpha_2 + \alpha_3 + \alpha_4) t^{-\frac{1}{2}} \tau_0 \tau_1 - \tau_4 s_4 s_2 s_3(\tau_3) + \tau_3 s_3 s_2 s_4(\tau_4) &= 0, \\ \alpha_2 + \alpha_3 + \alpha_4) t^{-\frac{1}{2}} s_4(\tau_4) \tau_0 - s_1(\tau_1) s_4 s_2 s_3(\tau_3) + \tau_3 s_3 s_4 s_2 s_1(\tau_1) &= 0, \end{aligned}$$
(A.30)

and

$$\alpha_{2}t^{-\frac{1}{2}}\tau_{0}\tau_{3} - s_{1}(\tau_{1})s_{2}s_{4}(\tau_{4}) + s_{4}(\tau_{4})s_{2}s_{1}(\tau_{1}) = 0,$$

$$(\alpha_{1} + \alpha_{4} + \alpha_{2})t^{-\frac{1}{2}}\tau_{0}\tau_{3} - \tau_{1}s_{1}s_{2}s_{4}(\tau_{4}) + \tau_{4}s_{4}s_{2}s_{1}(\tau_{1}) = 0.$$
(A.31)

For instance, the first equation in (A.28) can be obtained by applying  $s_0s_4$  on the first one in (A.27). We also get the second equation in (A.28) by applying  $s_0s_4s_6$  on the second one in (A.27). Other equations can be derived in a similar manner. By applying the translation  $T_{14}^{n'} \widehat{T}_{34}^{l} \widehat{T}_{40}^{l} \widehat{T}_{13}^{k}$  to the bilinear relations (A.28) and noticing (A.20), we get

$$(\alpha_{0} + \alpha_{2} + \alpha_{4} - m) t^{-\frac{1}{2}} \tau_{k,l-1,m-1,n'} \tau_{k-1,l-1,m-1,n'+1} + \tau_{k,l-1,m,n'+1} \tau_{k-1,l-1,m-2,n'} + \tau_{k,l,m,n'} \tau_{k-1,l-2,m-2,n'+1} = 0,$$

$$(\alpha_{0} + \alpha_{2} + \alpha_{4} - m) \tau_{k-1,l-1,m-1,n'} \tau_{k,l-1,m-1,n'} + \tau_{k,l-1,m,n'} \tau_{k-1,l-1,m-2,n'} - \tau_{k,l,m,n'} \tau_{k-1,l-2,m-2,n'} = 0,$$
(A.32)

and then (3.13) for the hypergeometric  $\tau$  functions. Similarly, we obtain for the hypergeometric  $\tau$ functions (3.14), (3.15) and (3.22) from (A.29), (A.30) and (A.31), respectively. The constraints

$$f_0 = f_4 - t, \quad f_3 = f_4 - 1,$$
 (A.33)

yield

$$\tau_0 s_4 s_2 s_0(\tau_0) = s_4(\tau_4) s_2 s_4(\tau_4) - t\tau_1 s_4 s_2 s_1(\tau_1),$$
  

$$\tau_0 s_1 s_2 s_0(\tau_0) = \tau_4 s_1 s_2 s_4(\tau_4) - t s_1(\tau_1) s_2 s_1(\tau_1),$$
  

$$\tau_3 s_4 s_2 s_3(\tau_3) = s_4(\tau_4) s_2 s_4(\tau_4) - \tau_1 s_4 s_2 s_1(\tau_1),$$
  
(A.34)

$$\tau_3 s_3(\tau_3) - \tau_0 s_0(\tau_0) = (t-1)\tau_1 s_1(\tau_1),$$
  

$$t\tau_3 s_3(\tau_3) - \tau_0 s_0(\tau_0) = (t-1)\tau_4 s_4(\tau_4),$$
(A.35)

from which we obtain (3.23) and (3.33), respectively. Due to (A.11) we have the relation

$$h_0 - h_3 = (t - 1) \left[ f_2 f_4 + \frac{1}{2} (1 - \alpha_3 - \alpha_4) \right].$$
 (A.36)

Then we get the bilinear relations

$$D\tau_{0} \cdot \tau_{3} = t^{\frac{1}{2}} s_{4}(\tau_{4}) s_{2} s_{1}(\tau_{1}) + \frac{1}{2} (1 - \alpha_{3} - \alpha_{4}) \tau_{0} \tau_{3},$$
  

$$D\tau_{0} \cdot \tau_{3} = t^{\frac{1}{2}} \tau_{4} s_{4} s_{2} s_{1}(\tau_{1}) + \frac{1}{2} (1 - \alpha_{3} + \alpha_{4}) \tau_{0} \tau_{3},$$
  

$$D\tau_{0} \cdot \tau_{3} = t^{\frac{1}{2}} \tau_{1} s_{1} s_{2} s_{4}(\tau_{4}) + \frac{1}{2} (\alpha_{0} - \alpha_{1}) \tau_{0} \tau_{3},$$
  
(A.37)

where *D* denotes Hirota's differential operator defined by  $Dg \cdot f = t \left(\frac{dg}{dt}f - g\frac{df}{dt}\right)$ . By applying the translation  $T_{14}^{n'} \widehat{T}_{34}^{l} \widehat{T}_{40}^{l} \widehat{T}_{13}^{k}$  to the first bilinear relation of (A.37), one gets

$$\left[D + \frac{1}{2}\left(\alpha_3 + \alpha_4 - k + l + n' - \frac{1}{2}\right)\right]\tau_{k,l,m,n'}\cdot\tau_{k,l-1,m-1,n'} = -t^{\frac{1}{2}}(t-1)^{-\frac{1}{2}}\tau_{k-1,l-1,m-1,n'+1}\tau_{k+1,l,m,n'-1},$$
(A.38)

which is reduced to the first relation of (3.38). The second and third relations of (A.37) also yield their counterparts in (3.38).

### References

- [1] Agafonov, S. I. "Imbedded circle patterns with the combinatorics of the square grid and discrete Painlevé equations." *Discrete Comput. Geom.* 29, no. 2 (2003): 305–319.
- [2] Agafonov, S. I. "Discrete Riccati equation, hypergeometric functions and circle patterns of Schramm type." *Glasg. Math. J.* 47, no. A (2005): 1–16.
- [3] Agafonov, S. I., and Bobenko, A. I. "Discrete  $Z^{\gamma}$  and Painlevé equations." *Internat. Math. Res. Notices* 2000, no. 4: 165–193.
- [4] Agafonov, S. I., and Bobenko, A. I. "Hexagonal circle patterns with constant intersection angles and discrete Painlevé and Riccati equations." J. Math. Phys. 44, no. 8 (2003): 3455– 3469.
- [5] Beardon, A. F., and Stephenson, K. "The uniformization theorem for circle packing." *Indiana Univ. Math. J.* 39, no. 4 (1990): 1383–1425.
- [6] Bobenko, A. I. "Discrete conformal maps and surfaces" in Symmetries and integrability of difference equations (Canterbury, 1996), 97–108, London Math. Soc. Lecture Note Ser. 255 (Cambridge Univ. Press, Cambridge, 1999).
- [7] Bobenko, A. I., and Hoffmann, T. "Hexagonal circle patterns and integrable systems: patterns with constant angles." *Duke Math. J.* 116, no. 3 (2003): 525–566.

- [8] Bobenko, A. I., Hoffmann, T., and Suris, Y. B. "Hexagonal circle patterns and integrable systems: patterns with the multi-ratio property and Lax equations on the regular triangular lattice." *Int. Math. Res. Not.* 2002, no. 3: 111–164.
- [9] Bobenko, A. I., and Pinkall, U. "Discrete isothermic surfaces." J. Reine Angew. Math. 475 (1996): 187–208.
- [10] Bobenko, A. I., and Pinkall, U. "Discretization of surfaces and integrable systems." Discrete integrable geometry and physics (Vienna, 1996), 3–58, Oxford Lecture Ser. Math. Appl., 16, Oxford Univ. Press, New York, 1999.
- [11] Hay, M., Kajiwara, K., and Masuda, T. "Bilinearization and special solutions to the discrete Schwarzian KdV equation." *J. Math-for-Ind.* 3, (2011): 53–62.
- [12] Masuda, T. "Classical transcendental solutions of the Painlevé equations and their degeneration." *Tohoku Math. J.* 56, no. 4 (2004): 467–490.
- [13] Nijhoff, F. W., Ramani, A., Grammaticos, B., and Ohta, Y. "On discrete Painlevé equations associated with the lattice KdV systems and the Painlevé VI equation." *Stud. Appl. Math.* 106, no. 3 (2001): 261–314.
- [14] Rodin, B. "Schwarz's lemma for circle packings." Invent. Math. 89, no. 2 (1987): 271–289.
- [15] Schramm, O. "Circle patterns with the combinatorics of the square grid." Duke Math. J. 86, no. 2 (1997): 347–389.
- [16] Stephenson, K. Introduction to circle packing, New York: Cambridge University Press, 2005.
- [17] Thurston, W. P. "The finite Riemann mapping theorem." Invited address, International Symposium in Celebration of the Proof of the Bieberbach Conjecture (Purdue University, 1985).

## List of MI Preprint Series, Kyushu University

### The Global COE Program Math-for-Industry Education & Research Hub

#### $\mathbf{MI}$

MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI Abstract collision systems simulated by cellular automata

#### MI2008-2 Eiji ONODERA

The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds

- MI2008-3 Hiroaki KIDO On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU Numerical computations of cavity flow problems by a pressure stabilized characteristiccurve finite element scheme
- MI2008-5 Yoshiyasu OZEKI Torsion points of abelian varieties with values in nfinite extensions over a padic field
- MI2008-6 Yoshiyuki TOMIYAMA Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA Alpha-determinant cyclic modules and Jacobi polynomials

- MI2008-10 Sangyeol LEE & Hiroki MASUDA Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO On the  $L^2$  a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials
- MI2008-14 Takashi NAKAMURA Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI Variable selection for functional regression model via the  $L_1$  regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCHI & Yuichiro TAGUCHII Flat modules and Groebner bases over truncated discrete valuation rings

- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA Hypergeometric -functions of the q-Painlevé system of type  $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA On Gaussian decay estimates of solutions to some linear elliptic equations and its applications
- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI Large time behavior of the semigroup on  $L^p$  spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain

- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO On very accurate enclosure of the optimal constant in the a priori error estimates for  $H_0^2$ -projection

- MI2009-26 Manabu YOSHIDA Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance
- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI Finite element computation for scattering problems of micro-hologram using DtN map

- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO Abstract collision systems on groups
- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA An algebraic approach to underdetermined experiments

MI2010-10 Kei HIROSE & Sadanori KONISHI Variable selection via the grouped weighted lasso for factor analysis models

- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE Lagrangian approach to weakly nonlinear stability of an elliptical flow

MI2010-21 Hiroki MASUDA

Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test

- MI2010-22 Toshimitsu TAKAESU A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiko FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra
- MI2010-25 Toshimitsu TAKAESU On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI Spin-spin correlation functions of the q-VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI Nonlinear regression modeling and spike detection via Gaussian basis expansions

- MI2010-32 Nobutaka NAKAZONO Hypergeometric  $\tau$  functions of the *q*-Painlevé systems of type  $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA CAP representations of inner forms of Sp(4) with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide FUKUMOTO& Alexander B. SAMOKHIN Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI Anderson-Darling test and the Malliavin calculus
- MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI Nonlinear regression modeling via Compressed Sensing
- MI2011-5 Hiroshi INOUE Implications in Compressed Sensing and the Restricted Isometry Property
- MI2011-6 Daeju KIM & Sadanori KONISHI Predictive information criterion for nonlinear regression model based on basis expansion methods
- MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI Group variable selection via relevance vector machine

MI2011-8 Jan BREZINA & Yoshiyuki KAGEI Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow Group variable selection via relevance vector machine

- MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC Recursive structures in the multispecies TASEP
- MI2011-10 Kazunori YASUTAKE On projective space bundle with nef normalized tautological line bundle
- MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA An explicit formula for the discrete power function associated with circle patterns of Schramm type