Some Remarks on Symbols, Multiple Allelomorphism, Crossing Over and Linkage Groups in The Silkworm

Tanaka, Yoshimaro Sericultural Laboratory, Kyushu Imperial University

https://doi.org/10.5109/22333

出版情報:九州大学大学院農学研究院紀要. 1 (8), pp.275-279, 1927-12. Kyushu Imperial University バージョン: 権利関係:

SOME REMARKS ON SYMBOLS, MULTIPLE ALLE-LOMORPHISM, CROSSING OVER AND LINKAGE GROUPS IN THE SILKWORM⁴

Yoshimaro TANAKA

I. REVISION OF SYMBOLS FOR TWO LARVAL MARKINGS

According to the nomenclature of genes in *Drosophila*, a recessive mutation is to be represented by a small letter or letters significant to the mutated character, and its normal allelomorph by the corresponding capital letter with or without the suffix of the small letters. Since 1917 I have followed this method of nomenclature, though a different scheme was used in my earlier works. For instance, the plain marking was represented by P, to which q was added whenever necessary to distinguish the plain from normal. Thus it is actually not P or p that distinguishes the plain from the other markings. This is at least confusing if not absurd especially when there are many recessive genes affecting the same organ as already pointed out by MORGAN (1913). It seems, therefore, desirable to revise the symbols for recessive larval markings, i. e. plain and quail. To make clear the history of changes, the symbols are listed below:

	Papers		Normal	Plain	Quail	Pale quail	
	Танака 1913a, b. 1914	. 1915.	N	n			
	TANAKA 1916.		\mathbf{PQ}	\mathbf{Pq}	$^{\rm pQ}$	Pq	
	New symbols		PQ	р	q	Pq	
	Consequently,		8 8		<u>*</u>)	2	
			Striped	Moricaud	Striped quail	Moricaud quail	
	TANAKA 1916.	32	SPQ	MPQ	S_{PQ}	MpQ	
	New symbols		S	М	Sq	Mq	
51 B.		- 1 H	272 D				

I Contributions from the Sericultural Laboratory, Kyushu Imperial University. No. 10.

Yoshimaro Tanaka

Bombyx mori has been domesticated for thousands of years, and no one can tell exactly the origin of our silkworms, though the wild mulberry silkworm is generally presumed to be their ancestral form. The wild mulberry silkworm is sometimes called *Theophila mandarina*, and sometimes *Bombyx mandarina*, but *Bombyx mori var. mandarina* seems to me better fitting as its scientific name. Besides the unsettled systematic position, the genetic characters of the wild silkworm are not yet fully worked out. It seems, therefore, more convenient, in *Bombyx*, to take the more common type as the standard, and the less common form as the mutant as BRIDGES and MORGAN propose (1923) for the cases where the wild type is unknown or not certainly known. Some important standard characters are compared to those of *mandarina*.

Standa	ard form of B. mori	Mandarina form
Larval marking	Normal	Moricaud
Blood color	Colorless	Light yellowish
Cocoon color	White	Light green
Voltinism	Univoltine	Polyvoltine
Number of molts before spinning	Four	Four
Skin	Opaque	Opaque

Determination of the standard form may sometimes be arbitrary, because we can not say which larval marking, "normal" or "plain," for example, is really more common in the world. It is, however, true that we can pick the more common type as the standard without difficulty in majority of cases.

II. MULTIPLE ALLELOMORPHISM

Special relations between four larval markings, striped, moricaud, normal and plain described in my previous papers (1913b, 1914) were pointed out by STURTEVANT (1915) as they form a series of multiple allelomorphs. Though I adopted the explanation by complete coupling and repulsion in my paper of 1916 as I did in the preceding work, I have later accepted the idea of multiple allelomorphism for this case. The students of *Drosophila* make use of exponents to denote multiple allelomorphs, as w^e, e^s etc. for example. But I am, for the time being, content with the use of the simpler expressions, S (striped) and M (moricaud) instead of symbols p⁸ and p^M.

III. CROSSING OVER VALUE

It may be somewhat useful that the phenotypic ratios published in my previous paper (1916) to be expressed in terms of the crossing over

276

value. As there is no crossing over in the female sex in *Bombyx*, the cross over value in diheterozygous males may be calculated from F_2 results by the following formula, in which k, l, m and n represent the numbers of individuals belonging to AB, Ab, aB and ab F_2 phenotypes.

Cross-over value =
$$1 \div \left\{ \frac{k+n}{2(l+m)} + 0.5 \right\} \times 100$$

This formula is available for F_2 ratios in "coupling" only, while in case of "repulsion" the F_2 ratio is uniformly 2 : I : I whatever the linkage value may be (TANAKA 1916), which makes caculation of cross-over per cents impossible. The crossing over per cents are to be obtained only through back-crossing, when two dominants are introduced from different parents.

a) Striped-yellow linkage.

2 D C (24)	SY	Sy	s¥	sy	Totals	c.o. %
"Coupling " Back cross	4421	1631	1735	4418	12205	27.58
"Coupling \r ₂	10958	1075	IIOI	3017	16151	26.95
" Repulsion " (back cross)	112	338	372	119	941	24.55
Weighted mean of c.o.	8					27.22
b) Moricaud-yello	w linkage	•				
	MY	Му	mY	my	Totals	c.o. %
"Compling" Back cross	3053	1030	1057	3129	8269	25.24
Coupring F2	10512	873	858	2949	15192	22.78
"Repulsion" (back cross)	86	453	483	104	1126	24.02
Weighted mean of c.o.						23.66
c) Plain-yellow li	nkage.					
	$\mathbf{P}\mathbf{Y}$	Ру	$\mathbf{P}\mathbf{Y}$	ру	Totals	c.o. %
"Coupling " Back cross	719	207	186	614	1726	22.77
lF2	3361	376	294	910	4941	27.12
"Repulsion" (back cross)	280	827	756	222	2085	24.08
Weighted mean of c.o.						25.54

As the genes S,M and p are multiple allelomorphs, they ought to be all the same in cross over percentage, but there is actually remarkable deviation from expectation. This is, however, not astonishing when we take into consideration the fact that very different numbers were obtained from back-crossing and F_2 results even between the same genes. For instance, crossing over value calculated from back-crossing in p-Y linkage is 22.77, while that derived from F_2 figures is 27.12, showing a difference of 4.35%. Kogure (1926) studied, in my laboratory, on the crossing over variation between S and Y, and discovered the existence of a Yoshimaro Tanaka

dominant modifier which partly suppresses crossing over between them and reduces it as low as 12%. He finds also that the crossing over percentage is influenced by the room temperature where the heterozygotes were reared: the average crossing over value of his "high crossing over" strain was 21.48 in about 30° C, while it was 25.86% in about 19°C, and some intermediate figures were obtained in intermediate thermometric degrees. The weighted mean of cross-over per cents between Y and the locus of p, S,M is 25.6 in my experiments.

IV. LINKAGE GROUPS AND CHROMOSOME MAPS

Four linkage groups have hitherto been described in the silkworm. They are as follow:



² The description in English will soon appear.

278

A few more possible cases of linkage are known to me at present, but they have not been fully worked out.

The chromosome maps of the silkworm are drawn on basis of published data, taking the cross-over per cents as units as such.

LITERATURE CITED

- BRIDGES, C. B. and MORGAN, T. H. -- 1923. The Third-Chromosome Group of Mutant Characters of Drosophila melanogaster. Carnegie Inst. Washington, Publication No.327.
- KOGURE, M.—1926. Modifying Factor Influencing the Striped-Yellow Crossing Over Value in the Silkworm. Bulteno Scienca de la Fakultato Terkultura, Kjuŝu Imperia Universitato. Vol.2, No.1, pp. 33-50. (In Japanese with English Résumé).
- MORGAN, T. H.—1913. Factors and Unit Characters in Mendelian Heredity. Amer. Nat. Vol.47, pp. 5-16.
- NISHIKAWA, H.-1922. On Sex-Linked Inheritance of Translucent Silkworm. (In Japanese.) Sericultural Experiment Station, Suigen, Korea.
- OGURA, S.-1922. Linkage of the Zebra and the Other Gene. Japanese Jour. Genetics. Vol.1. No.3. (In Japanese.)
- STURTEVANT, A. H.—1915. No Crossing Over in the Female of the Silkworm Moth. Amer. Nat. Vol.49, pp. 42-44.
- TANAKA, Y.-1913, a. A Study of Mendelian Factors in the Silkworm, Bombyx mori. Jour. Coll. Agric., Tohoku Imp. Univ. Vol. 5, pp. 61-113.
- ---- 1913 b. Gametic Coupling and Repulsion in the Silkworm, Bombyx mori. Ibid. pp. 115-148.
- ---- 1914. Further Data on Reduplication in Silkworms. Ibid. Vol.6, pp. 1-16.
- ----- 1915. Occurrence of Different Systems of Gametic Reduplication in Male and Female Hybrids. Zeits. ind. Abst. u. Vererb. Bd. 14, pp. 12-30.
- 1916. Genetic Studies on the Silkworm. Jour. Coll. Agric. Tohoku Imp. Univ. Vol. 7, pp. 129-255.
- --- 1921. Sex-Linkage in the Silkworm. Reports of Imp. Seric. Exp. Stat. Vol.6, No.1. (In Japanese.)
- ---- 1922. Sex-Linkage in the Silkworm. Jour. Genetics, Vol.12, No.2, pp. 163-172.
- ---- 1924. A New Sex-Linked Mutation in the Silkworm, Bombyx mori L. Jour. Dept. Agric. Kyushu Imp. Univ. Vol.1, No.2, pp. 135-150.
- _____ 1926. Somatic Mutation in Animals. Report of Japanese Assoc. for Advancem. Science for 1925. Vol.1, pp. 275-285. (In Japanese.)
- TANAKA, Y. and MATSUNO, S.-1927. "Stick" and "Multilunar," a Fourth Linkage Group in the Silkworm. Jour. Dept. Agric., Kyushu Imp. Univ. Vol.1, No.8, pp. 266-274