# Effect of Specific Ions in Agar on Antibiotic Production by Burkholderia glumae

Yamasaki, Shuichi Laboratory of Plant Pathology, Faculty of Agriculture, Kyushu University

Furuya, Naruto Laboratory of Plant Pathology, Faculty of Agriculture, Kyushu University

Matsuyama, Nobuaki Laboratory of Plant Pathology, Faculty of Agriculture, Kyushu University

https://doi.org/10.5109/24218

出版情報:九州大学大学院農学研究院紀要. 42 (3/4), pp.309-314, 1998-03. Kyushu University バージョン: 権利関係: J. Fac. Agr., Kyushu Univ., 43 (3.4), 309-314 (1998)

# Effect of Specific Ions in Agar on Antibiotic Production by *Burkholderia glumae*

# Shuichi Yamasaki\*, Naruto Furuya and Nobuaki Matsuyama

Laboratory of Plant Pathology, Faculty of Agriculture, Kyushu University 46-01, Fukuoka 812-8581, Japan (Received October 31, 1997 and accepted December 3, 1997)

Burkholderia glumae produced antibiotic substance(s) on agar plate media, but none in various liquid media. Some minor elements in agar were considered to be involved in the antibiotic production. K<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>3+</sup> and NH<sub>4</sub><sup>-</sup> were shown to play significant roles in the production by culturing the bacterium in liquid media supplemented with various inorganic chemicals. Highest production was observed when *B. glumae* was cultured in the liquid mediaum containing a mixture of MgCl<sub>2</sub>, KCl and CaCl<sub>2</sub>. No antibiotic production was observed in PGC broth containing MgSO<sub>4</sub>which was substituted for MgCl<sub>2</sub> or Mg(NO<sub>3</sub>)<sub>2</sub>. These results suggest that specific cations and anions are involved in antibiotic(s) production by *B. glumae*.

# INTRODUCTION

In previous papers (Wakimoto *et al.*, 1986; Furuya *et al.*, 1992a), it was shown that *Burkholderia glumae* has abilities to produce antibiotic substance(s) against various phytopathogenic bacteria including *Ralstonia solanacearum* and to suppress bacterial wilt of tomato under controlled conditions (Furuya *et al.*, 1991; 1992b). Attempts to obtain a large amount of the antibiotic substance failed because it was not produced in liquid media. Consequently, elements in agar were considered to be important factors in production of the antibiotic substance(s) (Yamasaki *et al.*, 1993; 1994). In this paper, qualitative and quantitative effects of some elements (cations and anions) on the antibiotic production were described.

# MATERIALS AND METHODS

## **Bacterial strains**

Bacterial strains used in this study were derived from the culture collection of the Laboratory of Plant Pathology, Faculty of Agriculture, Kyushu University. Seven strains of *B. glumae* were used as the producers of the antibiotics. *R. solanacearum* C319 was used as an indicator bacterium for the antibiotic substance(s).

# **Detection of antibiotic production**

*B. glumae* was cultured on YPDA (10g of yeast extract, 10g of peptone, 20g of Dglucose, 15g of agar, 1 liter of water, pH 7.0), PSA (2g of Na<sub>2</sub>HPO<sub>4</sub>  $\cdot$  12H<sub>2</sub>O, 0.5g of Ca(NO<sub>3</sub>)<sub>2</sub>  $\cdot$  4H<sub>2</sub>O, 5g of sucrose, 15g of agar, 1 liter of potato (300g) decoction, pH 7.0)

<sup>\*</sup> Present address: Oita Prefectural Agricultural Research Center, Oita 872-0103, Japan.

and TTC (10g of peptone, 5g of D-glucose, 1g of casein, 18g of agar, 50 ppm of 2,3,5triphenyltetrazorium chloride, 1 liter of water) media at 30 °C for 72 hr. After the bacteria were cultured on the media, the agar plates were frozen at -20 °C for longer than 2 hr and thawed at room temperature for 2hr. By these treatments, exudate was obtained from the culture plates. The exudate was sterilized by filtration through a Millipore filter ( $0.2 \mu$ m) and antibiotic activity was estimated by the penicillin-cup method.

To investigate the production of antibiotic substance in liquid media, *B. glumae* was cultured in YPD, PS and TTC liquid media at 30 °C for 6 days under still conditions. The cultured fluid was sterilized by filtration as described above and the presence of antibiotic was examined by the penicillin-cup method.

# Production of antibiotic substance(s) in the agar-extract amended (AePGC) broth

AePGC broth (10g of peptone, 5g of D-glucose, 1g of casein, 1 liter of agar-extract solution) was used to investigate the effect of agar-extract on the antibiotic production by *B. glumae*. Agar-extract solution was prepared by suspending a certain grams of agar powder (Katayama Chemical Co. Ltd.) in 1 liter of distilled water. Agar suspension was stirred overnight at 5°C, and filtered through a filter paper. Concentration of agar-extract solution was expressed as the amount (gram) of agar powder which was added to 1 liter of distilled water. *B. glumae* was cultured in AePGC broth at 30°C for 6 days under still conditions. Antibiotic production in the broth was examined by the penicillin-cup method.

#### Effects of minor elements in agar on antibiotic production

Since 30 g of agar powder was calculated to contain NaCl  $(3.2 \times 10^{\circ} \text{mM})$ , KCl  $(6.6 \times 10^{-2} \text{mM})$ , CaCl<sub>2</sub>  $(2.8 \times 10^{-1} \text{mM})$ , MgCl<sub>2</sub>  $(4.4 \times 10^{-1})$ , NH<sub>4</sub>Cl  $(6.2 \times 10^{-2} \text{mM})$  and Na<sub>3</sub>PO<sub>4</sub>  $(1.1 \times 10^{-1} \text{mM})$  according to Uzuka (1992), a liquid medium, PGC broth (10 g of peptone, 5 g of D-glucose, 1 g of casein, 1 liter of distilled water), was amended with each or mixtures of the inorganic chemicals so that the medium contained the same concentration of the respective elements. *B. glumae* was cultured in the amended PGC broth at 30 °C for 6 days under still conditions. The effect of elements on the antibiotic production was estimated by the penicillin-cup method.

# RESULTS

# Antibiotic production by B. glumae in various liquid media

The exudate from the culture plates showed anti-*R. solanacearum* activity, but the culture filtrates of YPD, PS and TTC liquid media did not show any anti-*R. solanacearum* activity (Table 1). As shown in Table 2, the antibiotic production was observed in AePGC liquid medium which contained the agar-extract from 5 g agar powder.

#### Effect of various elements in agar on antibiotic production by B. glumae

As can be seen in Table 3, the antibiotic production was observed in PGC broth supplemented with KCl,  $CaCl_2$ ,  $MgCl_2$  or  $NH_4Cl$ . Highest antibiotic production was recorded in PGC broth containing  $MgCl_2$ . However, the degree of production was lower than that in AePGC medium. A test was performed to examine cumulative effect of two or more kinds of elements on the production. The cumulative effect of cation on the

310

	Anti-Ralstonia solanacearum activity <sup>a</sup>						
<i>B. glumae</i> strain	shown by plate- chloroform method <sup>b)</sup>	of the exudate <sup>()</sup> from agar medium	of the culture filtrate <sup>d)</sup>				
N7401		~	_				
N7503	+	±	_				
805	+++	+	-				
N750	+++	+	-				
Ku8113	+	±	_				
Ku8117	_	~	-				
Ku8121	+++	+	-				

lable 1.	Production	of antibiotic	substance(s	) b	v Burkholderia	alumae
	1 I O G G G G G G G G G G G G G G G G G G	or anoiororo	oucounteers	10	1 19001 1010000001 000	grownood

a) Activity index (semidiameter of inhibition zone): -, not detected;  $\pm$ , faint activity; +, less than 5 mm; +++, greater than 10 mm.

b) *B. glumae* was spot-inoculated on YPDA, PSA and TTC plates and cultured at 30 °C for 3 days. Aftert treatment with chloroform vapor, an indicator bacterium, *R. solanacearum*, was overlaid.

c) YPDA, PSA and TTC media on which *B. glumae* was cultured at 30 °C for 3 days were frozen at -20 °C for above 2 hr and thawed at room temperature to obtain the exudate.

d) YPD, PS and TTC liquid media in which *B. glumae* was cultured at 30 °C for 6 days were filtered through Millipore filter  $(0.2 \mu m)$ .

	Width of inhibition zone <sup>a)</sup>					
B. glumae 🗍		Concentration of agar-extract <sup>b)</sup>			(g of agar/l of water)	
strain	0.0	1	5	15	30	50
N7401	_	NT	NT	NT	NT	_
N7503	-	NT	NT	NT	NT	+
805	-	-	+++	+++	NT	NT
N750	-	-	+++	+++	+++	+ + +
Ku8113	_	NT	NT	NT	NT	+
Ku8117	-	NT	NT	NT	NT	-
Ku8121	-	_	+++	+++	NT	NT

Table 2. Effect of agar-extract on the antibiotic production by Burkholderia glumae

a) Activity index (semidameter of inhibition zone): -, not detected; +, less than 5 mm; +++, greater than 10 mm; NT, not tested. *Ralstonia solanacearum* C319 was used as an indicator.

b) Concentration of agar-extract was expressed as the amount (gram) of agar powder which was added in 1 liter of distilled water.

#### S. Yamasaki et al.

production was remarkable when the mixture of  $MgCl_2$ , KCl and  $CaCl_2$  was added to PGC broth (Table 4). As shown in Table 5, the production was observed in PGC broth containing  $MgCl_2$  or  $Mg(NO_3)_2$ , but not  $MgSO_4$ .

	Cation			
R <sup>+b</sup>	R <sup>2+c)</sup>	R <sup>3+d)</sup>	Activity <sup>e)</sup>	pН
Na			0.0 h <sup>n</sup>	7.22 m
Κ			1.3 j	6.69 k
$NH_1$			0.9 hj	6.87 j
	Mg		3.7 k	6.27 m
	Ca		0.3 h	6.81 j
		Р	0.0 h	7.29 h
Na, K, NH₄	Mg, Ca	Р	5.4 m	6.33 m
	Cont. (AePGC)		5.6 m	6.24 m

 
 Table 3. Effect of various cations on the antibiotic production by Burkholderia glumae N750\*)

a) Cultured at 30 °C for 6 days. b) Monovalent cation. c) Divalent cation. d) Trivalent cation.

e) Semidiameter of inhibition zone (mm). Ralstonia solanacearum C319 was used as an indicator. f) Values within each column followed by the same letter are not significantly different (P = 0.05) according to Duncan's multiple range test.

Table 4.	Cumulative effects of various cations on the antibiotic production by Burkholderia	
	glumae N750 <sup>a)</sup>	

	Cation			
$\mathbf{R}^{+\mathrm{b}}$	R <sup>2 * c)</sup>	R <sup>3+d)</sup>	Activity <sup>e)</sup>	pН
K, NH4	Mg, Ca		6.5 h <sup>n</sup>	6.08 h
K, NH,	Mg		5.4 hj	6.25 jk
K	Mg, Ca		7.0 h	6.10 h
	Mg, Ca		4.4 j	6.21 j
К	Mg		5.5 hj	6.27 jk
$\rm NH_4$	Mg		1.6 k	6.31 k
K, NH₄			0.0 m	6.65 m
K	Ca		0.0 m	6.62 m
Na, K, NH4	Mg, Ca	Р	5.4 j	6.33 k
	Cont. (AePGC)		5.6 j	6.24 jk

a) Cultured at 30 °C for 6 days. b) Monovalent cation. c) Divalent cation. d) Trivalent cation. e) Semidiameter of inhibition zone (mm). *Ralstonia solanacearum* C319 was used as an indicator. f) Values within each column followed by the same letter are not significantly different (P = 0.05) according to Duncan's multiple range test.

	Anion			
R <sup>-b)</sup>	R <sup>2-c)</sup>	Activity	pН	
$MgCl_2$		3.7 h <sup>e)</sup>	6.27 h	
$Mg(NO_3)_2$		3.9 h	6.21 h	
	MgSO₄	0.0 j	6.35 h	

**Table 5.** Effect of various anions on the antibiotic production by Burkholderia glumae N750<sup>ai</sup>

a) Cultured at 30 °C for 6 days. b) Monovalent anion. c) Divalent anion. d) Semidiameter of inhibition zone (mm). *Ralstonia solanacearum* C319 was used as an indicator. e) Values within each column followed by the same letter are not significantly different (P = 0.05) according to Duncan's multiple range test.

#### DISCUSSION

Antibiotic production by *B. glumae* was observed in solid agar media, but not in liquid media. This phenomenon indicated an important role of element(s) in agar for the production (Furuya *et al*, 1992a). Since the existence of six ions (Na<sup>+</sup>, K<sup>+</sup>, NH<sub>4</sub><sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup> and P<sup>3+</sup>) in agar was formerly documented (Uzuka, 1992), the effect of these minor elements on the production was examined. Among these cations tested, K<sup>+</sup>, NH<sub>4</sub><sup>+</sup>, Mg<sup>2+</sup> and Ca<sup>2+</sup> were revealed to participate in the antibiotic production. The effect of Mg<sup>2+</sup> was particularly prominent. Moreover, highest production was recorded when the bacterium was cultured in PGC liquid medium containing the mixture of MgCl<sub>2</sub>, KCl and CaCl<sub>2</sub>. Thus, in liquid culture conditions, the presence of specific cations was necessary for the production of the antibacterial substance(s). The antibiotic production was observed in PGC broth containing MgCl<sub>2</sub> or Mg(NO<sub>3</sub>)<sub>2</sub>, but not in the broth supplemented with MgSO<sub>4</sub>, suggesting that anions also play an important role(s) for the production.

Like the case of *B. glumae*,  $Mg^{2+}$  was shown to affect magnesidin production by *P. magnesiorubra* nov. sp. ATCC21856 (Gandhi *et al.*, 1973) and *Vibrio gazogenes* ATCC29988 (Imamura *et al.*, 1994). The effect of minor element, ion, has also been reported in pyoveridine production by *P. fluorescens* (Loper and Lindow, 1994).

The mechanisms of the promotive effects on antibiotic production by amendment of the minor nutrients in agar have not been elucidated in the present study. This will be an important subject for future study.

# REFERENCES

- Furuya, N., Y. Kushima, K. Tsuchiya, N. Matsuyama and S. Wakimoto 1991 Protection of tomato seedlings by pre-treatment with *Pseudomonas glumae* from infection with *Pseudomonas solanacearum* and its mechanisms. Ann. Phytopathol. Soc. Jpn., 57: 363-370
- Furuya, N., Y. Kushima and N. Matsuyama 1992a Production of antibacterial substances by Pseudomonas glumae. J. Fac. Agr., Kyushu Univ., 37: 149-158
- Furuya, N. and N. Matsuyama 1992b Biological control of the bacterial wilt of tomato with antibioticproducing strains of *Pseudomonas glumae*. J. Fac. Agr., Kyushu Univ., **37**: 159-171
- Gandhi, M., J. Nazareth, P. Divekar, H. Kohl and N. De Souza 1973 Magnesidin, a novel magnesiumcontaining antibiotic. J. Antibiotics, **26**: 797-798

S. Yamasaki et al.

- Imamura, N., K. Asachi and H. Sano 1994 Magnesidin A, a component of marine antibiotic magnesidin, produced by *Vibrio gazogenes* ATCC 29988. *J. Antibiotics*, **47**: 257-261
- Loper, J.E. and S.E. Lindow 1994 A biological sensor for iron available to bacteria in their habitats on plant surfaces. *Appl. Environ. Microbiol.*, **60**: 1934-1941
- Uzuka, Y. 1992 Yeasts as oligotrophic microorganisms- a new concept derived from soil yeast physiology. *Kagaku To Seibutsu*, **30**: 431-440
- Wakimoto, S., K. Hirayae, K. Tsuchiya, Y. Kushima, N. Furuya and N. Matsuyama 1986 Production of antibiotics by plant pathogenic pseudomonads. Ann. Phytopathol. Soc. Jpn., 52: 835-842
- Yamasaki, S., N. Furuya, Y. Kushima, Y. Takanami and N. Matsuyama 1993 The antibiotics produced by *Pseudomonas glumae* (1). Conditions for antibiotic production (Abstr.). Ann. Phytopathol. Soc. Jpn., 59: 721-722
- Yamasaki, S., N. Furuya, Y. Kushima, Y. Takanami and N. Matsuyama 1994 The antibiotics produced by *Pseudomonas glumae* (2). Conditions for antibiotic production (2) (Abstr.). Ann. Phytopathol. Soc. Jpn., **60**: 370-371