Critical involvement of extracellular ATP acting on P2RX7 purinergic receptors in photoreceptor cell death

Notomi, Shoji Department of Ophthalmology, Graduate School of Medical Sciences, Kyushu University

Hisatomi, Toshio Department of Ophthalmology, National Hospital Organization Kyushu Medical Center | Department of Ophthalmology, Graduate School of Medical Sciences, Kyushu University

Kanemaru, Takaaki Morphology Core Unit, Graduate School of Medical Sciences, Kyushu University

Takeda, Atsunobu Department of Ophthalmology, Graduate School of Medical Sciences, Kyushu University

他

https://hdl.handle.net/2324/25676

出版情報:The American Journal of Pathology. 179 (6), pp.2798-2809, 2011-12. Elsevier バージョン: 権利関係:(C) 2011 American Society for Investigative Pathology

Critical involvement of extracellular ATP acting on P2RX7 purinergic receptors

in photoreceptor cell death

Shoji Notomi¹, Toshio Hisatomi^{1, 2}, Takaaki Kanemaru³, Atsunobu Takeda¹, Yasuhiro Ikeda¹, Hiroshi Enaida¹, Guido Kroemer⁴⁻⁸, and Tatsuro Ishibashi¹.

¹Department of Ophthalmology and ³Morphorogy Core Unit, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan, ²Department of Ophthalmology, National Hospital Organization Kyushu Medical Center, Fukuoka, Japan, ⁴INSERM, U848, F-94805 Villejuif, France; ⁵Metabolomics Platform, Institut Gustave Roussy, F-94805 Villejuif, France; ⁶Centre de Recherche des Cordeliers, F-75005 Paris, France; ⁷Pôle de Biologie, Hôpital Européen Georges Pompidou, AP-HP, F-75908 Paris, France; ⁸Université Paris Descartes, Paris 5, F-75270 Paris, France.

Manuscript information: the number of text pages: 46, of figures: 7, of table: 0.Short running head: ATP and P2RX7 receptors in photoreceptor deathThis study was supported by grants from the Ministry of Education, Science, Sports

and Culture, Japan (Grant-in-Aid for Scientific Research nos.21791690)

Correspondence to: Tatsuro Ishibashi, MD, PhD, Department of Ophthalmology, Graduate school of Medical Sciences, Kyushu University, 3-1-1 Maidashi Higashi-ku, Fukuoka, 812-8582, Japan. Phone: +81-92-642-5648; Fax: +81-92-642-5663, e-mail: ishi@eye.med.kyushu-u.ac.jp; or Guido Kroemer, MD, PhD, INSERM, U848, Institut Gustave Roussy, Pavillon de Recherche 1, 39 rue Camille-Desmoulins, F-94805 Villejuif, France. Phone: +33 1 4211 6046; Fax: +33 1 4211 6047; e-mail: kroemer@orange.fr

ABSTRACT

ATP can be released from stressed cells and participate in neurodegenerative processes through the specific ligation of P2RX7 purinergic receptors. Here we demonstrate that extracellular ATP and the more specific P2RX7 agonist, 2'- and

3'-O-(4-benzoylbenzoyl)-ATP (BzATP), both induce photoreceptor cell death when added to primary retinal cell cultures or when injected into the eyes from wild type mice, but not those from $P2RX7^{-/-}$ mice. The photoreceptor cell death was accompanied by the activation of caspase-8 and -9, translocation of apoptosis inducing factor (AIF) from mitochondria to nuclei, and TUNEL-detectable chromatin fragmentation. All hallmarks of photoreceptor apoptosis were prevented by pre-medication or co-application of brilliant blue G (BBG), a selective P2RX7 antagonist that is already approved for the staining of internal limiting membranes during ocular surgery. ATP release is upregulated by nutrient-starvation in primary retinal cell cultures and seems to be an initializing event that triggers primary and/or secondary cell death via the positive feedback loop on P2RX7. Our results encourage the potential application of BBG as a novel neuroprotective agent in retinal diseases, or similar neurodegenerative pathologies linked to excessive extracellular ATP.

Introduction

Photoreceptor degeneration involves the activation of several pathways of regulated cell death that may constitute potential therapeutic targets. Accordingly, attempts have been undertaken to inhibit caspases, which play a central role in the acquisition of the apoptotic morphology,^{1, 2} although pharmacological pan-caspase inhibitors have largely failed to preserve the structure and function of photoreceptors.^{3,4} Caspases can be activated as the result of mitochondrial outer membrane permeabilization (MOMP) in thus far that the mitochondrial release of cytochrome c results in the Apaf-1 apoptosome-dependent activation of caspase-9. MOMP also results in the mitochondrial release of apoptosis-inducing factor (AIF), which then translocates to the nucleus and participates in caspase-independent peripheral chromatin condensation and large-scale DNA fragmentation,⁵ suggesting the existence of redundant cell death mechanisms downstream of MOMP.^{4, 6, 7} Pharmacological inhibition of MOMP has indeed been shown to confer some degree of neuroprotection in a model of photoreceptor degeneration induced by retinal detachment.⁴

Upstream of or independently from MOMP, death receptors from the tumor necrosis factor receptor (TNF-R) family proteins including Fas receptor (Apo-1/CD95) or DR4

(death receptor 4) can transmit apoptotic and/or necrotic signals initiated by specific "death ligands".⁸ Increased TNF-R or CD95 signaling can participate in photoreceptor death induced by retinal detachment (RD),^{9, 10} in which case these receptors may either stimulate the initiation of apoptosis (via the activation of caspase-8) or programmed necrosis (via the activation of RIP1 kinase). Indeed, simultaneous inhibition of caspases (with Z-VAD-fmk) and RIP1 kinase (with necrostatin 1) has a more pronounced neuroprotective effect on photoreceptors in RD than either treatment alone.¹¹ Taken together, these results underscore the existence of multiple cell death mechanisms that have to be inhibited simultaneously to confer optimal neuroprotection.

An alternative to the inhibition of executioner pathways consists in intercepting the initiating events that account for upstream damage signals. Recently, adenosine-5'-triphosphate (ATP) has been discovered as a major extracellular messenger that can contribute to lethal signaling.¹² ATP, which can be released via exocytosis, anion channels or transporters, can act in autocrine and paracrine signaling pathways,¹³ for instance by evoking action potentials in brain slices, isolated nerves and glial cells,¹⁴ indicating that ATP acts as a physiological mediator of neurotransmission and neuron-glia communication.¹⁵ Moreover, extracellular ATP

concentrations increase in spinal cord injury,¹⁶ chronic neuropathic pain,¹⁷ and brain ischemia,¹⁸ suggesting a pathological role for extracellular ATP as well. Extracellular ATP can act on purinergic receptors, which are classified into two classes, the ionotropic, ligand-gated P2X receptors and the metabotropic, G protein-coupled P2Y receptors.¹⁹ Among the seven mammalian P2X receptors,²⁰ the P2X7 receptor (official name: P2RX7) has the highest affinity for ATP. P2RX7 differs from other P2X receptor subtypes by its long cytoplasmic, carboxy-terminal tail (240 amino acids). Brief application of agonists renders P2RX7 permeable to small cations (K^+ , Na^+ , Ca^{2+}) similar to other P2X receptors,¹² while repeated or prolonged exposure to agonists can lead to the formation of P2RX7-receptor-dependent pores that become permeable to solutes up to 900 daltons, hence triggering cell death.²¹ Thus, extracellular ATP can induce apoptotic and/or necrotic cell death by acting on P2RX7.²² Despite the great interest in the carboxyl-terminal region of P2RX7, the mechanism by which P2RX7 mediates apoptotic signaling is largely unknown. Of note, the residues 436-531 of P2RX7 are similar to a region of TNFR1 that overlaps its death domain,²³ which has led to the proposal of a potential mechanism for P2RX7-induced caspase-8 activity and apoptosis.²⁴ Thus, the plasma membrane receptor, P2RX7, may cleave and activate caspase-8 during extrinsic apoptotic pathway.

Importantly, P2RX7 is widely expressed in various organs including the immune system (thymus or spleen)²¹ and the central nervous system (CNS) (cortex, hippocampus,²⁵ and spinal cord²⁶). P2RX7 is expressed on astrocytes, microglial cells and neurons.¹⁴ In the retina, P2RX7 is expressed on Müller glia,²⁷ and in both inner and outer retinal neurons, including retinal ganglion cells^{28, 29} and photoreceptors.³⁰ Genetic or functional inactivation of P2RX7 can attenuate the development of several neurodegenerative diseases, including Alzheimer's disease³¹ and Huntington's disease.³² Thus, systemic administration of a pharmacological P2RX7 antagonist, brilliant blue G (BBG) can confer neuroprotective effects in models of Alzheimer's, Parkinson's disease, and spinal cord injury.³¹⁻³³

BBG is a triphenylmethane dye that has been approved for intraoperative use in ocular surgery, in the context of chromovitrectomy, which involves the use of vital dyes to improve the visualization of intraocular tissues during vitrectomy, thereby improving specific procedures such as internal limiting membrane (ILM) peeling.³⁴ Driven by the recent characterization of BBG as a P2RX7 antagonist,³¹⁻³³ we decided to investigate the pathogenic implications of P2RX7 in pathological photoreceptor loss, as well as the therapeutic utility of BBG in this context. As a result of these investigations, we report here that increased extracellular ATP levels contribute to pathological conditions

of photoreceptor loss and that BBG efficiently avoids photoreceptor cell death.

Materials and methods

Animals

All animal experiments were performed according to the guidelines of the Association for Research in Vision and Ophthalmology on adult (8 weeks of age) male C57BL6JJcl mice (CLEA, Tokyo, Japan) and isogenic P2RX7^{-/-} mice (B6.129P2-*P2rx7^{tm1Gab}*/J; stock number 005576, The Jackson Laboratory, Bar Harbor, ME).

Adult mouse primary retinal cell cultures

Adult primary retinal cell cultures were prepared as previously described with minor modifications.⁶ Primary retinal cells were cultured in 4-well chamber (NUNC; part of Thermo Fisher Scientific, Rochester, NY) with Neurobasal-A medium (Invitrogen, Carlsbad, CA) containing B27 supplement without antioxidants (NBA/B27AO–; Invitrogen), 1 μ g/ml insulin, and 12 μ g/ml gentamicin. To determine the number of adherent photoreceptor cells, immunofluorescent staining was performed with a rabbit anti-recoverin antibody (Millipore, Bedford, MA).

For nutrient-starvation, primary retinal cells were cultured for 3 hours (ATP

measurement) or 24 hours (immunocytochemistry and viability assay) with Neurobasal A medium without B27 supplement as 'starvation medium'. P2RX7 agonists (ATP or BzATP; Sigma-Aldrich, Saint Louis, MO) or P2RX7 antagonists (KN-62 or BBG; Sigma-Aldrich) were added to the medium and incubated for 24 hours. To investigate the effect of BBG pre-incubation, cells were cultured in the presence of BBG for 30 min, washed two times with culture medium, and incubated for 24hours. To examine the secondary effects of death ligands, rat anti-mouse TNF-alpha neutralizing antibody (1-10 ng/ml MP6-XT22; R&D systems, Minneapolis, MN)³⁵ or hamster anti-mouse CD95 ligand neutralizing antibody (1-10µg/ml MFL3; BD biosciences, San Jose, CA)³⁶ were added to culture medium, while isogenic rat or hamster IgG was added as control.

Viability assay in primary retinal cell cultures

To assess the viability of primary retinal cells, we used calcein AM (2 μ M; Invitrogen) or MitoTracker Orange CMTMRos (200 nM, M7510; Invitrogen) that were added for 34 min to primary retinal cell cultures. Then, cultured cells were fixed with 4% paraformaldehyde and photoreceptors were labeled with recoverin. Calcein⁺ or CMTMRos⁺ photoreceptors were counted in 10 random fields by blinded observers

using ImageJ software. Values are given as means \pm SD of 10 replicate wells.

Intraocular injections

For evaluation of the toxicity of BzATP and the effect of BBG in vivo, we took advantages of gas compression vitrectomy for uniform diffusion of intravitreally injected solutions. Mouse eyes were vitrectomized using SF6 gas as we and others previously described.³⁷⁻⁴⁰ To investigate the effects of gas compression vitrectomy on the viability of retinal cells, we observed ophthalmic examination every day up to 2 weeks and evaluated histochemical changes in hematoxylin eosin staining and TUNEL positive cells in the retina at 3 and 14 days after gas injections. Two weeks after gas injection, mice were anesthetized with an intraperitoneal injection of pentobarbital, and their pupils were dilated with topical 1% tropicamide and 2.5% phenylephrine hydrochloride. Then, 2 µl of liquid (PBS supplemented with 10-20 mM BzATP or BzATP plus 500 µM BBG) was injected into the vitreous cavity with a 32-gauge needle on a Hamilton syringe through posterior to the limbus. Intraocular injections were only performed in the right eye of each animal, and 5 eyes were examined in each group. The mice were sacrificed 24 hours after treatment, and their eyes were harvested, frozen at nitrogen liquid temperature and cryosectioned for histochemical or

ultrastructural examinations.

TUNEL

TUNEL analysis and quantification of TUNEL-positive cells were performed as previously described⁴ using the ApopTag Fluorescein In Situ Apoptosis Detection Kit (Millipore). Nuclei were counter-stained with propidium iodide or Hoechst 33342. TUNEL⁺ cells in the ONL were counted by two blinded observers, and results were presented as means \pm SD.

Immunohistochemistry

As previously reported,⁶ rabbit anti-AIF (R&D Systems, Minneapolis, MN), anti-mouse cleaved caspase-9 (Cell Signaling Technology, Beverly, MA), anti-mouse cleaved caspase-8 (Cell Signaling Technology) and anti-recoverin (Millipore) were used as primary antibodies and incubated at 4°C overnight. Goat anti-rabbit IgG conjugated to Alexa Fluor 546 or 647 (Invitrogen) were used as secondary antibodies and incubated at room temperature for 1 hour.

Electron Microscopy

The posterior segments of enucleated eyes were fixed in PBS containing 1% glutaraldehyde and 1% paraformaldehyde, postfixed in veronal acetate buffer osmium tetroxide (2%), dehydrated in ethanol and water, and embedded in Epon. Primary retinal cell cultures were similarly fixed, dehydrated, and embedded in Epon. Ultrathin sections were cut from blocks and mounted on copper grids. The specimens were observed with H-7650 transmission electron microscope (Hitachi, Tokyo, Japan).

Calcium imaging

Primary retinal cells were cultured on 3.5mm coverslip dishes (BD Biosciences, San Jose, CA) in Neurobasal A medium with B27 supplement. Then the calcium indicator, Fluo-4 AM (5µM; Invitrogen) was loaded onto primary cultures. The calcium-dependent fluorescence of Fluo-4 was captured by microscopy during 20 minutes incubation after addition of 1mM BzATP or vehicle PBS. To evaluate the effect of chelation of extracellular Ca²⁺ or P2RX7 antagonist (BBG), the fluorescence of Fluo-4 was measured in culture medium containing 1mM EGTA (Sigma-Aldrich) or 10 µM BBG, respectively.

ATP measurements

For ATP measurements of culture medium, primary retinal cells were starved for 3 hours with Neurobasal A medium without B27 supplement as 'starvation cultures' and cultures with Neurobasal A medium with B27 supplement were prepared as controls. Then 100µl culture medium were collected, centrifuged at 4°C, and subjected to ATP measurements as follows. The ATP levels of collected medium (100 µl in 96 well microplates; BD Biosciences) were immediately determined by adding luciferin-luciferase reaction buffer (ATP bioluminescent assay kit, FL-AA; Sigma-Aldrich) and a multi-mode microplate reader, Flex Station 3 (Molecular Devices, Sunnyvale, CA). The ATP levels in medium of 'control cultures' were determined by calibration of standard ATP in Neurobasal A medium with/without B27 supplement. To define the detection limit, ecto-nucleotidase (Apyrase; 10 unit /ml, Sigma-Aldrich) was added to culture medium. In control experiments, β , γ -methylene-ATP, a potent ecto-ATPase inhibitor (300 µM, Sigma-Aldrich), was added to control or starvation cultures. Values are given as mean \pm SD of 10 replicate wells.

Imaging of ATP release

Primary retinal cells were cultured on 3.5 mm coverslip dishes (BD Biosciences) in Neurobasal A medium with B27 supplement. ATP assay mix (a part of the FL-AA kit;

Sigma-Aldrich) was added to culture media (one vial per 1.5 ml culture medium with/without B27 supplement). Light production from the luciferin-luciferase reaction was captured by an electron multiplier charge-coupled camera (ImagEM; Hamamatsu Photonics, Shizuoka, Japan) with a 40 x oil lens (N.A. 1.35; Olympus, Tokyo, Japan) for 3 hours during 5 min exposure periods.

Statistical analysis

Statistical differences between two groups were analyzed by means of the Mann–Whitney U test. Multiple group comparison was performed by ANOVA followed by Tukey–Kramer adjustments. Differences were considered significance at P < 0.05 (*) and P < 0.01 (**). All values were expressed as means ± SD.

Results

P2RX7 mediated photoreceptor death in primary retinal cell cultures

To investigate the possible implication of P2RX7 in photoreceptor degeneration, we took advantage of primary retinal cell cultures.^{6, 41} Photoreceptor viability was assessed by means of two fluorescent sensors, calcein AM and MitoTracker CMTMRos, which only label intact and non-apoptotic, metabolically active cells, respectively. Calcein AM, which is non-fluorescent and cell permeable, becomes fluorescent and is trapped in live cells after removal of the AM moiety by cellular esterases. Cells that stain with this dye hence must possess an intact, impermeable plasma membrane. The lipophilic cation MitoTracker CMTMRos labels mitochondria, driven by the mitochondrial transmembrane potential (DY_m), and hence stains cells with intact mitochondrial membranes. Addition of ATP (Figure 1A) or 2'-and 3'-O-(4-benzoyl-benzoyl)-ATP (BzATP) (Figure 1B), a potent, more selective P2RX7 antagonist,²¹ resulted in a dose-dependent decline of viable, calcein⁺ or CMTMRos⁺ photoreceptors that were identified by immunofluorescence detection of recoverin. Before starting blocking experiments by BBG, we tested whether BBG has potential toxicity. Primary retinal cell cultures were incubated in the presence or absence of 1-10 µM BBG for 24 h. No

significant decline of photoreceptor viability was observed (Figure 2A). BBG showed minimal toxicity at those concentrations *in vitro*. BBG, a potent P2RX7 antagonist⁴² as well as another selective P2RX7 inhibitor,

1-[N,O-bis(5-isoquinolinesulphonyl)-N-methyl-L-tyrosyl]-4-phenylpiperazine (KN-62), reduced BzATP-induced photoreceptor death (Figure 2B). BBG is a non-competitive and slowly reversible P2RX7 inhibitor that antagonizes P2RX7 by allosteric regulation.⁴² Accordingly, a short (30 min) pre-incubation with BBG could reduce BzATP-induced photoreceptor death to some extent, though less efficiently than continuous co-incubation (Figure 2B). Of note, photoreceptors isolated from P2RX7^{-/-} mice readily survived exposure to high dose (1 mM) BzATP, under conditions in which large numbers of wild type photoreceptors died, underscoring that these effects are truly specific (Figure 2C).

These results indicated the specific effect of P2RX7 on photoreceptor death. However, we needed to exclude the secondary effect of death ligands, which could potentially be released by other cell types, on photoreceptor death. To this end, we performed the blocking experiment by neutralizing antibody against TNF-alpha and CD95 ligand. Of note, neither TNF-alpha (1-10 ng/ml MP6-XT22)³⁵ nor CD95 ligand (1-10 μ g/ml MFL3)³⁶ neutralizing antibody could prevent the photoreceptor death induced by 1mM

BzATP (Figure 2, D and E). These results suggest that P2RX7 stimulation could directly mediate the cell death pathway on photoreceptors, rather than an indirect pathway being mediated via TNF-alpha or CD95 ligands released by other cells.

P2RX7 induced retinal cell death via calcium influx and caspase-8 activation

Transmission electron microscopy revealed the relatively well-preserved organelles in the photoreceptors without BzATP administration, while P2RX7 stimulation by BzATP resulted in apoptotic cell death, characterized by chromatin condensation, as well as in some instances in necrotic cell death (Figure 3A). After culture in the presence of 1 mM BzATP, a significant percentage of photoreceptors exhibited a type of apoptotic DNA fragmentation that could be detected with the TUNEL. To examine caspase-8 activation by P2RX7, TUNEL-positive cells also were stained by immunofluorescence with an antibody that only recognizes active, proteolytically mature caspase-8 (Figure 3B). BBG largely prevented these BzATP-induced signs of the extrinsic cell death pathway (Figure 3, B and C).

As to be expected for an ionotropic receptor, BzATP also stimulated a rapid (20 min) increase in intracellular Ca^{2+} concentrations determined by means of Fluo-4 AM, a Ca^{2+} indicator that was loaded into primary cultures (Figure 3D; top). We performed

the extracellular Ca^{2+} chelating with EGTA to clarify the origin of the Ca^{2+} increase. This increase of intracellular Ca^{2+} was attenuated by addition of 1 mM EGTA into the culture medium (Figure 3D; middle), indicating that BzATP mediated Ca^{2+} influx mainly through the cation channel, not via release from intracellular binding proteins. Furthermore, co-incubation of 10 μ M BBG inhibited the Ca^{2+} influx by BzATP (Figure 3D; bottom), suggesting that BBG prevents intracellular calcium signaling by blocking the P2RX7 ion channel.

In conclusion, P2RX7 ligation kills photoreceptors via the influx of Ca^{2+} and caspase-8 activation, suggesting a potential mechanism for the P2RX7-mediated extrinsic pathway in line with previous reports.^{23, 24}

Photoreceptor apoptosis induced by intraocular administration of P2RX7 agonists *in vivo*

To determine whether P2RX7 ligation can trigger retinal cell death *in vivo*, we measured retinal cell apoptosis 24 h after intraocular injection of 20 mM BzATP into the eyes of C57BL/6 mice. The BzATP injections were performed 14 days after gas compression vitrectomy to obtain uniform diffusion of the injected solutions. Neither ophthalmic abnormality (i.e. corneal edema) nor signs of retinal cell loss in

hematoxylin eosin staining were observed up to 14 days after gas injection (data not shown), and no TUNEL-positive cells were observed in the retina at 3 and 14 days after injections (Figure 4A). The intraocular injection of BzATP induced TUNEL⁺ apoptotic events in the ganglion cell layer, which is notoriously susceptible to induction of apoptosis by stimulation of P2RX7 receptors,^{29, 43, 44} as well as in the inner nuclear layer (INL) and outer nuclear layer (ONL). These TUNEL-detectable DNA degradations accompanied the caspase-8 cleavage (Figure 4B, arrows). In vivo apoptosis induction by BzATP was blunted by co-injection of 500 µM BBG (Figure 4, C and D). As in the in vitro system, even high doses of BzATP (10-20 mM) largely failed to induce apoptosis when injected into P2RX7^{-/-} mice (Figure 4, C and D). Transmission electron microscopy confirmed that BzATP induced the death of photoreceptors with hallmarks of apoptosis, such as cell shrinkage and chromatin condensation in the ONL. Moreover, BzATP caused the degeneration of pedicles and spherules in the OPL. The frequency of apoptotic events and degenerative changes in rod spherules and cone pedicles was decreased when BzATP was combined with BBG (Figure 4E).

Taken together, these results confirmed that pharmacological P2RX7 agonists have the potential to induce retinal cell death *in vivo*, under conditions in which P2RX7 antagonists can preserve photoreceptors.

BBG attenuates starvation-induced photoreceptor death in primary retinal cell cultures

Primary retinal cells cultured under starvation conditions (Neurobasal A medium without B27 supplement, a serum substitute) exhibited a significantly elevated level of photoreceptor apoptosis as compared to control cells (cultured in complete, B27-supplemented Neurobasal A medium). AIF was detected in the cytoplasm of control cultures, yet translocated into nuclei containing fragmented DNA upon starvation in agreement with our previous report that starvation induced the mitochondrial cell death pathway in photoreceptors.⁶ BBG treatment attenuated the starvation-induced AIF translocation and TUNEL staining (Figure 5, A and C). The proteolytic maturation of caspase-9 was rarely detected in control cells, yet increased upon starvation, an effect that was again reduced by BBG (Figure 5, B and C). Similarly, a 24 h-starvation period led to a decrease in the frequency of viable (calcein⁺) and metabolically active (CMTMRos⁺) photoreceptors to approximately half

of that found in control cultures. This decline was again reversed by 30 min pre-incubation with 0.1-10 μ M BBG in a dose-dependent manner (Figure 5D). Moreover, P2RX7^{-/-} photoreceptors showed a significantly greater survival rate (p<0.05) under starvation conditions (ratio of calcein⁺ cells over controls= 0.73; ratio of CMTMRos⁺ cells over controls=0.64) compared to WT photoreceptors (ratio for calcein⁺ cells=0.52; ratio for CMTMRos⁺ cells=0.50).

Taken together, these results indicate that P2RX7-blockade inhibits the caspase-dependent and caspase-independent arms of the mitochondrial cell death pathway activated by starvation.

Extracellular ATP in stressed retinal cell cultures

Retinal cell cultures spontaneously released ATP (0.22±0.06 nM) under the control conditions. This release increased significantly (p<0.01) to 0.38±0.07 nM upon starvation, as quantified by a luciferase-based commercial assay (Figure 6). This difference persisted in the presence of β , γ -methylene-ATP (β , γ -Me-ATP), a potent inhibitor of ecto-ATPases⁴⁵ that degrade extracellular ATP, suggesting that the increase in ATP concentrations induced by starvation results from ATP release rather than from reduced ATP degradation. Of note, addition of BBG reduced the starvation-induced

ATP release (Figure 6), suggesting the existence of a positive feedback loop that links activation of P2RX7 receptors to nucleotide release.⁴⁶ For a more direct investigation of ATP release, we monitored retinal cell cultures by videomicroscopy to detect ATP-dependent bioluminescence revealed by a luciferin-luciferase assay. The frequency of ATP-releasing cells was low in control cultures, increased upon starvation, and was again decreased when starved cells were pre-incubated with BBG (Figure 7). Thus, BBG inhibits ATP released from starved retinal cells.

Discussion

In the present work, we provide compelling evidence that extracellular ATP and activation of P2RX7 may contribute to the pathological loss of photoreceptors. Thus, addition of P2RX7 agonists to primary retinal cultures or their intraocular injection causes the apoptotic death of photoreceptors in a fashion that is strictly dependent on P2RX7, as demonstrated by the use of P2RX7^{-/-} mice. This death is accompanied by hallmarks of apoptosis, including the activation of caspases, the mitochondrio-nuclear translocation of AIF, and TUNEL-detectable chromatin fragmentation. Finally, we demonstrate that a pharmacological P2RX7 antagonist, BBG, can prevent all stigmata of pathological loss of photoreceptors by inhibiting extracellular ATP acting on P2RX7.

BBG is reported to be a selective antagonist of P2RX7.⁴² In the current study, BBG substantially inhibited the effects of the selective P2RX7 agonist BzATP, namely, cell death, caspase-8 activation, and calcium influx. To clarify the specificity of the BBG, we examined another known selective P2RX7 blocker, KN-62. KN-62 showed an effect similar to BBG. Furthermore, the examination with P2RX7 knockout mice

revealed the specificity of BBG *in vitro* and *in vivo*. Thus, we concluded that BBG selectively blocked P2RX7 activation, thereby contributing a neuroprotective effect. Moreover, the blocking experiment with neutralizing antibodies for TNF-alpha or CD95 ligand revealed that P2RX7 could directly induce cell death on photoreceptors rather than indirectly via death ligands released by other cells.

ATP release appears to be (one of) the initializing event(s) that drives retinal cell death by nutrient starvation, a pathological setting *in vitro*. Furthermore, these extracellular ATP released from dying cells may worsen the pathology by promoting the death of neighboring cells, namely, secondary cell death. The following potential mechanisms may account for the increase in extracellular ATP in ocular pathologies. First, acute cell lysis may constitute one source of ATP, in line with the observations that ATP is released by acute stresses such as ischemia,¹⁸ hypotony,⁴⁷ or oxygen/glucose deprivation.⁴⁸ However, here we observed ATP release from retinal cells in a model of nutrient starvation well before the occurrence of cell death,^{6, 41} suggesting that ATP release occurred through a more controlled mechanism than acute cell lysis. Second, massive ATP release may occur from the usurpation of a physiological mechanisms of release (when ATP serves as a neurotransmitter), perhaps as the result of a pathogenic

feed-forward loop. Indeed, it has been shown that extracellular ATP can stimulate further ATP release through the opening of P2RX7 channels,⁴⁶ and – as shown here – BBG can reduce ATP release from starved retinal cultures by blocking the positive feedback.

The doses of ATP required to induce photoreceptor cell death are higher than those measured in starved cultures (which are rescued by addition of BBG or knockout of P2RX7). There are three possible explanations for this discrepancy. First, ATP released from cells may be diffused into bulk culture medium and continuously degraded through the action of ecto-ATPases, which are very active both in cultures⁴⁵ and *in* vivo⁴⁹ in the CNS. Indeed, the local concentration of ATP acting on the plasma membrane exceeds micromolar level, as detected using membrane-bound protein A-luciferase as a spatially localized probe of ATP levels at the extracellular cell surface.⁵⁰ Hence, the local increase of ATP in the intercellular space may activate P2RX7 in retinal disorders. Second, the injected solutions diffuse into the intraocular cavity and can be washed out by aqueous humor circulation in vivo, thereby lowering the local concentration in the retina. Third, the function of P2RX7 may be modulated not only by extracellular ATP but also by local receptor modifications. Nagasawa et al. reported that P2RX7 channel opening may occur without exogenous ATP

administration in cultured brain astrocytes, suggesting P2RX7 activation by endogenous ATP.⁵¹ Furthermore, P2RX7 may also be sensitized to ATP following ADP ribosylation at the arginine 125 position of P2RX7.⁵² These results may provide additional alternative pathways for the activation of P2RX7 under pathological conditions. Further investigations will be needed to elucidate the exact details of P2RX7 activation with respect to fluctuating local ATP concentrations.

In this study, we have shown that BBG may have important neuroprotective effects on the retina. Of note, BBG is already approved for intraocular use in patients, for the surgical procedure of chromovitrectomy. This procedure initially involved the administration of indocyanine green (ICG) as an intraocular surgical adjuvant in 2000.^{53, 54} However, upon our initial description that ICG has toxic side effects in 2002,³⁹ the following clinical and experimental studies reported that ICG induces visual field defects or retinal pigment epithelial (RPE) atrophy.⁵⁵⁻⁵⁷ Brilliant Blue G (BBG) has been introduced as an alternative staining dye for capsular staining in cataract surgery and chromovitrectomy with satisfactory biocompatibility.^{40, 58, 59} Moreover, whereas ICG exposure turned out to trigger apoptotic signaling with caspase activation, BBG has minimal cytotoxicity in retinal cells.⁵⁷ In the present study,

we further confirmed the minimal toxicity of BBG by means of primary culture. Recent reports have supported the safety of BBG as a surgical adjuvant,^{60, 61, 62} encouraging the ever expanding clinical applications for BBG in vitreo-retinal surgery. It should be noted that a substantial neuroprotective effect of BBG could be obtained at a dose of $\leq 10 \,\mu$ M, which is considerably lower than the concentration recommended for use during chromovitrectomy (293 μ M). Moreover, the neuroprotective effect of BBG could be obtained through transient exposures of primary cultures, probably due to the characteristics as a slowly reversible antagonist of P2RX7. Considering these results, one might expect BBG to have a broad, long-lasting neuroprotective effect in ocular pathologies. Taken together, our study clarifies (part of) the mechanisms through which BBG exerts its therapeutic effects in clinically relevant models of neurodegenerative diseases,³¹⁻³³ and thereby lend support to the potential application of BBG as a novel neuroprotective agent.

Acknowledgement

We thank Mari Imamura and Fumiyo Morikawa (Kyushu University) for technical assistance. We acknowledge Olympus Corporation, Tokyo, Japan for provision of electron multiplier CCD technology. This study was supported by grants from the Ministry of Education, Science, Sports and Culture, Japan (Grant-in-Aid for Scientific Research nos.21791690)

Reference

- Ellis HM, Horvitz HR: Genetic control of programmed cell death in the nematode C. elegans. Cell 1986, 44:817-829
- Yuan J, Shaham S, Ledoux S, Ellis HM, Horvitz HR: The C. elegans cell death gene ced-3 encodes a protein similar to mammalian interleukin-1 beta-converting enzyme. Cell 1993, 75:641-652
- Green DR, Kroemer G: Pharmacological manipulation of cell death: clinical applications in sight? J Clin Invest 2005, 115:2610-2617
- Hisatomi T, Sakamoto T, Murata T, Yamanaka I, Oshima Y, Hata Y, Ishibashi T, Inomata H, Susin SA, Kroemer G: Relocalization of apoptosis-inducing factor in photoreceptor apoptosis induced by retinal detachment in vivo. Am J Pathol 2001, 158:1271-1278
- Susin SA, Lorenzo HK, Zamzami N, Marzo I, Snow BE, Brothers GM, Mangion J, Jacotot E, Costantini P, Loeffler M, Larochette N, Goodlett DR, Aebersold R, Siderovski DP, Penninger JM, Kroemer G: Molecular characterization of mitochondrial apoptosis-inducing factor. Nature 1999, 397:441-446

- 6. Hisatomi T, Nakazawa T, Noda K, Almulki L, Miyahara S, Nakao S, Ito Y, She H, Kohno R, Michaud N, Ishibashi T, Hafezi-Moghadam A, Badley AD, Kroemer G, Miller JW: HIV protease inhibitors provide neuroprotection through inhibition of mitochondrial apoptosis in mice. J Clin Invest 2008, 118:2025-2038
- 7. Murakami Y, Ikeda Y, Yonemitsu Y, Onimaru M, Nakagawa K, Kohno R, Miyazaki M, Hisatomi T, Nakamura M, Yabe T, Hasegawa M, Ishibashi T, Sueishi K: Inhibition of nuclear translocation of apoptosis-inducing factor is an essential mechanism of the neuroprotective activity of pigment epithelium-derived factor in a rat model of retinal degeneration. Am J Pathol 2008, 173:1326-1338
- Tartaglia LA, Ayres TM, Wong GH, Goeddel DV: A novel domain within the 55 kd TNF receptor signals cell death. Cell 1993, 74:845-853
- Nakazawa T, Matsubara A, Noda K, Hisatomi T, She H, Skondra D, Miyahara S, Sobrin L, Thomas KL, Chen DF, Grosskreutz CL, Hafezi-Moghadam A, Miller JW: Characterization of cytokine responses to retinal detachment in rats. Mol Vis 2006, 12:867-878
- 10. Zacks DN, Zheng QD, Han Y, Bakhru R, Miller JW: FAS-mediated apoptosis

and its relation to intrinsic pathway activation in an experimental model of retinal detachment. Invest Ophthalmol Vis Sci 2004, 45:4563-4569

- 11. Trichonas G, Murakami Y, Thanos A, Morizane Y, Kayama M, Debouck CM, Hisatomi T, Miller JW, Vavvas DG: Receptor interacting protein kinases mediate retinal detachment-induced photoreceptor necrosis and compensate for inhibition of apoptosis. Proc Natl Acad Sci U S A 2010,
- 12. Dubyak GR, el-Moatassim C: Signal transduction via P2-purinergic receptors for extracellular ATP and other nucleotides. Am J Physiol 1993, 265:C577-606
- Sabirov RZ, Okada Y: ATP release via anion channels. Purinergic Signal 2005, 1:311-328
- Burnstock G: Purinergic Receptors in the Nervous System. In Current Topics in Membranes 2003, 54:307-368
- Abbracchio MP, Burnstock G, Verkhratsky A, Zimmermann H: Purinergic signalling in the nervous system: an overview. Trends Neurosci 2009, 32:19-29
- Wang X, Arcuino G, Takano T, Lin J, Peng WG, Wan P, Li P, Xu Q, Liu QS,
 Goldman SA, Nedergaard M: P2X7 receptor inhibition improves recovery after spinal cord injury. Nat Med 2004, 10:821-827
- 17. Khakh BS, North RA: P2X receptors as cell-surface ATP sensors in health and

disease. Nature 2006, 442:527-532

- Melani A, Turchi D, Vannucchi MG, Cipriani S, Gianfriddo M, Pedata F: ATP extracellular concentrations are increased in the rat striatum during in vivo ischemia. Neurochem Int 2005, 47:442-448
- Abbracchio MP, Burnstock G: Purinoceptors: are there families of P2X and P2Y purinoceptors? Pharmacol Ther 1994, 64:445-475
- 20. Ralevic V, Burnstock G: Receptors for purines and pyrimidines. Pharmacol Rev 1998, 50:413-492
- Surprenant A, Rassendren F, Kawashima E, North RA, Buell G: The cytolytic
 P2Z receptor for extracellular ATP identified as a P2X receptor (P2X7).
 Science 1996, 272:735-738
- Ferrari D, Los M, Bauer MK, Vandenabeele P, Wesselborg S, Schulze-Osthoff
 K: P2Z purinoreceptor ligation induces activation of caspases with distinct
 roles in apoptotic and necrotic alterations of cell death. FEBS Lett 1999,
 447:71-75
- 23. Denlinger LC, Fisette PL, Sommer JA, Watters JJ, Prabhu U, Dubyak GR, Proctor RA, Bertics PJ: Cutting edge: the nucleotide receptor P2X7 contains multiple protein- and lipid-interaction motifs including a potential binding site

for bacterial lipopolysaccharide. J Immunol 2001, 167:1871-1876

- 24. Kong Q, Wang M, Liao Z, Camden JM, Yu S, Simonyi A, Sun GY, Gonzalez FA, Erb L, Seye CI, Weisman GA: P2X(7) nucleotide receptors mediate caspase-8/9/3-dependent apoptosis in rat primary cortical neurons. Purinergic Signal 2005, 1:337-347
- 25. Sperlagh B, Szabo G, Erdelyi F, Baranyi M, Vizi ES: Homo- and heteroexchange of adenine nucleotides and nucleosides in rat hippocampal slices by the nucleoside transport system. Br J Pharmacol 2003, 139:623-633
- 26. Deuchars SA, Atkinson L, Brooke RE, Musa H, Milligan CJ, Batten TF, Buckley NJ, Parson SH, Deuchars J: Neuronal P2X7 receptors are targeted to presynaptic terminals in the central and peripheral nervous systems. J Neurosci 2001, 21:7143-7152
- 27. Pannicke T, Fischer W, Biedermann B, Schadlich H, Grosche J, Faude F,
 Wiedemann P, Allgaier C, Illes P, Burnstock G, Reichenbach A: P2X7
 receptors in Muller glial cells from the human retina. J Neurosci 2000,
 20:5965-5972
- 28. Brandle U, Kohler K, Wheeler-Schilling TH: Expression of the P2X7-receptor subunit in neurons of the rat retina. Brain Res Mol Brain Res 1998, 62:106-109

- Zhang X, Zhang M, Laties AM, Mitchell CH: Stimulation of P2X7 receptors elevates Ca2+ and kills retinal ganglion cells. Invest Ophthalmol Vis Sci 2005, 46:2183-2191
- Puthussery T, Fletcher EL: Synaptic localization of P2X7 receptors in the rat retina. J Comp Neurol 2004, 472:13-23
- 31. McLarnon JG, Ryu JK, Walker DG, Choi HB: Upregulated expression of purinergic P2X(7) receptor in Alzheimer disease and amyloid-beta peptide-treated microglia and in peptide-injected rat hippocampus. J Neuropathol Exp Neurol 2006, 65:1090-1097
- 32. Diaz-Hernandez M, Diez-Zaera M, Sanchez-Nogueiro J, Gomez-Villafuertes R, Canals JM, Alberch J, Miras-Portugal MT, Lucas JJ: Altered P2X7-receptor level and function in mouse models of Huntington's disease and therapeutic efficacy of antagonist administration. FASEB J 2009, 23:1893-1906
- 33. Peng W, Cotrina ML, Han X, Yu H, Bekar L, Blum L, Takano T, Tian GF, Goldman SA, Nedergaard M: Systemic administration of an antagonist of the ATP-sensitive receptor P2X7 improves recovery after spinal cord injury. Proc Natl Acad Sci U S A 2009, 106:12489-12493
- 34. Rodrigues EB, Meyer CH, Kroll P: Chromovitrectomy: a new field in

vitreoretinal surgery. Graefes Arch Clin Exp Ophthalmol 2005, 243:291-293

- 35. Abrams JS, Roncarolo MG, Yssel H, Andersson U, Gleich GJ, Silver JE:
 Strategies of anti-cytokine monoclonal antibody development: immunoassay of
 IL-10 and IL-5 in clinical samples. Immunol Rev 1992, 127:5-24
- 36. Kayagaki N, Yamaguchi N, Nagao F, Matsuo S, Maeda H, Okumura K, Yagita H: Polymorphism of murine Fas ligand that affects the biological activity. Proc Natl Acad Sci U S A 1997, 94:3914-3919
- 37. Thresher RJ, Ehrenberg M, Machemer R: Gas-mediated vitreous compression: an experimental alternative to mechanized vitrectomy. Graefes Arch Clin Exp Ophthalmol 1984, 221:192-198
- Irvine WD, Johnson MW, Hernandez E, Olsen KR: Retinal toxicity of human tissue plasminogen activator in vitrectomized rabbit eyes. Arch Ophthalmol 1991, 109:718-722
- 39. Enaida H, Sakamoto T, Hisatomi T, Goto Y, Ishibashi T: Morphological and functional damage of the retina caused by intravitreous indocyanine green in rat eyes. Graefes Arch Clin Exp Ophthalmol 2002, 240:209-213
- 40. Enaida H, Hisatomi T, Goto Y, Hata Y, Ueno A, Miura M, Kubota T, IshibashiT: Preclinical investigation of internal limiting membrane staining and peeling

using intravitreal brilliant blue G. Retina 2006, 26:623-630

- 41. Nakazawa T, Hisatomi T, Nakazawa C, Noda K, Maruyama K, She H,
 Matsubara A, Miyahara S, Nakao S, Yin Y, Benowitz L, Hafezi-Moghadam A,
 Miller JW: Monocyte chemoattractant protein 1 mediates retinal
 detachment-induced photoreceptor apoptosis. Proc Natl Acad Sci U S A 2007,
 104:2425-2430
- 42. Jiang LH, Mackenzie AB, North RA, Surprenant A: Brilliant blue G selectively blocks ATP-gated rat P2X(7) receptors. Mol Pharmacol 2000, 58:82-88
- Hu H, Lu W, Zhang M, Zhang X, Argall AJ, Patel S, Lee GE, Kim YC,
 Jacobson KA, Laties AM, Mitchell CH: Stimulation of the P2X7 receptor kills
 rat retinal ganglion cells in vivo. Exp Eye Res 2010, 91:425-432
- 44. Puthussery T, Fletcher E: Extracellular ATP induces retinal photoreceptor
 apoptosis through activation of purinoceptors in rodents. J Comp Neurol 2009,
 513:430-440
- 45. Joseph SM, Buchakjian MR, Dubyak GR: Colocalization of ATP release sites and ecto-ATPase activity at the extracellular surface of human astrocytes. J Biol Chem 2003, 278:23331-23342
- 46. Pellegatti P, Falzoni S, Pinton P, Rizzuto R, Di Virgilio F: A novel

recombinant plasma membrane-targeted luciferase reveals a new pathway for ATP secretion. Mol Biol Cell 2005, 16:3659-3665

- 47. Oike M, Kimura C, Koyama T, Yoshikawa M, Ito Y: Hypotonic stress-induced dual Ca(2+) responses in bovine aortic endothelial cells. Am J Physiol Heart Circ Physiol 2000, 279:H630-638
- Liu HT, Sabirov RZ, Okada Y: Oxygen-glucose deprivation induces ATP release via maxi-anion channels in astrocytes. Purinergic Signal 2008, 4:147-154
- Langer D, Hammer K, Koszalka P, Schrader J, Robson S, Zimmermann H: Distribution of ectonucleotidases in the rodent brain revisited. Cell Tissue Res 2008, 334:199-217
- 50. Beigi R, Kobatake E, Aizawa M, Dubyak GR: Detection of local ATP release from activated platelets using cell surface-attached firefly luciferase. Am J Physiol 1999, 276:C267-278
- Nagasawa K, Escartin C, Swanson RA: Astrocyte cultures exhibit P2X7 receptor channel opening in the absence of exogenous ligands. Glia 2009, 57:622-633
- 52. Adriouch S, Bannas P, Schwarz N, Fliegert R, Guse AH, Seman M, Haag F,

Koch-Nolte F: ADP-ribosylation at R125 gates the P2X7 ion channel by presenting a covalent ligand to its nucleotide binding site. FASEB J 2008, 22:861-869

- 53. Kadonosono K, Itoh N, Uchio E, Nakamura S, Ohno S: Staining of internal limiting membrane in macular hole surgery. Arch Ophthalmol 2000, 118:1116-1118
- 54. Burk SE, Da Mata AP, Snyder ME, Rosa RH, Jr., Foster RE: Indocyanine green-assisted peeling of the retinal internal limiting membrane.
 Ophthalmology 2000, 107:2010-2014
- 55. Engelbrecht NE, Freeman J, Sternberg P, Jr., Aaberg TM, Sr., Aaberg TM, Jr., Martin DF, Sippy BD: Retinal pigment epithelial changes after macular hole surgery with indocyanine green-assisted internal limiting membrane peeling. Am J Ophthalmol 2002, 133:89-94
- 56. Uemura A, Kanda S, Sakamoto Y, Kita H: Visual field defects after uneventful vitrectomy for epiretinal membrane with indocyanine green-assisted internal limiting membrane peeling. Am J Ophthalmol 2003, 136:252-257
- 57. Kawahara S, Hata Y, Miura M, Kita T, Sengoku A, Nakao S, Mochizuki Y, Enaida H, Ueno A, Hafezi-Moghadam A, Ishibashi T: Intracellular events in

retinal glial cells exposed to ICG and BBG. Invest Ophthalmol Vis Sci 2007, 48:4426-4432

- 58. Hisatomi T, Enaida H, Matsumoto H, Kagimoto T, Ueno A, Hata Y, Kubota T, Goto Y, Ishibashi T: Staining ability and biocompatibility of brilliant blue G: preclinical study of brilliant blue G as an adjunct for capsular staining. Arch Ophthalmol 2006, 124:514-519
- 59. Enaida H, Hisatomi T, Hata Y, Ueno A, Goto Y, Yamada T, Kubota T,
 Ishibashi T: Brilliant blue G selectively stains the internal limiting
 membrane/brilliant blue G-assisted membrane peeling. Retina 2006,
 26:631-636
- Farah ME, Maia M, Rodrigues EB: Dyes in ocular surgery: principles for use in chromovitrectomy. Am J Ophthalmol 2009, 148:332-340
- Rodrigues EB, Penha FM, de Paula Fiod Costa E, Maia M, Dib E, Moraes M,
 Jr., Meyer CH, Magalhaes O, Jr., Melo GB, Stefano V, Dias AB, Farah ME:
 Ability of new vital dyes to stain intraocular membranes and tissues in ocular
 surgery. Am J Ophthalmol 2010, 149:265-277
- 62. Kumar A, Gogia V, Shah VM, Nag TC: Comparative evaluation of anatomical and functional outcomes using brilliant blue G versus triamcinolone assisted

ILM peeling in macular hole surgery in Indian population. Graefes Arch Clin

Exp Ophthalmol 2011, 249:987-995

Legends to Figures

Figure 1. P2RX7 agonists induced retinal cell death in primary retinal cell cultures. The viability of retinal cells was assessed by incubation with calcein AM or MitoTracker CMTMRos after 24 h of culture with ATP or BzATP. Photoreceptors were labeled by immunocytochemistry for recoverin (in green). **A:** The frequency of calcein⁺ (**top**; in blue) or CMTMRos⁺ (**bottom**; in red) photoreceptors decreased after incubation with 200 μ M to 5 mM ATP in a dose-dependent manner. **B:** BzATP also reduced calcein⁺ (**top**) or CMTMRos⁺ (**bottom**) photoreceptors in a dose-dependent fashion. These quantifications are shown in the **right** panels in **A** and **B**.

Figure 2. BBG attenuated photoreceptor death as a P2RX7 antagonist.

A: The viability of photoreceptors after 24 h of continuous incubation with 1-10 μ M BBG. **B:** Photoreceptor death (induced by 1 mM BzATP) was partially reversed by 30 min transient incubation (BBG) or 24 h continuous incubation (co-BBG) of 1-10 μ M BBG, or 1-10 μ M KN-62. **C:** Photoreceptor viability was preserved in BzATP-treated (1 mM, 24 h) photoreceptors from P2RX7^{-/-} mice (KO) as compared to wild type (WT) controls. **D** and **E:** Neutralizing antibody against TNF-alpha (MP6-XT22) or CD95 ligand (MFL3) failed to prevent BzATP-induced photoreceptor death. *n* = 10 per

group; **P < 0.01. Scale bar= 20 µm.

Figure 3. P2RX7 induced caspase-8 cleavage and calcium influx in vitro.

A: Electron microscopy revealed the relatively well-preserved organelle of photoreceptors in primary cultures without BzATP administration (left). Apoptotic cell death was observed with chromatin condensation (middle) after BzATP administration. Necrotic cell death could also be detected after BzATP (**right**). Scale bar= $2 \mu m$. **B:** Representative images of immunocytochemistry for cleaved caspase-8 in primary retinal cell cultures (left; cleaved caspase-8 in red, middle; TUNEL in green, right; merged with Hoechst 33342 in blue). Cytoplasmic caspase-8 cleavage was detected in TUNEL-positive cells after 1 mM BzATP administration, and decreased by BBG treatment. C: Quantified results of cleaved caspase- 8^+ or TUNEL⁺ cells, n = 10 per group; *P < 0.05. **D:** The fluorescent images of calcium indicator Fluo-4 in primary retinal cell cultures before and 20 min after BzATP administration. The phase contrast images are shown in the left panel. The BzATP-induced increase of Fluo-4 fluorescence (top) was attenuated by addition of 1 mM EGTA (middle) or 10 µM BBG (**bottom**). Scale bar= $5 \mu m$.

Figure 4. P2RX7 stimulation induced photoreceptor apoptosis/necrosis *in vivo*.A: Representative TUNEL (in green) and propidium iodide (in red) staining of mouse

retina after 3 (top) and 14 (bottom) days of gas vitrectomy. B: Representative images of cleaved caspase-8 in the mouse retina at 24 h after the intraocular injection of 20 mM BzATP (top left; Hoechst 33342 in blue, top right; cleaved caspase-8 in red, bottom left; TUNEL in green, bottom right; merged). In wild type mice, cleaved caspase-8-positive staining was observed in TUNEL-positive apoptotic cells in the inner nuclear layer (INL) and the outer nuclear layer (ONL) after the vitreous injection of BzATP (white arrows). C: The BzATP injections dose-dependently induced photoreceptor apoptosis in the ONL, as detected by TUNEL (left), and this effect was decreased by co-injection of 500 µM BBG (middle). Only very few TUNEL-positive photoreceptors were detected upon injection of BzATP into P2RX7^{-/-} mice (right). TUNEL-stainings are shown in green and propidium iodide in red. The quantitations are shown in **D**. n = 6 per group; **P < 0.01. Scale bar= 20 µm. **E:** Transmission electron microscopy revealed BzATP-treated photoreceptors with characteristics of apoptosis, cell shrinkage and chromatin condensation (white arrow) in the ONL, and degenerated pedicles and spherules in the OPL (black arrows). Apoptotic photoreceptors were decreased, and rod spherules and cone pedicles were well preserved in mice receiving BzATP plus BBG. Scale bar= $2 \mu m$.

Figure 5. Neuroprotective effects of BBG in primary retinal cell cultures. A and B:

Representative images of immunofluorescence detection of AIF (AIF in red, TUNEL in green, Hoechst33342 in blue) and cleaved caspase-9 (cleaved caspase-9 in red, TUNEL in green, Hoechst33342 in blue) in control cultures and after starvation in the absence or presence of BBG. Scale bar = 5 μ m. The quantitations are shown in **C**. *n*=10 per group; **P* < 0.05. **D**: The dose-dependent neuroprotective effect of BBG. *n* = 10 per group; ***P* < 0.01.

Figure 6. Nutrient starvation-induced ATP release from primary retinal cells *in vitro*. The ATP levels in primary culture were measured by luciferin-luciferase assay. Total cellular levels of ATP were measured after permeabilization of plasma membranes, while the inferior detection limit of ATP levels was determined by adding apyrase to the culture supernatants. Starvation was induced as in Figure 4 in the presence or absence of BBG or the ecto-ATPase inhibitor, β , γ -Me-ATP (n = 10 per group; *P < 0.05, **P < 0.01).

Figure 7. Time-lapse imaging of ATP release induced by starvation. For dynamic imaging of the ATP release, light production from the luciferin-luciferase reaction was captured by an electron multiplier charge-coupled camera for 3 h). Representative images are shown (see Supplemental movies for total time-lapse images). Circles in the **middle** and **bottom** panel indicate events of ATP release induced by starvation and

arrows indicate the same location during the observation periods. Scale bar = $10 \ \mu m$.





Α





С

В

Cleaved caspase-8	TUNEL	Merge
		•
BzATP		
BzATP+BBG	-	•









Figure 7 Control 0:10 2:00 1:10 2:55 S Ł L L Starved 0:05 2:55 2:10 1:10 6 BBG 0:15 0:35 2:55 2:35