

Numerical verification methods of solutions for nonlinear singularly perturbed problem

Hashimoto, Kouji
Graduate School of Mathematics, Kyushu University

Abe, Ryohei
Graduate School of Mathematics, Kyushu University

Nakao, Mitsuhiro T.
Faculty of Mathematics, Kyushu University

Watanabe, Yoshitaka
Computing and Communications Center, Kyushu University

<https://hdl.handle.net/2324/11834>

出版情報 : Japan Journal of Industrial and Applied Mathematics. 22 (1), pp.111-131, 2005-02. 日本応用数理学会
バージョン :
権利関係 :



MHF Preprint Series

Kyushu University
21st Century COE Program
Development of Dynamic Mathematics with
High Functionality

Numerical verification methods of solutions for nonlinear singularly perturbed problem

K. Hashimoto, R. Abe
M.T. Nakao, Y. Watanabe

MHF 2004-15

(Received May 21, 2004)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

Numerical Verification Methods of Solutions for Nonlinear Singularly Perturbed Problem

K. Hashimoto¹ R. Abe¹ M. T. Nakao² Y. Watanabe³

¹*Graduate School of Mathematics, Kyushu University, Fukuoka 812-8581, Japan.*

²*Faculty of Mathematics, Kyushu University.*

³*Computing and Communications Center, Kyushu University.*

Abstract

In order to verify the solutions of nonlinear boundary value problems by Nakao's computer-assisted numerical method on computer, it is required to find a constant, as sharp as possible, in the a priori error estimates for the finite element approximation of some simple linear problems. For singularly perturbed problems, however, generally it is known that the perturbation term produces a bad effect on the a priori error estimates, i.e., leads to a large constant, if we use the usual approximation methods. In this paper, we propose some verification algorithms of solutions for nonlinear singularly perturbed problems by using the constant obtained in the a priori error estimates based on the exponential fitting method with Green's function. Some numerical examples which confirm us the effectiveness of our method are presented.

Key word numerical verification, singularly perturbed problem, finite element method, a priori constant.

1 Introduction

The numerical verification method for solutions of nonlinear differential equations realizes a mathematically rigorous analysis on computer, which is often effectively applied to the problem for which usual theoretical approaches no longer work. However, there have been no practical verification methods which are suitable to the nonlinear singularly perturbed problem up to the present. In this paper, we consider the verification method of solutions for the following nonlinear singularly perturbation problem including parameter ε ($1 \gg \varepsilon > 0$):

$$\begin{cases} Lu \equiv -\varepsilon u'' - b(x)u' + c(x)u = f(u) & \text{in } (0, 1), \\ u(0) = u(1) = 0, \end{cases} \quad (1.1)$$

where $f : L^\infty(0, 1) \rightarrow L^\infty(0, 1)$ is a bounded continuous map, and we suppose that $b(x), c(x) \in W_\infty^1(0, 1)$, and that there exists a constant γ such that $c(x) \geq \gamma > 0$.

Our arguments below are based on the finite element method on the interval $(0, 1)$ with mesh $: 0 = x_0 < x_1 < \cdots < x_n < x_{n+1} = 1$. Let h and h_{\min} denote the maximum and the minimum width of subintervals, respectively. We denote the trial and the test

spaces by S_h and $V_h \subset L^\infty(0,1) \cap H_0^1(0,1)$, respectively. And we define the piecewise constant functions $\bar{b} = \bar{b}(x)$, $\bar{c} = \bar{c}(x) \in L^\infty(0,1)$ by

$$\bar{b}(x) = \frac{b(x_{i-1}) + b(x_i)}{2}, \quad \bar{c}(x) = \frac{c(x_{i-1}) + c(x_i)}{2}, \quad x \in (x_{i-1}, x_i),$$

for $i = 1, \dots, n$.

Here, we try to extend and apply Nakao's verification method [2] for the application to the nonlinear singularly perturbed problem (1.1). In this method, it plays an important and essential role to get an a priori error estimation of the finite element solution of the linear problem, usually based on H_0^1 -projection. For the singularly perturbed problems, however, when the usual finite element schemes are applied, it is known that the constant in the a priori error estimates increases proportionally to the perturbation parameter ε . That is, from the error estimates of the H_0^1 -projection to the solution of $-\varepsilon\phi'' = g$ ($g \in L^\infty(0,1)$), it is known that, if we use piecewise polynomials then we have

$$\|\phi - P_{H_0^1}\phi\|_\infty \leq C(\varepsilon)h^2\|g\|_\infty, \quad (1.2)$$

where $C(\varepsilon) \rightarrow \infty$ if $\varepsilon \rightarrow 0$. The estimates (1.2) imply that using such an approximate method would lead the inefficient computational procedure for the verification for small ε . Therefore, in this paper we paid attention to the exponential fitting technique, \bar{L} -spline method, studied by M. Stynes and E. O'Riordan [4][6][7][8]. That is, we carefully and numerically estimated various kinds of constants appearing in the error analyses of their methods. And we succeed in estimating an a priori constant, such as in (1.2), numerically with independent of or less dependent on the perturbation parameter, and to apply it to the verification of solutions for the nonlinear singularly perturbed problem (1.1).

In Section 2, we show the computational result of the constructive a priori error estimation of constants to the linear problem. Especially, as for the convection-diffusion problem of the subsection 2.1, we obtained the error estimation of $O(h)$ which is independent of ε . In the subsection 2.2, the theoretical analysis yields some constructive error estimations to the reaction-diffusion problem, which can be more effectively used for the verification of the nonlinear problem compared with usual methods, even though it still depends on the perturbation parameter. In the section 3, an actual verification algorithm is shown, and some verification results are presented in the section 4.

2 Constructive a priori error estimates for a linear singularly perturbed problem using Green's function

In this section, we try to compute the constant in the a priori error estimation of finite element approximations for linear convection and reaction diffusion equations. We determine two different kinds of constants corresponding to the convection and the reaction diffusion problems, respectively.

2.1 Convection Diffusion Problem

We first consider the following linear convection diffusion problems.

$$\begin{cases} L\phi & \equiv & -\varepsilon\phi'' - b(x)\phi' + c(x)\phi & = & g & \text{in } (0,1), \\ \phi(0) = \phi(1) & = & 0, \end{cases} \quad (2.1)$$

where $g \in L^\infty(0, 1)$, and $b(x), c(x) \in W_\infty^1(0, 1)$ are given functions satisfying $b(x) \geq \beta > 0$ and $c(x) \geq \gamma > 0$.

Now we define the bilinear form of (2.1) by, for each $\varphi, \psi \in H_0^1(0, 1)$,

$$\begin{aligned} a(\varphi, \psi) &\equiv \varepsilon(\varphi', \psi') - (b\varphi', \psi) + (c\varphi, \psi), \\ a_h(\varphi, \psi) &\equiv \varepsilon(\varphi', \psi') - (\bar{b}\varphi', \psi) + (\bar{c}\varphi, \psi), \end{aligned}$$

where (\cdot, \cdot) denotes L^2 inner product on $(0, 1)$. Then, the projection $P_h : H_0^1 \rightarrow S_h$ is defined as

$$a(\phi - P_h\phi, \psi_h) = 0, \quad \text{for all } \psi_h \in V_h. \quad (2.2)$$

And we also define the approximation $P_h^\varepsilon \phi \equiv \phi_h^\varepsilon \in S_h$ of solution ϕ to (2.1), which we call the P_h^ε -projection, as follows :

$$a_h(\phi_h^\varepsilon, \psi_h) = a(\phi, \psi_h), \quad \text{for all } \psi_h \in V_h. \quad (2.3)$$

Now, M. Stynes and E. O'Riordan [7] introduced the following \bar{L} -spline $\{\varphi_i\}_{i=1}^n$ and \bar{L}^* -spline $\{\psi_i\}_{i=1}^n$ which constitute bases of S_h and V_h , respectively, satisfying for $i = 1, \dots, n$

$$\begin{aligned} \bar{L}\varphi_i &\equiv -\varepsilon\varphi_i'' - \bar{b}\varphi_i' + \bar{c}\varphi_i = 0 & \text{in } [0, 1] \setminus \{x_1, \dots, x_n\}, \\ \varphi_i(x_k) &= \delta_i^k & \text{for } k = 0, \dots, n+1, \\ \bar{L}^*\psi_i &\equiv -\varepsilon\psi_i'' + \bar{b}\psi_i' + \bar{c}\psi_i = 0 & \text{in } [0, 1] \setminus \{x_1, \dots, x_n\}, \\ \psi_i(x_k) &= \delta_i^k & \text{for } k = 0, \dots, n+1, \end{aligned}$$

where δ_i^k stands for Kronecker's delta.

We now define the Green function $G_i = G(x, x_i)$ by the following equation, which is spanned by $\{\psi_i\}_{i=1}^n$:

$$a_h(w, G_i) = w(x_i) \quad \text{for all } w \in H_0^1(0, 1).$$

Remark 1 [7] For each $i \in \{1, \dots, n\}$, the Green function $G_i(\cdot) \in C[0, 1]$ is characterized by

$$-\varepsilon G_i''(x) + \bar{b}G_i'(x) + \bar{c}G_i(x) = 0 \quad \text{in } [0, 1] \setminus \{x_1, \dots, x_n\},$$

$$G_i(0) = G_i(1) = 0,$$

$$\lim_{x \rightarrow x_k^+} (\varepsilon G_i'(x) - \bar{b}G_i(x)) - \lim_{x \rightarrow x_k^-} (\varepsilon G_i'(x) - \bar{b}G_i(x)) = -\delta_i^k,$$

where $x_k^- \equiv x_k - 0$ and $x_k^+ \equiv x_k + 0$. Moreover, $G_i(x)$ lies in V_h .

Lemma 2.1 For each $i \in \{1, \dots, n\}$, the Green function $G_i(\cdot)$ satisfies

$$\varepsilon G_i'(x) - \bar{b}G_i(x) = \varepsilon G_i'(0) + \int_0^x \bar{c}G_i(t) dt - H_i(x), \quad x \in [0, 1] \setminus \{x_1, \dots, x_n\},$$

where

$$H_i(x) = \begin{cases} 1 & \text{if } x \geq x_i, \\ 0 & \text{if } x < x_i. \end{cases}$$

Proof: By Remark 1, we have

$$\begin{aligned}
\int_0^x \bar{c}G_i(t) dt &= \sum_{k=1}^{j-1} \int_{x_{k-1}}^{x_k} (\varepsilon G_i''(t) - \bar{b}G_i'(t)) dt + \int_{x_{j-1}}^x (\varepsilon G_i''(t) - \bar{b}G_i'(t)) dt \\
&= \sum_{k=1}^{j-1} [\varepsilon G_i'(t) - \bar{b}G_i(t)]_{x_{k-1}}^{x_k} + [\varepsilon G_i'(t) - \bar{b}G_i(t)]_{x_{j-1}}^x \\
&= (\varepsilon G_i'(x) - \bar{b}G_i(x)) - \varepsilon G_i'(0) \\
&\quad + \sum_{k=1}^{j-1} \left[(\varepsilon G_i'(x_k^-) - \bar{b}G_i(x_k^-)) - (\varepsilon G_i'(x_k^+) - \bar{b}G_i(x_k^+)) \right],
\end{aligned}$$

which proves the lemma. ■

Lemma 2.2 For each $i \in \{1, \dots, n\}$ and for arbitrary $x \in (0, 1)$, we have

$$0 \leq G_i(x) \leq \frac{1}{\beta}.$$

Proof: From [7], it follows that $G_i(x) \geq 0$ for all $i \in \{1, \dots, n\}$ (cf. the proof of Lemma 2.4 in this paper). We assume that there exists $z \in [x_{j-1}, x_j]$ such that $G_i(z) > 1/\beta$. Then, from Lemma 2.1 we have

$$\begin{aligned}
\varepsilon G_i'(z) &= \bar{b}G_i(z) + \varepsilon G_i'(0) + \int_0^z \bar{c}G_i(t) dt - H_i(z) \\
&\geq \bar{b}G_i(z) - 1 \\
&\geq \beta G_i(z) - 1 \\
&> 0,
\end{aligned}$$

where we have used the fact that $G_i'(0) \geq 0$ because $G_i(x) \geq 0$ and $G_i(0) = 0$. Hence we obtain $1/\beta < G_i(z) < G_i(x_j)$. It means that $1/\beta < G_i(1)$ from the inductive argument, which contradicts with $G_i(1) = 0$. ■

Lemma 2.3 Let ϕ be a solution of (2.1). Then for all $x \in (0, 1)$

$$|\phi'(x)| \leq \left(\frac{1}{\varepsilon} C_1 e^{-\frac{\beta}{\varepsilon}x} + C_2 e^{-\frac{\beta}{\varepsilon}x} + C_3 \right) \|g\|_\infty,$$

where $K = 1/\max\{\beta, \gamma\}$ and

$$C_1 = \frac{2}{\beta} K \|b\|_\infty^2, \quad C_2 = \frac{1}{\beta} (1 + K \|c\|_\infty + 2K \|b'\|), \quad C_3 = \frac{1}{\beta} (1 + K \|c\|_\infty).$$

Proof: We first show that $\|\phi\|_\infty \leq 1/\max\{\beta, \gamma\} \|g\|_\infty$. Let $y_1(x) = K_1(1-x)\|g\|_\infty \pm \phi(x)$ and $y_2(x) = K_2\|g\|_\infty \pm \phi(x)$ where K_1, K_2 are positive constants. Then we have that $y_1(0) \geq y_1(1) = 0$ and $y_2(0) = y_2(1) \geq 0$. Hence if $K_1 = 1/\beta$ and $K_2 = 1/\gamma$ then we find

$$\begin{array}{lcl}
Ly_1 & = & K_1 [b(x) + (1-x)c(x)] \|g\|_\infty \pm g \\
& \geq & K_1 \beta \|g\|_\infty \pm g \\
& \geq & 0,
\end{array} \quad \left| \quad \begin{array}{lcl}
Ly_2 & = & K_2 c(x) \|g\|_\infty \pm g \\
& \geq & K_2 \gamma \|g\|_\infty \pm g \\
& \geq & 0.
\end{array} \right.$$

Therefore, if we choose $K := 1/\max\{\beta, \gamma\}$, then we obtain $\|\phi\|_\infty \leq K\|g\|_\infty$ by the maximum principle.

Now let $g_0(x) \equiv g(x) - c(x)\phi$, and let

$$g_1(x) \equiv \int_0^x g_0(t) e^{\frac{1}{\varepsilon} \hat{b}(t)} dt, \quad g_2(x) \equiv \int_x^1 g_0(t) e^{-\frac{1}{\varepsilon} \hat{b}(t)} dt,$$

where $\hat{b}(x) := \int_0^x b(t) dt > 0$. Then we get

$$(-\varepsilon \phi' e^{\frac{1}{\varepsilon} \hat{b}(x)})' = (-\varepsilon \phi'' - b(x)\phi') e^{\frac{1}{\varepsilon} \hat{b}(x)}, \quad (2.4)$$

$$(-\varepsilon \phi' e^{-\frac{1}{\varepsilon} \hat{b}(x)})' = (-\varepsilon \phi'' + b(x)\phi') e^{-\frac{1}{\varepsilon} \hat{b}(x)}. \quad (2.5)$$

From (2.4)-(2.5) we have

$$\int_0^x (-\varepsilon \phi' e^{\frac{1}{\varepsilon} \hat{b}(t)})' dt = -\varepsilon \phi'(x) e^{\frac{1}{\varepsilon} \hat{b}(x)} + \varepsilon \phi'(0) e^{\frac{1}{\varepsilon} \hat{b}(0)} = g_1(x), \quad (2.6)$$

$$\begin{aligned} \int_x^1 (-\varepsilon \phi' e^{-\frac{1}{\varepsilon} \hat{b}(t)})' dt &= -\varepsilon \phi'(1) e^{-\frac{1}{\varepsilon} \hat{b}(1)} + \varepsilon \phi'(x) e^{-\frac{1}{\varepsilon} \hat{b}(x)} \\ &= g_2(x) + 2 \int_x^1 b(t) \phi'(t) e^{-\frac{1}{\varepsilon} \hat{b}(t)} dt \\ &= g_2(x) + 2 \left[b(t) \phi(t) e^{-\frac{1}{\varepsilon} \hat{b}(t)} \right]_x^1 - 2\kappa(x), \end{aligned} \quad (2.7)$$

where

$$\kappa(x) \equiv \int_x^1 \phi(t) \left(b'(t) e^{-\frac{1}{\varepsilon} \hat{b}(t)} - \frac{b(t)^2}{\varepsilon} e^{-\frac{1}{\varepsilon} \hat{b}(t)} \right) dt.$$

Rewriting (2.6) as

$$-\varepsilon \phi'(x) + \varepsilon \phi'(0) e^{-\frac{1}{\varepsilon} \hat{b}(x)} = g_1(x) e^{-\frac{1}{\varepsilon} \hat{b}(x)}, \quad (2.8)$$

we have the following linear system from (2.7) and (2.8).

$$\begin{aligned} \varepsilon \phi'(1) - \varepsilon \phi'(0) e^{-\frac{1}{\varepsilon} \hat{b}(1)} &= -g_1(1) e^{-\frac{1}{\varepsilon} \hat{b}(1)}, \\ -\varepsilon \phi'(1) e^{-\frac{1}{\varepsilon} \hat{b}(1)} + \varepsilon \phi'(0) &= g_2(0) - 2\kappa(0). \end{aligned}$$

Since $\hat{b}(1) := \int_0^1 b(t) dt \geq \beta > 0$, this system is nonsingular so that

$$\begin{pmatrix} 1 & -e^{-\frac{1}{\varepsilon} \hat{b}(1)} \\ -e^{-\frac{1}{\varepsilon} \hat{b}(1)} & 1 \end{pmatrix}^{-1} = \frac{1}{1 - e^{-\frac{2}{\varepsilon} \hat{b}(1)}} \begin{pmatrix} 1 & e^{-\frac{1}{\varepsilon} \hat{b}(1)} \\ e^{-\frac{1}{\varepsilon} \hat{b}(1)} & 1 \end{pmatrix} \geq 0.$$

Therefore we obtain

$$\begin{aligned} \begin{pmatrix} |\phi'(1)| \\ |\phi'(0)| \end{pmatrix} &\leq \frac{1}{\varepsilon} \frac{1}{1 - e^{-\frac{2}{\varepsilon} \hat{b}(1)}} \begin{pmatrix} 1 & e^{-\frac{1}{\varepsilon} \hat{b}(1)} \\ e^{-\frac{1}{\varepsilon} \hat{b}(1)} & 1 \end{pmatrix} \begin{pmatrix} |g_1(1) e^{-\frac{1}{\varepsilon} \hat{b}(1)}| \\ |g_2(0) + 2|\kappa(0)|| \end{pmatrix} \\ &\leq \frac{1}{\varepsilon} \frac{1}{1 - e^{-\frac{2}{\varepsilon} \beta}} \begin{pmatrix} 1 & e^{-\frac{\beta}{\varepsilon}} \\ e^{-\frac{\beta}{\varepsilon}} & 1 \end{pmatrix} \begin{pmatrix} |g_1(1) e^{-\frac{1}{\varepsilon} \hat{b}(1)}| \\ |g_2(0) + 2|\kappa(0)|| \end{pmatrix}. \end{aligned}$$

And we have

$$\begin{aligned}
|g_1(1)e^{-\frac{1}{\varepsilon}\hat{b}(1)}| &= \left| \int_0^1 g_0(t)e^{-\frac{1}{\varepsilon}[\hat{b}(1)-\hat{b}(t)]} dt \right| \leq \|g_0\|_\infty \int_0^1 e^{-\frac{1}{\varepsilon}[\hat{b}(1)-\hat{b}(t)]} dt \\
&\leq \|g_0\|_\infty \int_0^1 e^{-\frac{\beta}{\varepsilon}(1-t)} dt \\
&= \frac{\varepsilon}{\beta} \|g_0\|_\infty \left(1 - e^{-\frac{\beta}{\varepsilon}}\right),
\end{aligned}$$

$$\begin{aligned}
|g_2(0)| &= \left| \int_0^1 g_0(t)e^{-\frac{1}{\varepsilon}\hat{b}(t)} dt \right| \leq \|g_0\|_\infty \int_0^1 e^{-\frac{1}{\varepsilon}\hat{b}(t)} dt \\
&\leq \|g_0\|_\infty \int_0^1 e^{-\frac{\beta}{\varepsilon}t} dt \\
&= \frac{\varepsilon}{\beta} \|g_0\|_\infty \left(1 - e^{-\frac{\beta}{\varepsilon}}\right).
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
|\kappa(0)| &= \left| \int_0^1 \phi(t) \left(b'(t)e^{-\frac{1}{\varepsilon}\hat{b}(t)} - \frac{b(t)^2}{\varepsilon}e^{-\frac{1}{\varepsilon}\hat{b}(t)} \right) dt \right| \\
&\leq \left| \int_0^1 b'(t)\phi(t)e^{-\frac{1}{\varepsilon}\hat{b}(t)} dt \right| + \frac{1}{\varepsilon} \left| \int_0^1 b(t)^2\phi(t)e^{-\frac{1}{\varepsilon}\hat{b}(t)} dt \right| \\
&\leq \|b'\|_\infty \|\phi\|_\infty \int_0^1 e^{-\frac{1}{\varepsilon}\hat{b}(t)} dt + \frac{1}{\varepsilon} \|b\|_\infty^2 \|\phi\|_\infty \int_0^1 e^{-\frac{1}{\varepsilon}\hat{b}(t)} dt \\
&\leq \|b'\|_\infty \|\phi\|_\infty \int_0^1 e^{-\frac{\beta}{\varepsilon}t} dt + \frac{1}{\varepsilon} \|b\|_\infty^2 \|\phi\|_\infty \int_0^1 e^{-\frac{\beta}{\varepsilon}t} dt \\
&= \|b'\|_\infty \|\phi\|_\infty \frac{\varepsilon}{\beta} (1 - e^{-\frac{\beta}{\varepsilon}}) + \|b\|_\infty^2 \|\phi\|_\infty \frac{1}{\beta} (1 - e^{-\frac{\beta}{\varepsilon}}).
\end{aligned}$$

Since $\|g_0\|_\infty \leq (1 + K\|c\|_\infty)\|g\|_\infty$, we obtain

$$\begin{aligned}
|\phi'(0)| &\leq \frac{1 - e^{-\frac{\beta}{\varepsilon}}}{1 - e^{-\frac{2\beta}{\varepsilon}}} \left[\frac{1 + e^{-\frac{\beta}{\varepsilon}}}{\beta} \|g_0\|_\infty + \frac{2}{\beta} \|b'\|_\infty \|\phi\|_\infty + \frac{2}{\varepsilon\beta} \|b\|_\infty^2 \|\phi\|_\infty \right] \\
&\leq \frac{1}{1 + e^{-\frac{\beta}{\varepsilon}}} \left[\frac{1 + e^{-\frac{\beta}{\varepsilon}}}{\beta} \|g_0\|_\infty + \frac{2}{\beta} \|b'\|_\infty \|\phi\|_\infty + \frac{2}{\varepsilon\beta} \|b\|_\infty^2 \|\phi\|_\infty \right] \\
&\leq \frac{1}{\beta} (1 + K\|c\|_\infty + 2K\|b'\|) \|g\|_\infty + \frac{2}{\varepsilon\beta} K \|b\|_\infty^2 \|g\|_\infty. \tag{2.9}
\end{aligned}$$

Finally, we have the following estimate

$$\begin{aligned}
|g_1(x)e^{-\frac{1}{\varepsilon}\hat{b}(x)}| &= \left| \int_0^x g_0(t)e^{-\frac{1}{\varepsilon}[\hat{b}(x)-\hat{b}(t)]} dt \right| \leq \|g_0\|_\infty \int_0^x e^{-\frac{1}{\varepsilon}[\hat{b}(x)-\hat{b}(t)]} dt \\
&\leq \|g_0\|_\infty \int_0^x e^{-\frac{\beta}{\varepsilon}(x-t)} dt \\
&\leq \frac{\varepsilon}{\beta} (1 + K\|c\|_\infty) \|g\|_\infty. \tag{2.10}
\end{aligned}$$

Combining (2.9) and (2.10) with (2.8), we obtain the desired conclusion. ■

Theorem 2.1 *Let ϕ and ϕ_h^ε be solutions of (2.1) and (2.3), respectively. Then*

$$|(\phi - \phi_h^\varepsilon)(x_i)| \leq C_0 h \|g\|_\infty, \quad i = 1, 2, \dots, n, \quad (2.11)$$

where

$$C_0 = \frac{1}{\beta^2} \|b'\|_\infty (C_1 + \varepsilon C_2 + \beta C_3) + \frac{1}{\beta} K \|c'\|_\infty.$$

Here, constants C_i , $1 \leq i \leq 3$, and K are defined in Lemma 2.3.

Proof: From the property of Green's function, we have

$$\begin{aligned} (\phi - \phi_h^\varepsilon)(x_i) &= a_h(\phi - \phi_h^\varepsilon, G_i) = a_h(\phi, G_i) - (g, G_i) \\ &= -(\phi', (\bar{b} - b)G_i) + (\phi, (\bar{c} - c)G_i). \end{aligned}$$

Hence, we have

$$|(\phi - \phi_h^\varepsilon)(x_i)| \leq \|\phi'\|_{L_1} \|\bar{b} - b\|_\infty \|G_i\|_\infty + \|\phi\|_\infty \|\bar{c} - c\|_{L_1} \|G_i\|_\infty.$$

Now we estimate each term in the above. First, note that

$$\begin{aligned} \|\bar{b} - b\|_\infty &\leq \max_i \int_{x_{i-1}}^{x_i} |b'(t)| dt \leq \|b'\|_\infty h, \\ \|\bar{c} - c\|_{L_1} &\leq \int_0^1 \|\bar{c} - c\|_\infty dt \leq \|c'\|_\infty h. \end{aligned}$$

Since the estimations of $\|\phi\|_\infty$ and $\|G_i\|_\infty$ follow from Lemma 2.2 and 2.3, respectively, it suffices to bound $\|\phi'\|_{L_1}$. From Lemma 2.3, we obtain

$$\begin{aligned} \|\phi'\|_{L_1} &= \int_0^1 |\phi'(x)| dx \\ &\leq \int_0^1 \left(\frac{1}{\varepsilon} C_1 e^{-\frac{\beta}{\varepsilon} x} + C_2 e^{-\frac{\beta}{\varepsilon} x} + C_3 \right) \|g\|_\infty dx \\ &\leq \left(\frac{1}{\beta} C_1 + \frac{\varepsilon}{\beta} C_2 + C_3 \right) \|g\|_\infty. \end{aligned}$$

Thus the conclusion follows immediately. ■

Theorem 2.2 *Let ϕ and ϕ_h^ε be solutions of (2.1) and (2.3), respectively. Then*

$$\|\phi - \phi_h^\varepsilon\|_\infty \leq C h \|g\|_\infty,$$

where $C \equiv C(\lambda) = \tau_1 + (2 - \lambda)\tau_2$ for $\lambda \in [0, 1)$ and τ_1, τ_2 are constants satisfying

$$\tau_1 = \frac{4}{\beta^2 + 4\varepsilon\gamma} \|b'\|_\infty (C_1 + \varepsilon C_2), \quad \tau_2 = \begin{cases} \tau & \text{if } C_0 = 0, \\ \max\{\tau, C_0/(1 - \lambda)\} & \text{otherwise,} \end{cases}$$

where

$$\tau \equiv \frac{1}{\beta + (1 - \lambda)\gamma h} [1 + (\|b'\|_\infty C_3 + K \|c'\|_\infty) h].$$

Moreover if $C_0 = 0$ then we can choose that $\lambda \in [0, 1]$.

(Constants C_i , $0 \leq i \leq 3$, and K are same as in Theorem 2.1.)

Proof: We prove the theorem using the maximum principle. First, we consider the case of $C_0 = 0$. Notice that, by the maximum principle, if the function $r(x) \geq 0$ satisfies $\bar{L}(\pm(\phi - \phi_h^\varepsilon)) \leq \bar{L}r(x)$ for each $x \in (x_{i-1}, x_i)$, then we have $\pm(\phi - \phi_h^\varepsilon) \leq r(x)$, for all $x \in [x_{i-1}, x_i]$, because $\phi(x_{i-1}) = \phi_h^\varepsilon(x_{i-1})$ and $\phi(x_i) = \phi_h^\varepsilon(x_i)$. From this property, we derive the maximum norm estimates of local error in each subinterval.

For each $x \in (x_{i-1}, x_i)$, we have

$$\begin{aligned} \bar{L}(\pm(\phi - \phi_h^\varepsilon)) &= \bar{L}(\pm\phi) - \bar{L}(\pm\phi_h^\varepsilon) \\ &= \pm(-\varepsilon\phi'' - \bar{b}\phi' + \bar{c}\phi) \\ &= \pm\{g - (\bar{b} - b)\phi' + (\bar{c} - c)\phi\} \\ &\leq \|g\|_\infty + \|\bar{b} - b\|_\infty|\phi'| + \|\bar{c} - c\|_\infty\|\phi\|_\infty \\ &\leq \|g\|_\infty + \|b'\|_\infty|\phi'|h + \|c'\|_\infty\|\phi\|_\infty h. \end{aligned}$$

Hence we get from Lemma 2.3

$$\begin{aligned} \bar{L}(\pm(\phi - \phi_h^\varepsilon)) &\leq \left(\frac{1}{\varepsilon}C_1 + C_2\right) e^{-\frac{\beta}{2\varepsilon}x} \|b'\|_\infty h \|g\|_\infty \\ &\quad + [1 + (C_3\|b'\|_\infty + K\|c'\|_\infty)h] \|g\|_\infty. \end{aligned}$$

Now, for each $\lambda \in [0, 1]$, define $r(x)$ by

$$r(x) \equiv \tau_1 e^{-\frac{\beta}{2\varepsilon}x} h \|g\|_\infty + \tau_2 [h + \lambda x_{i-1} + (1 - \lambda)x_i - x] \|g\|_\infty. \quad (2.12)$$

Then we have

$$\begin{aligned} \bar{L}r(x) &= \tau_1 \left(-\frac{\beta^2}{4\varepsilon} + \frac{\beta\bar{b}}{2\varepsilon} + \bar{c}\right) e^{-\frac{\beta}{2\varepsilon}x} h \|g\|_\infty \\ &\quad + \tau_2 [\bar{b} + \bar{c}(h + \lambda x_{i-1} + (1 - \lambda)x_i - x)] \|g\|_\infty \\ &\geq \tau_1 \left(\frac{\beta^2}{4\varepsilon} + \gamma\right) e^{-\frac{\beta}{2\varepsilon}x} h \|g\|_\infty + \tau_2 [\beta + (1 - \lambda)\gamma h] \|g\|_\infty. \end{aligned}$$

Hence if

$$\begin{aligned} \tau_1 &\geq \frac{4}{\beta^2 + 4\varepsilon\gamma} (C_1 + \varepsilon C_2) \|b'\|_\infty, \\ \tau_2 &\geq \frac{1}{\beta + (1 - \lambda)\gamma h} [1 + (\|b'\|_\infty C_3 + K\|c'\|_\infty)h], \end{aligned}$$

then, noting that $e^{-\frac{\beta}{\varepsilon}x} \leq e^{-\frac{\beta}{2\varepsilon}x}$, we have $\bar{L}(\pm(\phi - \phi_h^\varepsilon)) \leq \bar{L}r(x)$ on $[x_{i-1}, x_i]$. In this case, we can choose that $\lambda \in [0, 1]$ since $r(x) \geq 0$ if $\lambda = 1$.

Next, consider the case of $C_0 \neq 0$. In that case, we also define $r(x)$ by (2.12), then we obtain a positive lower bound of $r(x)$ satisfying $r(x) \geq \tau_2(1 - \lambda)h\|g\|_\infty$. Hence the function $r(x)$ satisfies the condition of the maximum principle for the corresponding error estimation if $\tau_2(1 - \lambda)h\|g\|_\infty \geq C_0 h\|g\|_\infty$. Thus, we obtain the following condition on τ_2 .

$$\tau_2 \geq \frac{C_0}{1 - \lambda}.$$

Therefore, the proof is completed since $|r(x)| \leq [\tau_1 + (2 - \lambda)\tau_2] h\|g\|_\infty$ in $[x_{i-1}, x_i]$. \blacksquare

Note that the a priori constant C in this theorem still includes the perturbation parameter ε , but one can readily see that it is essentially independent of this parameter. As a special case, we have the following corollary.

Corollary 1 *If $b(x) \equiv \beta$ and $c(x) \equiv \gamma$ are constant then*

$$\|\phi - \phi_h^\varepsilon\|_\infty \leq \frac{h}{\beta} \|g\|_\infty. \quad (2.13)$$

Moreover if $\beta \geq \gamma h$ then the estimation (2.13) attains the minimum value of $C(\lambda)$ in Theorem 2.2 with respect to all $\lambda \in [0, 1]$.

Proof: In this case, we have $\|b'\|_\infty = \|c'\|_\infty = 0$ in Theorem 2.2. Hence, we find

$$C(\lambda) = \frac{2 - \lambda}{\beta + (1 - \lambda)\gamma h}, \quad C'(\lambda) = \frac{\gamma h - \beta}{[\beta + (1 - \lambda)\gamma h]^2}.$$

Therefore $C(1) = 1/\beta$, and, moreover, $C(\lambda)$ is a monotone decreasing, if $\beta \geq \gamma h$. Thus, we have the desired conclusion immediately. \blacksquare

2.2 Reaction Diffusion Problem

We consider the following linear reaction diffusion problems.

$$\begin{cases} L\phi & \equiv & -\varepsilon\phi'' + c(x)\phi & = & g & \text{ in } & (0, 1), \\ \phi(0) = \phi(1) & = & 0, \end{cases} \quad (2.14)$$

where $g \in L^\infty(0, 1)$ and $c(x) \in W_\infty^1(0, 1)$ with $c(x) \geq \gamma > 0$.

For all $\varphi, \psi \in H_0^1(0, 1)$, we define the bilinear form associated with (2.14) by

$$\begin{aligned} a(\varphi, \psi) & \equiv \varepsilon(\varphi', \psi') + (c\varphi, \psi), \\ a_h(\varphi, \psi) & \equiv \varepsilon(\varphi', \psi') + (\bar{c}\varphi, \psi). \end{aligned}$$

Then, the projection $P_h : H_0^1 \rightarrow S_h$ is defined as

$$a(\phi - P_h\phi, \psi_h) = 0, \quad \text{for all } \psi_h \in S_h. \quad (2.15)$$

And we also define the approximation $P_h^\varepsilon\phi \equiv \phi_h^\varepsilon \in S_h$ of solution ϕ to (2.14), which we call the P_h^ε -projection, as follows :

$$a_h(\phi_h^\varepsilon, \psi_h) = a(\phi, \psi_h), \quad \text{for all } \psi_h \in S_h. \quad (2.16)$$

Then we define the basis $\{\varphi_i\}_{i=1}^n$ of S_h , \bar{L} -spline, by the solutions of the problems for $i = 1, \dots, n$

$$\begin{aligned} -\varepsilon\varphi_i'' + \bar{c}\varphi_i & = 0 & \text{ in } & [0, 1] \setminus \{x_1, \dots, x_n\}, \\ \varphi_i(x_k) & = \delta_i^k & \text{ for } & k = 0, \dots, n+1. \end{aligned}$$

Remark 2 *Note that the linear operator in (2.14) also satisfies the maximum principle. (see [4] as $b(x) \equiv 0$)*

As preciously, we define the Green function $G_i = G(x, x_i)$ which is spanned by $\{\varphi_i\}_{i=1}^n$ by the solution of following equation.

$$a_h(w, G_i) = w(x_i) \quad \text{for all } w \in H_0^1(0, 1). \quad (2.17)$$

We have the following equivalent formulation of G_i .

Lemma 2.4 For each $i \in \{1, \dots, n\}$, the Green function $G_i(\cdot) \in C[0, 1]$ is characterized by

$$-\varepsilon G_i''(x) + \bar{c} G_i(x) = 0 \quad \text{in } [0, 1] \setminus \{x_1, \dots, x_n\}, \quad (2.18)$$

$$G_i(0) = G_i(1) = 0, \quad (2.19)$$

$$\lim_{x \rightarrow x_k^+} (\varepsilon G_i'(x)) - \lim_{x \rightarrow x_k^-} (\varepsilon G_i'(x)) = -\delta_i^k, \quad (2.20)$$

where the notation that x_k^- and x_k^+ are the same as the one which was defined in Remark 1. The above $G_i(x)$ is well-defined and lies in S_h with $G_i(x) \geq 0$. Moreover let $R = (R_{k,j})$ be a matrix with $R_{k,j} = a_h(\varphi_k, \varphi_j)$, ($1 \leq k, j \leq N$) then $\|G_i\|_\infty \leq \|R^{-1}\|_{\ell_\infty}$ where $\|\cdot\|_{\ell_\infty}$ means matrix maximum norm.

Proof: For each $i \in \{1, \dots, n\}$, we set

$$G_i = \sum_{j=1}^n \alpha_j^i \varphi_j.$$

Then (2.17) is equivalent to the following linear system.

$$\sum_{j=1}^n \alpha_j^i a_h(\varphi_k, \varphi_j) = \delta_i^k, \quad k = 1, \dots, n. \quad (2.21)$$

Observing that

$$\begin{aligned} R_{k,j} = a_h(\varphi_k, \varphi_j) &= \varepsilon(\varphi_k', \varphi_j') + \bar{c}(\varphi_k, \varphi_j) \\ &= \sum_{l=1}^{n+1} \varepsilon[\varphi_k \varphi_j']_{x_{l-1}}^{x_l} + (\varphi_k, -\varepsilon \varphi_j'' + \bar{c} \varphi_j) \\ &= \sum_{l=1}^{n+1} \varepsilon[\varphi_k \varphi_j']_{x_{l-1}}^{x_l}, \end{aligned}$$

and noting that R is a tri-diagonal matrix from the property of base functions, we have

$$\begin{aligned} R_{k,k} &= \varepsilon[\varphi_k \varphi_k']_{x_{k-1}}^{x_k} + \varepsilon[\varphi_k \varphi_k']_{x_k}^{x_{k+1}} = \varepsilon \varphi_k'(x_k^-) - \varepsilon \varphi_k'(x_k^+) > 0, \\ R_{k,k-1} &= \varepsilon[\varphi_k \varphi_{k-1}']_{x_{k-1}}^{x_k} = \varepsilon \varphi_{k-1}'(x_k^-) < 0, \\ R_{k,k+1} &= \varepsilon[\varphi_k \varphi_{k+1}']_{x_k}^{x_{k+1}} = -\varepsilon \varphi_{k+1}'(x_k^+) < 0. \end{aligned} \quad (2.22)$$

From

$$\varphi_k(x) = \begin{cases} \sinh\left(\sqrt{\frac{\bar{c}}{\varepsilon}}(x - x_{k-1})\right) / \sinh\left(\sqrt{\frac{\bar{c}}{\varepsilon}}h_k\right) & \text{if } x \in [x_{k-1}, x_k] \\ \sinh\left(\sqrt{\frac{\bar{c}}{\varepsilon}}(x_{k+1} - x)\right) / \sinh\left(\sqrt{\frac{\bar{c}}{\varepsilon}}h_{k+1}\right) & \text{if } x \in [x_k, x_{k+1}] \\ 0 & \text{otherwise} \end{cases} \quad (2.23)$$

we have $\varphi'_{k+1}(x_k^-) = \varphi'_{k-1}(x_k^+) = 0$. Hence (2.21) can be rewritten as

$$\begin{aligned}
\delta_i^k &= \alpha_{k-1}^i R_{k,k-1} + \alpha_k^i R_{k,k} + \alpha_{k+1}^i R_{k,k+1} \\
&= \varepsilon \left(\alpha_{k-1}^i \varphi'_{k-1} + \alpha_k^i \varphi'_k \right) (x_k^-) - \varepsilon \left(\alpha_k^i \varphi'_k + \alpha_{k+1}^i \varphi'_{k+1} \right) (x_k^+) \\
&= \varepsilon \left(\alpha_{k-1}^i \varphi'_{k-1} + \alpha_k^i \varphi'_k + \alpha_{k+1}^i \varphi'_{k+1} \right) (x_k^-) - \varepsilon \left(\alpha_{k-1}^i \varphi'_{k-1} + \alpha_k^i \varphi'_k + \alpha_{k+1}^i \varphi'_{k+1} \right) (x_k^+) \\
&= \varepsilon G_i'(x_k^-) - \varepsilon G_i'(x_k^+).
\end{aligned}$$

Therefore it follows that conditions (2.18)-(2.20) are equivalent to (2.17).

Next, we show that the coefficient matrix R is an M-matrix [1]. This can be easily proved as below by the fact that R is a symmetric and Z-matrix (all off-diagonal elements are nonpositive) from the definition of $a_h(\cdot, \cdot)$ and (2.22). Since

$$R_{k-1,k} = -\varepsilon \varphi'_k(x_{k-1}^+), \quad R_{k+1,k} = \varepsilon \varphi'_k(x_{k+1}^-),$$

we obtain

$$\begin{aligned}
R_{k-1,k} + R_{k,k} + R_{k+1,k} &= -\varepsilon \varphi'_k(x_{k-1}^+) + \varepsilon \varphi'_k(x_k^-) - \varepsilon \varphi'_k(x_k^+) + \varepsilon \varphi'_k(x_{k+1}^-) \\
&= \left(\int_{x_k}^{x_{k+1}} + \int_{x_{k-1}}^{x_k} \right) \varepsilon \varphi_k''(t) dt \\
&= \left(\int_{x_k}^{x_{k+1}} + \int_{x_{k-1}}^{x_k} \right) \bar{c} \varphi_k(t) dt \\
&\geq \left(\int_{x_k}^{x_{k+1}} + \int_{x_{k-1}}^{x_k} \right) \gamma \varphi_k(t) dt \\
&> 0.
\end{aligned}$$

Hence we have that R is a strictly diagonally dominant matrix, which means R is an M-matrix. Thus, $G_i(x)$ is well-defined and lies in S_h with $G_i(x) \geq 0$, because M-matrix is nonsingular, all elements of its inverse matrix are nonnegative and the components of the right-hand side of (2.21) are nonnegative.

Next we show that $G_i(x)$ attains the maximum value at the end point on each subinterval.

Let

$$r_k(x) = \begin{cases} \frac{1}{h_k}(x - x_{k-1}) - \varphi_k(x) & \text{if } x \in [x_{k-1}, x_k] \\ \frac{1}{h_{k+1}}(x_{k+1} - x) - \varphi_k(x) & \text{if } x \in [x_k, x_{k+1}] \\ 0 & \text{otherwise} \end{cases}$$

for $k = 1, \dots, n$. Then, since $r_k(x_{i-1}) = r_k(x_i) = r_k(x_{i+1}) = 0$ and $\bar{L}r_k(x) > 0$ in each subinterval, we obtain $r_k(x) \geq 0$ from the maximum principle. And each Green's function satisfies $G_i(x) = \alpha_k^i \varphi_k(x) + \alpha_{k+1}^i \varphi_{k+1}(x)$ on each subinterval. Therefore it follows that

$$\begin{aligned}
|\alpha_k^i \varphi_k(x) + \alpha_{k+1}^i \varphi_{k+1}(x)| &\leq |\alpha_k^i| \varphi_k(x) + |\alpha_{k+1}^i| \varphi_{k+1}(x) \\
&\leq \max(|\alpha_k^i|, |\alpha_{k+1}^i|) \cdot (\varphi_k(x) + \varphi_{k+1}(x)) \\
&\leq \max(|\alpha_k^i|, |\alpha_{k+1}^i|),
\end{aligned}$$

because $r_k(x) \geq 0$. Thus, we get the following estimate.

$$\|G_i\|_\infty = \max_k |(R^{-1})_{k,i}| \leq \max_k \max_j |(R^{-1})_{k,j}| \leq \max_k \sum_{j=1}^n |(R^{-1})_{k,j}| = \|R^{-1}\|_{\ell_\infty},$$

Therefore the proof is completed. ■

Theorem 2.3 *Let ϕ and ϕ_h^ε be solutions of (2.14) and (2.16), respectively. Then*

$$|(\phi - \phi_h^\varepsilon)(x_i)| \leq C_0 h \|g\|_\infty, \quad i = 1, 2, \dots, n, \quad (2.24)$$

where $C_0 = 1/\gamma \|R^{-1}\|_{\ell_\infty} \|c'\|_\infty$, where R is the matrix same as in Lemma 2.4.

Proof: From the property of Green's function, we have

$$\begin{aligned} (\phi - \phi_h^\varepsilon)(x_i) &= a_h(\phi - \phi_h^\varepsilon, G_i) \\ &= (\phi, (\bar{c} - c)G_i). \end{aligned}$$

Hence the nodal errors can be estimated as

$$\begin{aligned} |(\phi - \phi_h^\varepsilon)(x_i)| &\leq \|\phi\|_\infty \|\bar{c} - c\|_{L_1} \|G_i\|_\infty \\ &\leq \frac{1}{\gamma} \|R^{-1}\|_{\ell_\infty} \|c'\|_\infty h \|g\|_\infty. \end{aligned}$$

■

For the numerical computation of $\|R^{-1}\|_{\ell_\infty}$ with guaranteed error bound, refer, e.g., [3].

Theorem 2.4 *Let ϕ and ϕ_h^ε be solutions of (2.14) and (2.16), respectively. Then*

$$\|\phi - \phi_h^\varepsilon\|_\infty \leq C(h, \varepsilon) \|g\|_\infty$$

where

$$\begin{aligned} C(h, \varepsilon) &= \max \left\{ \frac{1}{\gamma} \left(1 + \frac{1}{\gamma} \|c'\|_\infty h \right) \delta(h, \varepsilon) + 2e^{-\sqrt{\frac{\gamma}{\varepsilon}} \frac{h_{\min}}{2}} C_0 h, \ C_0 h \right\}, \\ \delta(h, \varepsilon) &\equiv \left(1 - e^{-\sqrt{\frac{\gamma}{\varepsilon}} \frac{h}{2}} \right)^2 < 1. \end{aligned}$$

Here, C_0 is same as in Theorem 2.3.

Proof: We use the maximum principle like the convection diffusion case. We now constitute a function $r(x)$ which satisfies with $\bar{L}(\pm(\phi - \phi_h^\varepsilon)) \leq \bar{L}r(x)$ and $\pm(\phi - \phi_h^\varepsilon)(x_{i-1}) \leq r(x_{i-1})$, $\pm(\phi - \phi_h^\varepsilon)(x_i) \leq r(x_i)$ in $[x_{i-1}, x_i]$ for $i = 1, \dots, n$ as below.

We first estimate $\bar{L}(\pm(\phi - \phi_h^\varepsilon))$ as follows.

$$\begin{aligned} \bar{L}(\pm(\phi - \phi_h^\varepsilon)) &= \bar{L}(\pm\phi) - \bar{L}(\pm\phi_h^\varepsilon) = \pm(-\varepsilon\phi'' + \bar{c}\phi) = \pm\{g + (\bar{c} - c)\phi\} \\ &\leq \|g\|_\infty + \|\bar{c} - c\|_\infty \|\phi\|_\infty \\ &\leq \|g\|_\infty + \|c'\|_\infty \|\phi\|_\infty h \\ &\leq \left(1 + \frac{1}{\gamma} \|c'\|_\infty h \right) \|g\|_\infty, \quad (=:\mathbf{g}) \end{aligned}$$

where we have used the estimate $\|\phi\|_\infty \leq 1/\gamma \|g\|_\infty$ which is following by the similar argument to that in the proof of Lemma 2.3. Here we solve the following ordinary differential equation with $\hat{C}_0 := C_0 h \|g\|_\infty$.

$$\begin{aligned} -\varepsilon r''(x) + \bar{c}r(x) &= \mathbf{g} \quad \text{in } (x_{i-1}, x_i), \\ r(x_{i-1}) = r(x_i) &= \hat{C}_0. \end{aligned} \quad (2.25)$$

Then the solution of (2.25) is written as

$$r(x) = \frac{1}{1 + e^{-\sqrt{\frac{\varepsilon}{h}}}} \left[\frac{\mathbf{g}}{\hat{c}} \left(1 - e^{\sqrt{\frac{\varepsilon}{h}}(x_{i-1}-x)} - e^{\sqrt{\frac{\varepsilon}{h}}(x-x_i)} + e^{-\sqrt{\frac{\varepsilon}{h}}h} \right) + \hat{C}_0 \left(e^{\sqrt{\frac{\varepsilon}{h}}(x_{i-1}-x)} + e^{\sqrt{\frac{\varepsilon}{h}}(x-x_i)} \right) \right].$$

Since $r'(x) = 0$ at $x = (x_{i-1} + x_i)/2$, we obtain

$$|r(x)| \leq \frac{1}{1 + e^{-\sqrt{\frac{\varepsilon}{h}}}} \max \left\{ \frac{\mathbf{g}}{\hat{c}} \left(1 - e^{-\sqrt{\frac{\varepsilon}{h}} \frac{h_i}{2}} \right)^2 + 2e^{-\sqrt{\frac{\varepsilon}{h}} \frac{h_i}{2}} \hat{C}_0, \hat{C}_0 \right\},$$

where $h_i = x_i - x_{i-1}$. Therefore, the maximum principle completes the proof. \blacksquare

3 Numerical Verification Algorithm

The nonlinear singularly perturbed problem (1.1) is transformed to fixed point equation such that $u = F(u)$ ($:= L^{-1}f(u)$), where F becomes a compact operator from $L^\infty(0, 1) \cap H_0^1(0, 1)$ to itself. We apply the verification method similar to that in [2].

Instead of the H_0^1 -projection in [2], using the P_h -Projection defined by (2.2) and (2.15), we have the following decomposition of the fixed point equation $u = F(u)$.

$$\begin{aligned} P_h u &= P_h F(u), \\ (I - P_h)u &= (I - P_h)F(u). \end{aligned}$$

Let \hat{u}_h^ε be an approximate solution of (1.1) which satisfies $a_h(\hat{u}_h^\varepsilon, \psi_h) = (f(\hat{u}_h^\varepsilon), \psi_h)$, for all $\psi_h \in V_h$, and let

$$N_h(u) \equiv P_h u - [I - F'(\hat{u}_h^\varepsilon)]_h^{-1} (P_h u - P_h F(u)). \quad (3.1)$$

Here $[I - F'(\hat{u}_h^\varepsilon)]_h^{-1}$ is an inverse operator of $P_h(I - F'(\hat{u}_h^\varepsilon))|_{S_h} : S_h \rightarrow S_h$ and $F'(\hat{u}_h^\varepsilon)$ is a Fréchet derivative of $F(u)$ at \hat{u}_h^ε . Then (1.1) is equivalent to the following fixed point equation.

$$u = Tu, \quad Tu \equiv N_h(u) + (I - P_h)F(u).$$

Defining the candidate set, a set expected to include the desired solutions, as $U \equiv \hat{u}_h^\varepsilon + W_h + [[\alpha]]_\infty \subset L^\infty(0, 1) \cap H_0^1(0, 1)$, where

$$\begin{aligned} W_h &\equiv \left\{ w_h \in S_h : w_h = \sum_{i=1}^n W_i \varphi_i, \ W_i = [-w_i, w_i], \ w_i \geq 0 \right\}, \\ [[\alpha]]_\infty &\equiv \left\{ \hat{\alpha} \in L^\infty(0, 1) \cap H_0^1(0, 1) : \|\hat{\alpha}\|_\infty \leq \alpha \right\}, \end{aligned}$$

if the condition

$$N_h(U) \subset U_h, \quad (3.2)$$

$$(I - P_h)F(U) \subset [[\alpha]]_\infty \quad (3.3)$$

holds, then, by Schauder's fixed point theorem, there exists a function $\hat{u} \in U$ such that $L\hat{u} = f(\hat{u})$.

In order to estimate the error $\|\phi - P_h\phi\|_\infty$, we use following triangle inequality

$$\|\phi - P_h\phi\|_\infty \leq \|\phi - P_h^\varepsilon\phi\|_\infty + \|P_h^\varepsilon\phi - P_h\phi\|_\infty. \quad (3.4)$$

For the first term, we use the estimation already discussed in the previous section. For the second term, by the fact that $a(P_h\phi, \psi_h) = a_h(P_h^\varepsilon\phi, \psi_h)$, for all $\psi_h \in V_h$, it is essentially independent of ε , because of Lemma 2.3, Theorems 2.1 and 2.3. Actually, it is easily seen that, by using matrices $A = (A_{i,j}) = [a(\varphi_j, \psi_i)]$ and $A^h = (A_{i,j}^h) = [a_h(\varphi_j, \psi_i)]$, we have $\|P_h^\varepsilon\phi - P_h\phi\|_\infty \leq \mathbf{C}h\|g\|_\infty$, where $\mathbf{C} := 3\|A^{-1} - (A^h)^{-1}\|_{\ell_\infty}$, which is followed by the properties of base functions of S_h, V_h .

From the viewpoint of the effectiveness of computational cost, usually, we use the residual form below instead of the original equation (1.1) [9].

That is, for the nonlinear singularly perturbed problem (1.1), we define the solution \bar{u} of the following *linear* singularly perturbed problem.

$$\begin{aligned} -\varepsilon\bar{u}'' - b(x)\bar{u}' + c(x)\bar{u} &= f(\hat{u}_h^\varepsilon) \quad \text{in } (0, 1), \\ \bar{u}(0) = \bar{u}(1) &= 0. \end{aligned}$$

Then, defining $v_0 \equiv \bar{u} - \hat{u}_h^\varepsilon$ we get the following estimates

$$\|v_0\|_\infty \leq Ch\|f(\hat{u}_h^\varepsilon)\|_\infty$$

by the same constant C in Theorems 2.2 and 2.4, because \hat{u}_h^ε coincides with P_h^ε -projection of \bar{u} . Thus the concerned problem is reduced to find $w := u - \bar{u}$ satisfying

$$\begin{aligned} -\varepsilon w'' - b(x)w' + c(x)w &= f(w + v_0 + \hat{u}_h^\varepsilon) - f(\hat{u}_h^\varepsilon) \quad \text{in } (0, 1), \\ w(0) = w(1) &= 0. \end{aligned} \quad (3.5)$$

Then, since the approximate solution of (3.5) is taken as 0, the candidate set for the solution is usually taken of the form $W \equiv W_h + [[\alpha]]_\infty$.

4 Numerical Examples

We now present some numerical examples, to show the effectiveness of the Theorems 2.2 and 2.4. In the below, the verifications were carried out by using the residual form (3.5).

We first consider the following example of the convection diffusion problem.

Example 4.1

$$\begin{aligned} L_1 u &\equiv -\varepsilon u'' - bu' + cu = 1 - u^3 \quad \text{in } (0, 1), \\ u(0) = u(1) &= 0, \end{aligned}$$

where $b = 1/(2\pi)^2$, $c = 1$.

We omit detailed and actual computational procedures for checking conditions (3.2) and (3.3) (see, e.g., [2][9] etc.). The approximate solutions are shown in Figure 1.

Table 1 shows the verification results for uniform mesh with $n = 999$ using \bar{L} -spline in this paper. We also illustrate, for comparison, in Tables 2 to 3 the results using the usual verification algorithm with piecewise linear functions as in [9]. We show that the distribution of mesh size for non-uniform mesh in Figure 3.

Example 4.1




Figure 1: The approximate solutions of Example 4.1.

In Tables 1 to 3 and 4 to 6, the exact solution is enclosed by $\hat{u}_h^\varepsilon + W_h + [[\alpha]]_\infty + v_0$, i.e., "**Total**"s mean the total error of the approximate solution \hat{u}_h^ε . "**Fail**"s in tables mean that we could not get the solution u with guaranteed error bound, in the sense of infinite dimension, but we could get the approximate solution with guaranteed error bound, in the sense of finite dimension.

We next consider the following example of the reaction diffusion problem.

Example 4.2 (Allen-Cahn equation)

$$\begin{aligned} L_2 u &\equiv -\varepsilon u'' + cu = (c+1)u^2 - u^3 & \text{in } (0,1), \\ u(0) &= u(1) = 0, \end{aligned}$$

where $c = 1/10$.

The approximate solutions are shown in Figure 2.

Tables 4 to 6 show the comparison of the effectiveness for \bar{L} -spline with uniform mesh and the usual piecewise linear finite element method with both uniform and nonuniform meshes for $n = 999$. Figure 3 shows that the distribution of mesh size for non-uniform mesh. By these tables, it is seen that if we use the uniformed mesh then \bar{L} -spline yields always better approximation than the usual piecewise linear finite element. "**Singular**" in Table 5 means that we could not get the approximate solution with guaranteed error bound, in the sense of finite dimension. As shown in Figure 4, the usefulness of \bar{L} -spline method should be more and more clear compared with the usual method when ε tends to be very small.

The numerical computations were carried out on a Dell Precision 650 Workstation by using INTLAB, a tool box in MATLAB developed by Rump [5] for self-validating

Table 1: \bar{L} -spline (Uniform) for Example 4.1

$1/\varepsilon$	Total	α	$\ W_h\ _\infty$	$\ v_0\ _\infty$
100	4.8819e-2	2.7863e-3	1.9576e-2	3.7979e-2
1000	4.9337e-2	2.7858e-3	1.9554e-2	3.7979e-2
10000	5.1568e-2	2.7913e-3	1.9621e-2	3.7979e-2
100000	6.0336e-2	3.0415e-3	2.0178e-2	3.7979e-2
1000000	6.3029e-2	3.7266e-3	2.1801e-2	3.7979e-2

Table 2: Piecewise Linear (Uniform) for Example 4.1

$1/\varepsilon$	Total	α	$\ W_h\ _\infty$	$\ v_0\ _\infty$
100	9.4484e-3	1.0157e-5	1.1025e-4	9.3539e-3
1000	6.4661e-2	6.6864e-4	6.1947e-4	6.3552e-2
10000	9.1327e-1	8.7964e-2	3.8504e-3	8.2299e-1
100000	Fail	∞	∞	2.0192e+1
1000000	Fail	∞	∞	5.4777e+2

Table 3: Piecewise Linear (Non-Uniform) for Example 4.1

$1/\varepsilon$	Total	α	$\ W_h\ _\infty$	$\ v_0\ _\infty$
100	5.1665e-3	5.2703e-6	8.1393e-5	5.0948e-3
1000	2.4911e-2	1.4200e-4	2.9216e-4	2.4568e-2
10000	2.7563e-1	9.4780e-3	1.2795e-3	2.6568e-1
100000	Fail	∞	∞	6.7784e-0
1000000	Fail	∞	∞	2.0396e+2

algorithms.

Acknowledgments.

This work is supported by Kyushu University 21st Century COE Program, *Development of Dynamic Mathematics with High Functionality*, of the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- [1] Axelsson O., Iterative Solution Methods, Cambridge University Press, (1994).
- [2] Nakao, M.T., Numerical verification methods for solutions of ordinary and partial differential equations, Numerical Functional Analysis and Optimization 22(3&4) (2001), 321-356.
- [3] Ogita T., Oishi S. and Ushiro Y., Fast verification of solutions for sparse monotone matrix equations, Computing [Suppl] 15 (2001), 175-187.
- [4] O’Riordan E., Stynes M., An analysis of a superconvergence result for a singularly perturbed boundary value problem, Mathematics of Computation vol. 46 number 173 (1986), 81-92.




Figure 2: The approximate solutions of Example 4.2.

- [5] Rump S. M., INTLAB-INTerval LABoratory, a Matlab toolbox for verified computations, version 4.1.2. Inst. Infomatik, Technical University Hamburg Harburg, (2002). <http://www.ti3.tu-harburg.de/rump/intlab/index.html/>
- [6] H. - G. Roos, M. Stynes, L. Tobiska, Numerical Methods for Singularly Perturbed Differential Equations, Springer, Berlin Heidelberg, (1996).
- [7] M. Stynes, E. O'Riordan, A finite element method for a singularly perturbed boundary value problem, Numerische Mathematik 50 (1986), 1-15.
- [8] M. Stynes, E. O'Riordan, An analysis of a singularly perturbed two-point boundary value problem using only finite element techniques, Mathematics of Computation vol. 56 number 194 (1991), 663-675.
- [9] Yamamoto, N., Nakao, M.T., Numerical verifications for solutions to elliptic equations using residual iterations with higher order finite element, Journal of Computational and Applied Mathematics 60 (1995), 271-279.

Table 4: \bar{L} -spline (Uniform) for Example 4.2

$1/\varepsilon$	Total	α	$\ w_h\ _\infty$	$\ v_0\ _\infty$
1000	4.1860e-5	2.8959e-9	1.7250e-5	2.4666e-5
10000	4.3532e-4	2.9477e-7	1.8943e-4	2.4645e-4
100000	4.9995e-3	3.2202e-5	2.5339e-3	2.4621e-3
400000	2.7049e-2	7.1905e-4	1.6925e-2	9.8181e-3
500000	3.9078e-2	1.3660e-3	2.6007e-2	1.2256e-2
600000	5.8141e-2	2.5107e-3	4.1667e-2	1.4691e-2

Table 5: Piecewise Linear (Uniform) for Example 4.2

$1/\varepsilon$	Total	α	$\ w_h\ _\infty$	$\ v_0\ _\infty$
1000	4.1922e-5	2.4793e-9	2.6180e-5	1.5934e-5
10000	4.6950e-4	2.8032e-7	3.0774e-4	1.6279e-4
100000	6.7719e-3	4.4062e-5	5.0283e-3	1.7376e-3
400000	Singular	—	—	—
500000	Fail	∞	∞	9.8976e-3
600000	Fail	∞	∞	1.2015e-2

Table 6: Piecewise Linear (Non-Uniform) for Example 4.2

$1/\varepsilon$	Total	α	$\ w_h\ _\infty$	$\ v_0\ _\infty$
1000	3.0688e-5	7.3430e-9	1.8028e-5	1.3748e-5
10000	5.5890e-5	4.3732e-9	3.5027e-5	2.1099e-5
100000	6.3696e-4	5.0772e-7	4.2204e-4	2.1622e-4
400000	3.0715e-3	1.0212e-5	2.1771e-3	8.9695e-4
500000	Fail	∞	∞	1.1304e-3
600000	Fail	∞	∞	1.3611e-3




Figure 3: The distribution of mesh sizes.




Figure 4: Constructive a priori constant $C(h, \varepsilon)$ for Example 4.2

List of MHF Preprint Series, Kyushu University

21st Century COE Program

Development of Dynamic Mathematics with High Functionality

- MHF2003-1 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Yoshitaka WATANABE
A numerical method to verify the invertibility of linear elliptic operators with applications to nonlinear problems
- MHF2003-2 Masahisa TABATA & Daisuke TAGAMI
Error estimates of finite element methods for nonstationary thermal convection problems with temperature-dependent coefficients
- MHF2003-3 Tomohiro ANDO, Sadanori KONISHI & Seiya IMOTO
Adaptive learning machines for nonlinear classification and Bayesian information criteria
- MHF2003-4 Kazuhiro YOKOYAMA
On systems of algebraic equations with parametric exponents
- MHF2003-5 Masao ISHIKAWA & Masato WAKAYAMA
Applications of Minor Summation Formulas III, Plücker relations, Lattice paths and Pfaffian identities
- MHF2003-6 Atsushi SUZUKI & Masahisa TABATA
Finite element matrices in congruent subdomains and their effective use for large-scale computations
- MHF2003-7 Setsuo TANIGUCHI
Stochastic oscillatory integrals - asymptotic and exact expressions for quadratic phase functions -
- MHF2003-8 Shoki MIYAMOTO & Atsushi YOSHIKAWA
Computable sequences in the Sobolev spaces
- MHF2003-9 Toru FUJII & Takashi YANAGAWA
Wavelet based estimate for non-linear and non-stationary auto-regressive model
- MHF2003-10 Atsushi YOSHIKAWA
Maple and wave-front tracking — an experiment
- MHF2003-11 Masanobu KANEKO
On the local factor of the zeta function of quadratic orders
- MHF2003-12 Hidefumi KAWASAKI
Conjugate-set game for a nonlinear programming problem

- MHF2004-1 Koji YONEMOTO & Takashi YANAGAWA
Estimating the Lyapunov exponent from chaotic time series with dynamic noise
- MHF2004-2 Rui YAMAGUCHI, Eiko TSUCHIYA & Tomoyuki HIGUCHI
State space modeling approach to decompose daily sales of a restaurant into time-dependent multi-factors
- MHF2004-3 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Cubic pencils and Painlevé Hamiltonians
- MHF2004-4 Atsushi KAWAGUCHI, Koji YONEMOTO & Takashi YANAGAWA
Estimating the correlation dimension from a chaotic system with dynamic noise
- MHF2004-5 Atsushi KAWAGUCHI, Kentarou KITAMURA, Koji YONEMOTO, Takashi YANAGAWA & Kiyofumi YUMOTO
Detection of auroral breakups using the correlation dimension
- MHF2004-6 Ryo IKOTA, Masayasu MIMURA & Tatsuyuki NAKAKI
A methodology for numerical simulations to a singular limit
- MHF2004-7 Ryo IKOTA & Eiji YANAGIDA
Stability of stationary interfaces of binary-tree type
- MHF2004-8 Yuko ARAKI, Sadanori KONISHI & Seiya IMOTO
Functional discriminant analysis for gene expression data via radial basis expansion
- MHF2004-9 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Hypergeometric solutions to the q -Painlevé equations
- MHF2004-10 Raimundas VIDŪNAS
Expressions for values of the gamma function
- MHF2004-11 Raimundas VIDŪNAS
Transformations of Gauss hypergeometric functions
- MHF2004-12 Koji NAKAGAWA & Masakazu SUZUKI
Mathematical knowledge browser
- MHF2004-13 Ken-ichi MARUNO, Wen-Xiu MA & Masayuki OIKAWA
Generalized Casorati determinant and Positon-Negaton-Type solutions of the Toda lattice equation
- MHF2004-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO
Generating function associated with the determinant formula for the solutions of the Painlevé II equation

MHF2004-15 Kouji HASHIMOTO, Ryohei ABE, Mitsuhiro T. NAKAO & Yoshitaka
WATANABE
Numerical verification methods of solutions for nonlinear singularly perturbed
problem