

## Generating function related to the Okamoto polynomials for the Painlevé IV equation

Goto, Hiromichi  
Graduate School of Mathematics, Kyushu University

Kajiwara, Kenji  
School of Mathematics and Statistics F07, The University of Sydney

<https://hdl.handle.net/2324/11850>

---

出版情報 : Bulletin of the Australian Mathematical Society. 71, pp.517-526, 2005. Australian Mathematical Society

バージョン :

権利関係 :



# MHF Preprint Series

Kyushu University  
21st Century COE Program  
Development of Dynamic Mathematics with  
High Functionality

## Generating function related to the Okamoto polynomials for the Painlevé IV equation

H. Goto & K. Kajiwara

MHF 2005-13

( Received March 29, 2005 )

Faculty of Mathematics  
Kyushu University  
Fukuoka, JAPAN

# Generating functions related to the Okamoto polynomials for the Painlevé IV equation

Hiromichi Goto<sup>1</sup> and Kenji Kajiwara<sup>1,2</sup>

<sup>1</sup> Graduate School of Mathematics, Kyushu University, 6-10-1 Hakozaki, Fukuoka 812-8581, Japan

<sup>2</sup> School of Mathematics and Statistics F07, The University of Sydney, Sydney 2006, Australia

March 23, 2005

## Abstract

We construct generating functions for the entries of Hankel determinant formula for the Okamoto polynomials which characterize a class of rational solutions to the Painlevé IV equation. Generating functions are characterized as asymptotic expansions of log derivative of Ai and Bi, which are solutions of the Airy equation.

2000 Mathematics Subject Classification: 34M55, 34E05, 33C10.

## 1 Introduction

In the theory of integrable systems, structure of exact solutions are considered to reflect essential mathematical structures behind the equations. In the soliton theory, as clarified by the Sato theory[12], the dependent variables which are introduced intuitively in the Hirota's method ( $\tau$  functions) are regarded as the most fundamental object, and determinant or Pfaffian structure of the  $\tau$  functions is reflection of the solution space and the transformation group acting on it. For the Painlevé equations, it is considered that the  $\tau$  functions and their determinant structure are of the same importance as the soliton equations, although the Painlevé equations are ordinary differential equations while the soliton equations are partial differential equations. In the series of works[15, 16, 17, 18], Okamoto has introduced the  $\tau$  functions through the Hamiltonians, and studied their properties. He has shown that the  $\tau$  functions satisfy the Toda equation in general, and that some class of transcendental classical solutions (hypergeometric solutions, or special function solutions) can be expressible in terms of determinants whose entries are given in terms of special functions of hypergeometric type. Moreover, for algebraic (or rational) solutions, which are another class of classical solutions to the Painlevé equations, it has been revealed that the special polynomials that characterize the solutions are specialization of the Schur functions or their generalization (universal characters) by studying the determinant formula of Jacobi-Trudi type [4, 7, 8, 10, 11, 14].

For example, the Painlevé II equation(P<sub>II</sub>)

$$\frac{d^2u}{dx^2} = 2u^3 - 4xu + 4\left(\alpha + \frac{1}{2}\right), \quad (1)$$

admits a one-parameter family of transcendental classical solution expressible in terms of the solution of the Airy equation when  $\alpha$  is an integer, and rational solutions when  $\alpha$  is a half-integer. As for the rational solutions, the following fact is known: let  $T_N$  ( $N \in \mathbb{Z}$ ) be the polynomials which are generated by the Toda equation

$$T_N''T_N - (T_N')^2 = T_{N+1}T_{N-1} - xT_N^2, \quad T_0 = 1, \quad T_1 = x, \quad ' = \frac{d}{dx}. \quad (2)$$

Then we have  $T_N = T_{-N-1}$ , and it is shown that the function

$$u = \frac{d}{dx} \log \frac{T_{N+1}}{T_N}, \quad (3)$$

satisfy  $P_{II}$  (1) with  $\alpha = N + \frac{1}{2}$ . The polynomials  $T_N$  is called the Yablonski-Vorob'ev (Y-V) polynomials. It is known that the Y-V polynomials admit the following determinant formula of Jacobi-Trudi type[7]:

$$T_N = c_N \begin{vmatrix} p_N & p_{N+1} & \cdots & p_{2N-1} \\ p_{N-2} & p_{N-1} & \cdots & p_{2N-3} \\ \vdots & \vdots & \ddots & \vdots \\ p_{-N+1} & p_{-N+2} & \cdots & p_1 \end{vmatrix}, \quad \sum_{k=0}^{\infty} p_k \lambda^k = \exp \left[ x\lambda + \frac{\lambda^3}{3} \right], \quad p_n = 0 \ (n < 0), \quad (4)$$

$$c_N = (2N - 1)!! = (2N - 1)!!(2N - 3)!! \cdots 3!! \quad (5)$$

Since the Schur function  $S_Y(t_1, t_2, t_3, \dots)$  associated with the 2-core Young diagram  $Y = (N, N - 1, \dots, 1)$  is given by the determinant in (4) with  $p_k$  replaced by  $\sum_{k=0}^{\infty} p_k \lambda^k = \exp \sum_{n=1}^{\infty} t_n \lambda^n$  [12], the above formula immediately implies that  $T_N$  are nothing but the specialization of 2-core Schur functions. Furthermore, since the 2-core Schur functions give rational solutions to KdV or modified KdV equation[12], this coincides with the fact that  $P_{II}$  can be derived from similarity reduction of the modified KdV equation(see, for example, [1]). In this manner, the determinant formula is useful to clarify the essential property of the object.

On the other hand, it is known that  $T_N$  admits the following Hankel determinant formula[7]:

$$T_N = \begin{vmatrix} a_0 & a_1 & \cdots & a_{N-1} \\ a_1 & a_2 & \cdots & a_N \\ \vdots & \vdots & \ddots & \vdots \\ a_{N-1} & a_N & \cdots & a_{2N-2} \end{vmatrix}, \quad a_0 = x, \quad a_n = a'_{n-1} + \sum_{k=0}^{n-2} a_k a_{n-2-k}. \quad (6)$$

If a determinant formula reflects some mathematical structure, what does this formula imply? In particular, what are the entries  $a_n$ ? In order to investigate this problem, a generating function for  $a_n$  was considered in [2], and the following result was obtained:

**Theorem 1.1** [2] *We define the generating function  $F(x, t)$  of  $a_n$  ( $n \geq 0$ ) by*

$$F(x, t) = \sum_{n=0}^{\infty} a_n (-2t)^{-n}. \quad (7)$$

*Then, the following asymptotic expansion around  $t \rightarrow \infty$  holds for any subsector of  $|\arg t| < \frac{\pi}{2}$ :*

$$F(x, t) \sim \frac{\partial}{\partial t} \log \theta(x, t), \quad \theta(x, t) = e^{\frac{2t^3}{3}} \text{Ai}(t^2 - x), \quad (8)$$

where  $\text{Ai}$  is the Airy function.

This result is quite suggestive, since log derivative of the Airy function is another particular solution to  $P_{II}$ . In fact,

$$u = \frac{d}{dx} \log \text{Ai}(2^{1/3}x), \quad (9)$$

satisfies  $P_{II}$  (1) with  $\alpha = 0$ . Is this phenomenon caused by an accident? In order to answer this question, it may be an important problem to investigate whether similar phenomena take place for other Painlevé equations or not.

In this article, we consider the Painlevé IV equation ( $P_{IV}$ )

$$\frac{d^2 y}{dx^2} = \frac{1}{2y} \left( \frac{dy}{dx} \right)^2 + \frac{3}{2} y^3 + 6xy + \frac{9}{2} \left( x^2 - \frac{4}{3} \alpha \right) y - \frac{\beta}{2y}, \quad (10)$$

where  $\alpha$  and  $\beta$  are parameters, and the Okamoto polynomials which characterize a class of rational solutions for  $P_{IV}$ . We construct the Hankel determinant formula for the Okamoto polynomials and discuss the generating functions for the entries of determinants.

## 2 Okamoto polynomials and Hankel determinant formula

Let  $Q_n$  ( $n \in \mathbb{Z}$ ) be polynomials generated by the following Toda equation:

$$Q_n'' Q_n - (Q_n')^2 = Q_{n+1} Q_{n-1} - (x^2 + 2n + 1) Q_n^2, \quad (11)$$

$$Q_0 = 1, \quad Q_1 = x^2 + 1. \quad (12)$$

Then, it is known that [17, 13]

$$y = \frac{d}{dx} \log \frac{Q_{n+1}}{Q_n} - x, \quad (13)$$

satisfies  $P_{IV}$  (10) with  $\alpha = n + 1, \beta = 1$ . The polynomials  $Q_n$  are called the Okamoto polynomials. A Jacobi-Trudi type determinant formula for  $Q_n$  is discussed in [8, 14, 13], and it is shown that  $Q_n$  are specialization of 3-core Schur functions.

Let us construct the Hankel determinant formula for  $Q_n$ . A Hankel determinant formula for the general solution of the Toda equation has been presented in [6] as follows:

**Theorem 2.1** [6] *Let  $\{\tau_n\}_{n \in \mathbb{Z}}$  functions satisfying the Toda equation:*

$$\tau_n'' \tau_n - \tau_n'^2 = \tau_{n+1} \tau_{n-1} - \psi \varphi \tau_n^2, \quad \tau_{-1} = \psi, \quad \tau_0 = 1, \quad \tau_1 = \varphi. \quad (14)$$

*Let  $\{a_n\}_{n \in \mathbb{N}}, \{b_n\}_{n \in \mathbb{N}}$  be sequences defined by the recursion relation*

$$a_n = a'_{n-1} + \psi \sum_{k=0}^{n-2} a_k a_{n-k}, \quad b_n = b'_{n-1} + \varphi \sum_{k=0}^{n-2} b_k b_{n-k}, \quad a_0 = \varphi, \quad b_0 = \psi, \quad (15)$$

*respectively. Then  $\tau_n$  is expressed as*

$$\tau_n = \begin{cases} \det(a_{i+j-2})_{i,j \leq n} & n > 0, \\ 1 & n = 0, \\ \det(b_{i+j-2})_{i,j \leq |n|} & n < 0. \end{cases} \quad (16)$$

Applying Theorem 2.1 to the Toda equation for  $Q_n$  (11), we have the following formula:

**Proposition 2.2**

$$Q_n = \begin{cases} \det(a_{i+j-2})_{1 \leq i,j \leq n}, & n \geq 1 \\ 1 & n = 0, -1 \\ \det(b_{i+j-2})_{1 \leq i,j \leq |n|-1}, & n \leq -1 \end{cases} \quad (17)$$

$$a_n = a'_{n-1} + 2x a_{n-1} + \sum_{k=0}^{n-2} a_k a_{n-2-k}, \quad a_0 = x^2 + 1, \quad a_1 = 2x^3 + 4x, \quad (18)$$

$$b_n = b'_{n-1} - 2x b_{n-1} + \sum_{k=0}^{n-2} b_k b_{n-2-k}, \quad b_0 = x^2 - 1, \quad b_1 = 2x^3 - 4x. \quad (19)$$

*Proof* For technical reason, we separate the cases  $n > 0$  and  $n < 0$ . Putting  $Q_n = e^{-nx^2} R_n$  in (11), we have

$$\begin{aligned} R_n'' R_n - (R_n')^2 &= R_{n+1} R_{n-1} - (x^2 + 1) R_n^2, \\ R_{-1} &= e^{-x^2}, \quad R_0 = 1, \quad R_1 = e^{x^2} (x^2 + 1). \end{aligned} \quad (20)$$

Applying Theorem 2.1 to (20) with  $n > 0$ , we have

$$R_n = \det(\tilde{a}_{i+j-2}), \quad \tilde{a}_n = \tilde{a}'_{n-1} + e^{-x^2} \sum_{k=0}^{n-2} \tilde{a}_k \tilde{a}_{n-k}, \quad \tilde{a}_0 = e^{x^2} (x^2 + 1). \quad (21)$$

Now putting  $\tilde{a}_n = e^{x^2} a_n$  and noticing that  $R_n = e^{nx^2} Q_n$ , we obtain the formula for  $n > 0$  as

$$Q_n = \det(a_{i+j-2}), \quad a_n = a'_{n-1} + 2x a_{n-1} + \sum_{k=0}^{n-2} a_k a_{n-k}, \quad a_0 = x^2 + 1. \quad (22)$$

For the case of  $n < 0$ , putting  $Q_n = e^{-(n+1)x^2} S_{n+1}$ , we have

$$\begin{aligned} S_n'' S_n - (S_n')^2 &= S_{n+1} S_{n-1} - (x^2 - 1) S_n^2, \\ S_{-1} &= e^{-x^2} (x^2 - 1), \quad S_0 = 1, \quad S_1 = e^{x^2}. \end{aligned} \quad (23)$$

Applying Theorem 2.1 to (23) with  $n < 0$ , we get the formula for  $n < 0$  by similar calculations.  $\square$

### 3 Generating functions

#### 3.1 Main result

Let  $I_\nu(z)$  be the modified Bessel function

$$I_\nu(z) = \sum_{k=0}^{\infty} \frac{(z/2)^{\nu+2k}}{\Gamma(k+1)\Gamma(k+\nu+1)}, \quad |z| < \infty, \quad |\arg z| < \pi. \quad (24)$$

We define the functions  $\text{Ai}(z)$  and  $\text{Bi}(z)$  by

$$\text{Ai}(z) = \frac{z^{1/2}}{3} \left[ I_{-1/3} \left( \frac{2z^{3/2}}{3} \right) - I_{1/3} \left( \frac{2z^{3/2}}{3} \right) \right], \quad (25)$$

$$\text{Bi}(z) = \frac{z^{1/2}}{3} \left[ I_{-1/3} \left( \frac{2z^{3/2}}{3} \right) + I_{1/3} \left( \frac{2z^{3/2}}{3} \right) \right], \quad (26)$$

respectively. It is known that  $\text{Ai}(z)$  and  $\text{Bi}(z)$  are independent solutions of the Airy equation

$$\frac{d^2 u}{dz^2} = zu, \quad (27)$$

and that they admit the following asymptotic expansion around  $z \sim \infty$  [9]:

$$\begin{aligned} \text{Ai}(z) &\sim U_-(z) \quad |\arg z| < \pi, \\ \text{Bi}(z) &\sim U_+(z) \quad |\arg z| < \frac{\pi}{3}, \end{aligned} \quad (28)$$

where

$$U_{\pm}(z) = \frac{1}{2\sqrt{\pi}} \exp\left(\pm \frac{2z^{3/2}}{3}\right) z^{-1/4} \sum_{n=0}^{\infty} \frac{\left(\frac{1}{6}\right)_n \left(\frac{5}{6}\right)_n}{n!} \left(\pm \frac{4z^{3/2}}{3}\right)^{-n}. \quad (29)$$

Now we define the generating functions for the entries  $a_n$  and  $b_n$  of the Hankel determinant formula of  $Q_n$  obtained in Proposition 2.2 by

$$F(x, t) = \sum_{n=0}^{\infty} a_n t^{-3n}, \quad G(x, t) = \sum_{n=0}^{\infty} b_n t^{-3n}. \quad (30)$$

Then the main result of this article is stated as follows:

**Theorem 3.1** *Define the functions  $\theta_1(x, t)$ ,  $\theta_2(x, t)$  by*

$$\begin{cases} \theta_1(x, t) = \text{Bi}\left(\frac{t^4}{4} - xt\right) \exp\left(-\frac{t^6}{12} + \frac{xt^3}{2}\right), \\ \theta_2(x, t) = \text{Ai}\left(\frac{t^4}{4} + xt\right) \exp\left(\frac{t^6}{12} + \frac{xt^3}{2}\right), \end{cases} \quad (31)$$

*respectively. Then, the following asymptotic expansions hold around  $t \sim \infty$ :*

$$F(x, t) \sim \frac{t^4}{x - t^3} \left[ \frac{\partial}{\partial t} \log \theta_1(x, t) - \frac{x^2}{t} \right], \quad |\arg t| < \frac{\pi}{12}, \quad (32)$$

$$G(x, t) \sim \frac{t^4}{x + t^3} \left[ \frac{\partial}{\partial t} \log \theta_2(x, t) + \frac{x^2}{t} \right], \quad |\arg t| < \frac{\pi}{4}. \quad (33)$$

### 3.2 Riccati equation

In the following we give the proof of Theorem 3.1. The strategy for proof is described as follows:

- (1) We derive Riccati equations for  $F$  and  $G$  from the recursion relations (18) and (19), respectively.
- (2) We then linearize the Riccati equations by a standard technique to yield the Airy equation.
- (3) Finally we identify the functions by investigating asymptotic behaviours.

**Proposition 3.2**  $F(x, t)$  and  $G(x, t)$  satisfy the following Riccati equations,

$$t^3 \frac{\partial F}{\partial x} = -F^2 + t^3(t^3 - 2x)F - t^6(x^2 + 1), \quad (34)$$

$$t^3 \frac{\partial G}{\partial x} = -G^2 + t^3(t^3 + 2x)G - t^6(x^2 - 1), \quad (35)$$

respectively.

*proof:* (34) can be derived as follows:

$$\begin{aligned} F^2 &= \sum_{n=0}^{\infty} \left( \sum_{k=0}^{\infty} a_k a_{n-k} \right) t^{-3n} = \sum_{n=0}^{\infty} (a_{n+2} - a'_{n+1} - 2xa_{n+1}) t^{-3n} \\ &= (t^6 F - t^6 a_0 - t^3 a_1) - \left( t^3 \frac{\partial F}{\partial x} - t^3 a'_0 \right) - 2x(t^3 F - t^3 a_0) \\ &= -t^3 \frac{\partial F}{\partial x} + t^3(t^3 - 2x)F - t^6(x^2 + 1), \end{aligned}$$

where we have used (18) to derive the second equality. We obtain (35) by the similar calculation.  $\square$

Since  $F(x, t)$  and  $G(x, t)$  are defined as formal series around  $t \sim \infty$ , it is useful to derive differential equations with respect to  $t$ . For this purpose we present the following lemma:

**Lemma 3.3** (1)  $a_n$  and  $b_n$  satisfy the following recursion relations,

$$a'_n = xa'_{n-1} + (3n + 1)a_{n-1}, \quad (36)$$

$$b'_n = -xb'_{n-1} - (3n + 1)b_{n-1}, \quad (37)$$

respectively.

- (2)  $F(x, t)$  and  $G(x, t)$  satisfy the following linear differential equations,

$$(x - t^3) \frac{\partial F}{\partial x} = t \frac{\partial F}{\partial t} - 4F - 2xt^3, \quad (38)$$

$$(x + t^3) \frac{\partial G}{\partial x} = t \frac{\partial G}{\partial t} - 4G + 2xt^3, \quad (39)$$

respectively.

The first statement can be proved by a simple induction. The second statement follows immediately from (36) and (37).  $\square$

From Lemma 3.3, we obtain the Riccati equations with respect to  $t$ .

**Proposition 3.4**  $F(x, t)$  and  $G(x, t)$  satisfy the following Riccati equations

$$t \frac{\partial F}{\partial t} = -\frac{x - t^3}{t^3} F^2 + (-t^6 + 3t^3 x - 2x^2 + 4)F + t^3(t^3 x^2 + t^3 - x^3 + x), \quad (40)$$

$$t \frac{\partial G}{\partial t} = -\frac{x + t^3}{t^3} G^2 + (t^6 + 3xt^3 + 2x^2 + 4)G + t^3(-t^3 x^2 + t^3 - x^3 - x), \quad (41)$$

respectively.

### 3.3 Linearization and identification of functions

We obtain the Airy equation by linearization of the Riccati equations by standard technique.

**Proposition 3.5** (1) Putting

$$F = t^3 \frac{v_x}{v} - xt^3 + \frac{t^6}{2}, \quad (42)$$

the Riccati equation (34) reduces to the following linear equation:

$$\frac{\partial^2 v}{\partial x^2} = t^3 \left( -x + \frac{t^3}{4} \right) v. \quad (43)$$

(2) Similarly, putting

$$F = \frac{t^4}{x - t^3} \frac{v_t}{v} - xt^3 + \frac{t^6}{2}, \quad (44)$$

the Riccati equation (40) reduces to the following linear equation:

$$\frac{\partial^2 v}{\partial t^2} = -\frac{3t^2}{x - t^3} \frac{\partial v}{\partial t} + t(x - t^3)^2 \left( -x + \frac{t^3}{4} \right) v. \quad (45)$$

(3) (43) and (45) are transformed into the Airy equation

$$\frac{d^2 v}{dz^2} = zv, \quad (46)$$

by the change of variable

$$z = t \left( -x + \frac{t^3}{4} \right). \quad (47)$$

A similar proposition holds for  $G(x, t)$ :

**Proposition 3.6** (1) Putting

$$G = t^3 \frac{w_x}{w} + xt^3 + \frac{t^6}{2}, \quad (48)$$

the Riccati equation (35) reduces to the following linear equation:

$$\frac{\partial^2 w}{\partial x^2} = t^3 \left( x + \frac{t^3}{4} \right) w. \quad (49)$$

(2) Similarly, putting

$$G = \frac{t^4}{x + t^3} \frac{w_t}{w} + xt^3 + \frac{t^6}{2}, \quad (50)$$

the Riccati equation (41) reduces to the following linear equation:

$$\frac{\partial^2 w}{\partial t^2} = \frac{3t^2}{x + t^3} \frac{\partial w}{\partial t} + t(x + t^3)^2 \left( x + \frac{t^3}{4} \right) w. \quad (51)$$

(3) (49) and (51) are transformed into the Airy equation

$$\frac{d^2 w}{dz^2} = zw, \quad (52)$$

by the change of variable,

$$z = t \left( x + \frac{t^3}{4} \right). \quad (53)$$

We omit the proof since it is done by elementary calculations.



**Remark 3.7** Assertions (1) and (2) in Proposition 3.5 are consistent. Namely, (1) and (2) can be transformed to each other under the assumption that  $(x, t)$  dependence of  $u$  is given by  $u = u(z)$ . A similar remark also holds for Proposition 3.6.

Let us finally identify the function by considering its asymptotic behaviour around  $t \sim \infty$ . Solving (44) in terms of  $v$ , we have from (30)

$$v(x, t) = \text{const.} \times \exp \left[ \frac{1}{12} t^6 - \frac{1}{2} x t^3 \right] \times \frac{1}{t} \times \exp \left[ - \sum_{n=0}^{\infty} \frac{x a_{n-1} - a_n}{3n} t^{-3n} \right]. \quad (54)$$

The function  $v$  is given as linear combination of  $\text{Ai}(z)$  and  $\text{Bi}(z)$ . We compare the above expression with (28) and (29). Noticing that we have from (47)

$$\frac{2}{3} z^{\frac{3}{2}} \underset{t \rightarrow \infty}{\sim} \frac{t^6}{12} - \frac{1}{2} x t^3 + \dots, \quad z^{-\frac{1}{4}} \underset{t \rightarrow \infty}{\sim} \frac{\sqrt{2}}{t} \left( 1 + \frac{x}{t^3} + \dots \right),$$

we find that  $u$  should be

$$u(x, t) \sim U_+(z), \quad (55)$$

by comparing the exponential factors. This proves the statement regarding  $F$  in Theorem 3.1. The statement for  $G$  can be proved in a similar manner by using Proposition 3.6.  $\square$

## 4 Concluding remarks

In this article, we have shown that the generating functions associated with the Hankel determinant formula for Okamoto polynomials can be characterized by the log derivative of solutions of the Airy equation. It is an interesting observation that  $\text{Bi}$  and  $\text{Ai}$ , the solutions of Airy equation with different asymptotic behaviours, appear in opposite directions with respect to  $n$ .

We have posed the question as to whether the appearance of the log derivative of the Airy function in the generating function associated with Y-V polynomials is an accident or not. The result of this article appears to imply that the answer is “No”. It may be an interesting problem to characterize generating functions associated with special polynomials for other Painlevé equations in terms of the log derivative of special functions of hypergeometric type.

If this phenomena is not caused by accident, we should clarify the reason and mechanism underlying it. Unfortunately we cannot give an answer to this problem yet. Recently, Hankel determinant formula for generic solutions to  $\text{P}_{\text{II}}$  and the generating functions of its entries have been discussed in [3]. It is shown that the generating functions are also characterized by log derivative of solutions of certain linear differential equations, which are nothing but the auxiliary linear problem for  $\text{P}_{\text{II}}$ . This result is not a conclusive answer to the problem, but it may imply some (unknown) correspondence between the Painlevé equations and their auxiliary linear problems. At least, it may be true that there is a nontrivial mathematical structure behind Hankel determinant formulae, which deserves further study.

**Acknowledgment** The authors would like to thank Prof. Nalini Joshi and Dr. Marta Mazzocco for discussion and encouragement. This work is partially supported by the JSPS Grant-in-Aid for Scientific Research.

## References

- [1] M. J. Ablowitz and H. Segur, *Solitons and the Inverse Scattering Transform*, SIAM Studies in Applied Mathematics **4** (SIAM, Philadelphia, 1981).
- [2] K. Iwasaki, K. Kajiwara and T. Nakamura, *Generating function associated with the rational solutions of the Painlevé II equation*, J. Phys. A: Math. Gen. **35**(2002) L207-L211.
- [3] N. Joshi, K. Kajiwara and M. Mazzocco, *Generating function associated with the determinant formula for solutions of the Painlevé II equation*, to appear in Astérisque, arXiv:nlin.SI/0406035.
- [4] K. Kajiwara and T. Masuda, *On the Umemura polynomials for the Painlevé III equation*, Phys. Lett. **A260**(1999) 462-467.

- [5] K. Kajiwara and T. Masuda, *A generalization of determinant formulae for the solutions of Painlevé II and XXXIV equations*, J. Phys. A **32**(1999) 3763-3778.
- [6] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, *Determinant formulas for the Toda and discrete Toda equations*, Funkcial. Ekvac. **44**(2001) 291-307.
- [7] K. Kajiwara and Y. Ohta, *Determinant structure of rational solutions for the Painlevé II equation*, J. Math. Phys. **37**(1996) 4162-4174.
- [8] K. Kajiwara and Y. Ohta, *Determinant structure of the rational solutions for the Painlevé IV equation*, J. Phys. A: Math. Gen. **31**(1998) 2431-2446.
- [9] N. N. Lebedev, *Special Functions and Their Applications* (Prentice-Hall, Englewood Cliffs, 1965).
- [10] T. Masuda, *On a class of algebraic solutions to the Painlevé VI equation, its determinant formula and coalescence cascade*, Funkcial. Ekvac. **46**(2003) 121-171.
- [11] T. Masuda, Y. Ohta and K. Kajiwara, *A determinant formula for a class of rational Solutions of Painlevé V equation*, Nagoya Math. J. **168**(2002) 1-25.
- [12] T. Miwa, M. Jimbo and E. Date, *Solitons. Differential equations, symmetries and infinite-dimensional algebras* Cambridge Tracts in Mathematics **135**(Cambridge University Press, Cambridge, 2000).
- [13] M. Noumi, *Painlevé equations through symmetry*, Translations of Mathematical Monographs **223**(American Mathematical Society, Providence, 2004).
- [14] M. Noumi and Y. Yamada, *Symmetries in the fourth Painlevé equation and Okamoto polynomials*, Nagoya Math. J. **153**(1999) 53-86.
- [15] K. Okamoto, *Studies on the Painlevé equations. I. Sixth Painlevé equation  $P_{VI}$* , Ann. Mat. Pura Appl. **146**(1987) 337-381.
- [16] K. Okamoto, *Studies on the Painlevé equations. II. Fifth Painlevé equation  $P_V$* , Japan. J. Math. **13**(1987) 47-76.
- [17] K. Okamoto, *Studies on the Painlevé equations. III. Second and fourth Painlevé equations,  $P_{II}$  and  $P_{IV}$* , Math. Ann. **275**(1986) 221-255.
- [18] K. Okamoto, *Studies on the Painlevé equations. IV. Third Painlevé equation  $P_{III}$* , Funkcial. Ekvac. **30** (1987) 305-332.

# List of MHF Preprint Series, Kyushu University

## 21st Century COE Program

### Development of Dynamic Mathematics with High Functionality

- MHF2003-1 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Yoshitaka WATANABE  
A numerical method to verify the invertibility of linear elliptic operators with applications to nonlinear problems
- MHF2003-2 Masahisa TABATA & Daisuke TAGAMI  
Error estimates of finite element methods for nonstationary thermal convection problems with temperature-dependent coefficients
- MHF2003-3 Tomohiro ANDO, Sadanori KONISHI & Seiya IMOTO  
Adaptive learning machines for nonlinear classification and Bayesian information criteria
- MHF2003-4 Kazuhiro YOKOYAMA  
On systems of algebraic equations with parametric exponents
- MHF2003-5 Masao ISHIKAWA & Masato WAKAYAMA  
Applications of Minor Summation Formulas III, Plücker relations, Lattice paths and Pfaffian identities
- MHF2003-6 Atsushi SUZUKI & Masahisa TABATA  
Finite element matrices in congruent subdomains and their effective use for large-scale computations
- MHF2003-7 Setsuo TANIGUCHI  
Stochastic oscillatory integrals - asymptotic and exact expressions for quadratic phase functions -
- MHF2003-8 Shoki MIYAMOTO & Atsushi YOSHIKAWA  
Computable sequences in the Sobolev spaces
- MHF2003-9 Toru FUJII & Takashi YANAGAWA  
Wavelet based estimate for non-linear and non-stationary auto-regressive model
- MHF2003-10 Atsushi YOSHIKAWA  
Maple and wave-front tracking — an experiment
- MHF2003-11 Masanobu KANEKO  
On the local factor of the zeta function of quadratic orders
- MHF2003-12 Hidefumi KAWASAKI  
Conjugate-set game for a nonlinear programming problem

- MHF2004-1 Koji YONEMOTO & Takashi YANAGAWA  
Estimating the Lyapunov exponent from chaotic time series with dynamic noise
- MHF2004-2 Rui YAMAGUCHI, Eiko TSUCHIYA & Tomoyuki HIGUCHI  
State space modeling approach to decompose daily sales of a restaurant into time-dependent multi-factors
- MHF2004-3 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA  
Cubic pencils and Painlevé Hamiltonians
- MHF2004-4 Atsushi KAWAGUCHI, Koji YONEMOTO & Takashi YANAGAWA  
Estimating the correlation dimension from a chaotic system with dynamic noise
- MHF2004-5 Atsushi KAWAGUCHI, Kentarou KITAMURA, Koji YONEMOTO, Takashi YANAGAWA & Kiyofumi YUMOTO  
Detection of auroral breakups using the correlation dimension
- MHF2004-6 Ryo IKOTA, Masayasu MIMURA & Tatsuyuki NAKAKI  
A methodology for numerical simulations to a singular limit
- MHF2004-7 Ryo IKOTA & Eiji YANAGIDA  
Stability of stationary interfaces of binary-tree type
- MHF2004-8 Yuko ARAKI, Sadanori KONISHI & Seiya IMOTO  
Functional discriminant analysis for gene expression data via radial basis expansion
- MHF2004-9 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA  
Hypergeometric solutions to the  $q$ -Painlevé equations
- MHF2004-10 Raimundas VIDŪNAS  
Expressions for values of the gamma function
- MHF2004-11 Raimundas VIDŪNAS  
Transformations of Gauss hypergeometric functions
- MHF2004-12 Koji NAKAGAWA & Masakazu SUZUKI  
Mathematical knowledge browser
- MHF2004-13 Ken-ichi MARUNO, Wen-Xiu MA & Masayuki OIKAWA  
Generalized Casorati determinant and Positon-Negaton-Type solutions of the Toda lattice equation
- MHF2004-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO  
Generating function associated with the determinant formula for the solutions of the Painlevé II equation

- MHF2004-15 Kouji HASHIMOTO, Ryohei ABE, Mitsuhiro T. NAKAO & Yoshitaka WATANABE  
Numerical verification methods of solutions for nonlinear singularly perturbed problem
- MHF2004-16 Ken-ichi MARUNO & Gino BIONDINI  
Resonance and web structure in discrete soliton systems: the two-dimensional Toda lattice and its fully discrete and ultra-discrete versions
- MHF2004-17 Ryuei NISHII & Shinto EGUCHI  
Supervised image classification in Markov random field models with Jeffreys divergence
- MHF2004-18 Kouji HASHIMOTO, Kenta KOBAYASHI & Mitsuhiro T. NAKAO  
Numerical verification methods of solutions for the free boundary problem
- MHF2004-19 Hiroki MASUDA  
Ergodicity and exponential  $\beta$ -mixing bounds for a strong solution of Lévy-driven stochastic differential equations
- MHF2004-20 Setsuo TANIGUCHI  
The Brownian sheet and the reflectionless potentials
- MHF2004-21 Ryuei NISHII & Shinto EGUCHI  
Supervised image classification based on AdaBoost with contextual weak classifiers
- MHF2004-22 Hideki KOSAKI  
On intersections of domains of unbounded positive operators
- MHF2004-23 Masahisa TABATA & Shoichi FUJIMA  
Robustness of a characteristic finite element scheme of second order in time increment
- MHF2004-24 Ken-ichi MARUNO, Adrian ANKIEWICZ & Nail AKHMEDIEV  
Dissipative solitons of the discrete complex cubic-quintic Ginzburg-Landau equation
- MHF2004-25 Raimundas VIDŪNAS  
Degenerate Gauss hypergeometric functions
- MHF2004-26 Ryo IKOTA  
The boundedness of propagation speeds of disturbances for reaction-diffusion systems
- MHF2004-27 Ryusuke KON  
Convex dominates concave: an exclusion principle in discrete-time Kolmogorov systems

- MHF2004-28 Ryusuke KON  
Multiple attractors in host-parasitoid interactions: coexistence and extinction
- MHF2004-29 Kentaro IHARA, Masanobu KANEKO & Don ZAGIER  
Derivation and double shuffle relations for multiple zeta values
- MHF2004-30 Shuichi INOKUCHI & Yoshihiro MIZOGUCHI  
Generalized partitioned quantum cellular automata and quantization of classical CA
- MHF2005-1 Hideki KOSAKI  
Matrix trace inequalities related to uncertainty principle
- MHF2005-2 Masahisa TABATA  
Discrepancy between theory and real computation on the stability of some finite element schemes
- MHF2005-3 Yuko ARAKI & Sadanori KONISHI  
Functional regression modeling via regularized basis expansions and model selection
- MHF2005-4 Yuko ARAKI & Sadanori KONISHI  
Functional discriminant analysis via regularized basis expansions
- MHF2005-5 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA  
Point configurations, Cremona transformations and the elliptic difference Painlevé equations
- MHF2005-6 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA  
Construction of hypergeometric solutions to the  $q$ -Painlevé equations
- MHF2005-7 Hiroki MASUDA  
Simple estimators for non-linear Markovian trend from sampled data:  
I. ergodic cases
- MHF2005-8 Hiroki MASUDA & Nakahiro YOSHIDA  
Edgeworth expansion for a class of Ornstein-Uhlenbeck-based models
- MHF2005-9 Masayuki UCHIDA  
Approximate martingale estimating functions under small perturbations of dynamical systems
- MHF2005-10 Ryo MATSUZAKI & Masayuki UCHIDA  
One-step estimators for diffusion processes with small dispersion parameters from discrete observations
- MHF2005-11 Junichi MATSUKUBO, Ryo MATSUZAKI & Masayuki UCHIDA  
Estimation for a discretely observed small diffusion process with a linear drift

MHF2005-12 Masayuki UCHIDA & Nakahiro YOSHIDA

AIC for ergodic diffusion processes from discrete observations

MHF2005-13 Hiromichi GOTO & Kenji KAJIWARA

Generating function related to the Okamoto polynomials for the Painlevé IV equation