Casorati Determinant Form of Dark Soliton Solutions of the Discrete Nonlinear Schrödinger Equation

Maruno, Ken-ichi
Faculty of Mathematics, Kyushu University

Ohta, Yasuhiro
Department of Mathematics, Kobe University

<http://hdl.handle.net/2324/11854>

出版情報 : Journal of the Physical society of Japan. 75 (5), pp.054002-, 2006-05-10. The
Physical Society of Japan

バージョン :

権利関係 :

MHF Preprint Series

Kyushu University
21st Century COE Program
Development of Dynamic Mathematics with
High Functionality

Determinant form of dark soliton solutions of the discrete nonlinear Schrödinger equation

K. Maruno & Y. Ohta

MHF 2005-24

(Received June 24, 2005)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

Determinant Form of Dark Soliton Solutions of the Discrete Nonlinear Schrödinger Equation

Ken-ichi Maruno^{1†} and Yasuhiro Ohta²

¹Faculty of Mathematics, Kyushu University, Hakozaki, Higashi-ku, Fukuoka, 812-8581, Japan

²Department of Mathematics, Kobe University, Rokko, Kobe 657-8501, Japan

Abstract. It is shown that the N -dark soliton solutions of the integrable discrete nonlinear Schrödinger (IDNLS) equation are given in terms of the Casorati determinant. The conditions for reduction, complex conjugacy and regularity for the Casorati determinant solution are also given explicitly. The relationship between the IDNLS and the relativistic Toda lattice is discussed.

24 June 2005

PACS numbers: 02.30.Ik, 05.45.Yv

To be submitted to : *J. Phys. A: Math. Gen.*

1. Introduction

The nonlinear Schrödinger (NLS) equation,

$$i\psi_t = \psi_{xx} + \alpha|\psi|^2\psi, \quad (1.1)$$

is one of most important soliton equations in mathematics and physics. The study of discrete analogues of the NLS equation has received considerable attention recently from both physical and mathematical point of view [1, 2]. The integrable discrete nonlinear Schrödinger (IDNLS) equation is given by

$$i\frac{d\psi_n}{dt} = \psi_{n+1} + \psi_{n-1} + \alpha|\psi_n|^2(\psi_{n+1} + \psi_{n-1}). \quad (1.2)$$

The IDNLS equation was originally derived by Ablowitz and Ladik using the Lax formulation [3, 4, 1]. Thus the IDNLS equation is often called the Ablowitz-Ladik lattice.

The IDNLS equation can be bilinearized into

$$D_t G_n \cdot F_n = G_{n+1} F_{n-1} - F_{n+1} G_{n-1}, \quad (1.3)$$

$$F_n^2 + \alpha G_n G_n^* = F_{n+1} F_{n-1}, \quad (1.4)$$

through the dependent variable transformation $\psi_n = (i)^n \frac{G_n}{F_n}$ where F_n is a real function, G_n is a complex function and G_n^* is its complex conjugate. With this bilinear forms the IDNLS

† E-mail: maruno@math.kyushu-u.ac.jp

equation in the case of $\alpha = 1$ can be solved by the Hirota bilinear method, i.e. N -bright soliton solutions are obtained [5, 6, 7, 8].

It is well known that the NLS equation with defocusing parameter ($\alpha < 0$) has dark soliton solutions. It is easily confirmed that there is a stationary dark soliton solution

$$\psi_n = e^{-2it \operatorname{sech}^2(k)} \frac{\tanh(k)}{\sqrt{-\alpha}} \tanh(kn) \quad (1.5)$$

by the direct substitution of an ansatz of stationary solution into eq.(1.2). Understanding various properties of dark solitons in discrete lattices is important from physical point of view [9, 10, 11]. There is a result of the IDNLS equation under non-vanishing boundary condition using the inverse scattering theory by Vekslerchik and Konotop [12]. However, it is difficult to understand the dependence on a lattice space parameter of the N -dark soliton solutions in the result. The N -dark soliton solutions of the IDNLS equation have not been studied much until now.

The relationship between the IDNLS and relativistic Toda lattice (RTL) equation was pointed out by several authors [13, 14, 8]. An interesting open problem was given in conclusion in Ref.[8]: The 2-component Casorati determinant solution of the IDNLS equation and RTL equation in Ref.[8] is different from the single-component Casorati determinant solution of the RTL equation which was derived in Ref.[15]. Is there any relationship between the 2-component Casorati determinant solution and single-component Casorati determinant solution?

Our goal in this paper is to construct Casorati determinant form of N -dark soliton solutions with a lattice space parameter and give an answer of the above open problem. In this article, we give an explicit formula of the dark soliton solutions. It is known that the dark soliton solutions for NLS are written in terms of Wronski determinants. We show that the solution for IDNLS is given in the Casorati determinant form which is a discrete analogue of the Wronskian.

This paper is organized as follows. In the next section, we discuss about the gauge transformation and bilinearization of IDNLS equation. In section 3, we give bilinear identities for Casorati determinant both for the Bäcklund transformation of Toda lattice and the discrete 2-dimensional Toda lattice equation. In the section 4, we perform a reduction of the Casorati determinant which derives the bilinear form for IDNLS given in section 2. The complex conjugate condition is considered in the section 5. In the section 6, we give examples of dark soliton solutions. We also clarify the relationship between the IDNLS and the relativistic Toda lattice in the final section.

2. Gauge Transformation of IDNLS

First of all, we should examine the transformation of IDNLS equation by gauge. Applying the gauge transformation,

$$\psi_n = u_n A e^{i(\theta_n - \omega t)},$$

eq. (1.2) is transformed as

$$i \frac{du_n}{dt} = -\omega u_n + (1 + \alpha A^2 |u_n|^2)(u_{n+1} e^{i\theta} + u_{n-1} e^{-i\theta}),$$

thus we can scale $|\alpha|$ by using the amplitude A . To obtain the dark soliton solutions, we set $\alpha = -(a^2 - 1)/a^2$, ($a > 1$). We also scale t by $t \rightarrow a^2 t$, then the IDNLS eq. (1.2) can be bilinearized into

$$(iD_t + \omega)G_n \cdot F_n = G_{n+1}F_{n-1}e^{i\theta} + G_{n-1}F_{n+1}e^{-i\theta}, \quad (2.1)$$

$$a^2 F_n^2 - (a^2 - 1)G_n G_n^* = F_{n+1}F_{n-1}, \quad (2.2)$$

through the dependent variable transformation,

$$\psi_n = \frac{G_n}{F_n} e^{i(\theta n - \omega t)},$$

where F_n is a real function, G_n is a complex function and G_n^* is its complex conjugate.

The above gauge factor $e^{i(\theta n - \omega t)}$ is nothing but the carrier wave of the dark soliton, and essentially $u_n = G_n/F_n$ gives the envelope of the soliton solution. In the context of NLS, in order to construct physical solutions, it is more convenient to consider the above bilinear form than the one in eqs. (1.3) and (1.4). Using the bilinear forms, eqs. (2.1) and (2.2), we present the Casorati determinant form of dark soliton solutions of the IDNLS equation. Instead of applying the Hirota bilinear method directly to eqs. (2.1) and (2.2), we rather start from the algebraic identities for Casorati determinant and derive the bilinear form of IDNLS equation by using the reduction technique.

3. Bilinear Identities for Casorati Determinant

3.1. Bäcklund Transformation of Toda Lattice

We have the bilinear forms for Bäcklund Transformation (BT) of Toda lattice (TL) equation,

$$(aD_x - 1)\tau_{n+1}(k+1, l) \cdot \tau_n(k, l) + \tau_n(k+1, l)\tau_{n+1}(k, l) = 0, \quad (3.1)$$

$$(bD_y - 1)\tau_{n-1}(k, l+1) \cdot \tau_n(k, l) + \tau_n(k, l+1)\tau_{n-1}(k, l) = 0, \quad (3.2)$$

where a and b are constants.

Consider the following Casorati determinant solution,

$$\tau_n(k, l) = \begin{vmatrix} \phi_1^{(n)}(k, l) & \phi_1^{(n+1)}(k, l) & \cdots & \phi_1^{(n+N-1)}(k, l) \\ \phi_2^{(n)}(k, l) & \phi_2^{(n+1)}(k, l) & \cdots & \phi_2^{(n+N-1)}(k, l) \\ \vdots & \vdots & & \vdots \\ \phi_N^{(n)}(k, l) & \phi_N^{(n+1)}(k, l) & \cdots & \phi_N^{(n+N-1)}(k, l) \end{vmatrix}, \quad (3.3)$$

where $\phi_i^{(n)}$'s are arbitrary functions of two continuous independent variables, x and y , and two discrete ones, k and l , which satisfy the dispersion relations,

$$\partial_x \phi_i^{(n)}(k, l) = \phi_i^{(n+1)}(k, l), \quad (3.4)$$

$$\partial_y \varphi_i^{(n)}(k, l) = \varphi_i^{(n-1)}(k, l), \quad (3.5)$$

$$\Delta_k \varphi_i^{(n)}(k, l) = \varphi_i^{(n+1)}(k, l), \quad (3.6)$$

$$\Delta_l \varphi_i^{(n)}(k, l) = \varphi_i^{(n-1)}(k, l). \quad (3.7)$$

Here Δ_k and Δ_l are the backward difference operators with difference intervals a and b , defined by

$$\Delta_k f(k, l) = \frac{f(k, l) - f(k-1, l)}{a}, \quad (3.8)$$

$$\Delta_l f(k, l) = \frac{f(k, l) - f(k, l-1)}{b}. \quad (3.9)$$

For simplicity we introduce a convenient notation,

$$|n_{1_{k_1}, n_{2_{k_2}}, \dots, n_{N_{k_N}}}| = \begin{vmatrix} \varphi_1^{(n_1)}(k_1, l_1) & \varphi_1^{(n_2)}(k_2, l_2) & \cdots & \varphi_1^{(n_N)}(k_N, l_N) \\ \varphi_2^{(n_1)}(k_1, l_1) & \varphi_2^{(n_2)}(k_2, l_2) & \cdots & \varphi_2^{(n_N)}(k_N, l_N) \\ \vdots & \vdots & & \vdots \\ \varphi_N^{(n_1)}(k_1, l_1) & \varphi_N^{(n_2)}(k_2, l_2) & \cdots & \varphi_N^{(n_N)}(k_N, l_N) \end{vmatrix}. \quad (3.10)$$

In this notation, the solution for BT of TL, $\tau_n(k, l)$ in eq. (3.3), is rewritten as

$$\tau_n(k, l) = |n_{\underset{l}{k}}, n+1_{\underset{l}{k}}, \dots, n+N-1_{\underset{l}{k}}|, \quad (3.11)$$

or suppressing the index k and l , we simply write as

$$\tau_n(k, l) = |n, n+1, \dots, n+N-1|.$$

We show that the above $\tau_n(k, l)$ actually satisfies the bilinear eqs. (3.1) and (3.2) by using the Laplace expansion technique[16, 17]. At first we investigate the difference formula for τ . From eq. (3.6) we have

$$a\varphi_i^{(n+1)}(k+1, l) = \varphi_i^{(n)}(k+1, l) - \varphi_i^{(n)}(k, l).$$

Noticing this relation, we get

$$\begin{aligned} \tau_n(k+1, l) &= |n_{\underset{l}{k+1}}, n+1_{\underset{l}{k+1}}, n+2_{\underset{l}{k+1}}, \dots, n+N-1_{\underset{l}{k+1}}| \\ &= |n_{\underset{l}{k}}, n+1_{\underset{l}{k+1}}, n+2_{\underset{l}{k+1}}, \dots, n+N-1_{\underset{l}{k+1}}| \end{aligned}$$

where we have subtracted the 2nd column multiplied by a from the 1st column,

$$= |n_{\underset{l}{k}}, n+1_{\underset{l}{k}}, n+2_{\underset{l}{k+1}}, \dots, n+N-1_{\underset{l}{k+1}}|$$

where we have subtracted the 3rd column multiplied by a from the 2nd column,

...

$$= |n_{\underset{l}{k}}, n+1_{\underset{l}{k}}, \dots, n+N-2_{\underset{l}{k}}, n+N-1_{\underset{l}{k+1}}|, \quad (3.12)$$

that is,

$$\tau_n(k+1, l) = |n_{\underset{l}{k}}, n+1_{\underset{l}{k}}, \dots, n+N-2_{\underset{l}{k}}, n+N-1_{\underset{l}{k+1}}|. \quad (3.13)$$

Moreover in eq. (3.13), multiplying the N -th column by a and adding the $(N-1)$ -th column to the N -th column, we get

$$a\tau_n(k+1, l) = |n_k, n+1_k, \dots, n+N-2_k, n+N-2_{k+1}|. \quad (3.14)$$

Differentiating eq.(3.14) with x and using eq.(3.4) we obtain

$$\begin{aligned} a\partial_x \tau_n(k+1, l) &= |n_k, n+1_k, \dots, n+N-3_k, n+N-1_k, n+N-2_{k+1}| \\ &\quad + |n_k, n+1_k, \dots, n+N-3_k, n+N-2_k, n+N-1_{k+1}| \\ &= |n_k, n+1_k, \dots, n+N-3_k, n+N-1_k, n+N-2_{k+1}| + \tau_n(k+1, l). \end{aligned} \quad (3.15)$$

We have also

$$\partial_x \tau_n(k, l) = |n_k, n+1_k, \dots, n+N-2_k, n+N_k|. \quad (3.16)$$

In short, we write

$$\tau_n(k+1, l) = |n, n+1, \dots, n+N-2, n+N-1_{k+1}|, \quad (3.17)$$

$$a\tau_n(k+1, l) = |n, n+1, \dots, n+N-2, n+N-2_{k+1}|, \quad (3.18)$$

$$(a\partial_x - 1)\tau_n(k+1, l) = |n, n+1, \dots, n+N-3, n+N-1, n+N-2_{k+1}|, \quad (3.19)$$

$$\partial_x \tau_n(k, l) = |n, n+1, \dots, n+N-2, n+N|. \quad (3.20)$$

Let us introduce an identity for $2N \times 2N$ determinant,

$$\begin{vmatrix} n+1 & \cdots & n+N-2 & \vdots & n+N & \vdots & n+N-1_{k+1} & \vdots & \emptyset & \vdots & n+N-1 \\ \hline \emptyset & \vdots & n+N & \vdots & n+N-1_{k+1} & \vdots & n & n+1 & \cdots & n+N-2 & \vdots & n+N-1 \end{vmatrix} = 0.$$

Applying the Laplace expansion to the left-hand side, we obtain the algebraic bilinear identity for determinants,

$$\begin{aligned} &|n+1, \dots, n+N-2, n+N, n+N-1_{k+1}| \times |n, n+1, \dots, n+N-2, n+N-1| \\ &- |n+1, \dots, n+N-2, n+N-1, n+N-1_{k+1}| \times |n, n+1, \dots, n+N-2, n+N| \\ &- |n+1, \dots, n+N-2, n+N, n+N-1| \times |n, n+1, \dots, n+N-2, n+N-1_{k+1}| = 0, \end{aligned} \quad (3.21)$$

which is rewritten by using eqs. (3.17)-(3.20), into the differential bilinear equation,

$$(a\partial_x - 1)\tau_{n+1}(k+1, l) \times \tau_n(k, l) - a\tau_{n+1}(k+1, l)\partial_x \tau_n(k, l) + \tau_{n+1}(k, l)\tau_n(k+1, l) = 0.$$

This equation is equal to eq. (3.1).

Similarly we obtain

$$(b\partial_y - 1)\tau_{n-1}(k, l+1) \times \tau_n(k, l) - b\tau_{n-1}(k, l+1)\partial_y \tau_n(k, l) + \tau_{n-1}(k, l)\tau_n(k, l+1) = 0,$$

because the role of y, l and b is parallel to that of x, k and a except that the ordering of index n is reversed. The above equation is equal to eq. (3.2). Hence we have proved that the Casorati determinant $\tau_n(k, l)$ in eq. (3.3) gives the solution for the BT of TL equation.

3.2. Discrete 2-dimensional Toda Lattice

Using the Casoratian technique, we show that the bilinear form of discrete 2-dimensional Toda lattice (D2DTL) equation,

$$\begin{aligned} & \tau_n(k+1, l+1)\tau_n(k, l) - \tau_n(k+1, l)\tau_n(k, l+1) \\ &= ab(\tau_n(k+1, l+1)\tau_n(k, l) - \tau_{n+1}(k+1, l)\tau_{n-1}(k, l+1)), \end{aligned} \quad (3.22)$$

is also satisfied by the Casorati determinant.

Let us further examine the difference formula for the Casorati determinant τ_n in eq. (3.3). Similarly to eqs. (3.17) and (3.18), we have

$$\tau_n(k, l+1) = |n_{l+1}, n+1, \dots, n+N-2, n+N-1|, \quad (3.23)$$

$$b\tau_n(k, l+1) = |n+1_{l+1}, n+1, \dots, n+N-2, n+N-1|. \quad (3.24)$$

Thus $\tau_n(k+1, l+1)$ is given by

$$\tau_n(k+1, l+1) = |n_{k+1, l+1}, n+1_{k+1, l+1}, \dots, n+N-2_{k+1, l+1}, n+N-1_{k+1, l+1}|. \quad (3.25)$$

In the following we show that the shifts of index k is condensed into only the right-most column of the determinant. From eqs. (3.6) and (3.7), $\phi_i^{(n)}$ satisfies

$$\Delta_l \Delta_k \phi_i^{(n)} = \phi_i^{(n)}, \quad (3.26)$$

that is,

$$\phi_i^{(n)}(k, l) - \phi_i^{(n)}(k-1, l) - \phi_i^{(n)}(k, l-1) + \phi_i^{(n)}(k-1, l-1) = ab\phi_i^{(n)}(k, l), \quad (3.27)$$

which is rewritten as

$$\begin{aligned} (1-ab)\phi_i^{(n)}(k+1, l+1) &= \phi_i^{(n)}(k, l+1) + \phi_i^{(n)}(k+1, l) - \phi_i^{(n)}(k, l) \\ &= \phi_i^{(n)}(k, l+1) + a\phi_i^{(n+1)}(k+1, l). \end{aligned} \quad (3.28)$$

Therefore multiplying the both hand sides of eq. (3.25) by $(1-ab)$ and rewriting the 1st column of the determinant by the use of eq. (3.28), we obtain

$$\begin{aligned} & (1-ab)\tau_n(k+1, l+1) \\ &= |n_{k, l+1}, n+1_{k, l+1}, \dots, n+N-2_{k, l+1}, n+N-1_{k, l+1}| \\ &+ a|n_{k+1, l+1}, n+1_{k+1, l+1}, \dots, n+N-2_{k+1, l+1}, n+N-1_{k+1, l+1}|. \end{aligned} \quad (3.29)$$

The second term in r.h.s. vanishes because its 1st and 2nd columns are the same. So we get

$$(1-ab)\tau_n(k+1, l+1) = |n_{l+1}, n+1_{k+1}, \dots, n+N-2_{k+1}, n+N-1_{k+1}|. \quad (3.30)$$

In the above determinant, by subtracting the $(j+1)$ -th column multiplied by a from the j -th column for $j = 2, 3, \dots, N-1$, $\tau_n(k+1, l+1)$ is given by

$$(1-ab)\tau_n(k+1, l+1) = |n_{l+1}, n+1, \dots, n+N-2, n+N-1_{k+1}|. \quad (3.31)$$

As is shown, even if the two variables k and l are shifted, τ_n is also expressed in the determinant form whose columns are almost unchanged and only edges are varied. Thus we can use the Laplace expansion technique.

We consider an identity for $2N \times 2N$ determinant,

$$\begin{vmatrix} n_{l+1} & n+1 & \cdots & n+N-2 & n+N-1_{k+1} & n & \emptyset & n+N-1 \\ \hline n_{l+1} & \emptyset & & & & n & n+1 & \cdots & n+N-2 & n+N-1 \end{vmatrix} = 0.$$

By the Laplace expansion, we have

$$\begin{aligned} & |n_{l+1}, n+1, \dots, n+N-2, n+N-1_{k+1}| \times |n, n+1, \dots, n+N-2, n+N-1| \\ & - |n, n+1, \dots, n+N-2, n+N-1_{k+1}| \times |n_{l+1}, n+1, \dots, n+N-2, n+N-1| \\ & + |n+1, \dots, n+N-2, n+N-1, n+N-1_{k+1}| \times |n_{l+1}, n, n+1, \dots, n+N-2| = 0. \end{aligned} \quad (3.32)$$

Using eqs. (3.17), (3.18), (3.23), (3.24) and (3.31), we obtain from eq. (3.32),

$$\begin{aligned} & (1-ab)\tau_n(k+1, l+1)\tau_n(k, l) - \tau_n(k+1, l)\tau_n(k, l+1) \\ & + a\tau_{n+1}(k+1, l)b\tau_{n-1}(k, l+1) = 0, \end{aligned} \quad (3.33)$$

which recovers eq. (3.22). This completes the proof that the Casorati determinant is indeed the solution for D2DTL equation.

For example, we can take $\varphi_i^{(n)}$ as

$$\begin{aligned} \varphi_i^{(n)}(k, l) &= p_i^n (1 - ap_i)^{-k} \left(1 - b \frac{1}{p_i}\right)^{-l} e^{\xi_i} + q_i^n (1 - aq_i)^{-k} \left(1 - b \frac{1}{q_i}\right)^{-l} e^{\eta_i}, \\ \xi_i &= p_i x + \frac{1}{p_i} y + \xi_{i0}, \quad \eta_i = q_i x + \frac{1}{q_i} y + \eta_{i0}, \end{aligned} \quad (3.34)$$

where p_i , q_i and ξ_{i0} , η_{i0} are arbitrary constants which correspond to the wave numbers and phase parameters of solitons, respectively. It can easily be seen that $\varphi_i^{(n)}$ in eq. (3.34) actually satisfies the dispersion relations, eqs. (3.4)-(3.7).

4. Reduction to IDNLS

In this section, we give the reduction technique in order to derive the IDNLS equation from the system of BT of TL and D2DTL. In eq. (3.34), we apply a condition for the wave numbers,

$$q_i = -p_i \frac{1 - b \frac{1}{p_i}}{1 - ap_i}. \quad (4.1)$$

Then we get

$$p_i = -q_i \frac{1 - b \frac{1}{q_i}}{1 - aq_i}, \quad (4.2)$$

and thus

$$p_i^2 \frac{1 - b \frac{1}{p_i}}{1 - ap_i} = q_i^2 \frac{1 - b \frac{1}{q_i}}{1 - aq_i}. \quad (4.3)$$

On this condition, $\varphi_i^{(n)}$ in eq. (3.34) satisfies

$$\varphi_i^{(n+2)}(k+1, l-1) = p_i^2 \frac{1 - b \frac{1}{p_i}}{1 - ap_i} \varphi_i^{(n)}(k, l). \quad (4.4)$$

Hence we obtain

$$\tau_{n+2}(k+1, l-1) = \left(\prod_{i=1}^N p_i^2 \frac{1 - b \frac{1}{p_i}}{1 - ap_i} \right) \tau_n(k, l). \quad (4.5)$$

By using eq. (4.5), the bilinear forms for BT of TL and D2DTL, eqs. (3.1), (3.2) and (3.22), are rewritten as

$$(aD_x - 1)\tau_{n+1}(k+1, l) \cdot \tau_n(k, l) + \tau_n(k+1, l)\tau_{n+1}(k, l) = 0, \quad (4.6)$$

$$(bD_y - 1)\tau_{n+1}(k+1, l) \cdot \tau_n(k, l) + \tau_{n+2}(k+1, l)\tau_{n-1}(k, l) = 0, \quad (4.7)$$

$$\begin{aligned} & \tau_{n+1}(k+1, l)\tau_{n-1}(k-1, l) - \tau_{n+1}(k, l)\tau_{n-1}(k, l) \\ & = ab(\tau_{n+1}(k+1, l)\tau_{n-1}(k-1, l) - \tau_n(k, l)\tau_n(k, l)). \end{aligned} \quad (4.8)$$

Here we may drop l -dependence, thus for simplicity we take $l = 0$ hereafter. Equation (4.8) is nothing but the bilinear form of the discrete one-dimensional TL equation.

Let us substitute

$$x = iact, \quad y = ibdt, \quad (4.9)$$

where c and d are constants. Then we have

$$-i\partial_t = ac\partial_x + bd\partial_y, \quad (4.10)$$

and from the bilinear eqs. (4.6) and (4.7), we get

$$(-iD_t - c - d)\tau_{n+1}(k+1) \cdot \tau_n(k) + c\tau_n(k+1)\tau_{n+1}(k) + d\tau_{n+2}(k+1)\tau_{n-1}(k) = 0. \quad (4.11)$$

The Casorati determinant solution of eqs. (4.11) and (4.8) (with $l = 0$) is given by

$$\tau_n(k) = \begin{vmatrix} \varphi_1^{(n)}(k) & \varphi_1^{(n+1)}(k) & \cdots & \varphi_1^{(n+N-1)}(k) \\ \varphi_2^{(n)}(k) & \varphi_2^{(n+1)}(k) & \cdots & \varphi_2^{(n+N-1)}(k) \\ \vdots & \vdots & \ddots & \vdots \\ \varphi_N^{(n)}(k) & \varphi_N^{(n+1)}(k) & \cdots & \varphi_N^{(n+N-1)}(k) \end{vmatrix},$$

$$\varphi_i^{(n)}(k) = p_i^n (1 - ap_i)^{-k} e^{\xi_i} + q_i^n (1 - aq_i)^{-k} e^{\eta_i},$$

$$\xi_i = i \left(acp_i + \frac{bd}{p_i} \right) t + \xi_{i0}, \quad \eta_i = i \left(acq_i + \frac{bd}{q_i} \right) t + \eta_{i0}.$$

Here p_i and q_i are related by eq. (4.1).

By defining

$$f_n = \tau_n(0), \quad g_n = \tau_{n+1}(1), \quad h_n = \tau_{n-1}(-1), \quad (4.12)$$

from eqs. (4.8) and (4.11), we obtain the bilinear form of IDNLS equation,

$$(-iD_t - c - d)g_n \cdot f_n + dg_{n+1}f_{n-1} + cg_{n-1}f_{n+1} = 0, \quad (4.13)$$

$$(iD_t - c - d)h_n \cdot f_n + ch_{n+1}f_{n-1} + dh_{n-1}f_{n+1} = 0, \quad (4.14)$$

$$f_{n+1}f_{n-1} - f_nf_n = (ab - 1)(f_nf_n - g_nh_n). \quad (4.15)$$

By using the dependent variable transformation,

$$u_n = \frac{g_n}{f_n}, \quad v_n = \frac{h_n}{f_n},$$

the above bilinear equations are transformed into

$$i \frac{du_n}{dt} = -(c+d)u_n + (ab - (ab-1)u_nv_n)(du_{n+1} + cu_{n-1}), \quad (4.16)$$

$$-i \frac{dv_n}{dt} = -(c+d)v_n + (ab - (ab-1)u_nv_n)(cv_{n+1} + dv_{n-1}). \quad (4.17)$$

5. Complex Conjugate Condition

In order to take v_n to be the complex conjugate of u_n , we need to restrict f_n as real and h_n as the complex conjugate of g_n up to gauge freedom. Moreover to take u_n to be regular (i.e., u_n not to diverge for real t), we need $f_n \neq 0$ for real t . In this section, we give the conditions for the complex conjugacy and regularity for the Casorati determinant solution.

Firstly we take

$$b = a,$$

for simplicity, and next we take

$$a > 1.$$

Then the conditions for complex conjugate are given as follows:

$$p_i = a + \sqrt{a^2 - 1}r_i, \quad |r_i| = 1, \quad (5.1)$$

$$e^{\eta_{i0}} = \left(\prod_{\substack{k=1 \\ k \neq i}}^N \frac{p_i - q_k}{q_i - q_k} \right) e^{-\xi_{i0}^*}, \quad (5.2)$$

$$d = c^*, \quad (5.3)$$

where r_i is a complex parameter of absolute value 1, and $*$ means the complex conjugate. On the condition (5.1), eq. (4.1) turns to be

$$q_i = \frac{1}{p_i^*},$$

and eqs. (5.2) and (5.3) gives

$$e^{\eta_i} = \left(\prod_{\substack{k=1 \\ k \neq i}}^N \frac{p_i - q_k}{q_i - q_k} \right) e^{-\xi_i^*},$$

for real t .

Now we summarize the final result of the Casorati determinant solution for IDNLS equation. The bilinear form of IDNLS equation is given as

$$(-iD_t - c - c^*)g_n \cdot f_n + c^*g_{n+1}f_{n-1} + cg_{n-1}f_{n+1} = 0, \quad (5.4)$$

$$(iD_t - c - c^*)h_n \cdot f_n + ch_{n+1}f_{n-1} + c^*h_{n-1}f_{n+1} = 0, \quad (5.5)$$

$$f_{n+1}f_{n-1} - f_nf_n = (a^2 - 1)(f_nf_n - g_nh_n), \quad (5.6)$$

where $a > 1$ and t is real. The Casorati determinant solution for the above bilinear equations is given as

$$f_n = \begin{vmatrix} \varphi_1^{(n)}(0) & \varphi_1^{(n+1)}(0) & \cdots & \varphi_1^{(n+N-1)}(0) \\ \varphi_2^{(n)}(0) & \varphi_2^{(n+1)}(0) & \cdots & \varphi_2^{(n+N-1)}(0) \\ \vdots & \vdots & \ddots & \vdots \\ \varphi_N^{(n)}(0) & \varphi_N^{(n+1)}(0) & \cdots & \varphi_N^{(n+N-1)}(0) \end{vmatrix}, \quad (5.7)$$

$$g_n = \begin{vmatrix} \varphi_1^{(n+1)}(1) & \varphi_1^{(n+2)}(1) & \cdots & \varphi_1^{(n+N)}(1) \\ \varphi_2^{(n+1)}(1) & \varphi_2^{(n+2)}(1) & \cdots & \varphi_2^{(n+N)}(1) \\ \vdots & \vdots & \ddots & \vdots \\ \varphi_N^{(n+1)}(1) & \varphi_N^{(n+2)}(1) & \cdots & \varphi_N^{(n+N)}(1) \end{vmatrix}, \quad (5.8)$$

$$h_n = \begin{vmatrix} \varphi_1^{(n-1)}(-1) & \varphi_1^{(n)}(-1) & \cdots & \varphi_1^{(n+N-2)}(-1) \\ \varphi_2^{(n-1)}(-1) & \varphi_2^{(n)}(-1) & \cdots & \varphi_2^{(n+N-2)}(-1) \\ \vdots & \vdots & \ddots & \vdots \\ \varphi_N^{(n-1)}(-1) & \varphi_N^{(n)}(-1) & \cdots & \varphi_N^{(n+N-2)}(-1) \end{vmatrix}, \quad (5.9)$$

where

$$\varphi_i^{(n)}(k) = p_i^n (-\sqrt{a^2 - 1} p_i^* r_i)^{-k} e^{\xi_i} + \left(\frac{1}{p_i^*} \right)^n \left(\frac{\sqrt{a^2 - 1}}{p_i^* r_i} \right)^{-k} \left(\prod_{\substack{k=1 \\ k \neq i}}^N \frac{p_k^* p_i - 1}{p_k^* / p_i^* - 1} \right) e^{-\xi_i^*},$$

$$\xi_i = ia \left(cp_i + \frac{c^*}{p_i} \right) t + \xi_{i0}, \quad p_i = a + \sqrt{a^2 - 1} r_i, \quad |r_i| = 1,$$

where r_i is a complex parameter of absolute value 1 and ξ_{i0} is an arbitrary complex parameter. This Casorati determinant solution gives the N -dark soliton solution for IDNLS equation. r_i and ξ_{i0} parametrize the wave number and phase constant of i -th soliton, respectively.

After a straightforward and tedious calculation of expanding the determinants, f_n , g_n and h_n can be expressed in an explicit way as follows:

$$f_n = F_n \mathcal{G}, \quad g_n = G_n \mathcal{G} \frac{r_1 r_2 \cdots r_N}{(\sqrt{a^2 - 1})^N}, \quad h_n = H_n \mathcal{G} \frac{(\sqrt{a^2 - 1})^N}{r_1 r_2 \cdots r_N}, \quad (5.10)$$

where $\mathcal{G}$ is the gauge factor defined by

$$\mathcal{G} = \prod_{N \geq i > j \geq 1} (q_i - q_j) \prod_{i=1}^N q_i^n e^{\eta_i},$$

and F_n , G_n and H_n are given as

$$F_n = \sum_{M=0}^N \sum_{1 \leq i_1 < i_2 < \cdots < i_M \leq N} \left| \prod_{1 \leq \mu < \nu \leq M} \frac{p_{i_\mu} - p_{i_\nu}}{p_{i_\mu} p_{i_\nu}^* - 1} \right|^2 \prod_{\nu=1}^M |p_{i_\nu}|^{2n} e^{\xi_{i_\nu} + \xi_{i_\nu}^*}, \quad (5.11)$$

$$G_n = \sum_{M=0}^N \sum_{1 \leq i_1 < i_2 < \cdots < i_M \leq N} \left| \prod_{1 \leq \mu < \nu \leq M} \frac{p_{i_\mu} - p_{i_\nu}}{p_{i_\mu} p_{i_\nu}^* - 1} \right|^2 \prod_{\nu=1}^M \left(-\frac{p_{i_\nu}}{p_{i_\nu}^* r_{i_\nu}^2} \right) |p_{i_\nu}|^{2n} e^{\xi_{i_\nu} + \xi_{i_\nu}^*}, \quad (5.12)$$

$$H_n = \sum_{M=0}^N \sum_{1 \leq i_1 < i_2 < \cdots < i_M \leq N} \left| \prod_{1 \leq \mu < \nu \leq M} \frac{p_{i_\mu} - p_{i_\nu}}{p_{i_\mu} p_{i_\nu}^* - 1} \right|^2 \prod_{\nu=1}^M \left(-\frac{p_{i_\nu}^* r_{i_\nu}^2}{p_{i_\nu}} \right) |p_{i_\nu}|^{2n} e^{\xi_{i_\nu} + \xi_{i_\nu}^*}. \quad (5.13)$$

Now it is clear that F_n is real and $H_n = G_n^*$, thus v_n is the complex conjugate of u_n up to multiplication factor. Moreover since $F_n > 1$, the regularity of u_n is also satisfied.

From the bilinear eqs. (5.4)-(5.6), and the gauge transformation (5.10), the above F_n and G_n satisfy the bilinear equations,

$$(-iD_t - c - c^*)G_n \cdot F_n + c^*G_{n+1}F_{n-1} + cG_{n-1}F_{n+1} = 0, \quad (5.14)$$

$$F_{n+1}F_{n-1} - F_nF_n = (a^2 - 1)(F_nF_n - G_nG_n^*). \quad (5.15)$$

Let us take

$$c = e^{-i\theta},$$

where θ is a real parameter. By the variable transformation,

$$\psi_n = \frac{G_n}{F_n} e^{i(\theta n - \omega t)},$$

that is,

$$\psi_n = u_n \frac{(\sqrt{a^2 - 1})^N}{r_1 r_2 \cdots r_N} e^{i(\theta n - \omega t)},$$

where

$$\omega = e^{i\theta} + e^{-i\theta} = 2 \cos \theta,$$

we finally obtain the IDNLS equation,

$$\frac{i}{a^2} \frac{d\psi_n}{dt} = \psi_{n+1} + \psi_{n-1} - \frac{a^2 - 1}{a^2} |\psi_n|^2 (\psi_{n+1} + \psi_{n-1}). \quad (5.16)$$

Thus we have completed the proof that the N -dark soliton solutions for the IDNLS equation are given in terms of the Casorati determinant. Here we remark that a parameter a is related to a lattice space parameter $\delta = 1/a (< 1)$. Using a lattice parameter δ , we can rewrite the IDNLS equation (5.16) into

$$i \frac{d\psi_n}{dt} = \frac{\psi_{n+1} + \psi_{n-1}}{\delta^2} - \frac{1 - \delta^2}{\delta^2} |\psi_n|^2 (\psi_{n+1} + \psi_{n-1}). \quad (5.17)$$

We see that the IDNLS equation is defocusing whenever a lattice parameter δ is smaller than 1. When $\delta = 1$, the IDNLS equation reduces to a linear equation.

6. Soliton Solutions

In this section we give examples of dark soliton solutions for the IDNLS equation. By taking $N = 1$ in eqs. (5.11) and (5.12), we get the 1-soliton solution for the IDNLS eq. (5.16),

$$\psi_n = \frac{G_n}{F_n} e^{i(\theta n - 2t \cos \theta)},$$

where θ is a real parameter and

$$F_n = 1 + |p_1|^{2n} e^{\xi_1 + \xi_1^*}, \quad G_n = 1 - \frac{p_1}{p_1^* r_1^2} |p_1|^{2n} e^{\xi_1 + \xi_1^*},$$

Figure 1. Plots of 1-dark soliton. The vertical axis is $|\psi_n|$. ((a) $a = 2, r = \exp(i\pi), \theta = \pi$, (b) $a = 1.1, r = \exp(i\pi), \theta = \pi$.)

where

$$\xi_1 = ia \left(\frac{p_1}{e^{i\theta}} + \frac{e^{i\theta}}{p_1} \right) t + \xi_{10}, \quad p_1 = a + \sqrt{a^2 - 1}r_1, \quad |r_1| = 1.$$

By rewriting as $p_1 = pe^{i\theta}$ and $r_1 = re^{i\theta}$, we get a slightly simpler expression,

$$F_n = 1 + |p|^{2n} e^{\xi + \xi^*}, \quad G_n = 1 - \frac{p}{p^* r^2} |p|^{2n} e^{\xi + \xi^*},$$

where

$$\xi = ia \left(p + \frac{1}{p} \right) t + \xi_0, \quad p = ae^{-i\theta} + \sqrt{a^2 - 1}r, \quad |r| = 1.$$

Here $\arg(r)$ parametrizes the wave number of soliton, and $\Re(\xi_0)$ gives the phase constant which parametrizes the position of soliton. Figure 1 shows a stationary 1-dark (black) soliton. When $a(> 1)$ is getting closer to 1, the width of a dip is getting wider. Figure 2 and 3 show travelling 1-dark (gray and black) solitons.

Here we should comment about the difference of travelling velocity between black and gray solitons. In the continuous case, the NLS equation (1.1) is invariant under the gauge and Galilean transformations,

$$\psi(x, t) \rightarrow \psi(x - 2kt, t) e^{i(kx - k^2 t)},$$

thus the travelling velocity of any solution can be shifted by $2k$ by using the above transformation. If we normalize the freedom of Galilean transformation by requiring that the carrier wave disappears (i.e. $k = 0$) and ψ represents only the envelope, then the velocity of the envelope soliton is 0 if and only if the soliton is black. In the case of IDNLS, the situation is almost same. When $\theta = 0$ (i.e. the case of carrier-wave-less) or $\theta = \pi$ (i.e. $\psi_n = (-1)^n \times (\text{envelope soliton})$), then the velocity of black soliton is 0 and the travelling dark solitons are always gray solitons. We see this fact from Figure 1 and 3.

The 2-soliton solution is obtained by taking $N = 2$ in eqs. (5.11) and (5.12). Similarly by rewriting as $p_i \rightarrow p_i e^{i\theta}$ and $r_i \rightarrow r_i e^{i\theta}$, we get an explicit expression of the 2-soliton solution for eq. (5.16),

$$\psi_n = \frac{G_n}{F_n} e^{i(\theta n - 2t \cos \theta)},$$

$$F_n = 1 + |p_1|^{2n} e^{\xi_1 + \xi_1^*} + |p_2|^{2n} e^{\xi_2 + \xi_2^*} + \left| \frac{p_1 - p_2}{p_1 p_2^* - 1} \right|^2 |p_1 p_2|^{2n} e^{\xi_1 + \xi_1^* + \xi_2 + \xi_2^*},$$

Figure 2. Plots of travelling 1-gray soliton. The vertical axis is $|\psi_n|$. ((a) $a = 2, r = \exp(3i\pi/4), \theta = 5\pi/3$.)

Figure 3. Plots of travelling 1-black soliton. The vertical axis is $|\psi_n|$. ($a = 2, r = \exp(-5i\pi/3), \theta = 5\pi/3$.)

$$G_n = 1 - \frac{p_1}{p_1^* r_1^2} |p_1|^{2n} e^{\xi_1 + \xi_1^*} - \frac{p_2}{p_2^* r_2^2} |p_2|^{2n} e^{\xi_2 + \xi_2^*} \\ + \left| \frac{p_1 - p_2}{p_1 p_2^* - 1} \right|^2 \frac{p_1}{p_1^* r_1^2} \frac{p_2}{p_2^* r_2^2} |p_1 p_2|^{2n} e^{\xi_1 + \xi_1^* + \xi_2 + \xi_2^*},$$

where

$$\xi_i = i a \left(p_i + \frac{1}{p_i} \right) t + \xi_{i0}, \quad p_i = a e^{-i\theta} + \sqrt{a^2 - 1} r_i, \quad |r_i| = 1.$$

Here the carrier wave of the dark soliton solution is given by the exponential factor, $e^{i(\theta n - 2t \cos \theta)}$. Figure 4 shows 2-dark soliton interaction.

Figure 4. Plots of 2-dark soliton interaction. The vertical axis is $|\psi_n|$. ($a = 2, r_1 = \exp(3i\pi/4), r_2 = \exp(i\pi/4), \theta = 5\pi/4$)

7. Concluding Remarks

We have shown that the N -dark soliton solutions of the IDNLS equation are given by the Casorati determinant. From the above derivation of the N -dark soliton solutions for the IDNLS equation, we notice that there is correspondence between the Casorati determinant solutions of the IDNLS equation and the RTL equation [15]. The RTL equation is decomposed into three bilinear equations, i.e. the Toda lattice (TL) equation, the Bäcklund transformation (BT) for TL and the DTL equation (See appendix). The IDNLS equation is also decomposed into three bilinear equations, i.e. two types of the BT for TL and the DTL equation.

This fact reminds us several works on the relationship between IDNLS and RTL [13, 14, 8]. Moreover, our result is an answer to an open problem of Ref.[8], i.e., the N -bright soliton solutions of IDNLS are written by the 2-component Casorati determinant in Ref.[8], and the N -dark soliton solution are written by the single-component Casorati determinant which is given in this paper and corresponds to the solution of RTL in Ref.[15].

It is interesting to investigate N -dark soliton solutions of the discrete time IDNLS equation. It will be reported in the forthcoming paper.

The authors gratefully acknowledge valuable discussions with Prof. M. J. Ablowitz, Prof. Y. Kodama, Prof. K. Yoshimura, Prof. B. F. Feng, Prof. K. Kajiwara and Dr. A. Mukaihira. K.M. acknowledges support from the 21st Century COE program “Development of Dynamic Mathematics with High Functionality” at Faculty of Mathematics, Kyushu University.

8. Appendix

In this appendix we briefly explain τ -functions of the RTL equation. The RTL equation

$$\begin{aligned} \frac{d^2 q_n}{dt^2} = & (1 + \frac{1}{c} \dot{q}_{n-1}) (1 + \frac{1}{c} \dot{q}_n) \frac{\exp(q_{n-1} - q_n)}{1 + \frac{1}{c^2} \exp(q_{n-1} - q_n)} \\ & - (1 + \frac{1}{c} \dot{q}_n) (1 + \frac{1}{c} \dot{q}_{n+1}) \frac{\exp(q_n - q_{n+1})}{1 + \frac{1}{c^2} \exp(q_n - q_{n+1})}, \end{aligned} \quad (8.1)$$

where q_n is the coordinates of n -th lattice point and c is the light speed, was introduced and studied by Ruijsenaars [18]. The RTL equation (8.1) is transformed into the three bilinear equations,

$$\begin{aligned} D_x^2 f_n \cdot f_n &= 2(\bar{g}_n g_n - f_n^2), \\ (aD_x - 1)f_n \cdot f_{n-1} + \bar{g}_{n-1} g_n &= 0, \\ \bar{g}_{n-1} g_{n+1} - f_n^2 &= a^2(f_{n+1} f_{n-1} - f_n^2), \end{aligned} \quad (8.2)$$

through the variable transformations,

$$q_n = \log \frac{f_{n-1}}{f_n}, \quad (8.3)$$

$$t = \frac{\sqrt{1+c^2}}{c} x, \quad (8.4)$$

$$c = \frac{\sqrt{1-a^2}}{a}. \quad (8.5)$$

In eqs. (8.2), g and $\bar{g}$ are the auxiliary variables and D is the bilinear differential operator defined by

$$D_x^k f \cdot g = (\partial_x - \partial_y)^k f(x)g(y)|_{y=x}.$$

Letting

$$f_n = \tau_n(n), \quad g_n = \tau_{n-1}(n), \quad \bar{g}_n = \tau_{n+1}(n), \quad (8.6)$$

we have three bilinear equations, the TL equation, the Bäcklund transformation (BT) for TL and the DTL equation,

$$\begin{aligned} D_x^2 \tau_n(k) \cdot \tau_n(k) &= 2(\tau_{n+1}(k) \tau_{n-1}(k) - \tau_n(k) \tau_n(k)), \\ (aD_x - 1) \tau_n(k) \cdot \tau_{n-1}(k-1) + \tau_n(k-1) \tau_{n-1}(k) &= 0, \\ \tau_n(k+1) \tau_n(k-1) - \tau_n(k) \tau_n(k) &= a^2(\tau_{n+1}(k+1) \tau_{n-1}(k-1) - \tau_n(k) \tau_n(k)). \end{aligned} \quad (8.7)$$

References

- [1] Ablowitz M J, Prinari B and Trubatch A D 2004 *Discrete and Continuous Nonlinear Schrödinger Systems* (Cambridge University Press)
- [2] Kevrekidis P G, Rasmussen K O and Bishop A R 2001 *Int. J. Mod. Phys. B*, **15**, 2833
- [3] Ablowitz M J and Ladik J F 1975 *J. Math. Phys.* **16**, 598
- [4] Ablowitz M J and Ladik J F 1976 *J. Math. Phys.* **17**, 1011

- [5] Hirota R 2004 *The Direct Method in Soliton Theory* (Cambridge University Press)
- [6] Hirota R, unpublished work
- [7] Ohta Y 2000 *Chaos, Solitons & Fractals*, **11**, 91
- [8] Sadakane T 2003 *J. Phys. A: Math. Gen.* **36**, 87
- [9] Kivshar Y and Kr'olikowski 1994 *Phys. Rev. E* **50**, 5020
- [10] Sanchez-Rey B and Johansson M 2005 *Phys. Rev. E* **71**, 036627
- [11] Kevrekidis P G, Carretero-Gonzalez R, Theocharis G, Frantzeskakis D J and Malomed B A 2003 *Phys. Rev. E* **68**, 035602
- [12] Vekslerchik V E and Konotop V V 1992 *Inv. Prob.* **8**, 889
- [13] Vekslerchik V E 1995 *Inv. Prob.* **11**, 463
- [14] Kharchev S, Mironov A and Zhedanov A 1997 *Int. J. Mod. Phys. A*, **12**, 2675
- [15] Ohta Y, Kajiwara K, Matsukidaira J and Satsuma J 1993 *J. Math. Phys.* **34**, 5190
- [16] Freeman N C and Nimmo J J C 1983 *Phys. Lett. A* **95**, 1
- [17] Freeman N C and Nimmo J J C 1983 *Proc. Roy. Soc. London Ser. A* **389**, 319
- [18] Ruijsenaars S N M 1990 *Commun. Math. Phys.* **133**, 217

List of MHF Preprint Series, Kyushu University

21st Century COE Program

Development of Dynamic Mathematics with High Functionality

- MHF2003-1 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Yoshitaka WATANABE
A numerical method to verify the invertibility of linear elliptic operators with applications to nonlinear problems
- MHF2003-2 Masahisa TABATA & Daisuke TAGAMI
Error estimates of finite element methods for nonstationary thermal convection problems with temperature-dependent coefficients
- MHF2003-3 Tomohiro ANDO, Sadanori KONISHI & Seiya IMOTO
Adaptive learning machines for nonlinear classification and Bayesian information criteria
- MHF2003-4 Kazuhiro YOKOYAMA
On systems of algebraic equations with parametric exponents
- MHF2003-5 Masao ISHIKAWA & Masato WAKAYAMA
Applications of Minor Summation Formulas III, Plücker relations, Lattice paths and Pfaffian identities
- MHF2003-6 Atsushi SUZUKI & Masahisa TABATA
Finite element matrices in congruent subdomains and their effective use for large-scale computations
- MHF2003-7 Setsuo TANIGUCHI
Stochastic oscillatory integrals - asymptotic and exact expressions for quadratic phase functions -
- MHF2003-8 Shoki MIYAMOTO & Atsushi YOSHIKAWA
Computable sequences in the Sobolev spaces
- MHF2003-9 Toru FUJII & Takashi YANAGAWA
Wavelet based estimate for non-linear and non-stationary auto-regressive model
- MHF2003-10 Atsushi YOSHIKAWA
Maple and wave-front tracking — an experiment
- MHF2003-11 Masanobu KANEKO
On the local factor of the zeta function of quadratic orders
- MHF2003-12 Hidefumi KAWASAKI
Conjugate-set game for a nonlinear programming problem

- MHF2004-1 Koji YONEMOTO & Takashi YANAGAWA
Estimating the Lyapunov exponent from chaotic time series with dynamic noise
- MHF2004-2 Rui YAMAGUCHI, Eiko TSUCHIYA & Tomoyuki HIGUCHI
State space modeling approach to decompose daily sales of a restaurant into time-dependent multi-factors
- MHF2004-3 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Cubic pencils and Painlevé Hamiltonians
- MHF2004-4 Atsushi KAWAGUCHI, Koji YONEMOTO & Takashi YANAGAWA
Estimating the correlation dimension from a chaotic system with dynamic noise
- MHF2004-5 Atsushi KAWAGUCHI, Kentarou KITAMURA, Koji YONEMOTO, Takashi YANAGAWA & Kiyofumi YUMOTO
Detection of auroral breakups using the correlation dimension
- MHF2004-6 Ryo IKOTA, Masayasu MIMURA & Tatsuyuki NAKAKI
A methodology for numerical simulations to a singular limit
- MHF2004-7 Ryo IKOTA & Eiji YANAGIDA
Stability of stationary interfaces of binary-tree type
- MHF2004-8 Yuko ARAKI, Sadanori KONISHI & Seiya IMOTO
Functional discriminant analysis for gene expression data via radial basis expansion
- MHF2004-9 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Hypergeometric solutions to the q -Painlevé equations
- MHF2004-10 Raimundas VIDŪNAS
Expressions for values of the gamma function
- MHF2004-11 Raimundas VIDŪNAS
Transformations of Gauss hypergeometric functions
- MHF2004-12 Koji NAKAGAWA & Masakazu SUZUKI
Mathematical knowledge browser
- MHF2004-13 Ken-ichi MARUNO, Wen-Xiu MA & Masayuki OIKAWA
Generalized Casorati determinant and Positon-Negaton-Type solutions of the Toda lattice equation
- MHF2004-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO
Generating function associated with the determinant formula for the solutions of the Painlevé II equation

- MHF2004-15 Kouji HASHIMOTO, Ryohei ABE, Mitsuhiro T. NAKAO & Yoshitaka WATANABE
Numerical verification methods of solutions for nonlinear singularly perturbed problem
- MHF2004-16 Ken-ichi MARUNO & Gino BIONDINI
Resonance and web structure in discrete soliton systems: the two-dimensional Toda lattice and its fully discrete and ultra-discrete versions
- MHF2004-17 Ryuei NISHII & Shinto EGUCHI
Supervised image classification in Markov random field models with Jeffreys divergence
- MHF2004-18 Kouji HASHIMOTO, Kenta KOBAYASHI & Mitsuhiro T. NAKAO
Numerical verification methods of solutions for the free boundary problem
- MHF2004-19 Hiroki MASUDA
Ergodicity and exponential β -mixing bounds for a strong solution of Lévy-driven stochastic differential equations
- MHF2004-20 Setsuo TANIGUCHI
The Brownian sheet and the reflectionless potentials
- MHF2004-21 Ryuei NISHII & Shinto EGUCHI
Supervised image classification based on AdaBoost with contextual weak classifiers
- MHF2004-22 Hideki KOSAKI
On intersections of domains of unbounded positive operators
- MHF2004-23 Masahisa TABATA & Shoichi FUJIMA
Robustness of a characteristic finite element scheme of second order in time increment
- MHF2004-24 Ken-ichi MARUNO, Adrian ANKIEWICZ & Nail AKHMEDIEV
Dissipative solitons of the discrete complex cubic-quintic Ginzburg-Landau equation
- MHF2004-25 Raimundas VIDŪNAS
Degenerate Gauss hypergeometric functions
- MHF2004-26 Ryo IKOTA
The boundedness of propagation speeds of disturbances for reaction-diffusion systems
- MHF2004-27 Ryusuke KON
Convex dominates concave: an exclusion principle in discrete-time Kolmogorov systems

- MHF2004-28 Ryusuke KON
Multiple attractors in host-parasitoid interactions: coexistence and extinction
- MHF2004-29 Kentaro IHARA, Masanobu KANEKO & Don ZAGIER
Derivation and double shuffle relations for multiple zeta values
- MHF2004-30 Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Generalized partitioned quantum cellular automata and quantization of classical CA
- MHF2005-1 Hideki KOSAKI
Matrix trace inequalities related to uncertainty principle
- MHF2005-2 Masahisa TABATA
Discrepancy between theory and real computation on the stability of some finite element schemes
- MHF2005-3 Yuko ARAKI & Sadanori KONISHI
Functional regression modeling via regularized basis expansions and model selection
- MHF2005-4 Yuko ARAKI & Sadanori KONISHI
Functional discriminant analysis via regularized basis expansions
- MHF2005-5 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Point configurations, Cremona transformations and the elliptic difference Painlevé equations
- MHF2005-6 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Construction of hypergeometric solutions to the q -Painlevé equations
- MHF2005-7 Hiroki MASUDA
Simple estimators for non-linear Markovian trend from sampled data:
I. ergodic cases
- MHF2005-8 Hiroki MASUDA & Nakahiro YOSHIDA
Edgeworth expansion for a class of Ornstein-Uhlenbeck-based models
- MHF2005-9 Masayuki UCHIDA
Approximate martingale estimating functions under small perturbations of dynamical systems
- MHF2005-10 Ryo MATSUZAKI & Masayuki UCHIDA
One-step estimators for diffusion processes with small dispersion parameters from discrete observations
- MHF2005-11 Junichi MATSUKUBO, Ryo MATSUZAKI & Masayuki UCHIDA
Estimation for a discretely observed small diffusion process with a linear drift

- MHF2005-12 Masayuki UCHIDA & Nakahiro YOSHIDA
AIC for ergodic diffusion processes from discrete observations
- MHF2005-13 Hiromichi GOTO & Kenji KAJIWARA
Generating function related to the Okamoto polynomials for the Painlevé IV equation
- MHF2005-14 Masato KIMURA & Shin-ichi NAGATA
Precise asymptotic behaviour of the first eigenvalue of Sturm-Liouville problems with large drift
- MHF2005-15 Daisuke TAGAMI & Masahisa TABATA
Numerical computations of a melting glass convection in the furnace
- MHF2005-16 Raimundas VIDŪNAS
Normalized Leonard pairs and Askey-Wilson relations
- MHF2005-17 Raimundas VIDŪNAS
Askey-Wilson relations and Leonard pairs
- MHF2005-18 Kenji KAJIWARA & Atsushi MUKAIHIRA
Soliton solutions for the non-autonomous discrete-time Toda lattice equation
- MHF2005-19 Yuu HARIYA
Construction of Gibbs measures for 1-dimensional continuum fields
- MHF2005-20 Yuu HARIYA
Integration by parts formulae for the Wiener measure restricted to subsets in $\mathbb{R}^d$
- MHF2005-21 Yuu HARIYA
A time-change approach to Kotani's extension of Yor's formula
- MHF2005-22 Tadahisa FUNAKI, Yuu HARIYA & Mark YOR
Wiener integrals for centered powers of Bessel processes, I
- MHF2005-23 Masahisa TABATA & Satoshi KAIZU
Finite element schemes for two-fluids flow problems
- MHF2005-24 Ken-ichi MARUNO & Yasuhiro OHTA
Determinant form of dark soliton solutions of the discrete nonlinear Schrödinger equation