Convergence of integral and its applications

Izaki, Mamoru
Electrical Communication Laboratory, Ministry of Telecommunications

<https://doi.org/10.5109/12954>

出版情報 : 統計数理研究. 5 (1/2), pp.31-34, 1952-09. Research Association of Statistical Sciences
バージョン :
権利関係 :

CONVERGENCE OF INTEGRAL AND ITS APPLICATIONS

Mamoru IZAKI

(Electrical Communication Laboratory,
Ministry of Telecommunications)

1. Introduction

It is a matter of common knowledge that if $f(t)$ be a continuous function defined over the closed interval $0 \leq t \leq l$, the following sum (1) converges to a finite value as $\max(t_k - t_{k-1}) \rightarrow 0$.

$$(1) \quad S_{(1)} = \sum_{k=1}^n f(\xi_k)(t_k - t_{k-1}),$$

where $t_0 = 0$, $t_n = l$ and $n - 1$ t -points are randomly chosen in the interval $0 \leq t \leq l$, and ξ_k is randomly drawn from the sub-interval $t_{k-1} \leq \xi_k \leq t_k$.

The random variable defined by the sum (1) will converge in probability as merely $n \rightarrow \infty$. We will investigate the aspect of convergence and its applications will be described.

2. Mean value

Since the random variables $(t_1, t_2, \dots, t_{n-1})$ and $(\xi_1, \xi_2, \dots, \xi_n)$ are mutually independent and $(t_1, t_2, \dots, t_{n-1})$ are the order statistics, the mean value of random variable defined by the sum (1) is given by the integral

$$(2) \quad E(S_{(1)}) = \frac{(n-1)!}{l^{n-1}} \int_{0 \leq t_1 \leq t_2 \leq \dots \leq t_{n-1} \leq l} \left(\prod_{k=1}^n \int_{t_{k-1}}^{t_k} \frac{d\xi_k}{t_k - t_{k-1}} \right) \\ \times \left[\sum_{k=1}^n (t_k - t_{k-1}) f(\xi_k) \right] dt_1 dt_2 \dots dt_{n-1}.$$

This integration is easily performed and its result is given by

$$(3) \quad E(S_{(1)}) = \int_0^l f(t) dt.$$

This equation shows that the sum (1) is an unbiased estimator.

3. Variance

Similary the variance of random variable (1) is given by

$$(4) \quad var.(S_{(1)}) = \frac{(n-1)!}{l^{n-1}} \int_{0 \leq t_1 \leq \dots \leq t_{n-1} \leq l} \left(\prod_{k=1}^n \left(\int_{t_{k-1}}^{t_k} \frac{d\xi_k}{t_k - t_{k-1}} \right) \right) \\ \times \left[\sum_{k=1}^n (t_k - t_{k-1}) \{ f(\xi_k) \}^2 + \sum_{i \neq j} (t_i - t_{i-1})(t_j - t_{j-1}) f(\xi_i) f(\xi_j) \right]$$

$$\begin{aligned}
& \times dt_1 dt_2 \cdots dt_{n-1} - \left\{ \int_0^l f(t) dt \right\}^2 \\
& = \frac{(n-1)!}{l^{n-1}} \int_{0 \leq t_1 \leq \cdots \leq t_{n-1} \leq l} \cdots \int_{k=1}^n \left[(t_k - t_{k-1}) \int_{t_{k-1}}^{t_k} \{f(t)\}^2 dt \right. \\
& \quad \left. - \left\{ \int_{t_{k-1}}^{t_k} f(t) dt \right\}^2 \right] dt_1 dt_2 \cdots dt_{n-1},
\end{aligned}$$

but in this case, under the condition that $f(t)$ is continuous, it is difficult that the integral is led to a simple formula. But if $f(t)$ be differentiable, we shall have

$$(5) \quad \text{var.}(S_{(1)}) = 2 \left\{ \frac{1}{l} \int_0^l \left(\frac{df}{dt} \right)^2 dt \right\} \frac{l}{\left(\frac{n}{l} \right)^3} + O\left(\frac{1}{n^4} \right).$$

Hence the variance of random variable (1) is proportional to the average of square of tangent of the function $f(t)$ and the length of interval l , and inversely proportional to the cube of number of the divided strata per unit length.

4. A special case

Here, we will consider a special case such that the interval $0 \leq t \leq l$ is divided into n strata of the equal length l/n by t_k ($k = 1, 2, \dots, n-1$). A sample $(\xi_1, \xi_2, \dots, \xi_n)$ is such that each ξ_k is randomly drawn from the sub-interval $(k-1) \frac{l}{n} \leq t \leq k \frac{l}{n}$. Thus we shall have the finite sum of the following form corresponding to the equation (1)

$$(6) \quad S_{(2)} = \frac{l}{n} \sum_{k=1}^n f(\xi_k).$$

Also in this case, its mean value is the unbiased estimator⁽¹⁾, and when $f(t)$ be differentiable, then the variance is given by

$$(7) \quad \text{var.}(S_{(2)}) = \frac{1}{12} \left\{ \frac{1}{l} \int_0^l \left(\frac{df}{dt} \right)^2 dt \right\} \frac{l}{\left(\frac{l}{n} \right)^3} + O\left(\frac{1}{n^5} \right).$$

It is worthy of notice that the equations (7) and (5) are the similar formula.

5. Some applications

The errors in the integrator, correlator etc. will be estimated by the results of (3), (5) etc. and this paper deals with integrator which was just made at our laboratory. The integration by this computer is carried out in such a way that the integrand is substituted by a step function, hence

it appears that the integration is just the evaluation of the $S_{(1)}$ (eq.(1)), but strictly speaking, the calculation is performed in such a way that the point t is drawn sequentially, so it should be treated as process. If the random variable t is Poisson's process, we shall have the same formulae with eq. (3) and (5) as follows.

Since t is the Poisson's process, we shall have

$$P_r(t_k \leq t_k \leq t_k + dt_k | t_{k-1}) = a e^{-a(t_k - t_{k-1})} dt_k,$$

where a is the mean value of the points taken in a unit length. When a sample $(t_1, t_2, \dots, t_{n-1})$ was taken in the interval $0 \leq t \leq l$ and t'_n in the interval $l \leq t' \leq \infty$, the integration is performed.

While, it may be permissible that ξ is mutually independent of t in our case.

Thus we shall have the following equation corresponding to equation (2)

$$(8) \quad a^n \int_{0 \leq t_1 \leq \dots \leq t_{n-1} \leq l \leq t'_n < \infty} \left[\prod_{k=1}^n \int_{t_{k-1}}^{t_k} \frac{d\xi_k}{t_k - t_{k-1}} \right] \\ \times \left[\sum_{k=1}^n (t_k - t_{k-1}) f(\xi_k) \right] dt_1 dt_2 \dots dt_{n-1} dt'_n.$$

And by comparison with equation (8) and (2), we shall easily have

$$(9) \quad \frac{(al)^{n-1}}{(n-1)!} e^{-al} \int_0^l f(t) dt,$$

but in this case we must treat so as n is a random variable, then the mean value is

$$(10) \quad E(S_{(1)}) = \sum_{n=1}^{\infty} \frac{(al)^{n-1}}{(n-1)!} e^{-al} \int_0^l f(t) dt \\ = \int_0^l f(t) dt.$$

Simiraly, if the integrand $f(t)$ be differentiable, we shall have

$$(11) \quad var(S_{(1)}) = 2 \left\{ \frac{1}{l} \int_0^l \left(\frac{df}{dt} \right)^2 dt \right\} l^4 \sum_{n=1}^{\infty} \frac{1}{n^3} \frac{(al)^{n-1}}{(n-1)!} e^{-al} + O\left(\frac{1}{n^4}\right) \\ \equiv 2 \left\{ \frac{1}{l} \int_0^l \left(\frac{df}{dt} \right)^2 dt \right\} l^4 E \left[\left(\frac{1}{n+1} \right)^3 \right] + O\left(\frac{1}{n^4}\right),$$

and since we can estimate the value of $\{ \}$ in the equation (11) by an appropriate expedient, the estimation of error in the integrator will be performed numerically.

References

- (1) Tosio KITAGAWA: Random Integrations, Bull. Math. Stat. Vol. 4, No. 1 ~ 2, (1950).

Added in proof: To prove (5) and (7), we have assumed that $f(t)$ be analytic in $0 \leq t \leq l$, and expansible in a power series of t whose convergence-radius R is greater than l :

$$f(t) = \sum_{k=0}^{\infty} a_n t^n, \quad |t| < R.$$