On a Non-parametric Test in Life Test

Ishii, Goro
Ikuno Senior High School

<https://doi.org/10.5109/12985>

出版情報 : 統計数理研究. 8 (3/4), pp.73-79, 1959-03. Research Association of Statistical Sciences

バージョン :

権利関係 :

ON A NON-PARAMETRIC TEST IN LIFE TEST

By

Gorô ISHII

(Received November 30, 1958)

§ 1. Introduction. There are various topics in the theory of life test. The purpose of this paper is to treat the problem from the standpoint of "test of fit." In life test we start with M items and stop the test at a certain preassigned time T or stop when the s -th death has occurred, because if we wait till all the death occur we have to wait for a long time, c.f. B. Epstein [2], [3], G. Ishii [5]. In a previous paper, we had treated the test of fit in life test in the case that we stop when the s -th death has occurred, G. Ishii [4]. In the present paper we treat the case when we stop the life test at a certain preassigned time T .

Let X be a random variable having the continuous distribution function. In order to test the hypothesis H_0 that the distribution function is a known function $F(x)$, F. N. David [1] and M. Okamoto [6] have proposed the following non-parametric test:

Let x_i ($i=1, 2, \dots, M$) be M independent observation of a random variable X . There are real numbers $\{a_i\}$ $i=1, 2, \dots, m-1$ such that $F(a_i) - F(a_{i-1}) = 1/m$, $i=1, 2, \dots, m$ where $a_0 = -\infty$, $a_m = +\infty$. $(a_{i-1}, a_i]$ will be called "part." Let v be the number of parts which contain no x 's. If v is too large, we reject H_0 .

Now we shall apply the above non-parametric test to life test.

§ 2. Distribution of v in life test. In the following, we shall treat the life test in the case that we stop the test at a certain preassigned time T , where $F(T) = t$. We call T 'stop time.' We take the stop time T as is equal to a divided point a_n such that $F(T) = F(a_n) = t = n/m$. We start the life test with M items and stop at time T . Suppose that there are N deaths. Then N is a random variable which follows the binomial distribution.

$$(1) \quad P(\mathfrak{N} = N) = \binom{M}{N} t^N (1-t)^{M-N}.$$

N observations are situated in some $(-\infty, a_1]$, $(a_1, a_2]$, $\dots$, $(a_{n-1}, T]$. Let v be the number of parts which contain no x 's in the above n parts.

For fixed N , we have

$$(2) \quad P(v = \nu | \mathfrak{N} = N) = n^{-N} \binom{n}{\nu} \sum_{k=1}^{n-\nu} (-1)^{n-\nu-k} \binom{n-\nu}{k} k^\nu.$$

Then we have

$$(3) \quad P(v = \nu, \mathfrak{N} = N) = P(v = \nu | \mathfrak{N} = N) \cdot P(\mathfrak{N} = N).$$

We put $v^{(s)} = v(v-1) \cdots (v-s+1)$.

Then the s -th factorial moment is written as

$$(4) \quad E(v^{(s)}) = \frac{n!}{(n-s)!} \left(1 - \frac{s}{m}\right)^M.$$

Putting $s = 1, 2$,

$$(5) \quad E(v) = n(1 - 1/m)^M,$$

$$(6) \quad E(v(v-1)) = n(n-1)(1 - 2/m)^M.$$

If $M \rightarrow \infty$, $m \rightarrow \infty$, $n \rightarrow \infty$ under the restriction of $M = mr$ (r is a constant) and $n = mt$ (t is a constant),

$$(7) \quad E(v/n) = e^{-r}(1 - r/2m) + O(m^{-2}),$$

$$(8) \quad D^2(v/n) = e^{-2r}(e^r - 1 - tr)/n + O(m^{-2}).$$

Under the above conditions we have next theorem.

Theorem 1. v/n is asymptotically normally distributed with mean e^{-r} and variance $e^{-2r}(e^r - 1 - tr)/n$, where $M = mr$, $n = mt$, (r and t are constants).

Proof. It is sufficient to prove that the moments of $(n/c)^{1/2}(v/n - e^{-r})$ tend to the moments of the standard normal distribution, where $c = e^{-2r}(e^r - 1 - tr)$. The proof is almost parallel to that of theorem 1 of M. Okamoto [6], and theorem 2 of B. Sherman [7].

Denoting by $B_r^{(n)}$ the Bernoulli's number of order n and degree r .

$$\begin{aligned} (9) \quad E \left\{ \left(\frac{n}{c} \right)^{1/2} \left(\frac{v}{n} - e^{-r} \right) \right\}^l &= \left(\frac{n}{c} \right)^{l/2} \sum_{k=0}^l \binom{l}{k} (-e^{-r})^{l-k} E \left(\frac{v}{n} \right)^k \\ &= \left(\frac{n}{c} \right)^{l/2} \sum_{k=0}^l \binom{l}{k} (-e^{-r})^{l-k} n^{-k} \sum_{q=0}^k \binom{k}{q} B_q^{(q-k)} E(v^{(k-q)}) \\ &= \left(\frac{n}{c} \right)^{l/2} \sum_{k=0}^l \sum_{q=0}^k \frac{l! (-1)^{l-k} e^{-r(l-k)}}{k! (l-k)!} \frac{k!}{q! (k-q)!} B_q^{(q-k)} \frac{n! (m-k+q)^M}{(n-k+q)! m^M n^k} \\ &= \frac{(-1)^l n^{l/2} l!}{(e^r - 1 - tr)^{l/2}} \left\{ a_0 + \frac{a_1}{n} + \frac{a_2}{n^2} + \cdots + \frac{a_\alpha}{n^\alpha} + \cdots \right\}. \end{aligned}$$

We have to evaluate these a_i . In the above expansion

$$\begin{aligned} (10) \quad \frac{n! (m-k+q)^M}{(n-k+q)! m^M n^k} &= \frac{1}{n^k} \left(1 - \frac{k-q}{m} \right)^M n(n-1) \cdots (n-k+q+1) \\ &= n^{-q} \left(1 - \frac{k-q}{m} \right)^M \left(1 - \frac{1}{n} \right) \left(1 - \frac{2}{n} \right) \cdots \left(1 - \frac{k-q-1}{n} \right) \end{aligned}$$

$$\begin{aligned}
 &= x^q \{1 - (k-q)tx\}^M \prod_{j=1}^{k-q-1} (1-jx) \\
 &= x^q \cdot G(x),
 \end{aligned}$$

where

$$\begin{aligned}
 x &= 1/n \\
 G(x) &= \{1 - (k-q)tx\}^{\frac{r}{xt}} \prod_{j=1}^{k-q-1} (1-jx).
 \end{aligned}$$

Expanding this, we have

$$G(x) = a_{kq0} + a_{kq1}x + a_{kq2}x^2 + \dots,$$

where

$$\begin{aligned}
 (11) \quad a_{kq0} &= e^{-(k-q)r} \\
 a_{kqp} &= \frac{1}{p!} \left\{ \frac{d^p G(x)}{dx^p} \right\}_{x=0} \\
 &= \frac{1}{p!} \sum_{s=0}^{p-1} \binom{p-1}{s} \frac{d^{p-1} G(x)}{dx^{p-1}} \frac{d^{s+1} \log G(x)}{dx^{s+1}} \Big|_{x=0} \\
 \log G(x) &= - \sum_{j=1}^{k-q-1} \left(jx + \frac{j^2 x^2}{2} + \frac{j^3 x^3}{3} + \dots + \frac{j^{s+1} x^{s+1}}{s+1} + \dots \right) \\
 &\quad - \frac{r}{xt} \left((k-q)tx + \frac{(k-q)^2 t^2 x^2}{2} + \dots + \frac{(k-q)^{s+2} t^{s+2} x^{s+2}}{s+2} + \dots \right).
 \end{aligned}$$

Therefore

$$\begin{aligned}
 (12) \quad \frac{d^{s+1} \log G(x)}{dx^{s+1}} \Big|_{x=0} &= - \sum_{j=1}^{k-q-1} s! j^{s+1} - \frac{(s+1)! (k-q)^{s+2}}{s+2} r t^{s+1} \\
 &= s! b_{kqs} \quad (\text{say}) \\
 b_{kqs} &= - \sum j^{s+1} - \frac{s+1}{s+2} r t^{s+1} (k-q)^{s+2}.
 \end{aligned}$$

Then b_{kqs} is a polynomial in k of degree $s+2$.

Putting $s=0$,

$$\begin{aligned}
 b_{kq0} &= - \frac{(k-q)(k-q-1)}{2} - \frac{1}{2} r t (k-q)^2 \\
 &= - (1+rt) \frac{k^2}{2} + Ak + B \quad (\text{say}).
 \end{aligned}$$

From (11),

$$\begin{aligned}
 (13) \quad a_{kpq} &= \frac{1}{p!} \sum_{s=0}^{p-1} \frac{(p-1)!}{s! (p-s-1)!} a_{kq(p-1-s)} (p-s-1)! s! b_{kqs} \\
 &= \frac{1}{p} \sum_{s=0}^{p-1} a_{kq(p-1-s)} b_{kqs}.
 \end{aligned}$$

Then we have next equations

$$\begin{aligned}
 a_{kq0} &= e^{-(k-q)s} \\
 a_{kq0}b_{kq0} - a_{kq1} &= O \\
 a_{kq0}b_{kq1} + a_{kq1}b_{kq0} - 2a_{kq2} &= O \\
 &\dots\dots\dots \\
 a_{kq0}b_{kq(i-q-1)} + a_{kq1}b_{kq(i-q-2)} + \dots + a_{kq(i-q-1)}b_{kq0} - (i-q)a_{kq(i-q)} &= O.
 \end{aligned}$$

Hence we have

$$(14) \quad a_{kq(i-q)} = e^{-(k-q)r} B_{kqi},$$

where

$$\begin{aligned}
 B_{kqi} &= \frac{(b_{kq0})^{i-q}}{(i-q)!} + (\text{terms of lower degree in } k) \\
 &= \frac{1}{(i-q)!} \left(-\frac{1+rt}{2} \right)^{i-q} k^{2(i-q)} + \sum_{j=0}^{2(i-q)-1} A_j k^j.
 \end{aligned}$$

In (9) we put

$$\begin{aligned}
 (15) \quad & \sum_{q=0}^l \sum_{k=q}^l \frac{(-1)^k e^{rk}}{q! (l-k)! (k-q)!} B_q^{(q-k)} \frac{n! (m-k+q)^M}{(n-k+q)! m^M n^k} \\
 &= a_0 + \frac{a_1}{n} + \frac{a_2}{n^2} + \dots + \frac{a_\alpha}{n^\alpha} + \dots.
 \end{aligned}$$

If we denote by a_{iq} the coefficient of n^{-i} in the expansion in the power of n^{-1} of

$$\begin{aligned}
 & \sum_{k=q}^l \frac{(-1)^k e^{rk}}{(l-k)! (k-q)!} B_q^{(q-k)} \frac{n! (m-k+q)^M}{(n-k+q)! m^M n^k} \\
 &= \sum_{k=q}^l \frac{(-1)^k e^{rk}}{(l-k)! (k-q)!} B_q^{(q-k)} x^q \{a_{kq0} + a_{kq1}x + \dots + a_{kq(i-q)}x^{i-q} + \dots\},
 \end{aligned}$$

we have

$$(16) \quad a_i = \sum_{q=0}^l \frac{a_{iq}}{q!}$$

and

$$\begin{aligned}
 a_{iq} &= \sum_{k=q}^l \frac{(-1)^k e^{rk}}{(l-k)! (k-q)!} B_q^{(q-k)} a_{i,q(i-q)} \\
 &= e^{qr} \sum_{k=q}^l \frac{(-1)^k}{(l-k)! (k-q)!} B_q^{(k-b)} B_{kqi}.
 \end{aligned}$$

$B_q^{(q-k)}$ is the coefficient of the expansion of $(q/(e^t - 1))^{q-k}$, then $B_q^{(q-k)}$ is the polynomial in k of degree q with 2^{-q} as the coefficient of the term of highest degree. Therefore

$$\begin{aligned}
 (17) \quad a_{iq} &= e^{qr} \sum_{k=q}^l \frac{(-1)^k}{(l-k)! (k-q)!} \left\{ \frac{k^q}{2^q} + \dots \right\} \left\{ \frac{k^{2(i-q)}}{(i-q)!} \left(-\frac{1+tr}{2} \right)^{i-q} + \sum A_j k^j \right\} \\
 &= \frac{e^{qr}}{2^q (i-q)!} \left(-\frac{1+tr}{2} \right)^{i-q} \sum_{k=q}^l \frac{(-1)^k}{(l-k)! (k-q)!} \left\{ k^{2i-q} + \text{terms of lower degree in } k \right\}.
 \end{aligned}$$

As

$$\sum_{k=q}^l \frac{(-1)^k k^{l'}}{(l-k)! (k-q)!} = \begin{cases} 0 & l' < l-q \\ 1 & l' = l-q \end{cases}$$

we obtain

$$\begin{aligned}
 a_{iq} &= 0 \quad i < \frac{l}{2} \\
 a_{hq} &= \frac{e^{qr} \left(-\frac{1+tr}{2} \right)^{i-q}}{2^q (h-q)!} \quad i = \frac{l}{2} = h \quad (\text{when } l \text{ is even}).
 \end{aligned}$$

Thus from (16)

$$\begin{aligned}
 a_i &= 0 \quad \text{when } i < \frac{l}{2} \\
 a_h &= \sum_{q=0}^{2h} \frac{a_{hq}}{q!} = \sum_{q=0}^{2h} \frac{1}{q! (h-q)!} \left(\frac{e^r}{2} \right)^q \left(-\frac{1+tr}{2} \right)^{h-q} \\
 &= \frac{1}{h!} \frac{(e^r - 1 - tr)^h}{2^h}.
 \end{aligned}$$

Then we have

$$\begin{aligned}
 \lim_{n \rightarrow \infty} E \left\{ \left(\frac{n}{c} \right)^{1/2} \left(\frac{v}{n} - e^{-r} \right) \right\}^l &= 0 \quad (\text{when } l \text{ is odd}) \\
 \lim_{n \rightarrow \infty} E \left\{ \left(\frac{n}{c} \right)^{1/2} \left(\frac{v}{n} - e^{-r} \right) \right\}^{2h} &= \frac{a_h (2h)!}{(e^r - 1 - tr)^h} \\
 &= \frac{(2h)!}{2^h h!} \quad (\text{when } l \text{ is even, } l = 2h).
 \end{aligned}$$

This completes the proof.

§ 3. The power function. The power function of the test with respect to the alternative hypothesis H_1 is

$$P = P(v \geq l | H_1).$$

Under H_1 : distribution function is $F_1(x)$, and the range of distribution is equal to that of $F(x)$, where $F(T) = \tau$

$$\int_{t-1/m}^{t/m} dH(x) = p_i, \quad i = 1, 2, \dots, n$$

$$\text{where } H(x) = F_1(F^{-1}(x)), \quad \sum_{i=1}^n p_i = \tau.$$

We put $p_i/\tau = q_i$, then $\sum_{i=1}^n q_i = 1$

$$P(\mathfrak{N} = N | H_1) = \binom{M}{N} \tau^N (1 - \tau)^{M-N}$$

$$P(v = k | \mathfrak{N} = N | H_1) = \sum_{j=1}^{n-k} (-1)^{n-k-j} \binom{n-j}{k} \sum_{(i_1, \dots, i_j)}^n (q_{i_1} + \dots + q_{i_j})^N,$$

where $\sum_{(i_1, \dots, i_j)}^n$ denotes the summation over all combinations $(i_1, \dots, i_j)$ drawn from $(1, 2, \dots, j)$.

$$P(v = k, \mathfrak{N} = N | H_1) = P(v = k | \mathfrak{N} = N | H_1) \cdot P(\mathfrak{N} = N | H_1)$$

$$\begin{aligned} E(v^{(s)} | H_1) &= \sum_N s! \sum_{(i_1, \dots, i_s)}^n (1 - q_{i_1} - \dots - q_{i_s})^N \binom{M}{N} \tau^N (1 - \tau)^{M-N} \\ &= s! \sum_{(i_1, \dots, i_s)}^n (1 - p_{i_1} - \dots - p_{i_s})^M. \end{aligned}$$

Putting $s = 1, 2$, we get

$$E(v | H_1) = \sum_{i=1}^n (1 - p_i)^M$$

$$E(v(v-1) | H_1) = \sum_{i \neq j}^n (1 - p_i - p_j)^M.$$

If $F_1(x)$ is absolutely continuous with respect to $F(x)$ and its relative density is differentiable, putting $H'(x) = h(x)$,

$$p_i = \frac{1}{m} h\left(\frac{i}{m}\right) - \frac{1}{2m^2} h'\left(\frac{i}{m}\right) + O(m^{-3})$$

$$(1 - p_i)^M = e^{-r h\left(\frac{i}{m}\right)} \left[1 + \frac{r}{2m} \left\{ h'\left(\frac{i}{m}\right) - h^2\left(\frac{i}{m}\right) \right\} + O(m^{-2}) \right]$$

$$E(v/n | H_1) = \frac{1}{t} \int_0^t e^{-r h(x)} dx - \frac{r}{2mt} \int_0^t e^{-r h(x)} h^2(x) dx + O(m^{-2}).$$

The power function

$$P = \sum_{k \geq l} \sum_{N=0}^M P(v = k, \mathfrak{N} = N | H_1)$$

satisfies the next condition when $\tau \leq t$

$$P \geq P(v \geq l | H_0).$$

This inequality can be proved by the same method in theorem 3 of Okamoto [6].

Thus we have the next theorem.

Theorem 2. *The test based on v is unbiased against the class of alternative hypotheses, $\tau \leq t$.*

Acknowledgement. The author is deeply indebted to Mr. M. Okamoto whose helpful guidance made this work possible.

IKUNO SENIOR HIGH SCHOOL*

References

- [1] F. N. DAVID; "*Two combinatorial tests of whether a sample has come from a given population*" *Biometrika*, vol. **37** (1950), pp. 97-110.
- [2] B. EPSTEIN; "*Truncated life tests in the exponential case*" *Ann. Math. Stat.*, vol. **25** (1954), pp. 555-64.
- [3] B. EPSTEIN; "*Simple estimator of the parameter of exponential distributions when sample are censored*" *Ann. Inst. Stat. Math.*, vol. **8** (1956-7), pp. 15-26.
- [4] G. ISHII; "*Test of fit in life test*" *Ann. Inst. Stat. Math.*, vol. **9** (1957-8) pp. 117-125.
- [5] G. ISHII; "*Life test in two dimensional exponential distribution*" in Japanese. *Tokei Kagaku Kenkyu*, vol. **2** (1958).
- [6] M. OKAMOTO; "*On a non-parametric test*" *Osaka Math. J.*, vol. **4** (1952), pp. 77-85.
- [7] B. SHERMAN; "*A random variable related to the spacing of sample values*" *Ann. Math. Stat.*, vol. **22** (1950), pp. 339-361.
- [8] G. ISHII; "*Kolmogorov-Smirnov test in life test*" *Ann. Inst. Stat. Math.*, vol. **10** (1958-9) pp. 37-46.
- [9] I. WEISS; "*Limiting distribution in some occupancy problem*" *Ann. Math. Stat.*, vol. **29** (1958) pp. 878-884.

* Present address: Atomic Bomb Casualty Commission, Hiroshima.