ON A STATISTICAL ANALYSIS OF HOMOGENEOUS RANDOM FIELDS ON A METRIC SPACE ACTED UPON BY A COMPACT METRIC GROUP

Nagai, Takeaki
Faculty of Science, Kagoshima University

<https://doi.org/10.5109/13037>

出版情報 : 統計数理研究. 13 (3/4), pp.65-92, 1969-03. Research Association of Statistical
Sciences
バージョン :
権利関係 :

ON A STATISTICAL ANALYSIS OF HOMOGENEOUS RANDOM FIELDS ON A METRIC SPACE ACTED UPON BY A COMPACT METRIC GROUP

By

Takeaki NAGAI*

(Received Jan. 5, 1969)

- § 1. Introduction and summary.
- § 2. Preliminaries.
- § 3. On the optimal linear interpolation of homogeneous random fields I.
- § 4. On the optimal linear interpolation of homogeneous random fields II.
- § 5. On the uniformly minimum variance unbiased linear estimates of mean value functions of homogeneous random fields.
- § 6. Appendix.
- § 7. Acknowledgment.

§ 1. Introduction and summary.

In this paper, we shall consider a problem on statistical analysis of second order homogeneous random fields $\{X(t), t \in T\}$ on a metric space T acted upon by a compact metric group G such that each $g \in G$ is a homeomorphism acting on T and the mapping $(g, t) \rightarrow gt$ from $G \times T$ into T is continuous.

The fields $\{X(t), t \in T\}$ are observable on a compact set $A \subset T$ invariant under every $g \in G$.

As preliminaries, we shall consider in § 2 a measure μ on a compact metric space A being a topological subspace of T invariant under every $g \in G$ such that for every $g \in G$ and any Borel set A of A , $\mu(gA) = \mu(A)$. By a measure on a metric space we always mean a non-negative countably additive set-function defined on the σ -field of Borel subsets of the metric space.

It is shown in § 2 that there exists a Borel set $V \subset A$ such that (i) $G \cdot V = A$ and (ii) $Gv_1 \cap Gv_2 = \emptyset$ if $v_1 \neq v_2$ ($v_1, v_2 \in V$), (lemma 2.1) and that a mapping $\xi: (g, v) \rightarrow gv$ is a Borel isomorphism between $G \times V$ and A , (lemma 2.2). We shall call V the Borel cross section of A with respect to G or simply the Borel cross section of A .

It is shown in theorem 2.1 that the induced measure $\mu^* = \mu\xi$ on $G \times V$ is decomposed into a direct product measure of the normalized Haar measure on G and a measure m on the Borel cross section V in such a way that for any Borel set $A \times B \subset G \times V$,

$$\mu^*(A \times B) = \nu(A) \cdot m(B)$$

* Faculty of Science, Kagoshima University, Kagoshima.

where ν is the normalized Haar measure on G and m is a measure on V . This result will be used in § 4.

We shall consider in § 3 the optimal linear estimator $\hat{Z}$ of a random variable $Z \in L_2(X(t), t \in T)$ such that $Z \in L_2(X(t), t \in A)$ and $U(g)Z = Z$ for all $g \in G$ where for any $S \subset T$ $L_2(X(t), t \in S)$ is a Hilbert space spanned by linear combinations of the field $\{X(t), t \in S\}$ under the scalar product (U, W) defined by $(U, W) = E\{U \cdot \overline{W}\}$ and $\{U(g), g \in G\}$ are unitary operators on $L_2(X(t), t \in T)$ defined by writing $U(g)X(t) = X(gt), t \in T$.

It is shown in theorem 3.1 that for any small $\varepsilon > 0$ there exists a random variable $Y \in L_2(X(t), t \in V)$ such that

$$E\left\{\left|\hat{Z} - \int_G U(g)dg \cdot Y\right|^2\right\} < \varepsilon$$

where V is a Borel cross section of A . The operator $\int_G U(g)dg$ is well-defined on $L_2(X(t), t \in A)$ and actually the projection operator onto the closed subspace V_1 consisting of all random variables invariant under every $U(g)$, $g \in G$ in $L_2(X(t), t \in A)$.

The situations being the same as in § 3, we shall proceed in § 4 further in details to construct more practical approximating sequence $\{Y_n, n = 1, 2, 3, \dots\}$ for the optimal linear estimator $\hat{Z}$ of the random variable Z invariant under every $U(g)$, $g \in G$ such that

$$\hat{Z} = \text{l.i.m}_{n \rightarrow \infty} \int_G U(g)dg \cdot Y_n,$$

where $Y_n \in L_2(X(t), t \in V)$, $n = 1, 2, 3, \dots$, are given by the followings:

$$Y_n = \int_V X(s)H_n(s)dm(s), \quad n = 1, 2, 3, \dots,$$

where $\{H_n(s), s \in V, n = 1, 2, 3, \dots\}$ is a sequence of continuous bounded functions on V which are determined by recursively and the measure m is determined by a measure on A invariant under every $g \in G$.

In § 5 the metric space T has the same properties as in the previous sections except for T being compact and T itself being invariant under every $g \in G$. We shall consider a second order homogeneous random field $\{X(t), t \in T\}$ with a mean value function $m(t) = E\{X(t)\}$, $t \in T$. We assume that the field is observable on T . Let V be a Borel cross section of T with respect to G , that is, V is a Borel set of T such that $G \cdot V = T$ and $Gv_1 \cap Gv_2 = \emptyset$ if $v_1 \neq v_2$ ($v_1, v_2 \in V$).

Then it is shown in theorem 5.3 that when the mean value function $m(t)$, $t \in T$, is invariant under every $g \in G$, that is, $m(t) = m(gt)$, for all $g \in G$, and all $t \in T$, the uniformly minimum variance unbiased linear estimator $\hat{m}(t)$ of the mean value function $m(t)$ at $t \in T$ is given by $\hat{m}(t) = \int_G U(g)dg \cdot X(v)$, where for some $h \in G$ the point t is written as $t = hv$.

As an appendix, we shall write in § 6 several theorems which are referred to in our discussions.

§ 2. Preliminaries.

In this section, we shall consider on decomposability of a measure μ on a compact set A of a metric space T acted upon by a compact metric group G .

Let T be a metric space. Let G be a compact metric group of homeomorphisms acting on T such that the mapping $(g, t) \rightarrow gt$ from $G \times T$ into T is continuous.

For any metric space X , we shall denote by $\mathfrak{B}_X$ the σ -field of Borel subsets of X , that is, $\mathfrak{B}_X$ is the smallest σ -algebra of subsets of X containing all open sets.

Lemma 2.1. *Let $A \subset T$ be a compact set invariant under every $g \in G$. Then, there exists a Borel set $V \subset A$ such that*

$$(1) \quad G \cdot V = A,$$

$$(2) \quad G \cdot v_1 \cap G \cdot v_2 = \emptyset \quad \text{if} \quad v_1 \neq v_2 \quad (v_1, v_2 \in V).$$

We shall call a Borel set V satisfying (1) and (2) a Borel cross section of A with respect to G or simply a Borel cross section of A .

Proof. For any two points t_1, t_2 in A we shall say that $t_1 \sim t_2$ if there exists $g \in G$ such that $t_1 = gt_2$. " $\sim$ " is an equivalence relation. Let $[t]$ denote the equivalence class containing $t \in A$. Let M be the space of all such equivalence classes. The mapping $t \rightarrow [t]$ from A into M is continuous under the quotient topology and onto. Thus, the space M is a continuous image of the compact set A under the mapping and hence M is a compact metric space.

Thus, by a theorem of Federer and Morse (see theorem 6.1 in § 6) it follows that there exists a Borel set $V \subset A$ satisfying (1) and (2). The proof of lemma 2.1 is complete.

We shall show that the space A may be identified with the product space $G \times V$.

Lemma 2.2. *Suppose that for any $g \in G$, $g \neq e$, there is no fixed point in A .*

Then, the mapping

$$(3) \quad \xi: (g, v) \rightarrow gv$$

is a Borel isomorphism between $G \times V$ and A .

Proof. From lemma 2.1 the mapping $\xi: (g, v) \rightarrow gv$ from $G \times V$ into A is onto. Since there is no fixed point in A for any $g \in G$, $g \neq e$, the mapping ξ is one-one. From our assumption that the mapping $(g, t) \rightarrow gt$ from $G \times T$ into T is continuous, it is clear that the mapping ξ is continuous. In particular, ξ is measurable.

Hence, by a theorem of Kuratowski (see theorem 6.2 in § 6) the inverse mapping ξ^{-1} is also measurable. Thus, we have proved lemma 2.2.

Remark. Lemma 2.2 may be proved alternatively by making use of a theorem 6.3 in § 6 in the virtue of the space A being compact.

Let μ be a measure on A invariant under every $g \in G$, that is, for each $g \in G$, $\mu(gA) = \mu(A)$, $A \in \mathfrak{B}_A$.

Without loss of generality, we assume that $\mu(A) = 1$, in other words, μ is a probability measure on A .

Let μ^* be the induced measure on $G \times V$ by the Borel isomorphism ξ from the measure μ on A .

Then, we have the following:

Theorem 2.1. *The measure μ^* on $G \times V$ is decomposed into a direct product measure in such a way that for any Borel set $A \times B \subset G \times V$,*

$$\mu^*(A \times B) \equiv \mu(\xi(A \times B)) = \nu(A) \cdot m(B),$$

where ν is the normalized Haar measure on G and m a probability measure on the Borel cross section V of A .

Proof. Both $G \times V$ and G are compact metric spaces and hence these are complete separable metric spaces. Therefore, these are automatically separable standard Borel spaces (see [10], page 133).

Let us consider the mapping $\pi: (g, v) \rightarrow g$ from $G \times V$ onto G . Then it is clear that π is measurable since the mapping π is a projection from the product space $G \times V$ onto its coordinate space G .

Thus, it follows that there exists a regular conditional probability distribution of μ^* given π , which we shall denote by $\tilde{m}_g(A \times B)$, $g \in G$, $A \times B \in \mathfrak{B}_{G \times V}$.

This satisfies the following conditions:

- (i) For each $g \in G$, $\tilde{m}_g(\cdot)$ is a probability measure on $G \times V$.
- (ii) For each Borel set $A \times B \subset G \times V$, the mapping $g \rightarrow \tilde{m}_g(A \times B)$ is $\mathfrak{B}_G$ -measurable.
- (iii) For any Borel set $A \times B \subset G \times V$,

$$\mu^*(A \times B) = \int_G \tilde{m}_g(A \times B) d\nu(g),$$

where ν is a probability measure on G such that $\nu(E) = \mu^*(\pi^{-1}(E)) = \mu^*(E \times V)$, $E \in \mathfrak{B}_G$. (See [10], page 146).

In particular, we have for any $E \in \mathfrak{B}_G$,

$$(4) \quad \begin{aligned} \mu^*(A \times B \cap \pi^{-1}(E)) &\equiv \mu^*((A \cap E) \times B) \\ &= \int_E \tilde{m}_g(A \times B) d\nu(g), \quad A \times B \in \mathfrak{B}_{G \times V}. \end{aligned}$$

Let us write

$$m_g(B) = \tilde{m}_g(G \times B), \quad B \in \mathfrak{B}_V.$$

Then we have by letting $A = G$ in (4),

$$(5) \quad \mu^*(E \times B) = \int_E m_g(B) d\nu(g),$$

for any $E \in \mathfrak{B}_G$ and any $B \in \mathfrak{B}_V$, where $m_g(\cdot)$ is a probability measure on V .

Since for each $g \in G$ and any Borel sets A and B of G and V respectively, $\mu^*(gA \times B) = \mu(\xi(gA \times B)) = \mu(\xi(A \times B)) = \mu^*(A \times B)$, it follows by letting particularly $B = V$ in (5) that for any $A \in \mathfrak{B}_G$,

$$\nu(gA) = \mu^*(gA \times V) = \mu^*(A \times V) = \nu(A).$$

This equality and the uniqueness of the invariant measure on G imply that the measure ν is the normalized Haar measure on G .

Thus, we may write

$$\mu^*(A \times B) = \int_A m(B) dg, \quad A \times B \in \mathfrak{B}_{G \times V}.$$

Hence we have for any $A \in \mathfrak{B}_G$, $B \in \mathfrak{B}_V$ and $h \in G$,

$$\begin{aligned}\mu^*(A \times B) &= \int_A m_g(B) dg \\ &= \mu^*(hA \times B) \\ &= \int_{hA} m_g(B) dg \\ &= \int_A m_{h^{-1} \cdot g}(B) dg.\end{aligned}$$

This implies that for any $B \in \mathfrak{B}_V$ and any $g, h \in G$,

$$m_g(B) = m_h(B),$$

in other words, $m_g(\cdot)$ is a probability measure on V independent of $g \in G$.

Let us write

$$m(B) = m_g(B), \quad B \in \mathfrak{B}_V.$$

Then we have for any Borel set $A \times B \subset G \times V$,

$$\begin{aligned}\mu^*(A \times B) &= \int_A m(B) dg \\ &= \nu(A) \cdot m(B),\end{aligned}$$

where ν is the normalized Haar measure on G and m a probability measure on V . Thus, we have proved theorem 2.1.

Example 2.1. Let $T = R_2 = \{(x, y) \mid -\infty < x < \infty, -\infty < y < \infty\}$. Let $G = SO(2)$, the group consisting of all rotations on T around the origin $(0, 0)$. Let us consider a closed disk A with a finite radius r , that is, $A = \{(x, y) \in R_2 \mid 0 \leq x^2 + y^2 \leq r^2\}$.

It is obvious that as a Borel cross section of A , we may take a set $V = \{(x, 0) \mid 0 \leq x \leq r\}$.

Since $SO(2)$ is isomorphic to the torus group $G_0 = \{\theta \mid 0 \leq \theta < 2\pi\}$ with the group operation $\theta_1 + \theta_2 = \theta_3$ where $\theta_1 + \theta_2 \equiv \theta_3 \pmod{2\pi}$, $0 \leq \theta_3 < 2\pi$, we may identify the space A with the product space $V \times G_0$.

Let μ be a probability measure on A such that for any $A \in \mathfrak{B}_A$, $\mu(A)$ is proportional to the "area" of the set A , in other words, μ is a uniform distribution on A .

Then μ is invariant under every $g \in SO(2)$ and it is written as follows:

For any Borel set $A \subset G_0$ and $B \subset V$,

$$\int_{B \times A} d\mu(x, \theta) = \int_B \frac{2}{r^2} x dx \cdot \int_A \frac{1}{2\pi} d\theta.$$

Example 2.2. Let $L_3 = \{(t, x, y, z) \mid t^2 - (x^2 + y^2 + z^2) = c^2\}$ where t, x, y, z and c are real numbers. Let G be a subgroup of the proper Lorentz group of order (3.1) such that its elements are of the following form:

$$g = \begin{pmatrix} 1 & 0 \\ 0' & h \end{pmatrix}, \quad h \in SO(3),$$

where 0 denotes a zero vector $(0, 0, 0)$ and $0'$ its transposed one.

Let $A = \{(t, x, y, z) \in L_3 \mid 0 \leq x^2 + y^2 + z^2 \leq \alpha^2\}$ where α is a finite real number.

Since A is invariant under every $g \in G$, we may take a set

$$V = \{(\sqrt{c^2+u^2}, u, 0, 0) \mid 0 \leq u \leq \alpha\} \subset A$$

as a Borel cross section of A .

Let μ be a probability measure on A such that for any $C \in \mathfrak{B}_A$, $\mu(C)$ is proportional to the "volume" of C , in other words, μ is uniformly distributed on A . Then μ is invariant under every $g \in G$. Thus, the measure μ is decomposed into a direct product measure as follows:

$$\begin{aligned} \int_C d\mu(x, y, z) &= \int_{B \times (A_1 \times A_2)} d\mu(u, \theta, \varphi) \\ &= \int_B \frac{\beta u^2}{\sqrt{c^2+u^2}} du \cdot \frac{1}{4\pi} \int_{A_1 \times A_2} \sin \theta d\theta d\varphi \end{aligned}$$

where C is a set of all points (t, x, y, z) in A such that

$$\begin{aligned} x &= u \cdot \sin \theta \cdot \cos \varphi, \\ y &= u \cdot \sin \theta \cdot \sin \varphi, \\ z &= u \cdot \cos \theta, \\ t &= \sqrt{c^2+u^2}, \end{aligned}$$

$(\theta, \varphi) \in A_1 \times A_2 \in \mathfrak{B}_G$ and $u \in B \in \mathfrak{B}_V$ (u is identified with a point $(\sqrt{c^2+u^2}, u, 0, 0)$ in V) and

$$\beta = 2 \cdot [\alpha \cdot \sqrt{\alpha^2+c^2} + c^2 \cdot \log(|c| - |\alpha + \sqrt{\alpha^2+c^2}|)]^{-1}.$$

§ 3. On the optimal linear interpolation of homogeneous random fields I.

3.1. Summary.

In this section we shall consider an interpolation problem of a homogeneous random field $\{X(t), t \in T\}$ on a metric space T acted upon by a compact metric group G of homeomorphisms acting on T such that the mapping $(g, t) \rightarrow gt$ from $G \times T$ into T is continuous.

The field $\{X(t), t \in T\}$ is observable on a compact set $A \subset T$ invariant under every $g \in G$.

Let $Z \in L_2(X(t), t \in T)$ be a random variable invariant under all unitary operators $U(g)$, $g \in G$ defined on $L_2(X(t), t \in T)$ by writing $U(g)X(t) = X(gt)$, $t \in T$. Let $\hat{Z} \in L_2(X(t), t \in A)$ be the best linear estimator of Z on the basis of the realization of $X(t)$ on A .

Let V be a subset of A such that $A = G \cdot V$ and $G \cdot v_1 \cap G \cdot v_2 = \emptyset$ if $v_1 \neq v_2$ ($v_1, v_2 \in V$).

Then it will be shown the following result:

For each $\varepsilon > 0$, there exists a random variable $Y \in L_2(X(t), t \in V)$ such that $E\{|\hat{Z} - AY|^2\} < \varepsilon$ where $A = \int_G U(g) dg$ whose restriction to $L_2(X(t), t \in A)$ is a projection operator onto the maximal closed subspace V_1 invariant under all $U(g)$, $g \in G$, that is, $V_1 = \{u \mid U(g)u = u, \text{ for all } g \in G\} \subset L_2(X(t), t \in T)$.

3.2. An interpolation problem of homogeneous random fields.

Let T be a metric space. Let G be a compact metric group of homeomorphisms acting on T such that the mapping $(g, t) \rightarrow gt$ from $G \times T$ into T is continuous.

Let $\{X(t), t \in T\}$ be a complex-valued homogeneous random field satisfying the conditions:

- (1) $E\{X(t)\} = 0$, for all $t \in T$.
- (2) $E\{|X(t)|^2\} < \infty$, for all $t \in T$.
- (3) The covariance function of the field $K(t, s) = E\{X(t) \cdot \overline{X(s)}\}$ is a continuous positive definite function on $T \times T$.
- (4) For each $g \in G$,

$$K(t, s) = K(gt, gs), \quad t, s \in T.$$

Let $A \subset T$ be a compact set invariant under every $g \in G$ and the field $\{X(t), t \in T\}$ be observable on A .

For a set $S \subset T$ we shall denote by $L_2(X(t), t \in S)$ a Hilbert space consisting of all random variables which may be represented either as a finite linear combinations

$$U = \sum_{i=1}^n c_i \cdot X(t_i)$$

for some integer n , points $t_1, t_2, \dots, t_n$ in S and complex numbers $c_1, c_2, \dots, c_n$ or as a limit in quadratic mean of such finite linear combinations under the scalar product (U, W) defined by $(U, W) = E\{U \cdot \overline{W}\}$.

It is well-known that for each $g \in G$ one can define a unitary operator $U(g)$ on $L_2(X(t), t \in T)$ such that

- (5) $U(g)X(t) = X(gt)$, $g \in G$,
- (6) $U(g) \cdot U(h) = U(gh)$, $g, h \in G$.

Let H_0 be a subspace of $L_2(X(t), t \in T)$ consisting of all random variables u such that $U(g)u = u$ for all $g \in G$.

Let $\{u_n, n = 1, 2, 3, \dots\}$ be a convergent sequence of random variables in H_0 such that $\lim_{n \rightarrow \infty} u_n = u$. Then, since $\|u - U(g)u\| \leq \|u - u_n\| + \|U(g)u_n - U(g)u\| = 2 \cdot \|u - u_n\|$, for all $g \in G$, $U(g)u = u$, for all $g \in G$. This implies that the subspace H_0 is closed.

Let us write $H_1 = H_0 \cap L_2(X(t), t \in A)$.

For any closed subspace H of $L_2(X(t), t \in T)$ and any random variable u in $L_2(X(t), t \in T)$, we shall denote by $\text{Proj.}(u|H)$ (or simply $u|H$) the projection of u onto H .

Let $Z \in L_2(X(t), t \in T)$ be such that $Z \in H_0$ but $Z \notin L_2(X(t), t \in A)$, that is, $U(g)Z = Z$, for all $g \in G$ but $Z \notin L_2(X(t), t \in A)$.

Let $\hat{Z} = \text{Proj.}(Z|L_2(X(t), t \in A))$. Then $\hat{Z}$ is the best linear estimator of Z on the basis of the realization of $\{X(t), t \in A\}$ in such a sense that $E\{|Z - \hat{Z}|^2\} \leq E\{|Z - u|^2\}$ for any $u \in L_2(X(t), t \in A)$.

We shall prepare the following:

Lemma 3.1. For each $g \in G$, $U(g)\hat{Z} = \hat{Z}$.

Proof. Z is decomposed uniquely in such a way that $Z = \hat{Z} + W$ where W is a

component of Z orthogonal to $L_2(X(t), t \in A)$. Since for all $g \in G$, $Z = \hat{Z} + W = U(g)\hat{Z} + U(g)W$, we have $\hat{Z} - U(g)\hat{Z} = U(g)W - W$ and hence $\|\hat{Z} - U(g)\hat{Z}\|^2 = (\hat{Z} - U(g)\hat{Z}, U(g)W - W) = (\hat{Z}, U(g)W) = (U(g)^{-1} \cdot \hat{Z}, W) = 0$. Thus, we have proved lemma 3.1.

Let $C(G)$ be a set of all bounded continuous functions on G .

Then we have the following lemma:

Lemma 3.2. *For each pair of random variables y, u in $L_2(X(t), t \in T)$*

$$(U(g)y, u) \in C(G).$$

Proof. Let $u_n = \sum_{k=1}^{N_n} c_{k,n} \cdot X(t_{k,n})$, $n = 1, 2, 3, \dots$, be such that $\lim_{n \rightarrow \infty} u_n = u$. Then $(U(g)y, u_n)$, $n = 1, 2, 3, \dots$, converges uniformly to $(U(g)y, u)$ since $|(U(g)y, u) - (U(g)y, u_n)| = |(U(g)y, u - u_n)| \leq \|y\| \cdot \|u - u_n\|$.

Hence it is sufficient for us to prove that for all $t \in T$, $(U(g)y, X(t)) \in C(G)$.

For each $t \in T$, we have

$$\begin{aligned} & |(U(g)y, X(t)) - (U(h)y, X(t))| \\ &= |(y, U(g^{-1})X(t) - U(h^{-1})X(t))| \\ &= |(y, X(g^{-1} \cdot t) - X(h^{-1} \cdot t))| \\ &\leq \|y\| \cdot \|X(g^{-1} \cdot t) - X(h^{-1} \cdot t)\| \\ &= 2 \cdot \|y\| \cdot (K(t, t) - K(t, gh^{-1} \cdot t)). \end{aligned}$$

Since $K(t, s)$ is continuous and $gh^{-1} \cdot t \rightarrow t$ as $g \rightarrow h$, we see that $(U(g)y, X(t)) \in C(G)$ for all $t \in T$. The boundedness of $(U(g)y, u)$ is obvious since G is compact.

Let $V \subset A$ be a Borel set such that $G \cdot V = A$ and $G \cdot v_1 \cap G \cdot v_2 = \emptyset$ if $v_1 \neq v_2$ ($v_1, v_2 \in V$). Since A is a compact metric space invariant under every $g \in G$, we can choose such a set V in A (lemma 2.1 in §2).

Now we have the following:

Theorem 3.1. *For each $\varepsilon > 0$, there is a random variable $Y \in L_2(X(t), t \in V)$ such that*

$$E\{|\hat{Z} - A \cdot Y|^2\} < \varepsilon,$$

where $A = \int_G U(g)dg$ whose restriction to $L_2(X(t), t \in A)$ is a projection operator onto H_1 .

Proof. From lemma 3.2, it follows that for any fixed $y \in L_2(X(t), t \in T)$, $J(u) = \int_G (U(g)y, u)dg$, $u \in L_2(X(t), t \in T)$, is a well-defined bounded linear functional on $L_2(X(t), t \in T)$ where by $\int \cdot dg$ we mean the integration with respect to the normalized Haar measure on G .

From a theorem of Riesz it follows that there exists a unique random variable $y^* = \int_G U(g)dg \cdot y$ in $L_2(X(t), t \in T)$ such that $J(u) = (y^*, u)$.

For any $\varepsilon > 0$, there exist $t_1 = g_1 v_1, t_2 = g_2 v_2, \dots, t_n = g_n v_n$ ($g_i \in G, v_i \in V$) and scalars $a_1, a_2, \dots, a_n$ such that

$$E\left\{\left|\hat{Z} - \sum_{i=1}^n a_i \cdot X(t_i)\right|^2\right\} < \varepsilon.$$

Let us put

$$Y = \sum_{i=1}^n a_i \cdot X(v_i).$$

Then, $Y \in L_2(X(t), t \in V)$.

We shall now evaluate $E\{|\hat{Z} - A \cdot Y|^2\}$.

We have

$$\begin{aligned} E\{|\hat{Z} - A \cdot Y|^2\} &= E\left\{\left|\hat{Z} - \int_G U(g)dg \cdot Y\right|^2\right\} \\ &= E\left\{\left|\hat{Z} - \sum_{i=1}^n a_i \int_G U(g) \cdot X(v_i)dg\right|^2\right\} \\ &= E\left\{\left|\hat{Z} - \sum_{i=1}^n a_i \int_G U(gg_i) \cdot X(v_i)dg\right|^2\right\} \end{aligned}$$

(here we have used the invariant measure dg).

Since the random variable $\hat{Z}$ is invariant with respect to all operators $U(g)$, $g \in G$ and $\int_G dG = 1$, we have

$$\begin{aligned} E\{|\hat{Z} - A \cdot Y|^2\} &= E\left\{\int_G U(g)\left(\hat{Z} - \sum_{i=1}^n a_i U(g_i)X(v_i)\right)dg\right\}^2 \\ &\leq E\left\{\int_G |U(g)\left(\hat{Z} - \sum_{i=1}^n a_i X(t_i)\right)|^2 dg\right\} \\ &= \int_G E\left\{|U(g)\left(\hat{Z} - \sum_{i=1}^n a_i X(t_i)\right)|^2\right\} dg \\ &= \int_G E\left\{\left|\hat{Z} - \sum_{i=1}^n a_i \cdot X(t_i)\right|^2\right\} dg \\ &= E\left\{\left|\hat{Z} - \sum_{i=1}^n a_i X(t_i)\right|^2\right\} < \varepsilon. \end{aligned}$$

Now, we shall show that for any $u \in L_2(X(t), t \in A)$,

$$\int_G U(g)dg \cdot u = \text{Proj}(u|H_1).$$

From our assumptions, the operator $U(\cdot)$ is a continuous unitary representation of G on $L_2(X(t), t \in T)$.

Hence, the representation $(U(g), L_2(X(t), t \in T))$ is completely reducible.

It is well-known that equivalent classes of irreducible representations for a compact group is only countably many and each irreducible representation is finite dimensional.

Thus, $L_2(X(t), t \in T)$ is decomposed into the direct sum of closed subspaces V_j , $j=1, 2, 3, \dots$ such that

$$(7) \quad L_2(X(t), t \in T) = V_1 + V_2 + \dots + V_n + \dots$$

and also for each j ($j=1, 2, 3, \dots$), V_j is a direct sum of d_j -dimensional (d_j is finite) subspaces V_{jk} , $k=1, 2, 3, \dots$ such that

$$(8) \quad V_j = V_{j1} + V_{j2} + \dots + V_{jk} + \dots$$

where (i) the decomposition (7) is unique, (ii) for each j ($j=1, 2, 3, \dots$), the dimension of V_{jk} , $k=1, 2, 2, \dots$, are identical and d_j , (iii) for each j, k ($j, k=1, 2, 3, \dots$), V_{jk} is

an invariant subspace under the operators $U(g)$, $g \in G$, and for each k ($k=1, 2, 3, \dots$) the restriction of $U(\cdot)$ onto V_{jk} (in symbols $M^{(j)}(\cdot) = U(\cdot)|V_{jk}$) is an irreducible unitary representation of G with dimension d_j and identical for each k ($k=1, 2, 3, \dots$), (iv) $M^{(j_1)}(g)$ and $M^{(j_2)}(g)$ are inequivalent if $j_1 \neq j_2$.

Let us choose an arbitrary orthonormal basis $(e_{jk}^{(1)}, e_{jk}^{(2)}, \dots, e_{jk}^{(d_j)})$ of the subspace V_{jk} and fix it. Then the operator $M^{(j)}(g)$ is written by $d_j \times d_j$ matrix

$$M^{(j)}(g) = \{m_{lk}^{(j)}(g)\}, \quad l, k = 1, 2, 3, \dots, d_j.$$

It is well-known that for each l, k, j ($l, k = 1, 2, 3, \dots, d_j, j = 1, 2, 3, \dots$), $m_{lk}^{(j)}(g)$ is a bounded continuous function on G and hence square-integrable with respect to the invariant measure on G .

The following orthogonality relations are well-known:

$$\int_G m_{lk}^{(j)}(g) \overline{m_{l'k'}^{(j)}(g)} dg = \frac{1}{d_j} \cdot \delta_{l \cdot l'} \cdot \delta_{k \cdot k'} \cdot \delta_{j \cdot j'},$$

where $\delta_{i,j} = 1$ if $i=j$; $=0$ otherwise.

These relations are independent of the choice of coordinate systems.

Let $M^{(1)}(g) = U(g)|V_{1k}$, $k=1, 2, 3, \dots$, be the irreducible identity representation of G , that is, $U(g)|V_{1k} = I$. Then, from Schur's lemma the irreducible identity representation of G is one-dimensional.

Hence, $d_1 = 1$, $M^{(1)}(g) = 1$ and

$$V_1 = H_0 = V_{11} + V_{12} + \dots + V_{1k} + \dots,$$

where $H_0 = \{u | U(g)u = u, \text{ for all } g \in G\} \subset L_2(X(t), t \in T)$.

Thus, from the orthogonality relations it follows that for each j ($j=2, 3, 4, \dots$), for all $\nu, \mu = 1, 2, 3, \dots, d_j$,

$$\int_G m_{\nu\mu}^{(j)}(g) dg = 0.$$

Let $y, u \in L_2(X(t), t \in T)$ be any elements but fixed. Let $y_j = \text{Proj}(y|V_j)$, $u_j = \text{Proj}(u|V_j)$ and $y_{jk} = \text{Proj}(y|V_{jk})$, $u_{jk} = \text{Proj}(u|V_{jk})$ and

$$y_{jk} = y_{jk}^{(1)} e_{jk}^{(1)} + y_{jk}^{(2)} e_{jk}^{(2)} + \dots + y_{jk}^{(d_j)} e_{jk}^{(d_j)},$$

$$u_{jk} = u_{jk}^{(1)} e_{jk}^{(1)} + u_{jk}^{(2)} e_{jk}^{(2)} + \dots + u_{jk}^{(d_j)} e_{jk}^{(d_j)}.$$

Then we have

$$\begin{aligned} \int_G (U(g)y, u) dg &= \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \int_G (M^{(j)}(g)y_{jk}, u_{jk}) dg \\ &= \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \sum_{\nu=1}^{\infty} \sum_{\mu=1}^{\infty} y_{jk}^{(\nu)} \cdot \overline{u_{jk}^{(\mu)}} \cdot \int_G m_{\nu\mu}^{(j)}(g) dg \\ &= \sum_{k=1}^{\infty} y_{jk}^{(1)} \cdot \overline{u_{1k}^{(1)}} = (y_1, u_1) \\ &= (y_1, u) \\ &= (\text{Proj}(y|V_1), u) \\ &= \left(\int_G U(g) dg \cdot y, u \right). \end{aligned}$$

Thus, the operator $\int_G U(g)dg$ is a projection onto the closed subspace H_0 . It is obvious that

$$\begin{aligned} \int_G U(g)dg \cdot L_2(X(t), t \in A) &= H_0 \cap L_2(X(t), t \in A) \\ &= H_1. \end{aligned}$$

This completes the proof.

Example 3.1. Let R_2 be the 2-dimensional Euclidean space. Let each point in R_2 be written by the polar coordinate (r, θ) , $0 \leq r < \infty$, $0 \leq \theta \leq 2\pi$.

Let $G = SO(2)$ and A be a compact set such that

$$A = \{(r, \theta) \in R_2 \mid 0 < \rho_1 \leq r \leq \rho_2 < \infty, 0 \leq \theta \leq 2\pi\}.$$

Then $G \cdot A = A$. Thus, as a Borel cross section of A , we may choose a set $V = \{(r, 0) \mid \rho_1 \leq r \leq \rho_2\}$, since V obviously satisfies the conditions:

- (i) $A = G \cdot V$,
- (ii) $Gv_1 \cap Gv_2 = \emptyset$ if $v_1 \neq v_2$ ($v_1, v_2 \in V$).

Let $\{X(r, \theta), (r, \theta) \in R_2\}$ be a complex-valued homogeneous random field such that $E\{X(r, \theta)\} = 0$, for all $(r, \theta) \in R_2$, $E\{|X(r, \theta)|^2\} < \infty$, for all $(r, \theta) \in R_2$, the covariance function $K((r, \theta), (r', \theta')) = E\{X(r, \theta) \cdot \overline{X(r', \theta')}\} = K((r, 0), (r', \theta' - \theta))$, for all $(r, \theta), (r', \theta') \in R_2$ and $K((r, \theta), (r', \theta'))$ is a continuous positive definite function on $R_2 \times R_2$.

For each $g \in SO(2)$ such that

$$g = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}, \quad 0 \leq \theta < 2\pi,$$

we shall define a unitary operator $U(\theta)$, $0 \leq \theta < 2\pi$, by writing for each $(r, \theta^*) \in R_2$,

$$U(\theta) \cdot X(r, \theta^*) = X(r, \theta^{**}), \quad 0 \leq \theta^{**} < 2\pi,$$

where $\theta + \theta^* \equiv \theta^{**}, \text{ mod } 2\pi$.

Let $Z = X(0, 0)$. Then, $U(\theta)Z = Z$, for all θ ($0 \leq \theta \leq 2\pi$).

Thus, from theorem 3.1, it follows that for each $\varepsilon > 0$ there exists a random variable Y such that

$$E\left\{\left|\hat{Z} - \frac{1}{2\pi} \int_0^{2\pi} U(\theta)d\theta \cdot Y\right|^2\right\} < \varepsilon,$$

where (a) Y is a finite linear combination of the field on V such that for some integer N , points $(r_1, 0), (r_2, 0), \dots, (r_N, 0)$ in V and complex numbers $c_1, c_2, \dots, c_N$

$$Y = \sum_{k=1}^N c_k X(r_k, 0),$$

(b) $\hat{Z}$ is the best linear estimator of $Z = X(0, 0)$ on the basis of the realizations of $\{X(t), t \in A\}$.

§ 4. On the optimal linear interpolation of homogeneous random fields II.

4.1. Summary.

In this section we shall continue considering a problem of interpolation for a homogeneous random field $\{X(t), t \in T\}$ on a metric space T acted upon by a compact

metric group G .

The field $\{X(t), t \in T\}$ is observable on a compact set A invariant under every $g \in G$.

Let $Z \in L_2(X(t), t \in T)$ be a random variable invariant with respect to all unitary operators $U(g)$, $g \in G$, defined on $L_2(X(t), t \in T)$ by writing $U(g)X(t) = X(gt)$, $t \in T$. Let $\hat{Z} \in L_2(X(t), t \in A)$ be the best linear estimator of Z based on realizations of $X(t)$ on A .

We shall proceed further in details to construct a sequence of random variables $\{Y_n, n = 1, 2, 3, \dots\}$ such that

$$(i) \quad Y_n \in L_2(X(t), t \in V), \quad n = 1, 2, 3, 4, \dots,$$

$$(ii) \quad \hat{Z} = \lim_{n \rightarrow \infty} \int_G U(g) dg \cdot Y_n,$$

where Y_n , $n = 1, 2, 3, \dots$, are concretely calculated on the basis of the realizations of the field on a Borel cross section V of A by making use of the theory of the reproducing kernel Hilbert space generated by the covariance function of the field and the previous result in theorem 2.1 of § 2 on decomposability of measures on a compact metric space.

Definitions and notations appeared in the previous sections are used in this section as well with the same meanings.

4.2. An approximating sequence for the optimal linear interpolation.

Let T and G be the spaces identical with those defined in § 3.

For any metric space X , we shall denote by $\mathfrak{B}_X$ the σ -field of Borel subsets of X and $C(X)$ a set of all bounded continuous functions on X .

Let $\{X(t), t \in T\}$ be a complex-valued homogeneous random field measurable with respect to $\mathfrak{B}_T$ satisfying the conditions:

$$(1) \quad E\{X(t)\} = 0, \quad \text{for all } t \in T.$$

$$(2) \quad E\{|X(t)|^2\} < \infty, \quad \text{for all } t \in T.$$

$$(3) \quad \text{The covariance function } K(t, s) = E\{X(t)\overline{X(s)}\} \text{ is a continuous positive definite function on } T \times T.$$

$$(4) \quad \text{For each } g \in G, K(t, s) = K(gt, gs), \quad t, s \in T.$$

The field $\{X(t), t \in T\}$ is observable on a compact set $A \subset T$ such that for every $g \in G$, $g \cdot A = A$.

For a set $S \subset T$, $L_2(X(t), t \in S)$ denotes the Hilbert space same as defined in § 3.

By $U(g)$, $g \in G$ we denote unitary operators defined on $L_2(X(t), t \in T)$ by writing $U(g)X(t) = X(gt)$, $t \in T$.

Let $V \subset A$ be a Borel cross section of A , that is, V is a Borel subset of A such that

$$(i) \quad G \cdot V = A,$$

$$(ii) \quad G \cdot v_1 \cap G \cdot v_2 = \emptyset \text{ if } v_1 \neq v_2 \text{ } (v_1, v_2 \in V).$$

Suppose that for any $g \in G$, $g \neq e$, there is no fixed point in A . Then, the mapping

$$\xi: (g, v) \rightarrow gv$$

is a Borel isomorphism between $G \times V$ and A (see lemma 2.2 in § 2).

Let μ be a measure on A invariant under every $g \in G$ such that the support of μ is A . Let us denote by μ^* the induced measure on $G \times V$ from μ on A by the Borel isomorphism ξ , that is, μ^* is a measure on $G \times V$ such that for any $A \times B \in \mathfrak{B}_{G \times V}$,

$$\mu^*(A \times B) = \mu(\xi(A \times B)).$$

Without loss of generality, we assume that $\mu^*(G \times V) = 1$, in other words, μ^* is a probability measure on $G \times V$. From theorem 2.1 in § 2, it follows that μ^* may be decomposed into a direct product measure of the normalized Haar measure of G and a probability measure m on V such that for any $A \times B \in \mathfrak{B}_{G \times V}$,

$$\mu^*(A \times B) = \nu(A) \cdot m(B).$$

Thus, we may and do identify the space A with the product space $G \times V$ and the measure μ on A with the measure μ^* on $G \times V$.

Let $Z \in L_2(X(t), t \in T)$ be such that

- (i) For all $g \in G$, $U(g)Z = Z$,
- (ii) $Z \in L_2(X(t), t \in A)$.

Let $\hat{Z}$ be the best linear estimator of Z on the basis of the realizations of the field $X(t)$ on A . Actually, $\hat{Z}$ is the projection of Z onto $L_2(X(t), t \in A)$, that is,

$$\hat{Z} = \text{Proj}(Z | L_2(X(t), t \in A)).$$

It has been shown in lemma 3.1 in § 3 that for each $g \in G$, $U(g)\hat{Z} = \hat{Z}$.

Since the covariance function $K(t, s)$ restricted to $A \times A$ is also continuous and positive definite, it follows from a theorem of Aronszajn (see theorem 6.4 in § 6) that $K(t, s)$, $t, s \in A$, generates a unique reproducing kernel Hilbert space $H(K)$.

$H(K)$ is actually a Hilbert space consisting of functions on A satisfying the conditions:

- (K.1) For each $t \in A$, $K(t, \cdot) \in H(K)$.
- (K.2) For any $f \in H(K)$, $(f, K(t, \cdot))_K = f(t)$, $t \in A$,

where by $(f_1, f_2)_K$ we denote the scalar product of every pair of elements f_1, f_2 in $H(K)$.

We shall make use of the following theorem:

Theorem 4.1. ([12], page 10, THEOREM 2D).

There is a one-one, scalar product preserving linear mapping ϕ from $H(K)$ onto $L_2(X(t), t \in A)$ such that

$$(5) \quad \phi(K(t, \cdot)) = X(t), \quad t \in A.$$

Since the mapping ϕ is an isometric isomorphism between $H(K)$ and $L_2(X(t), t \in A)$, one can define for each $g \in G$ a unitary operator $U^*(g)$ on $H(K)$ by writing

$$(6) \quad U(g) \cdot \phi(f) = \phi(U^*(g)f), \quad \text{for each } f \in H(K).$$

The operators $\{U^*(g), g \in G\}$ obviously satisfy the following:

$$U^*(g) \cdot U^*(h) = U^*(gh), \quad \text{for all } g, h \in G.$$

We shall prepare the following:

Lemma 4.1. For any $u \in L_2(X(t), t \in A)$,

$$(\phi^{-1}(u))(t) = E\{u \cdot \overline{X(t)}\}, \quad t \in A.$$

Proof. Since ϕ is an isometric isomorphism between $H(K)$ and $L_2(X(t), t \in A)$, there always exists a unique function $h \in H(K)$ such that $\phi(h) = u$. Since ϕ is scalar product preserving,

$$\begin{aligned} E\{u \cdot \overline{X(t)}\} &= (u, X(t)) = (\phi(h), \phi(K(t, \cdot))) \\ &= (h, K(t, \cdot))_K = h(t), \quad t \in A. \end{aligned}$$

This shows that $(\phi^{-1}(u))(t) = E\{u \cdot \overline{X(t)}\}$, $t \in A$.

Let us denote by Σ a set of all functions in $H(K)$ invariant under every $U^*(g)$, $g \in G$, that is,

$$\Sigma = \{f \in H(K) \mid U^*(g)f = f, \text{ for all } g \in G\}.$$

Since $H_1 = \{u \in L_2(X(t), t \in A) \mid U(g)u = u, \text{ for all } g \in G\}$ is a closed subspace in $L_2(X(t), t \in A)$ and Σ is an inverse image of H_1 under ϕ , Σ is a closed subspace of $H(K)$.

Now, let us define an operator K^* on $C(V)$ in such a manner that for any $f \in C(V)$,

$$(K^*f)(t) = \int_V K(v, t)f(v)dm(v), \quad t \in A.$$

Then, we have the following:

Lemma 4.2. (i) $K^* \cdot C(V) \subset H(K) \subset C(A)$.

(ii) For any $f \in C(V)$,

$$\phi(K^*f) = \int_V X(v)f(v)dm(v).$$

Thus, for any $f \in C(V)$, $\phi(K^*f) \in L_2(X(t), t \in V)$.

Proof. Let us consider a random variable

$$u = \int_V X(v)f(v)dm(v), \quad \text{for } f \in C(V).$$

Since $E\{|u|^2\} < \infty$, the random variable u is well-defined and $u \in L_2(X(t), t \in V)$. From lemma 4.1, it follows that

$$\begin{aligned} (K^*f)(t) &= \int_V K(v, t)f(v)dm(v) \\ &= \int_V E\{X(v)\overline{X(t)}\}f(v)dm(v) \\ &= E\{u \cdot \overline{X(t)}\} \\ &= (\phi^{-1}(u))(t), \quad t \in A. \end{aligned}$$

Thus $\phi(K^* \cdot f) = \int_V X(v)f(v)dm(v)$.

Let $h \in H(K)$. Then, for any $t, s \in A$,

$$\begin{aligned} |h(t) - h(s)| &= |(h, K(t, \cdot) - K(s, \cdot))_K| \\ &\leq \|h\|_K \cdot \sqrt{K(t, t) + K(s, s) - 2K(t, s)}. \end{aligned}$$

Thus, h is continuous. Since A is compact, h is bounded. Thus, $h \in C(A)$. This completes the proof of lemma 4.2.

Since for each pair of random variables y, u in $L_2(X(t), t \in A)$, $(U(g)y, u) \in C(G)$, (lemma 3.2 in § 3), the operator

$$A = \int_G U(g) dg$$

is well-defined on $L_2(X(t), t \in A)$. A is a projection operator on $L_2(X(t), t \in A)$ onto the closed subspace H_1 (theorem 3.1 in § 3).

Let us define an operator A^* on $H(K)$ in such a manner that for each $h \in H(K)$,

$$A \cdot \phi(h) = \phi(A^* \cdot h).$$

Then, $A^* = \int_G U^*(g) dg$, since for any $u \in L_2(X(t), t \in A)$,

$$\begin{aligned} (A \cdot \phi(h), u) &= \int_G (U(g) \cdot \phi(h), u) dg \\ &= \int_G (\phi(U^*(g)h), u) dg \\ &= \int_G (U^*(g)h, \phi^{-1}(u))_K dg \\ &= \left(\int_G U^*(g) dg \cdot h, \phi^{-1}(u) \right)_K \\ &= \left(\phi \left(\int_G U^*(g) dg \cdot h \right), u \right) \\ &= (\phi(A^* \cdot h), u). \end{aligned}$$

This implies that $A^* = \int_G U^*(g) dg$.

Lemma 4.3. For any $f \in H(K)$ and any $g \in G$,

$$(U^*(g)f)(t) = f(g^{-1} \cdot t), \quad t \in A.$$

In particular, if $f \in \Sigma$, then for all $g \in G$ and $t \in A$, $f(gt) = f(t)$.

Proof. From (5) and (6), it follows that $U(g)X(t) = X(gt) = \phi(K(gt, \cdot)) = \phi(U^*(g)K(t, \cdot))$. Hence, we have for any $g \in G$, and any $t, s \in A$,

$$\begin{aligned} (U^*(g)K(t, \cdot))(s) &= K(gt, s) \\ &= K(t, g^{-1} \cdot s). \end{aligned}$$

Thus, it follows that for any $f \in H(K)$,

$$\begin{aligned} (U^*(g)f)(t) &= (U^*(g)f, K(t, \cdot))_K \\ &= (f, U^*(g^{-1})K(t, \cdot))_K \\ &= (f, K(g^{-1} \cdot t, \cdot))_K \\ &= f(g^{-1} \cdot t), \quad t \in A. \end{aligned}$$

If $f \in \Sigma$, then $(U^*(g)f)(t) = f(t)$, $t \in A$. Thus, $f(g^{-1} \cdot t) = f(t)$, for all $g \in G$. This is equivalent to the statement that $f(gt) = f(t)$ for all $g \in G$.

We shall define an operator K on a Hilbert space $L_2(A, \mathfrak{B}_A, \mu)$ consisting of all

square integrable functions on A with respect to the measure μ in such a way that for any $f \in L_2(A, \mathfrak{B}_A, \mu)$,

$$(Kf)(t) = \int_A K(s, t)f(s)d\mu(s), \quad t \in A.$$

We have the following:

Lemma 4.4. *For any $h \in \Sigma$, $A^*K^*h = Kh$ and $K \cdot \Sigma \subset H(K)$.*

Proof. Suppose $h \in \Sigma$, and let us consider the following random variables:

$$u_1 = \int_V X(v)h(v)dm(v),$$

$$u_2 = \int_A X(t)h(t)d\mu(t).$$

Then, it is obvious that u_1 and u_2 are well-defined and $u_1 \in L_2(X(t), t \in V)$ and $u_2 \in L_2(X(t), t \in A)$. From lemma 4.2, it follows that $u_1 = \phi(K^* \cdot h)$.

Now, we have

$$\begin{aligned} (\phi^{-1}(u_2), K(t, \cdot))_K &= E\{u_2 \overline{X(t)}\} \\ &= \int_A E\{X(s) \overline{X(t)}\} h(s)d\mu(s) \\ &= \int_A K(s, t)h(s)d\mu(s) \\ &= (K \cdot h)(t), \quad t \in A. \end{aligned}$$

This implies that $K \cdot \Sigma \subset H(K)$ and

$$\phi(K \cdot h) = \int_A X(t)h(t)d\mu(t).$$

Thus, for any $g \in G$, any $f \in H(K)$ and any $h \in \Sigma$,

$$\begin{aligned} (f, U^*(g)K^* \cdot h)_K &= (\phi(f), \phi(U^*(g)K^* \cdot h)) \\ &= (\phi(f), U(g) \cdot \phi(K^* \cdot h)) \\ &= (\phi(f), U(g) \cdot \int_V X(s)h(s)dm(s)) \\ &= (\phi(f), \int_V X(gs)h(s)dm(s)) \\ &= \int_V E\{\phi(f) \cdot \overline{X(gs)}\} \overline{h(s)}dm(s) \\ &= \int_V f(gs)\overline{h(s)}dm(s). \end{aligned}$$

Hence, we have

$$\begin{aligned} (f, A^* \cdot K^* \cdot h)_K &= \int_G (f, U^*(g)K^* \cdot h)_K dg \\ &= \int_G dg \cdot \int_V f(gs)\overline{h(s)}dm(s) \\ &= \int_G \int_V f(gs)\overline{h(s)}dm(s)dg. \end{aligned}$$

Since for any $h \in \Sigma$, $h(t) = h(gt)$, $t \in A$,

$$\begin{aligned}
 (f, A^* \cdot K^* \cdot h)_K &= \int_G \int_V f(gs) \overline{h(gs)} dm(s) dg \\
 &= \int_A f(t) \overline{h(t)} d\mu(t) \\
 &= \int_A E\{\phi(f) \cdot \overline{X(t)}\} \overline{h(t)} d\mu(t) \\
 &= E\left\{\phi(f) \cdot \overline{\int_A X(t) h(t) d\mu(t)}\right\} \\
 &= (\phi(f), \phi(Kh)) \\
 &= (f, Kh)_K.
 \end{aligned}$$

Thus, for any $h \in \Sigma$, $A^* \cdot K^* \cdot h = K \cdot h$. The proof of lemma 4.4 is now complete.

Lemma 4.5. Let $\rho_z(t) = E\{Z \cdot \overline{X(t)}\}$, $t \in A$. Then, $\rho_z \in \Sigma$.

Proof. Since for any $t \in A$, $\rho_z(t) = E\{Z \cdot \overline{X(t)}\} = E\{\hat{Z} \cdot \overline{X(t)}\}$, $\rho_z \in H(K)$. Since $\hat{Z}$ is invariant with respect to all $U(g)$, $g \in G$, we have for any $g \in G$, $\hat{Z} = \phi(\rho_z) = U(g)\hat{Z} = U(g) \cdot \phi(\rho_z) = \phi(U^*(g) \cdot \rho_z)$. Thus, $\rho_z = U^*(g) \cdot \rho_z$, that is, $\rho_z \in \Sigma$.

Lemma 4.6.

(i) For any $H \in C(V)$,

$$A^*K^*H - \rho_z \in \Sigma.$$

(ii) For any constant α ,

$$(I^* - \alpha A^*K^*)\Sigma \subset \Sigma,$$

where I^* is the identity operator on $H(K)$.

Proof. First, we shall prove that for any $g \in G$, $U^*(g)A^* = A^*$. For any $f, h \in H(K)$ and $g \in G$,

$$\begin{aligned}
 (f, U^*(g)A^*h)_K &= (U^*(g^{-1})f, A^*h)_K \\
 &= \int_G (U^*(g^{-1})f, U^*(g^*)h)_K dg^* \\
 &= \int_G (f, U^*(g)U^*(g^*)h)_K dg^* \\
 &= \int_G (f, U^*(gg^*)h)_K dg^* \\
 &= \int_G (f, U^*(g')h)_K dg' \\
 &= (f, A^*h)_K.
 \end{aligned}$$

Since $K^*C(V) \subset H(K)$, A^*K^*H is well-defined and an element in $H(K)$. Thus, for any $g \in G$,

$$\begin{aligned}
 U^*(g)(A^*K^*H - \rho_z) &= U^*(g)A^*K^*H - U^*(g)\rho_z \\
 &= A^*K^*H - \rho_z.
 \end{aligned}$$

This implies that $A^*K^*H - \rho_z \in \Sigma$.

Now, we shall prove the second half of lemma 4.6.

For any $h \in \Sigma$ and any $g \in G$,

$$\begin{aligned} U^*(g)(I^* - \alpha \cdot A^*K^*)h &= U^*(g)h - \alpha \cdot U^*(g)A^*K^*h \\ &= h - \alpha \cdot A^*K^*h \\ &= (I^* - \alpha \cdot A^*K^*) \cdot h. \end{aligned}$$

Thus, the proof of lemma 4.6 is now complete.

Here, we have the following:

Theorem 4.2. *Let $\{H_n, n=1, 2, 3, \dots\}$ be a sequence of functions in $C(V)$ defined recursively in such a way that*

$$(7) \quad H_n(t) = H_{n-1}(t) - \alpha \cdot (A^*K^*H_{n-1}(t) - \rho_z(t)), \quad t \in V,$$

where α is a constant and $H_0(t)$ is an arbitrary function in $C(V)$.

Let us define a sequence of random variables $\{Y_n, n=1, 2, 3, \dots\}$ in such a way that

$$Y_n = \int_V X(s)H_n(s)dm(s), \quad n=1, 2, 3, \dots$$

Then, (i) $Y_n \in L_2(X(t), t \in V)$, for all n ($n=1, 2, 3, \dots$).

(ii) For a constant α such that $0 < \alpha < 2/D$, where $D = \int_A K(t, t)d\mu(t)$,

$$\hat{Z} = \text{l.i.m}_{n \rightarrow \infty} \int_G U(g)dg \cdot Y_n.$$

In order to prove theorem 4.2, we shall first need to derive several lemmas.

Since $\int_A \int_A |K(t, s)|^2 d\mu(t)d\mu(s) < \infty$, that is, the operator K is compact on $L_2(A, \mathfrak{B}_A, \mu)$, the eigen-values of the operator K are only countably many.

Let λ_ν , $\nu=1, 2, 3, \dots$, be non-trivial eigen-values and $\varphi_\nu(t)$, $t \in A$, $\nu=1, 2, 3, \dots$, be the corresponding normalized eigen-functions of the covariance function $K(t, s)$ such that

$$\lambda_\nu \cdot \varphi_\nu(t) = \int_A K(s, t) \cdot \varphi_\nu(s) d\mu(s), \quad t \in A.$$

Since $K(t, s)$, $t, s \in A$, is positive definite, the eigen-values λ_ν , $\nu=1, 2, 3, \dots$, are positive real numbers.

It is well-known that the covariance function $K(t, s)$ may be expanded as follows:

$$(8) \quad K(t, s) = \sum_{\nu=1}^{\infty} \lambda_\nu \cdot \overline{\varphi_\nu(t)} \cdot \varphi_\nu(s), \quad t, s \in A.$$

(Cf. [14], page 278, Mercer's theorem).

Lemma 4.7. *The family of all eigen-functions $\{\varphi_\nu, \nu=1, 2, 3, \dots\}$ spans the Hilbert space $L_2(A, \mathfrak{B}_A, \mu)$.*

Proof. Let $f \in L_2(A, \mathfrak{B}_A, \mu)$ be such that for all $\nu=1, 2, 3, \dots$,

$$(f, \varphi_\nu)_A = \int_A f(t) \cdot \overline{\varphi_\nu(t)} d\mu(t) = 0,$$

where we denote by $(f, h)_A$ the scalar product of each pair of functions f, h in $L_2(A, \mathfrak{B}_A, \mu)$.

Then, from (8), it follows that

$$\begin{aligned} & \int_A \int_A K(t, s) f(t) \overline{f(s)} d\mu(t) d\mu(s) \\ &= \sum_{\nu=1}^{\infty} \lambda_{\nu} \cdot |(f, \varphi_{\nu})_A|^2 = 0. \end{aligned}$$

Since $K(t, s)$ is positive definite, f must be a zero element in $L_2(A, \mathfrak{B}_A, \mu)$. This implies that $L_2(A, \mathfrak{B}_A, \mu)$ is spanned by $\{\varphi_{\nu}, \nu = 1, 2, 3, \dots\}$.

Lemma 4.8.

- (i) For each ν ($\nu = 1, 2, 3, \dots$), $\varphi_{\nu} \in H(K)$.
- (ii) $(\varphi_{\nu}, \varphi_{\mu})_K = \lambda_{\nu}^{-1}$, if $\nu = \mu$,
 $= 0$, if $\nu \neq \mu$.
- (iii) $\phi(\varphi_{\nu}) = \lambda_{\nu}^{-1} \cdot \int_A X(s) \cdot \varphi_{\nu}(s) d\mu(s)$.

Proof. Let us write

$$X_k = \int_A X(s) \cdot \varphi_k(s) d\mu(s), \quad k = 1, 2, 3, \dots$$

It is easily seen that $E\{|X_k|^2\} = \lambda_k < \infty$ and $X_k \in L_2(X(t), t \in A)$. Since $(\phi^{-1}(X_k))(t) = E\{X_k \cdot \overline{X(t)}\} = \lambda_k \cdot \varphi_k(t)$, $t \in A$,

$$X_k = \phi(\lambda_k \cdot \varphi_k) = \lambda_k \cdot \phi(\varphi_k).$$

Thus, $\varphi_k \in H(K)$ and $\phi(\varphi_k) = \lambda_k^{-1} \cdot X_k$.

Since $E\{X_k \cdot \overline{X_n}\} = \lambda_k$ if $k = n$; $= 0$ if $k \neq n$,

$$\begin{aligned} (\varphi_k, \varphi_n)_K &= (\phi(\varphi_k), \phi(\varphi_n)) \\ &= \lambda_k^{-1} \cdot \lambda_n^{-1} \cdot E\{X_k \cdot \overline{X_n}\} \\ &= \lambda_k^{-1}, \quad \text{if } k = n, \\ &= 0, \quad \text{if } k \neq n. \end{aligned}$$

Thus, we have proved lemma 4.8.

Lemma 4.9. For any $h \in H(K)$,

$$\|h\|_K^2 = \sum_{\nu=1}^{\infty} \lambda_{\nu}^{-1} \cdot |(h, \varphi_{\nu})_A|^2 < \infty.$$

Proof. Since $H(K) \subset L_2(A, \mathfrak{B}_A, \mu)$, h may be expanded as follows:

$$h(t) = \sum_{\nu=1}^{\infty} (h, \varphi_{\nu})_A \cdot \varphi_{\nu}(t), \quad t \in A.$$

Thus, we have

$$\begin{aligned} \|h\|_K^2 &= \sum_{\nu=1}^{\infty} \sum_{\mu=1}^{\infty} (h, \varphi_{\nu})_A \cdot \overline{(h, \varphi_{\mu})_A} \cdot (\varphi_{\nu}, \varphi_{\mu})_K \\ &= \sum_{\nu=1}^{\infty} \lambda_{\nu}^{-1} \cdot |(h, \varphi_{\nu})_A|^2. \end{aligned}$$

This completes the proof of lemma 4.9.

Lemma 4.10. Let $B(\alpha)$ be such that

$$B(\alpha) = \sup_{\substack{\|f\|_K=1 \\ f \in \Sigma}} \|(I^* - \alpha \cdot A^* \cdot K^*)f\|_K$$

where α is a constant.

Then, $B(\alpha) < 1$ if $0 < \alpha < 2/D$, where $D = \int_A K(t, t) d\mu(t)$.

Proof. For any $h \in \Sigma$, we have the following expansion:

$$h(t) = \sum_{\nu=1}^{\infty} (h, \varphi_{\nu})_A \cdot \varphi_{\nu}(t), \quad t \in A.$$

Thus, from lemma 4.4, it follows that

$$A^* \cdot K^* \cdot h = K \cdot h = \sum_{\nu=1}^{\infty} \lambda_{\nu} \cdot (h, \varphi_{\nu})_A \cdot \varphi_{\nu}.$$

Hence, we have

$$(I^* - \alpha \cdot A^* \cdot K^*)h = \sum_{\nu=1}^{\infty} (1 - \alpha \cdot \lambda_{\nu}) (h, \varphi_{\nu})_A \cdot \varphi_{\nu}.$$

Thus, we have the following inequality:

$$\begin{aligned} \|(I^* - \alpha \cdot A^* \cdot K^*)h\|_K^2 &= \sum_{\nu=1}^{\infty} \lambda_{\nu}^{-1} \cdot |1 - \alpha \cdot \lambda_{\nu}|^2 \cdot |(h, \varphi_{\nu})_A|^2 \\ &< \sum_{\nu=1}^{\infty} \lambda_{\nu}^{-1} \cdot |(h, \varphi_{\nu})_A|^2 = \|h\|_K^2, \end{aligned}$$

if for all $\nu = 1, 2, 3, \dots$, $|1 - \alpha \cdot \lambda_{\nu}| < 1$.

This condition can be satisfied by choosing the constant α such that $2/\lambda^* > \alpha > 0$, where $\lambda^* = \text{Max}(\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_n, \dots)$.

Since $D = \int_A K(t, t) d\mu(t) = \sum_{\nu=1}^{\infty} \lambda_{\nu} > \lambda^*$, if α is such that $0 < \alpha < 2/D$,

$$\|(I^* - \alpha \cdot A^* \cdot K^*)h\|_K^2 < \|h\|_K^2, \quad \text{for any } h \in \Sigma.$$

Thus, we have proved lemma 4.10.

We are now ready to derive the proof of theorem 4.2.

Let us apply the operator $A^* \cdot K^*$ on both sides of (7). Then, we have

$$A^* \cdot K^* \cdot H_n = A^* \cdot K^* \cdot H_{n-1} - \alpha \cdot A^* \cdot K^* (A^* \cdot K^* \cdot H_{n-1} - \rho_z).$$

Thus, for each n ($n = 1, 2, 3, \dots$),

$$\begin{aligned} A^* \cdot K^* \cdot H_n - \rho_z &= (I^* - \alpha \cdot A^* \cdot K^*) (A^* \cdot K^* \cdot H_{n-1} - \rho_z) \\ &= (I^* - \alpha \cdot A^* \cdot K^*)^n \cdot (A^* \cdot K^* \cdot H_0 - \rho_z). \end{aligned}$$

Since $A^* \cdot K^* \cdot H_0 - \rho_z \in \Sigma$, we readily find the following inequality:

$$\begin{aligned} \|A^* \cdot K^* \cdot H_n - \rho_z\|_K &= \|(I^* - \alpha \cdot A^* \cdot K^*)^n \cdot (A^* \cdot K^* \cdot H_0 - \rho_z)\|_K \\ &\leq B(\alpha)^n \cdot \|A^* \cdot K^* \cdot H_0 - \rho_z\|_K. \end{aligned}$$

Hence, by choosing the constant α such that $B(\alpha) < 1$, we have

$$\|A^* \cdot K^* \cdot H_n - \rho_z\|_K \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

From lemma 4.2, it follows that for each n ($n = 1, 2, 3, \dots$),

$$Y_n = \int_V X(s) H_n(s) dm(s),$$

is well-defined and $Y_n \in L_2(X(t), t \in V)$.

Since $Y_n = \phi(K^* \cdot H_n)$, $A \cdot Y_n = \phi(A^* \cdot K^* \cdot H_n)$, where $A = \int_G U(g) dg$ and $A^* = \int_G U^*(g) dg$.

Thus, it follows that

$$\begin{aligned} \|A^* \cdot K^* \cdot H_n - \rho_z\|_K &= \|\phi(A^* \cdot K^* \cdot H_n) - \phi(\rho_z)\| \\ &= \|A \cdot Y_n - \hat{Z}\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

This completes the proof of theorem 4.2.

Example 4.1. Let T be the unit sphere with radius c in R_3 , that is, $T = \{(x, y, z) \in R_3 \mid x^2 + y^2 + z^2 = c^2\}$.

Let (θ, φ) , $0 \leq \theta \leq \pi$, $0 \leq \varphi < 2\pi$, be such that

$$\begin{aligned} x &= c \cdot \sin \theta \cdot \cos \varphi, \\ y &= c \cdot \sin \theta \cdot \sin \varphi, \\ z &= c \cdot \cos \theta, \end{aligned}$$

where $(x, y, z) \in T$.

Let G be a subgroup of $SO(3)$ such that each element of G is of the form :

$$g = \begin{pmatrix} \cos \eta, & \sin \eta, & 0 \\ -\sin \eta, & \cos \eta, & 0 \\ 0, & 0, & 1 \end{pmatrix}, \quad 0 \leq \eta \leq 2\pi,$$

that is, G is a group consisting of all rotations around the z -axis.

Let $\{X(\theta, \varphi), (\theta, \varphi) \in T\}$ be a homogeneous random field on T such that

(i) $E\{X(\theta, \varphi)\} = 0$, for all $(\theta, \varphi) \in T$,

(ii) $E\{|X(\theta, \varphi)|^2\} < \infty$, for all $(\theta, \varphi) \in T$,

(iii) $K((\theta, \varphi), (\theta', \varphi')) = E\{X(\theta, \varphi) \cdot \overline{X(\theta', \varphi')}\}$

$$= K((\theta, 0), (\theta', \varphi' - \varphi)), \quad \text{for all } (\theta, \varphi), (\theta', \varphi') \in T.$$

Let us define unitary operators $U(\varphi)$, $0 \leq \varphi < 2\pi$, on $L_2(X(t), t \in T)$ by writing

$$U(\varphi), X(\theta, \varphi') = X(\theta, \varphi''),$$

where $0 \leq \varphi'' < 2\pi$, $\varphi + \varphi' \equiv \varphi'' \pmod{2\pi}$.

Let A be such that

$$A = \{(\theta, \varphi) \mid 0 < \rho_1 \leq \theta \leq \rho_2 \leq \pi, 0 \leq \varphi < 2\pi\}.$$

Then, we may choose as a Borel cross section of A a set V such that

$$V = \{(\theta, 0) \mid \rho_1 \leq \theta \leq \rho_2\}.$$

Suppose that the field $\{X(\theta, \varphi), (\theta, \varphi) \in T\}$ is observable on A .

Let μ be a probability measure distributing uniformly on T . Then, it is written as follows:

$$d\mu(\theta, \varphi) = \frac{1}{2} \sin \theta \, d\theta \cdot \frac{1}{2\pi} d\varphi.$$

Let $Z = X(0, 0)$. Then, Z is invariant with respect to every $U(\varphi)$, $0 \leq \varphi < 2\pi$.

Now, let a sequence of functions $\{H_n, n=1, 2, 3, \dots\}$ be defined recursively as follows:

$$H_n(t) = H_{n-1}(t) - \alpha \cdot \left(A^* \cdot \int_0^\pi K((\theta, 0), t) H_{n-1}(\theta, 0) \, dm(\theta) - \rho_z(t) \right)$$

where $\rho_z(t) = E\{X(0, 0) \cdot \overline{X(t)}\}$, $t \in A$,

$$dm(\theta) = \frac{1}{2} \sin \theta \, d\theta,$$

$$H_0(t) \equiv 1, \quad t \in A.$$

Then, from theorem 4.2, it follows that

$$\hat{Z} = \text{l.i.m}_{n \rightarrow \infty} \int_0^{2\pi} U(\varphi) d\varphi \cdot Y_n,$$

where $Y_n = \int_0^\pi X(\theta, 0) H_n(\theta, 0) dm(\theta)$.

§5. On the uniformly minimum variance unbiased linear estimates of mean value functions of homogeneous random fields.

5.1. Summary.

In this section we shall consider an estimation problem of mean value functions $m(t)$ of homogeneous random fields $\{X(t), t \in T\}$ on a compact metric space T acted upon by a compact metric group G .

We assume that mean value functions $m(t)$ are G -invariant, that is, $m(t) = m(gt)$, for all $g \in G$ and $t \in T$.

Let V be a Borel cross section of T , in other words, V be a subset of T such that $G \cdot V = T$ and $G \cdot v_1 \cap G \cdot v_2 = \emptyset$ if $v_1 \neq v_2$ ($v_1, v_2 \in V$).

Then, it is shown that the uniformly minimum variance unbiased linear estimate $\widehat{m}(t)$ of the mean value function $m(t)$ admits the expression

$$\widehat{m}(t) = \int_G U(g) dg \cdot X(v), \quad v \in V,$$

where $t = g' \cdot v$ for some $g' \in G$ and $v \in V$, and $\{U(g), g \in G\}$ are unitary operators on $L_2(X(t), t \in T)$ defined by writing $U(g)X(t) = X(gt)$.

5.2. An estimation problem of mean value functions.

Let T be a compact metric space acted upon by a compact metric group G such that (i) $G \cdot T = T$, (ii) each $g \in G$ is a homeomorphism acting on T and (iii) the mapping $(g, t) \rightarrow gt$ from $G \times T$ onto T is continuous.

From our assumptions, it follows that there exists a Borel subset V of T such that $G \cdot V = T$ and $G \cdot v_1 \cap G \cdot v_2 = \emptyset$ if $v_1 \neq v_2$ ($v_1, v_2 \in V$), (see lemma 2.1 in §2).

Let $\{X(t), t \in T\}$ be a homogeneous random field on T with the following properties:

- (1) $E\{|X(t)|^2\} < \infty$, for all $t \in T$.

(2) The covariance function $K(t, s) = \text{Cov}(X(t), X(s))$ is continuous positive definite function on $T \times T$.

(3) For each $g \in G$, $K(t, s) = K(gt, gs)$, $t, s \in T$.

Let us denote the mean value function of the field by

$$m(t) = E\{X(t)\}, \quad t \in T.$$

Let $L_2(X(t), t \in T)$ be the Hilbert space consisting of all random variables which may be represented either as a finite linear combination

$$u = \sum_{i=1}^n c_i \cdot X(t_i)$$

for some integer n , points $t_1, t_2, \dots, t_n$ in T and scalars $c_1, c_2, \dots, c_n$ or as a limit in quadratic mean of such finite linear combinations under the scalar product (u, z) defined by

$$(u, z) = E_m\{u \cdot \bar{z}\} = \text{Cov}(u, z) + E_m\{u\} \cdot E_m\{\bar{z}\}.$$

The subscript m on an expectation operator E is written to indicate that the expectation is computed under the assumption that $m(\cdot)$ is the true mean value function.

Since the covariance function $K(t, s)$, $t, s \in T$, is continuous positive definite, it follows from a theorem of Aronszajn (see theorem 6.4 in § 6) that $K(t, s)$ generates a unique reproducing kernel Hilbert space $H(K)$.

$H(K)$ is actually a Hilbert space consisting of functions on T satisfying the conditions:

(K.1) For each $t \in T$, $K(t, \cdot) \in H(K)$.

(K.2) For any $f \in H(K)$,

$$(f, K(t, \cdot))_K = f(t), \quad t \in T,$$

where by $(f, h)_K$ we denote the scalar product of each pair of functions f, h in $H(K)$.

Let M be an arbitrarily given subset of $H(K)$ and let us assume that $m \in M$.

We shall make use of the following theorem:

Theorem 5.1. ([11], page 29, Theorem 4A). *There is a linear one-one mapping ϕ from $H(K)$ onto $L_2(X(t), t \in T)$ with the following properties:*

(L.1) For each $t \in T$, $\phi(K(t, \cdot)) = X(t)$.

(L.2) For any $f \in H(K)$,

$$E_m\{\phi(f)\} = (f, m)_K, \quad \text{for all } m \in M.$$

(L.3) For any $f, h \in H(K)$,

$$\text{Cov}(\phi(f), \phi(h)) = (f, h)_K.$$

We shall make use of the definition of an unbiased linear estimate of the mean value function $m(t)$, $t \in T$, introduced by E. Parzen [11].

A random variable $\phi(h)$, $h \in H(K)$, is said to be an unbiased linear estimate of the value $m(t)$ at a particular point $t \in T$ of the mean value function $m(\cdot)$ if

$$E_m\{\phi(h)\} = (h, m)_K = m(t), \quad \text{for all } m \in M.$$

For any Hilbert space H and its closed subspace H^* and any element $u \in H$, we shall denote by

$$\text{Proj}(u|H^*)$$

the projection of u onto H^* .

We shall use the following:

Theorem 5.2. ([11], page 29, theorem 4A). *The uniformly minimum variance unbiased linear estimate $\widehat{m}(t)$ of $m(t) \in M$ is given by*

$$\widehat{m}(t) = \phi(\text{Proj}(K(t, \cdot)|\bar{M}))$$

where $\bar{M}$ is the smallest closed subspace of $H(K)$ containing M .

Now, let us write

$$\Sigma = \{f \in H(K) \mid f(t) = f(gt), \text{ for all } g \in G\}.$$

Hereafter, we shall assume that the mean value function $m(t)$ is invariant under every $g \in G$, in other words,

$$m(\cdot) \in \Sigma.$$

For each $g \in G$, let us define a linear operator $U(g)$ on $L_2(X(t), t \in T)$ by writing $U(g)X(t) = X(gt)$, $t \in T$. Then, since $m \in \Sigma$ and

$$\begin{aligned} (U(g)X(t), U(g)X(s)) &= \text{Cov}(X(gt), X(gs)) + m(gt)\overline{m(gs)} \\ &= K(t, s) + m(t)\overline{m(s)} \\ &= (X(t), X(s)), \end{aligned}$$

the operators $U(g)$, $g \in G$, are unitary.

Since the following inequality holds:

$$\begin{aligned} \|f - h\|_K^2 &\leq \|\phi(f) - \phi(h)\|^2 \\ &\leq (1 + \|m\|_K^2) \cdot \|f - h\|_K^2, \quad \text{for any } f, h \in H(K), \end{aligned}$$

the mapping ϕ is a linear homeomorphism between $H(K)$ and $L_2(X(t), t \in T)$.

Lemma 5.1. *Let B be a closed linear subspace of $H(K)$ and the mean value function $m \in B$. Then, for any $f \in H(K)$,*

$$\phi(\text{Proj}(f|B)) = \text{Proj}(\phi(f)|\phi(B)).$$

Proof. Since ϕ is a homeomorphism between $H(K)$ and $L_2(X(t), t \in T)$, $\phi(B)$ is closed.

For any $f \in H(K)$, there is a unique decomposition such that

$$f = f_1 + f_2$$

where $f_1 = \text{Proj}(f|B)$ and $f_2 = \text{Proj}(f|B^\perp)$.

Let $u_1 = \text{Proj}(\phi(f)|\phi(B))$ and $u_2 = \text{Proj}(\phi(f)|\phi(B)^\perp)$. Then, $\phi(f) = u_1 + u_2 = \phi(f_1) + \phi(f_2)$.

Since $u_1 \in \phi(B)$, there is a unique function $k \in B$ such that $\phi(k) = u_1$. Thus, we have

$$\begin{aligned} \|\phi(f_1) - u_1\|^2 &= (u_1, \phi(f_2)) - (\phi(f_1), \phi(f_2)) \\ &= (m, k)_K(m, f_2)_K - (m, f_1)_K(m, f_2)_K \\ &= 0, \end{aligned}$$

since $m \in B$ and $f_2 \in B^\perp$.

Lemma 5.2. *Let $V_1 = \{u \mid U(g)u = u, \text{ for all } g \in G\} \subset L_2(X(t), t \in T)$. Then, V_1 is a closed subspace of $L_2(X(t), t \in T)$ and $\phi(\Sigma) = V_1$. Thus, Σ is a closed subspace of $H(K)$.*

Proof. For any $u_1, u_2 \in V_1$ and any scalars α, β ,

$$\begin{aligned} U(g)(\alpha u_1 + \beta u_2) &= \alpha \cdot U(g)u_1 + \beta U(g)u_2 \\ &= \alpha u_1 + \beta u_2. \end{aligned}$$

Let $\{u_n, n = 1, 2, 3, \dots\}$ be a sequence in V_1 such that for $u^* \in L_2(X(t), t \in T)$,

$$\|u_n - u^*\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Then, for any $g \in G$,

$$\begin{aligned} \|u^* - U(g)u^*\| &\leq \|u^* - u_n\| + \|U(g)u_n - U(g)u^*\| \\ &= 2 \cdot \|u_n - u^*\| \rightarrow 0, \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Thus, $u^* \in V_1$ and hence V_1 is closed.

For any $h \in \Sigma$ and any $g \in G$,

$$\begin{aligned} (U(g)\phi(h), X(t)) &= (\phi(h), U(g^{-1})X(t)) \\ &= (\phi(h), X(g^{-1} \cdot t)) \\ &= (h, K(g^{-1} \cdot t, \cdot))_K + (h, m)_K (m, K(g^{-1}t, \cdot))_K \\ &= h(g^{-1}t) + (h, m)_K m(g^{-1}t) \\ &= h(t) + (h, m)_K m(t) \\ &= (\phi(h), X(t)), \quad \text{for all } t \in T. \end{aligned}$$

This implies that $U(g)\phi(h) = \phi(h)$, for all $g \in G$. Thus, $\phi(\Sigma) \subset V_1$.

Conversely, for any $z \in V_1$, there exists a unique function $f \in H(K)$ such that $\phi(f) = z$.

Since $U(g)\phi(f) = \phi(f)$, for all $g \in G$, we have

$$\begin{aligned} (\phi(f), X(t)) &= f(t) + (f, m)_K m(t) \\ &= (U(g)\phi(f), X(t)) \\ &= (\phi(f), X(g^{-1}t)) \\ &= f(g^{-1}t) + (f, m)_K \cdot m(t), \quad \text{for all } t \in T. \end{aligned}$$

Thus, it follows that $f(t) = f(g^{-1}t)$, for all $g \in G$ and $t \in T$.

This implies that $f \in \Sigma$. Thus, we have proved that

$$V_1 = \phi(\Sigma).$$

Since V_1 is closed and ϕ is a homeomorphism, Σ is closed.

We have now the following:

Theorem 5.3. *Let $\hat{m}(t)$ be the uniformly minimum variance unbiased linear estimate of the value $m(t)$ at the point $t \in T$ of the mean value function $m \in \Sigma$.*

Then, $\hat{m}(t)$ is written as follows:

$$\widehat{m}(t) = \int_G U(g)dg \cdot X(v),$$

where $t = g^*v$ for some $g^* \in G$ and $v \in V$.

If T is a homogeneous space, that is, G is transitive on T , then the Borel cross section V consisting of a single point $v_0 \in T$ and

$$\widehat{m}(t) = \int_G U(g)dg \cdot X(v_0).$$

Proof. First, we shall show that the operator $\int_G U(g)dg$ is well-defined.

Since for any fixed $y \in L_2(X(t), t \in T)$ and for all $t \in T$,

$$\begin{aligned} & |(U(g)y, X(t)) - (U(h)y, X(t))| \\ & \leq 2 \cdot \|y\| \cdot (K(t, t) - K(t, gh^{-1}t)), \quad g, h \in G, \end{aligned}$$

$(U(g)y, z) \in C(G)$, for any fixed $z \in L_2(X(t), t \in T)$, where $C(G)$ is a set of all bounded continuous functions on G . Hence, it is clear that the operator $\int_G U(g)dg$ is well-defined on $L_2(X(t), t \in T)$.

By the same arguments in the proof of theorem 3.1 in §3, we see that the operator $\int_G U(g)dg$ is the projection operator onto the closed subspace V_1 of $L_2(X(t), t \in T)$.

From theorem 5.2 and lemma 5.1, it follows that the uniformly minimum variance unbiased linear estimate $\widehat{m}(t)$ of $m(t)$, $t = g^*v \in T$ ($g^* \in G$, $v \in V$), is given by

$$\begin{aligned} \widehat{m}(t) &= \phi(\text{Proj}(K(t, \cdot) | \Sigma)) \\ &= \text{Proj}(\phi(K(t, \cdot)) | \phi(\Sigma)) \\ &= \text{Proj}(X(t) | V_1) \\ &= \int_G U(g)dg \cdot X(t). \end{aligned}$$

Since $X(t) = X(g^*v) = U(g^*)X(v)$, it is written as follows:

$$\begin{aligned} \widehat{m}(t) &= \int_G U(g)dg \cdot U(g^*)X(v) \\ &= \int_G U(g)U(g^*)dg \cdot X(v) \\ &= \int_G U(gg^*)dg \cdot X(v) \\ &= \int_G U(g)dg \cdot X(v), \end{aligned}$$

(here we have used the invariant measure dg).

If G is transitive on T , then for any $t \in T$ and any $v_0 \in T$, there exists $g^* \in G$ such that $t = g^*v_0$.

Thus, we have

$$\begin{aligned}
\widehat{m}(t) &= \int_G U(g) dg \cdot U(g^*) X(v_0) \\
&= \int_G U(gg^*) dg \cdot X(v_0) \\
&= \int_G U(g) dg \cdot X(v_0).
\end{aligned}$$

Now, the proof of theorem 5.3 is complete.

§ 6. Appendix.

Theorem 6.1. ([10], page 23, theorem 4.2)

Let X and Y be compact metric spaces and g a continuous map of X onto Y . Then there is a Borel set $B \subset X$ such that $g(B) = Y$ and g is one-one on B .

Theorem 6.2. ([10], page 21, theorem 3.9)

Let X_1, X_2 be complete separable metric spaces and $E_1 \subset X_1, E_2 \subset X_2$ two sets, E_1 being a Borel set. Let φ be a measurable one-one map of E_1 into X_2 such that $\varphi(E_1) = E_2$. Then E_2 is a Borel set.

Theorem 6.3. ([17], page 131)

A one-to-one continuous mapping of a compact space onto a Hausdorff space is a homeomorphism.

Theorem 6.4. ([12], page 7, theorem 2B)

A symmetric non-negative kernel K generates a unique Hilbert space, which we denote by $H(K)$, of which K is the reproducing kernel.

§ 7. Acknowledgment.

The author would like to express his heartiest thanks to Professor K. R. Parthasarathy of University of Manchester and Professor J. Gani of Sheffield University for their kind suggestions and encouragement while this paper was being prepared.

References

- [1] ARONSZAJN, N.: *Theory of reproducing kernels*, Trans. Amer. Math. Soc., **68** (1950), 337-404.
- [2] FEDERER, H. AND A. P. MORSE: *Some properties of measurable functions*, Bull. Amer. Math. Soc., **49** (1943), 270-277.
- [3] HALMOS, P.: *"Introduction to Hilbert Space,"* Chelsea, N. Y., 1951.
- [4] HALMOS, P. R.: *"Measure Theory,"* Van Nostrand, Princeton, New Jersey, 1962.
- [5] HANNAN, E. J.: *"Time Series Analysis,"* Methuen, London, 1960.
- [6] HANNAN, E. J.: *Group representations and applied probability*, J. Appl. Prob., **2** (1965), 1-68.
- [7] KELLEY, J. L.: *"General Topology,"* Van Nostrand, N. Y., 1961.
- [8] LOEVE, M.: *"Probability Theory,"* Van Nostrand, N. Y., 1955.
- [9] NAGAI, T.: *On decomposability of probability measures on a separable metric space acted upon by a compact metric group*, (to be published).
- [10] PARTHASARATHY, K. R.: *"Probability measures on Metric Spaces*, Academic Press, N. Y.,

- 1967.
- [11] PARZEN, E.: *A new approach to the synthesis of optimal smoothing and prediction systems*, Tech. Report, No. 34, Stanford Univ. 1960.
 - [12] PARZEN, E.: *Regression analysis of continuous parameter time series, II*, Tech. Report, No. 35, Stanford Univ., 1960.
 - [13] PARZEN, E.: *An approach to time series analysis*, Ann. Math. Statist., 32 (1961), 951-989.
 - [14] PARZEN, E.: *"Time series analysis papers,"* Holden-Day, Calif., 1967.
 - [15] PONTRJAGIN, L.S.: *"Topological Groups,"* Princeton Univ. Press, Princeton, New Jersey, 1946 (Translated from Russian).
 - [16] ROSENBLATT, M.: *"Random Processes,"* Oxford Univ. Press, N. Y., 1962.
 - [17] SIMMONS, G.F.: *"Introduction to Topology and Modern Analysis,"* McGraw-Hill, N. Y., 1963.
 - [18] YAGLOM, A.M.: *Second order homogeneous random fields*, Fourth Berkeley Symposium on Math. Statist. and Probability, 2 (1961), 593-622.