

ON EXPLICIT BOUNDS IN THE ERROR FOR THE \$ H_o^1 \$-PROJECTION INTO PIECEWISE POLYNOMIAL SPACES

Kimura, Seiji

Sony Oita, DA Section, Fukuoka System LSI Development Center

Yamamoto, Nobito

The University of Electro-Communications

<https://doi.org/10.5109/13484>

出版情報 : Bulletin of informatics and cybernetics. 31 (2), pp.109-115, 1999-12. Research
Association of Statistical Sciences

バージョン :

権利関係 :



ON EXPLICIT BOUNDS IN THE ERROR FOR THE H_0^1 -PROJECTION INTO PIECEWISE POLYNOMIAL SPACES

By

Seiji KIMURA * and Nobito YAMAMOTO †

Abstract

The values of constants appearing in error estimates of approximations by finite element methods play an important role in numerical verification methods for elliptic equations (Nakao and Yamamoto(1998), Yamamoto and Nakao(1995), etc.). For efficient implementation of the verification algorithms on computers, it is necessary that these constants can be estimated as close as possible to their optimal values. In Nakao, Yamamoto and Kimura(1998), the optimal constant was derived for quadratic elements as well as a nearly optimal value for cubic elements. In this paper, we establish a method to calculate the values of constants for approximation by piecewise polynomials of arbitrary degree and to give bounds on the difference between the constants so calculated and optimal values.

1. Introduction

We consider an interval $I \subset \mathbf{R}$ and Sobolev spaces on it, denoted by $L^2(I)$, $H^1(I)$, $H_0^1(I)$, and so on, as usual. The inner product in $L^2(I)$ is represented by (\cdot, \cdot) , and the norm in the same space by $\|\cdot\|$.

We divide the interval I into subintervals and consider a piecewise polynomial space $S_h(I) \subset H_0^1(I)$ of arbitrary degree. Let u be a function in $H_0^1(I) \cap H^2(I)$ and $u_h \in S_h$ its approximation which satisfies

$$(u' - u_h', \psi') = 0 \quad \text{for } \forall \psi \in S_h(I).$$

Since u_h is the orthogonal projection of u into S_h with respect to the inner-product in H_0^1 ,

$$\|u' - u_h'\| \leq \|u' - \psi'\| \quad \text{for } \forall \psi \in S_h(I),$$

holds, and we refer to u_h as the H_0^1 -projection of u into $S_h(I)$.

For u and u_h , the following error estimate holds :

$$\|u' - u_h'\| \leq C_0 h \|u''\|, \tag{1.1}$$

* Sony Oita, DA Section, Fukuoka System LSI Development Center, Fukuoka, Japan

† The University of Electro-Communications, Chofu-shi, Tokyo, 182-8585, Japan

where h is the maximal size of subintervals, and C_0 is a constant independent of u and h . The purpose of the present paper is to calculate the value of C_0 .

As is well known, an H_0^1 -projection into a piecewise polynomial space coincides with the interpolation of the function at each node (e.g., Schultz, 1973). Therefore, we shall reduce our problem to the error estimation for local approximations of functions on an interval $e \equiv [0, h]$, vanishing at the terminals.

Let $Q_N(e)$ with $N \geq 0$ be a polynomial space defined by

$$Q_N(e) = \left\{ p(x) \mid p \text{ is a polynomial of degree } N+1, p(0) = p(h) = 0 \right\},$$

and, for an arbitrary function $u \in H_0^1(e) \cap H^2(e)$, let $P_N u$ be the local H_0^1 -projection of u into $Q_N(e)$ defined by

$$(u' - (P_N u)', v')_e = 0 \quad \text{for } \forall v \in Q_N(e),$$

where $(\cdot, \cdot)_e$ denotes the inner-product of $L^2(e)$. We look for the minimal constant C_0 which satisfies an error estimate of the form:

$$\|u' - (P_N u)'\|_e \leq C_0 h \|u''\|_e, \quad (1.2)$$

where $\|\cdot\|_e$ is the norm of $L^2(e)$. Note that the constant in (1.2) gives an upper bound in (1.1) when $S_h(I)$ is a piecewise polynomial space of degree $N+1$, and that it can also be extended into multi-dimensional problems (cf. Nakao, Yamamoto and Kimura, 1998).

The following lemma provides a basis for error estimates of the approximation by piecewise linear functions.

LEMMA 1.1. *Wirtinger's inequality (Theorem 1.2 in Schultz, 1973)*

If $u \in H_0^1(e)$, then

$$\|u\|_e \leq \frac{h}{\pi} \|u'\|_e.$$

Moreover, the equality holds if $u(x) = \sin(\pi x/h)$.

From Lemma 1.1, it can be seen that the optimal value for the constant C_0 in (1.2) for the error estimate of linear interpolant ($N = 0$) is $1/\pi$. Therefore, it is natural to expect that the corresponding constant is smaller than $1/\pi$ when we use higher degree polynomials. Actually, this is proved in Nakao, Yamamoto and Kimura (1998) in the quadratic and cubic cases. In this paper, we are concerned with an arbitrary degree.

2. Explicit Estimate of Constants

Now we introduce a set of polynomials $\{g_i\}_{i=1}^\infty$ of the form

$$g_i(x) = \frac{i+1/2}{i! h^{i+1/2}} \frac{d^{i-1}}{dx^{i-1}} \left[x^i (h-x)^i \right].$$

The followings are well known properties of these polynomials:

$$(g_i, g_j)_e = \begin{cases} \frac{h^2}{2(2i-1)(2i+3)} & (i=j) \\ -\frac{h^2}{4(i+j+1)(2i+1)^{1/2}(2j+1)^{1/2}} & (i=j \pm 2) \\ 0 & (\text{otherwise}) \end{cases} \quad (2.1)$$

and

$$(g'_i, g'_j)_e = \begin{cases} 1 & (i=j) \\ 0 & (i \neq j). \end{cases} \quad (2.2)$$

The fact that $\{g'_i\}_{i=1}^\infty$ gives a complete orthonormal system in $L^2(e)$ enables us to expand an arbitrary function $u \in H_0^1(e)$ as

$$u(x) \sim \sum_{j=1}^\infty u_j g_j(x),$$

where

$$u_j \equiv (u', g'_j)_e.$$

Moreover, we can represent the H_0^1 -projection of $u \in H_0^1(e)$ into $Q_N(e)$ as

$$P_N u(x) = \sum_{j=1}^N u_j g_j(x),$$

owing to the uniqueness of the H_0^1 -projection for fixed $Q_N(e)$.

The following theorem gives a rough estimation for the case of an arbitrary integer $N \geq 0$.

THEOREM 2.1. *If $u \in H_0^1(e) \cap H^2(e)$, it follows that*

$$\|u' - (P_N u)'\|_e^2 \leq \tilde{C}_N h^2 \|u''\|_e^2,$$

where

$$\tilde{C}_N \equiv \frac{1}{8} \left(\frac{1}{2N+1} + \frac{1}{2N+3} \right).$$

PROOF.

$$\begin{aligned} \|u' - (P_N u)'\|_e^2 &= \sum_{i=N+1}^\infty (u', g'_i)_e^2 \\ &= \sum_{i=N+1}^\infty (u'', g_i)_e^2 \quad (\text{integrating by parts}) \end{aligned}$$

$$\begin{aligned}
&\leq \|u''\|_e^2 \sum_{i=N+1}^{\infty} \|g_i\|_e^2 \quad (\text{by Schwarz' inequality}) \\
&= h^2 \|u''\|_e^2 \sum_{i=N+1}^{\infty} \frac{1}{2(2i-1)(2i+3)} \quad (\text{from (2.1)}) \\
&= \frac{h^2}{8} \|u''\|_e^2 \sum_{i=N+1}^{\infty} \left(\frac{1}{2i-1} - \frac{1}{2i+3} \right) \\
&= \frac{h^2}{8} \left(\frac{1}{2N+1} + \frac{1}{2N+3} \right) \|u''\|_e^2.
\end{aligned}$$

□

In the following lemma, a smaller constant is derived by considering a finite dimensional eigenvalue problem in the case that u is a polynomial.

LEMMA 2.2. *Let $M \geq 1$ be an arbitrary integer. If $q \in Q_{N+M}(e)$, then the following holds:*

$$\|q - P_N q\|_e^2 \leq \Lambda_N h^2 \|q' - (P_N q)'\|_e^2,$$

where

$$\begin{aligned}
\Lambda_N \equiv \max \left\{ \frac{1}{2(2N+1)(2N+5)} + \frac{1}{4(2N+5)\sqrt{2N+3}\sqrt{2N+7}}, \right. \\
\left. \frac{1}{4(2N+5)\sqrt{2N+3}\sqrt{2N+7}} + \frac{1}{2(2N+5)(2N+9)}, \right. \\
\left. \frac{1}{4(2N+9)\sqrt{2N+7}\sqrt{2N+11}} \right\}. \quad (2.3)
\end{aligned}$$

PROOF.

Each polynomial $q \in Q_{N+M}(e)$ can be represented as

$$q = \sum_{i=1}^{N+M} q_i g_i,$$

and we have

$$\begin{aligned}
\|q - P_N q\|_e^2 &= \sum_{i=N+1}^{N+M} \sum_{j=N+1}^{N+M} q_i (g_i, g_j)_e q_j, \\
\|q' - (P_N q)'\|_e^2 &= \sum_{i=N+1}^{N+M} q_i^2.
\end{aligned}$$

Now let us consider an M -by- M matrix:

$$A_N^M \equiv \left(a_{ij}^N \right)_{1 \leq i, j \leq M},$$

where

$$a_{ij}^N \equiv \frac{1}{h^2} (g_{N+i}, g_{N+j})_\epsilon,$$

and let q_N^M be an M -dimensional vector:

$$q_N^M \equiv \left(q_i \right)_{1 \leq i \leq M}$$

Then we have

$$\sup_{q \neq 0} \frac{\|q - P_N q\|_\epsilon^2}{h^2 \|q' - (P_N q)'\|_\epsilon^2} = \sup_{q_N^M \neq 0} \frac{q_N^{M^T} A_N^M q_N^M}{q_N^{M^T} q_N^M}.$$

The right-hand side equals the largest eigenvalue of A_N^M , denoted by λ_N^M . From Gerschgorin's Theorem, λ_N^M can be estimated by

$$\lambda_N^M \leq \max_{1 \leq i \leq M} \sum_{j=1}^M |a_{ij}^N|.$$

Using (2.1), the right-hand side of the above inequality is calculated as

$$\begin{aligned} \lambda_N^M &\leq \max \left\{ |a_{11}^N| + |a_{13}^N|, |a_{22}^N| + |a_{24}^N|, \max_{3 \leq i \leq M} \{ |a_{ii}^N| + |a_{i-2}^N| + |a_{i+2}^N| \} \right\} \\ &= \max \{ |a_{11}^N| + |a_{13}^N|, |a_{31}^N| + |a_{33}^N| + |a_{35}^N| \} \\ &= \Lambda_N. \end{aligned}$$

This completes the proof. □

The following theorem, the main result of this article, is an extension of Lemma 2.2.

THEOREM 2.3. *For any function $u \in H_0^1(e) \cap H^2(e)$, we have*

$$\|u' - (P_N u)'\|_\epsilon^2 \leq \Lambda_N h^2 \|u''\|_\epsilon^2, \quad (2.4)$$

where the constant Λ_N is defined by (2.3).

PROOF.

Using Lemma 1.1, Lemma 2.2, Theorem 2.1, and the relation $\|(P_{N+M} u)' - (P_N u)'\|_\epsilon \leq \|u' - (P_N u)'\|_\epsilon$, we have the following for an arbitrary integer $M \geq 1$:

$$\begin{aligned} \|u - P_N u\|_\epsilon^2 &\leq (\|u - P_{N+M} u\|_\epsilon + \|P_{N+M} u - P_N u\|_\epsilon)^2 \\ &\leq \left\{ \left(\frac{h}{\pi} \right) \|u' - (P_{N+M} u)'\|_\epsilon + \Lambda_N^{1/2} h \|(P_{N+M} u)' - (P_N u)'\|_\epsilon \right\}^2 \\ &\leq \left\{ \left(\frac{h}{\pi} \right) \frac{h}{2\sqrt{2}} \left(\frac{1}{2N+2M+1} + \frac{1}{2N+2M+3} \right)^{1/2} \|u''\|_\epsilon \right. \\ &\quad \left. + \Lambda_N^{1/2} h \|u' - (P_N u)'\|_\epsilon \right\}^2. \end{aligned}$$

Letting M tend to infinity, we obtain

$$\|u - P_N u\|_\epsilon^2 \leq \Lambda_N h^2 \|u' - (P_N u)'\|_\epsilon^2. \quad (2.5)$$

Moreover, from (2.5),

$$\begin{aligned} \|u' - (P_N u)'\|_\epsilon^2 &= (u' - (P_N u)', u' - (P_N u)')_\epsilon \\ &= (u' - (P_N u)', u')_\epsilon \\ &= -(u - P_N u, u'')_\epsilon \\ &\leq \|u - P_N u\|_\epsilon \|u''\|_\epsilon \\ &\leq \Lambda_N^{1/2} h \|u' - (P_N u)'\|_\epsilon \|u''\|_\epsilon \end{aligned} \quad (2.6)$$

holds. Dividing both sides of (2.6) by $\|u' - (P_N u)'\|_\epsilon$, we have (2.4). \square

In Bernardi(1996), similar arguments (but not numerical) are presented. The above theorem assures (1.5) of Bernardi(1996) in the case of $s = 2$:

$$\|u - P_N u\|_{H^1(\epsilon)} \leq \frac{c}{N+1} \|u\|_{H^2(\epsilon)}$$

with numerical estimate of the constant c . For example, the theorem tells us that we can choose $c = 1/2$.

Moreover, we can estimate an upper bound of the difference between Λ_N and the optimal constant C_0^2 . Since $\lambda_N^M \leq C_0^2 \leq \Lambda_N$ holds for an arbitrary integer $M \geq 1$, we have

$$|\Lambda_N - C_0^2| \leq |\Lambda_N - \lambda_N^M|.$$

In the following table, we estimate relative errors of Λ_N on C_0^2 by $\frac{|\Lambda_N - \lambda_N^M|}{\Lambda_N}$ taking $M = 4$ for $N = 0, \dots, 10$. The exact values of C_0^2 obtained in Nakao, Yamamoto and Kimura(1998) are shown for $N = 0, 1$. For $N \geq 2$, exact values of C_0^2 are not yet known, which are represented by '?' in the table.

From this table, it is seen that the relative errors of Λ_N are less than 15% for $N = 0, \dots, 10$.

N	λ_N^4	C_0^2	Λ_N	$\frac{ \Lambda_N - \lambda_N^4 }{\Lambda_N}$
0	0.1013197	$(\frac{1}{\pi})^2 \cong 0.10132$	0.110911	0.0864766
1	0.0253155	$(\frac{1}{2\pi})^2 \cong 0.02533$	0.0291335	0.131053
2	0.0123516	?	0.0142767	0.134841
3	0.00748463	?	0.00859464	0.129152
4	0.00507143	?	0.00577062	0.121164
5	0.00368227	?	0.00415142	0.113011
6	0.00280337	?	0.00313354	0.105368
7	0.00220957	?	0.00245078	0.0984193
8	0.00178846	?	0.00197004	0.0921739
9	0.00147842	?	0.00162342	0.0893178
10	0.00124326	?	0.00138892	0.104871

3. Conclusion

We have obtained a method to get numerical estimates of the constants appearing in (1.1) for the approximation by polynomials of arbitrary degree. Since the estimates derived here are sharp enough for actual application, we believe that they will play an important role in the error estimates for the finite element method and in the numerical verification of solutions to nonlinear elliptic equations.

From the mathematical point of view, the relation between the degree of polynomials and the constant in (1.2) is very interesting. We leave it as an open problem to decide the optimal constants.

References

- Beckenbach, E.F., and Bellman, R. (1965), *Inequalities*, Springer-Verlag, Berlin.
- Bernardi, C. (1996), Inducteurs d'erreur en $h - N$ version des éléments spectraux, *Modélisation Mathématique et Analyse Numérique*, **30**, 1-38.
- Ciarlet, P.G. (1978), *The finite element method for elliptic problems*, North-Holland Publishing Company, Amsterdam.
- Grisvard, P. (1985), *Elliptic problems in nonsmooth domain*, Pitman, London.
- Johnson, C. (1987), *Numerical Solution of partial differential equations by the finite element method*, Cambridge University Press, Cambridge.
- Nakao, M.T., and Yamamoto, N. (1998), Numerical verification of solutions for nonlinear elliptic problems using an L^∞ residual method, *Journal of Mathematical Analysis and Applications*, **217**, 246-262.
- Nakao, M.T., Yamamoto, N. and Kimura, S. (1998), On the best constant in the error bound for the H_0^1 -projection into piecewise polynomial spaces, *Journal of Approximation Theory*, **93**, 491-500.
- Schultz, M.H. (1973), *Spline Analysis*, Prentice-Hall, London.
- Yamamoto, N. and Nakao, M.T. (1995), Numerical verifications for solutions to elliptic equations using residual iterations with higher order finite element, *Journal of Computational and Applied Mathematics*, **60**, 271-279.

Received September 7, 1998

Revised March 31, 1999

Re-revised July 12, 1999