MULTI-VALUED OPTIMAL STOPPING PROBLEM WITH MONOTONICITY

Ohtsubo, Yoshio
Department of Mathematics, Faculty of Science, Kochi University

<https://doi.org/10.5109/13486>

出版情報 : Bulletin of informatics and cybernetics. 31 (2), pp.137-146, 1999-12. Research
Association of Statistical Sciences
バージョン :
権利関係 :

MULTI-VALUED OPTIMAL STOPPING PROBLEM WITH MONOTONICITY*

By

Yoshio OHTSUBO†

Abstract

We consider a multi-valued optimal stopping problem for indexed random sequences. We perform a scalarization of the random sequences and give characterization of optimal value process for scalarized problem. We also find optimal stopping time in the sense of Pareto optimality for original problem. We apply its result to monotone cases, in which it is given an explicit optimal stopping time. It is given sufficient conditions for our problem to be a monotone case. One of conditions is a kind of concavity with regard to time parameter.

1. Introduction

We often happen to meet optimization problem with a lot of objectives. In this paper we formulate such a problem as optimal stopping problem for indexed random sequences and give a solution to the problem.

Let $(\Omega, \mathcal{F}, P)$ be a probability space and $(\mathcal{F}_n)_{n \in N}$ a filtration of $\mathcal{F}$, where $N = \{0, 1, 2, \dots\}$ is a discrete time space. Let T be an index set of objects in our problem. For each t in the index set T , let $(X_n^t)_{n \in N}$ be a random sequence defined on $(\Omega, \mathcal{F}, P)$ and adapted to $(\mathcal{F}_n)$ such that random variables $\sup_{n \in N} \sup_{t \in T} (X_n^t)^+$ and $\sup_{t \in T} (X_n^t)^-$ are integrable with regard to P , where $x^+ = \max(0, x)$ and $x^- = (-x)^+$. We assume that the set $\cup_{t \in T_1} B_t$ is $\mathcal{F}_n$ -measurable for any subset T_1 of T and for any family $\{B_t : t \in T_1\}$ of $\mathcal{F}_n$ -measurable sets. This assumption is clearly satisfied if T is a countable set. For each $n \in N$, we also denote by Λ_n the class of $(\mathcal{F}_n)$ -stopping times τ such that $n \leq \tau < \infty$ almost surely.

Our aim is to maximize the expectation $E[X_\tau^t]$ with respect to τ in Λ_0 for every $t \in T$. Especially, when the index set T consists of an element it is the classical optimal stopping problem (cf. Chow *et al.*(1971)), and when $T = \{1, 2, \dots, p\}$ it is a multi-objective stopping problem (cf. Gugerli(1987), Ohtsubo(1997)).

To accomplish our aim, we shall generalize the concept of Pareto optimality (cf. Aubin(1979, p.295), Yu(1985, p.22)). A Pareto optimal solution is also called an efficient

---

* This work was partly supported by Grant-in-Aid for Scientific Research C-2-10640124 from Ministry of Education, Science, Sports and Culture

† Department of Mathematics, Faculty of Science, Kochi University, Kochi 780-8520, Japan
e-mail: ohtsubo@math.kochi-u.ac.jp

or noninferior solution in the literature (see, for example, Yu(1985)). For the sake of general discussion we define a conditional expectation of X_τ^t at $n \in N$ by

$$G_n(\tau; t) = E[X_\tau^t | \mathcal{F}_n]$$

for each $t \in T$ and $\tau \in \Lambda_n$. For $n \in N$, $\varepsilon \geq 0$ and probability measure μ on $(T, 2^T)$, we say that a stopping time τ_ε in Λ_n is (ε, μ) -Pareto optimal at n , if there exists no stopping time τ in Λ_n such that almost surely

$$G_n(\tau; t) \geq G_n(\tau_\varepsilon; t) + \varepsilon$$

for every $t \in T_0$ and

$$G_n(\tau; t) > G_n(\tau_\varepsilon; t) + \varepsilon$$

for every $t \in T_1$, where T_0 and T_1 are some subsets of T satisfying $\mu(T_0) = 1$ and $\mu(T_1) > 0$.

When T is a finite set $\{t_1, t_2, \dots, t_N\}$, and we define μ to be probability measure such that $\mu(t_k) > 0$ for every k , $(0, \mu)$ -Pareto optimal stopping time τ_ε at n satisfies the following condition: there exists no stopping time τ in Λ_n such that almost surely

$$G_n(\tau; t) \geq G_n(\tau_\varepsilon; t)$$

for every $t \in T$ and

$$G_n(\tau; t) > G_n(\tau_\varepsilon; t)$$

for some $t \in T$. This means well-known Pareto optimal solution in mathematical programming (cf. Aubin(1979), Yu(1985)).

Similarly we shall define weak Pareto optimal stopping time as follows: For $n \in N$, $\varepsilon \geq 0$ and probability measure μ on $(T, 2^T)$, it is said for a stopping time τ_ε in Λ_n to be (ε, μ) -weak Pareto optimal at n , if there exists no stopping time τ in Λ_n such that almost surely

$$G_n(\tau; t) > G_n(\tau_\varepsilon; t) + \varepsilon$$

for every $t \in T_0$, where T_0 is some subset of T satisfying $\mu(T_0) = 1$.

We easily see that if τ_ε is (ε, μ) -Pareto optimal at n , it is (ε, μ) -weak Pareto optimal at n . In this paper we shall investigate only (ε, μ) -Pareto optimal stopping time.

For the sake of simplicity, without further comments we assume that all inequalities and equalities between random variables hold in the sense of "almost surely".

In the second section, we consider a scalarization of our problem, give fundamental properties of optimal value process for the scalarized problem and find an optimal stopping time on the scalarized problem. By the results of the scalarization we find an (ε, μ) -Pareto optimal stopping time for the original problem. In the third section, we consider the classical monotone case on the scalarized problem as a special model and we prove that a one-step-look-ahead (OLA) stopping time is $(0, \mu)$ -Pareto optimal. It is given sufficient conditions for the problem to be a monotone case. The first condition is a kind of concavity with regard to time parameter on the original process and the second is a generalization of classical stopping problem. Finally in Section 4 we give

examples of monotone case, including economic model which is a withdrawal problem from a market.

Such a problem is a generalization of multi-objective stopping problem, which has been investigated in Hisano(1980), Gugerli(1987) and Ohtsubo(1997). Hisano has introduced a partial order defined by a convex cone and give an existence theorem of optimal stopping time with regard to the order. Gugerli has considered such a problem for the class of all randomized stopping times on discrete time Markov chain and has characterized the set of all Pareto optimal stopping times. Ohtsubo(1997) has given a sufficient condition for multi-objective stopping problem to be a monotone case, and find 0-Pareto optimal stopping time which is OLA rule. This paper is not only a generalization of Ohtsubo(1997) but also contains many useful examples. In Stadje(1980), Preater(1993) and Gnedin(1994), multi-criteria best-choice problems have been investigated in different forms.

2. Scalarization and Pareto Optimality

In this section, we consider a scalarization of our problem with regard to probability measure on the set T , give fundamental properties of optimal value process for the scalarized problem and find an Pareto optimal stopping times on the original problem.

Let $\mathcal{M}$ denote the set of probability measures μ on the measurable space $(T, 2^T)$. This μ is called weighting function in mathematical programming. For μ in $\mathcal{M}$, we define a scalarized reward process $(X_n(\mu))$ by

$$X_n(\mu) = \int_T X_n^t d\mu(t),$$

where X_n^t is integrable from the conditions given in Section 1, and define the conditional expectation by

$$G_n(\tau; \mu) = \int_T G_n(\tau; t) d\mu(t) \equiv E[X_\tau(\mu) | \mathcal{F}_n].$$

The last equality is obtained from the boundedness (from above) of X_τ^t on $\Omega \times T$ and from Fubini's theorem for conditional expectation. The optimal value process $(V_n(\mu))$ for the scalarized problem is defined by

$$V_n(\mu) = \operatorname{ess\,sup}_{\tau \in \Lambda_n} G_n(\tau; \mu), \quad n \in N$$

for a given $\mu \in \mathcal{M}$.

For each $n \in N, \varepsilon \geq 0$ and $\mu \in \mathcal{M}$, we define a stopping time $\tau_n^\varepsilon(\mu)$ by

$$\tau_n^\varepsilon(\mu) = \inf\{k \geq n | V_k(\mu) \leq X_k(\mu) + \varepsilon\},$$

where $\inf \emptyset = +\infty$.

The scalarized problem is one of classical stopping problems and hence it follows the theorem below, the proof of which refers to Theorems 4.1, 4.2 and 4.3 in Chow *et al.*(1971) or Proposition VI-1-2 and VI-1-3 in Neveu(1975).

PROPOSITION 2.1. *Let μ in $\mathcal{M}$ be arbitrary.*

(i) *The optimal value process $V(\mu) = (V_n(\mu))$ satisfies the following recursive relation:*

$$V_n(\mu) = \max(X_n(\mu), E[V_{n+1}(\mu)|\mathcal{F}_n]), \quad n \in N.$$

(ii) *$V(\mu)$ is the smallest supermartingale dominating the process $X(\mu) = (X_n(\mu))$.*

(iii) $\limsup_n V_n(\mu) = \limsup_n X_n(\mu)$.

(iv) *For every $n \in N$ any $\varepsilon > 0$, the stopping time $\tau_n^\varepsilon(\mu)$ is $(\varepsilon, V(\mu))$ -optimal at n in the sense that $V_n(\mu) \leq G_n(\tau_n^\varepsilon(\mu); \mu) + \varepsilon$. If the stopping time $\tau_n^0(\mu)$ is a.s. finite, $\tau_n^0(\mu)$ is $(0, V(\mu))$ -optimal at n .*

The general lemma below is an important result and an optimal stopping version of a well-known result in multi-objective problem on mathematical programming (cf. Proposition 3 in Aubin(1979, p.297)).

LEMMA 2.2. *Let $n \in N, \varepsilon \geq 0$ and $\mu \in \mathcal{M}$ be arbitrary. If a stopping time τ_ε in Λ_n satisfies inequality $V_n(\mu) \leq G_n(\tau_\varepsilon; \mu) + \varepsilon$, then τ_ε is (ε, μ) -Pareto optimal at n .*

PROOF. We suppose that the stopping time τ_ε is not (ε, μ) -Pareto optimal at n . Then there exists a stopping time τ in Λ_n such that $G_n(\tau; t) \geq G_n(\tau_\varepsilon; t) + \varepsilon$ for every $t \in T_0$ and $G_n(\tau; t) > G_n(\tau_\varepsilon; t) + \varepsilon$ for every $t \in T_1$ where T_0 and T_1 satisfy $\mu(T_0) = 1$ and $\mu(T_1) > 0$. Thus since $(T - T_1) \cap T_0 \subset T_0$, $\mu(T - (T_0 \cup T_1)) = 0$ and $\mu(T_0 \cup T_1) = 1$, we have

$$\begin{aligned} G_n(\tau; \mu) &= \int_T G_n(\tau; t) d\mu(t) \\ &= \int_{T_1} G_n(\tau; t) d\mu(t) + \int_{(T-T_1) \cap T_0} G_n(\tau; t) d\mu(t) + \int_{T-(T_0 \cup T_1)} G_n(\tau; t) d\mu(t) \\ &> \int_T G_n(\tau_\varepsilon; t) d\mu(t) + \varepsilon \mu(T_0 \cup T_1) \\ &= G_n(\tau_\varepsilon; \mu) + \varepsilon, \end{aligned}$$

and hence $V_n(\mu) > G_n(\tau_\varepsilon; \mu) + \varepsilon$, which is a contradiction. Hence the stopping time τ_ε is (ε, μ) -Pareto optimal at n .

Proposition 2.1 and Lemma 2.2 immediately imply the following theorem.

THEOREM 2.3. *Let $n \in N$ and $\mu \in \mathcal{M}$ be arbitrary.*

(i) *For each $\varepsilon > 0$, the stopping time $\tau_n^\varepsilon(\mu)$ is (ε, μ) -Pareto optimal at n .*

(ii) *If the stopping time $\tau_n^0(\mu)$ is a.s. finite, $\tau_n^0(\mu)$ is $(0, \mu)$ -Pareto optimal at n .*

3. Monotone Case

For the scalarized reward process $(X_n(\mu))$ defined in Section 2 where $\mu \in \mathcal{M}$, we define subsets $A_n(\mu)$ of Ω by

$$A_n(\mu) = \{X_n(\mu) \geq E[X_{n+1}(\mu)|\mathcal{F}_n]\}, \quad n \in N,$$

and define a stopping time $\sigma_n(\mu)$ by

$$\sigma_n(\mu) = \inf\{k \geq n | X_k(\mu) \geq E[X_{k+1}(\mu)|\mathcal{F}_k]\}, \quad n \in N,$$

that is,

$$\sigma_n(\mu)(\omega) = \inf\{k \geq n | \omega \in A_k(\mu)\}, \quad \omega \in \Omega, n \in N,$$

where $\inf \phi = +\infty$. $\sigma_n(\mu)$ is called one-step-look-ahead (OLA) rule or myopic rule.

For each μ in $\mathcal{M}$, we introduce the following condition :

CONDITION $C(\mu)$. For every $n \in N$, $A_n(\mu) \subset A_{n+1}(\mu)$ and $\lim_{n \rightarrow \infty} P(A_n(\mu)) = 1$.

When the condition $C(\mu)$ is satisfied for a given $\mu \in \mathcal{M}$, the scalarized stopping problem corresponding to μ is in a well-known monotone case (cf. Chow *et al.*(1971, p.54)).

THEOREM 3.1. *Suppose that Condition $C(\mu)$ is satisfied for a given μ in $\mathcal{M}$. Then for each $n \in N$ $\sigma_n(\mu)$ is a.s. equal to $\tau_n^0(\mu)$ and is a.s. finite, and hence $\sigma_n(\mu)$ is $(0, \mu)$ -Pareto optimal at n .*

PROOF. The first part : $\sigma_n(\mu) = \tau_n^0(\mu) < \infty$ a.s. is proved similarly to Theorem 3.3 in Chow *et al.*(1971). Hence Theorem 2.3 implies that $\sigma_n(\mu)$ is $(0, \mu)$ -Pareto optimal at n .

Next we shall give two sufficient conditions for Condition $C(\mu)$ to be satisfied. For each $n \in N$ and $t \in T$, we define a stochastic process (Y_n^t) by

$$Y_n^t = X_n^t - E[X_{n+1}^t|\mathcal{F}_n]$$

and a subset A_n^t of Ω by

$$A_n^t = \{Y_n^t \geq 0\} \equiv \{X_n^t \geq E[X_{n+1}^t|\mathcal{F}_n]\}.$$

ASSUMPTION 3.1. For $\mu \in \mathcal{M}$, there exists a subset T_0 of T satisfying $\mu(T_0) = 1$ such that for all $t \in T_0$ (Y_n^t) is a submartingale and $\lim_{n \rightarrow \infty} P(\cap_{t \in T_0} A_n^t) = 1$.

When (Y_n^t) is a submartingale, we have the relation

$$E[X_{n+1}^t|\mathcal{F}_n] \geq \frac{1}{2}X_n^t + \frac{1}{2}E[X_{n+2}^t|\mathcal{F}_n],$$

which means that (X_n^t) is concave with regard to time parameter n . If T_0 is a finite set, (A_n^t) is increasing with regard to n and $\lim_{n \rightarrow \infty} P(A_n^t) = 1$ for every $t \in T_0$, then $\lim_{n \rightarrow \infty} P(\cap_{t \in T_0} A_n^t) = 1$.

We define a new scalarized process $(Y_n(\mu))$ by

$$Y_n(\mu) = \int_T Y_n^t d\mu(t) \equiv X_n(\mu) - E[X_{n+1}(\mu) | \mathcal{F}_n].$$

Then we notice that $A_n(\mu) = \{Y_n(\mu) \geq 0\}$.

LEMMA 3.2. *If Assumption 3.1 is satisfied for some $\mu \in \mathcal{M}$, then $(Y_n(\mu))$ is a submartingale.*

PROOF. By the definition of $Y_n(\mu)$ and the fact that $\mu(T - T_0) = 0$, it follows that

$$Y_n(\mu) - E[Y_{n+1}(\mu) | \mathcal{F}_n] = \int_{T_1} (Y_n^t - E[Y_{n+1}^t | \mathcal{F}_n]) d\mu(t).$$

Since (Y_n^t) is a submartingale for any $t \in T_0$, that is, $Y_n^t \leq E[Y_{n+1}^t | \mathcal{F}_n]$, $n \in N$, we have $Y_n(\mu) \leq E[Y_{n+1}(\mu) | \mathcal{F}_n]$, and so $(Y_n(\mu))$ is a submartingale for given $\mu \in \mathcal{M}$.

THEOREM 3.3. *If Assumption 3.1 is satisfied for some $\mu \in \mathcal{M}$, then Condition C(μ) holds, and hence for each $n \in N$ the OLA rule $\sigma_n(\mu)$ is $(0, \mu)$ -Pareto optimal at n .*

PROOF. In order to show that $A_n(\mu) \subset A_{n+1}(\mu)$, it suffices to prove that $X_{n+1}(\mu) \geq E[X_{n+2}(\mu) | \mathcal{F}_{n+1}]$ on $A_n(\mu)$. Since $Y_n(\mu)$ is a submartingale from Lemma 3.2 and $Y_n(\mu) \geq 0$ on $A_n(\mu)$, it follows that for any $B \in \mathcal{F}_n$,

$$\begin{aligned} & \int_{A_n(\mu) \cap B} (X_{n+1}(\mu) - E[X_{n+2}(\mu) | \mathcal{F}_{n+1}]) dP \\ &= \int_{A_n(\mu) \cap B} Y_{n+1}(\mu) dP \\ &\geq \int_{A_n(\mu) \cap B} Y_n(\mu) dP \\ &\geq 0 \end{aligned}$$

and hence $X_{n+1}(\mu) \geq E[X_{n+2}(\mu) | \mathcal{F}_{n+1}]$ on $A_n(\mu)$. Similarly, by using the fact that (Y_n^t) is a submartingale for any $t \in T_0$, it follows that $A_n^t \subset A_{n+1}^t$ for every $n \in N$ and $t \in T_0$, so $\cap_{t \in T_0} A_n^t \subset \cap_{t \in T_0} A_{n+1}^t$ for every $n \in N$.

Next we see that for all $t \in T_0$ $X_n^t \geq E[X_{n+1}^t | \mathcal{F}_n]$ on $\cap_{t \in T_0} A_n^t$. Since $\mu(T_0) = 1$ for a given $\mu \in \mathcal{M}$, it follows from Fubini's theorem for conditional expectation that

$$\begin{aligned} X_n(\mu) &= \int_{T_0} X_n^t d\mu(t) \\ &\geq \int_{T_0} E[X_{n+1}^t | \mathcal{F}_n] d\mu(t) \\ &= E\left[\int_{T_0} X_{n+1}^t d\mu(t) \middle| \mathcal{F}_n\right] \\ &= E[X_{n+1}(\mu) | \mathcal{F}_n] \end{aligned}$$

on $\cap_{t \in T_0} A_n^t$. Hence $\cap_{t \in T_0} A_n^t \subset A_n(\mu)$ and so $\cup_n \cap_{t \in T_0} A_n^t \subset \cup_n A_n(\mu)$. Since $P(\cup_n \cap_{t \in T_0} A_n^t) = \lim_{n \rightarrow \infty} P(\cap_{t \in T_0} A_n^t) = 1$, we have $\lim_{n \rightarrow \infty} P(A_n(\mu)) = P(\cup_n A_n(\mu)) = 1$. Thus Condition $C(\mu)$ is satisfied for a given $\mu \in \mathcal{M}$. From Theorem 3.1 it follows that the OLA rule $\sigma_n(\mu)$ is $(0, \mu)$ -Pareto optimal at n .

Next we shall give another sufficient condition for Condition $C(\mu)$ to be satisfied.

ASSUMPTION 3.2. For $\mu \in \mathcal{M}$, there exists a subset T_0 of T satisfying $\mu(T_0) = 1$ such that $A_n^s = A_n^t$ and $A_n^t \subset A_{n+1}^t$ for each $n \in N$ and any $s, t \in T_0$, and $\lim_{n \rightarrow \infty} P(A_n^t) = 1$ for any $t \in T_0$.

THEOREM 3.4. If Assumption 3.2 is satisfied for some $\mu \in \mathcal{M}$, then Condition $C(\mu)$ holds, and hence the OLA rule $\sigma_n(\mu)$ is $(0, \mu)$ -Pareto optimal at n .

PROOF. We see from Assumption 3.2 and by the way similar to the proof of Theorem 3.3 that $A_n^t = \cap_{s \in T_0} A_n^s \subset A_n(\mu)$ for each $n \in N$ and all $t \in T_0$. Conversely, since we have $X_n(\mu) \geq E[X_{n+1}(\mu) | \mathcal{F}_n]$ on $A_n(\mu)$, there exists $t \in T_0$ such that $X_n^t \geq E[X_{n+1}^t | \mathcal{F}_n]$, and so we have $A_n(\mu) \subset A_n^t = \cap_{s \in T_0} A_n^s$. Thus it follows that $A_n^t = \cap_{s \in T_0} A_n^s = A_n(\mu)$ for each $n \in N$ and all $t \in T_0$. From Assumption 3.2 the monotonicity of $\{A_n^t\}$ implies that of $\{A_n(\mu)\}$ and we have $\lim_{n \rightarrow \infty} P(A_n(\mu)) = 1$, that is, Condition $C(\mu)$ is satisfied for a given $\mu \in \mathcal{M}$. Hence it follows from Theorem 3.1 that the OLA rule $\sigma_n(\mu)$ is $(0, \mu)$ -Pareto optimal at n .

4. Examples

In this section we shall give four examples as applications of the previous section.

EXAMPLE 4.1 (WITHDRAWAL PROBLEM I). Let (W_n) be a random sequence which satisfies a regular condition, and assume that there is a subset T_0 of T such that for each $t \in T_0$ and all $n \in N$ $X_n^t = E[W_{n+a_t} | \mathcal{F}_n]$, where a parameter a_t of time-shift is a nonnegative integer.

We define random sequence (Z_n) and a subset B_n of $\mathcal{F}_n$ by

$$\begin{aligned} Z_n &= W_n - E[W_{n+1} | \mathcal{F}_n], \\ B_n &= \{W_n \geq E[W_{n+1} | \mathcal{F}_n]\} = \{Z_n \geq 0\}, \end{aligned}$$

respectively. We then have the relation $Y_n^t = E[Z_{n+a_t} | \mathcal{F}_n]$. We assume that (Z_n) is a submartingale and $\lim_{n \rightarrow \infty} P(B_n) = 1$. Here we notice that (B_n) is an increasing sequence of sets. Then we easily see that (Y_n^t) is a submartingale, so $A_n^t \subset A_{n+1}^t$, $n \in N$ for each $t \in T_0$. Also since $B_n \subset B_{n+a_t}$, we have $B_n \subset A_n^t$ for each $n \in N$ and all $t \in T_0$, so we have $\lim_{n \rightarrow \infty} P(\cap_{t \in T_0} A_n^t) = 1$. Hence Assumption 3.1 is satisfied for any $\mu \in \mathcal{M}$ satisfying $\mu(T_0) = 1$. Thus Theorem 3.3 implies that for such a μ the OLA rule $\sigma_n(\mu)$ is $(0, \mu)$ -Pareto optimal at n for every $n \in N$.

In a withdrawal problem from the market, we can explain this model as follows. There is a company which owns a lot of factories producing a common article at many

countries in the world. Let T be a set of all factories. W_n represents demands for the article on USA at time n and is susceptible to the influences of the market. For a part of factories T_0 , X_n^t represents demands at the factory $t \in T_0$ and is influenced from W_n with time delay a_t . The aim of the company is to find a withdrawal time τ from the market, at which all factories simultaneously stop to produce the article.

EXAMPLE 4.2 (WITHDRAWAL PROBLEM II). For $t \in T$, let f_t be a strictly increasing real-valued function on real line. For (W_n) given in Example 4.1, there is a subset T_0 of T such that for each $t \in T_0$ and all $n \in N$ $X_n^t = f_t(W_n)$, for example, $X_n^t = a_t W_n + b_t$, where a_t and b_t are constants such that $a_t > 0$. If (W_n) is in monotone case, i.e. $B_n \subset B_{n+1}$, $n \in N$ and $\lim_{n \rightarrow \infty} P(B_n) = 1$, we can easily check that Assumption 3.2 is satisfied. Thus it follows from Theorem 3.4 that for any $\mu \in \mathcal{M}$ satisfying $\mu(T_0) = 1$, the OLA rule $\sigma_n(\mu)$ is $(0, \mu)$ -Pareto optimal at n for every $n \in N$. This model is also applicable to a withdrawal problem.

EXAMPLE 4.3 (WITHDRAWAL PROBLEM III). For each $t \in T_0 \subset T$, let (U_n^t) be a bounded sequence of random variables defined on $(\Omega, \mathcal{F}, P)$ and adapted to $(\mathcal{F}_n)$. For a given constant $\beta_t (0 < \beta_t < 1)$: discount factor, we define random sequence (X_n^t) by

$$X_n^t = \sum_{k=0}^n \beta_t^k U_k^t,$$

for $t \in T_0$. We assume that for each $t \in T_0$ and any $n \in N$

$$\begin{aligned} E[U_{n+1}^t | U_n^t > 0] &> 0, \\ P(U_{n+1}^t \leq 0 | U_n^t \leq 0) &= 1 \end{aligned}$$

and

$$P(\cup_n \{U_n^t \leq 0\}) = 1.$$

Then we easily see that $U_n^t > 0$ if and only if $X_n^t < E[X_{n+1}^t | \mathcal{F}_n]$, and that if $U_n^t \leq 0$ then $X_k^t \geq E[X_{k+1}^t | \mathcal{F}_k]$ for every $k \geq n$. Thus it follows that $A_n^t = \{U_n^t \leq 0\}$, $A_n^t \subset A_{n+1}^t$, $t \in T_0$, $n \in N$ and $\lim_{n \rightarrow \infty} P(A_n^t) = 1$.

Since

$$Y_n^t = -\beta_t^{n+1} E[U_{n+1}^t | \mathcal{F}_n],$$

it follows that if

$$E[U_{n+1}^t | \mathcal{F}_n] \geq \beta_t E[U_{n+2}^t | \mathcal{F}_n]$$

for each $n \in N$ and any $t \in T_0$, (Y_n^t) is a submartingale, and hence that for any $\mu \in \mathcal{M}$ satisfying $\mu(T_0) = 1$ Assumption 3.1 holds under condition that $\lim_{n \rightarrow \infty} P(\cap_{t \in T_0} A_n^t) = 1$ or that T_0 is finite.

If $\{U_n^s \leq 0\} = \{U_n^t \leq 0\}$ for each $n \in N$ and any $s, t \in T_0$, Assumption 3.2 is satisfied for such a μ .

We can explain this model as other withdrawal problem similarly to Huang and Li(1990) and Ohtsubo(1991).

EXAMPLE 4.4 (CF. CHOW *et al.*(1971)). For each $t \in T_0 \subset T$, let (W_n^t) be a sequence of independent and identically distributed random variables with finite mean, and let

$$m_n^t = \max_{0 \leq k \leq n} W_k^t, \quad X_n^t = m_n^t - c_n^t$$

for each $n \in N$, where $(c_n^t)_{n=0}^\infty$ is any strictly increasing sequence of positive constants. Then we have

$$X_{n+1}^t - X_n^t = (W_{n+1}^t - m_n^t)^+ - b_n^t,$$

where $b_n^t = c_{n+1}^t - c_n^t$. From Chow *et al.*(1971, p.56), it follows that if $b_{n+1}^t \geq b_n^t$ for all $n \in N$, that is, (c_n^t) is convex with regard to n , then $A_n^t \subset A_{n+1}^t$ for any $n \in N$ and

$$\lim_{n \rightarrow \infty} P(A_n^t) = P(\sigma_t < \infty) = 1,$$

where

$$\sigma_t = \inf\{n \geq 0 | X_n^t \geq E[X_{n+1}^t | \mathcal{F}_n]\} = \inf\{n \geq 0 | m_n^t \geq \beta_n^t\}$$

and β_n^t is the unique solution of the equation

$$E[(W_n^t - \beta_n^t)^+] = b_n^t.$$

Now in order to check that Assumption 3.1 is satisfied for any $\mu \in \mathcal{M}$ satisfying $\mu(T_0) = 1$, it suffices to show that (Y_n^t) is a submartingale. We have easily

$$\begin{aligned} E[Y_{n+1}^t - Y_n^t | \mathcal{F}_n] &= E[(X_{n+1}^t - X_{n+2}^t) - (X_n^t - X_{n+1}^t) | \mathcal{F}_n] \\ &= E[(W_{n+1}^t - m_n^t)^+ - (W_{n+2}^t - m_{n+1}^t)^+ | \mathcal{F}_n] - (b_n^t - b_{n+1}^t). \end{aligned}$$

Since $m_n^t \leq m_{n+1}^t$, $n \in N$ and (W_n^t) is i.i.d.,

$$E[(W_{n+1}^t - m_n^t)^+ | \mathcal{F}_n] \geq E[(W_{n+2}^t - m_{n+1}^t)^+ | \mathcal{F}_n].$$

Thus when $(b_n^t)_{n=0}^\infty$ is an increasing sequence for every $t \in T_0$, it follows that $E[Y_{n+1}^t | \mathcal{F}_n] \geq Y_n^t$, that is, (Y_n^t) is a submartingale.

References

- Aubin, J.P.(1979). *Mathematical Methods of Game and Economic Theory*, North-Holland, Amsterdam.
- Chow, Y.S., Robbins H. and Siegmund D.(1971). *Great Expectations : The Theory of Optimal Stopping*, Houghton Mifflin, Boston.
- Gnedin, A.V.(1994). On a best-choice problem with dependent criteria, *J. Appl. Prob.* **31**, 221-234.
- Gugerli, U.S.(1987). Optimal stopping of a Markov chain with vector-valued gain function, *Proc. 4th Vilnius Conference Prob. Theory Math. Statist.* **2**, VNU Sci. Press, Utrecht, 523-528.

- Hisano, H.(1980). An existence theorem in a vector-valued optimal stopping problem, *Mem. Fac. Sci. Kyushu Univ., Ser.A* **34**, 99-106.
- Huang, C.-F. and Li, L.(1990). Continuous time stopping games with monotone reward structures, *Math. Operat. Res.* **15**, 496-507.
- Neveu, J.(1975). *Discrete-Parameter Martingales*, North-Holland, Amsterdam.
- Ohtsubo, Y.(1991). On a discrete-time non-zero-sum Dynkin problem with monotonicity, *J. Appl. Prob.* **28**, 466-472.
- Ohtsubo, Y.(1997). Multi-objective stopping problem for a monotone case, *Mem. Fac. Sci. Kochi Univ. Ser.A* **18**, 99-104.
- Preater, J.(1993). A note on monotonicity in optimal multiple stopping problems, *Stat. Probab. Lett.* **16**, 407-410.
- Stadje, W.(1980). Efficient stopping of a random series of partially ordered points, *Springer Lecture Notes in Economics and Mathematical Systems.* **177**, 430-477.
- Yu, P.-L.(1985). *Multiple-criteria Decision Making*, Plenum Press, New York.

Received October 9, 1998

Revised April 23, 1999