ON THE DYNAMIC PROBABILISTIC INVENTORY PROBLEMS WITH PIECEWISE COST FUNCTIONS WHICH MAY NOT BE PIECEWISE SMOOTH

Sakaguchi, Michinori
Faculty of Economic Sciences , Hiroshima Shudo University

Kodama, Masanori
Faculty of Economic Sciences , Hiroshima Shudo University

<https://doi.org/10.5109/13510>

出版情報 : Bulletin of informatics and cybernetics. 34 (1), pp.75-90, 2002-10. Research
Association of Statistical Sciences

バージョン :

権利関係 :

ON THE DYNAMIC PROBABILISTIC INVENTORY PROBLEMS WITH PIECEWISE COST FUNCTIONS WHICH MAY NOT BE PIECEWISE SMOOTH

By

Michinori SAKAGUCHI* and Masanori KODAMA†

Abstract

A mathematical model of the probabilistic inventory problems with piecewise cost functions which may not be piecewise smooth is presented and various properties in this model are studied. Also some sufficient conditions on cost functions are found to ensure simple treatment on an optimal policy.

Key Words and Phrases: Dynamic inventory problem, Dynamic programming, Piecewise cost function.

1. Introduction

We consider the optimal policy of a probabilistic inventory problem with a piecewise cost function. The decision criterion is the maximization of expected profit or the minimization of expected costs which include the ordering, holding, and shortage costs. A typical example in single period problem is as follows. Let x be the amount on hand before an order is placed and let $\phi(b)$ be the probability density function of demand B . Let h and p be the holding and shortage costs per unit per period. Further, let c be the purchasing cost per unit and let z be the amount on hand in initial period after an order is received, which means that the initial regular order is $z - x$. If we assume no setup cost is occurred, the expected cost for the period is given by

$$E\{C(B, z)\} = c(z - x) + hE\{\text{holding cost}\} + pE\{\text{shortage cost}\}.$$

We define the function $H(z)$ by the equation $E\{C(B, z)\} = -cx + H(z)$ and let $f_1(x)$ be the minimal expectation of the total cost. Then we may write

$$f_1(x) = \min_{z \geq x} \{-cx + H(z)\}.$$

In one period problem the maximization (Kabak, 1984, Sorai, Arizono and Ohta, 1986) and the minimization (Kodama and Kitahara, 1983, Kodama, 1986) of the expected function in inventory models are considered and the special inventory problems are solved in Kodama (1990a) and Kodama (1991a). Moreover, a mathematical generalization of

* Faculty of Economic Sciences, Hiroshima Shudo University 1-1 Ohtsukahigasi-1-chome, Asaminami-ku, Hiroshima-si, 731-3195, Japan. tel +81-82-830-1237 sakaguti@shudo-u.ac.jp

† Faculty of Economic Sciences, Hiroshima Shudo University 1-1 Ohtsukahigasi-1-chome, Asaminami-ku, Hiroshima-si, 731-3195, Japan. tel +81-82-830-1942 kodama@shudo-u.ac.jp

the cost functions in the inventory problem is defined and analyzed in Kodama (1990b), Kodama (1992) and Kodama (1996).

In multiperiod models we suppose that $\phi(b)$ remains unchanged from period to period and demands in each period are independent. We should take in the discounted value of money in this case. That is, if $\alpha (< 1)$ is the discount factor per period and $f_n(x)$ is the discount expected loss for n -period inventory model when an optimal policy is used at each purchasing opportunity, then

$$f_n(x) = \min_{z \geq x} \left\{ -cx + H(z) + \alpha \int_0^\infty f_{n-1}(z-b)\phi(b)db \right\}.$$

Various properties of the optimal policy in multi-stage model are studied in Kodama (1991b) and Kodama (1993). In recent papers (Kodama, 1998a and Kodama, 1998b), we have attempted to express the optimal function by closed forms with known cost functions in the multi-stage model. Furthermore, some sufficient conditions on cost functions are found to ensure simple treatments on an optimal policy and specializations of cost functions are made and many examples are analyzed. In Kodama and Sakaguchi (2001c) we studied inventory models with a piecewise cost function that was piecewise smooth and discussed the properties of the optimal policy. We define functions $F_k(z)$ by the equations

$$F_{k-1}(z) = H(z) + \alpha \int_0^\infty f_{k-1}(z-b)\phi(b)db, \quad f_0(\cdot) = 0, \quad k = 1, 2, \dots, N$$

and let

$$\bar{x}_n = \inf\{z \mid F'_{n-1}(z) \geq 0\}.$$

Then the *optimal policy* in the probabilistic dynamic inventory problem was given as follows:

If $x < \bar{x}_n$, then order $(\bar{x}_n - x)$, otherwise do not order.

In this paper we give a mathematical model of dynamic inventory problems with a piecewise cost function which may not be piecewise smooth. That is to say, we assume that the function $H(z)$ is a piecewise function which may not be piecewise smooth. In this case the fundamental properties of the functions $F_k(z)$ are given in Theorem 2.6 and we would set

$$\bar{x}_n = \inf\{z \mid F'_{(n-1)+}(z) \geq 0\}.$$

We assume that $H_i(z)$ ($1 \leq i \leq m+1$) has a continuous second derivative and is a convex function on $[R_{i-1}, R_i]$ such that $H(z) = H_i(z)$ for $z \in (R_{i-1}, R_i]$. It is effective to decide the sign of $F'_{(n-1)+}(R_i)$ in order to get $\bar{x}_n$ and we show some sufficient conditions for this in terms of the discount factor α and $H(R_i)$ in Corollary to Theorem 3.1. Further we discuss the method to obtain $\bar{x}_n$ in some cases expressing the equation to solve by the known functions $H_i(z)$.

2. A mathematical model

Let c, α be real numbers with $0 < c$, $0 < \alpha < 1$. Let $R_1, \dots, R_m$ be a sequence of real numbers such that $R_1 < \dots < R_m$. Let $H_i(z)$ ($1 \leq i \leq m+1$) be real valued-functions defined on $[R_{i-1}, R_i]$, where we respectively abbreviate $(-\infty, R_1]$ to $[R_0, R_1]$ and $[R_m, \infty)$ to $[R_m, R_{m+1}]$. Throughout this paper we assume that for all i with $1 \leq i \leq m+1$, $H_i(z)$ has a continuous second derivative on $[R_{i-1}, R_i]$, and it is a convex function on $[R_{i-1}, R_i]$ which means by the condition that $H_i''(z) \geq 0$ on (R_{i-1}, R_i) . We also assume that

$$H_i(R_i) = H_{i+1}(R_i) \quad \text{and} \quad H_i'(R_i) \leq H_{i+1}'(R_i) \quad \text{for all } i \quad 1 \leq i \leq m \quad (2.1)$$

$$\lim_{z \rightarrow -\infty} H_1'(z) < 0, \quad \lim_{z \rightarrow \infty} H_{m+1}'(z) > c. \quad (2.2)$$

Now set

$$H(z) = H_i(z) \quad \text{for } z \in (R_{i-1}, R_i]. \quad (2.3)$$

Then $H(z)$ has a derivative at every point in the set $\mathbf{R}$ of all real numbers except $z = R_i \quad 1 \leq i \leq m$. We know that $H'_-(R_i)$ and $H'_+(R_i)$ exist, and that the inequality

$$H'_-(R_i) \leq H'_+(R_i) \quad (2.4)$$

holds for all i and we see that $H'_+(z)$ is non-decreasing on $(-\infty, \infty)$.

Let $\phi(b)$ be the density function of a real random variable B and we assume that $\phi(b)$ is a piecewise continuous function with $\phi(b) = 0$ for $b < 0$. For a given real number x and z we define functions $f_k(x)$, $F_k(x)$ ($k = 1, \dots, N$) as follows.

$$f_1(x) = \min_{z \geq x} \{-cx + H(z)\}, \quad (2.5)$$

$$f_k(x) = \min_{z \geq x} \left\{ -cx + H(z) + \alpha \int_0^\infty f_{k-1}(z-b)\phi(b)db \right\}, \quad k = 2, 3, \dots, N, \quad (2.6)$$

$$F_{k-1}(z) = H(z) + \alpha \int_0^\infty f_{k-1}(z-b)\phi(b)db, \quad f_0(\cdot) = 0, \quad k = 1, 2, \dots, N. \quad (2.7)$$

By (2.1), (2.2), (2.3) and the fact that $H_i(z)$ is a convex function on $[R_{i-1}, R_i]$ for all i ($1 \leq i \leq m+1$), there exists $\bar{x}_1$ such that

$$\bar{x}_1 = \inf\{z \mid H'_+(z) \geq 0\}. \quad (2.8)$$

Note that if $z \neq R_i$, then we have $H'_+(z) = H'(z)$.

LEMMA 2.1. *We have*

$$f_1(x) = \begin{cases} -cx + H(\bar{x}_1), & \text{if } x \leq \bar{x}_1, \\ -cx + H(x), & \text{if } x \geq \bar{x}_1. \end{cases}$$

In particular $f_1(x)$ is continuous on $\mathbf{R}$.

PROOF. By the assumption (2.1), we know that $H'_+(z)$ is a non-decreasing function on $(-\infty, \infty)$. By (2.8) we see that $H(\bar{x}_1)$ is the minimum of $H(z)$ on the interval $(-\infty, \infty)$. It thus follows that

$$\min_{z \geq x} H(z) = \begin{cases} H(\bar{x}_1), & \text{if } x \leq \bar{x}_1, \\ H(x), & \text{if } x > \bar{x}_1 \end{cases}$$

and hence we get the equation in this lemma. $\square$

Now we shall discuss some of the basic properties of the functions $f_k(x)$, $F_k(x)$ ($k = 1, \dots, N$). At first it will be shown that the function $F_1(z)$ has the similar properties to the function $H(z)$.

LEMMA 2.2. *We have*

$$F_1(z) = \begin{cases} H(z) - \alpha cz + \alpha \{H(\bar{x}_1) + cE(B)\}, & \text{if } z \leq \bar{x}_1, \\ H(z) - \alpha cz + \alpha \int_0^{z-\bar{x}_1} \{H(z-b) - H(\bar{x}_1)\} \phi(b) db \\ \quad + \alpha \{H(\bar{x}_1) + cE(B)\}, & \text{if } z > \bar{x}_1. \end{cases}$$

Furthermore $F_1(z)$ is continuous on R .

PROOF. Assume first that $z \leq \bar{x}_1$. Then it is shown that if $b \geq 0$, then $z - b \leq \bar{x}_1$ and hence $f_1(z - b) = -c(z - b) + H(\bar{x}_1)$ by Lemma 2.1. We therefore obtain

$$\begin{aligned} F_1(z) &= H(z) + \alpha \int_0^\infty \{-c(z - b) + H(\bar{x}_1)\} \phi(b) db \\ &= H(z) - \alpha cz + \alpha \{H(\bar{x}_1) + cE(B)\}. \end{aligned}$$

Next suppose that $z \geq \bar{x}_1$. Then $z - b \leq \bar{x}_1$ if and only if $z - \bar{x}_1 \leq b$. Thus we have

$$\begin{aligned} F_1(z) &= H(z) + \alpha \int_0^\infty f_1(z - b) \phi(b) db \\ &= H(z) + \alpha \int_0^{z-\bar{x}_1} \{-c(z - b) + H(z - b)\} \phi(b) db \\ &\quad + \alpha \int_{z-\bar{x}_1}^\infty \{-c(z - b) + H(\bar{x}_1)\} \phi(b) db \\ &= H(z) - \alpha cz + \alpha \int_0^{z-\bar{x}_1} \{H(z - b) - H(\bar{x}_1)\} \phi(b) db + \alpha \{H(\bar{x}_1) + cE(B)\}, \end{aligned}$$

and we complete the proof because it is clear that $F_1(z)$ is continuous. $\square$

LEMMA 2.3. $F_1(z)$ has a derivative on $[R_{i-1}, R_i]$ for all i ($1 \leq i \leq m + 1$), given by

$$F_1'(z) = \begin{cases} H'(z) - \alpha c, & \text{if } z \leq \bar{x}_1 \text{ and } z \neq R_j, \\ H'(z) - \alpha c + \alpha \int_0^{z-\bar{x}_1} H'(z - b) \phi(b) db, & \text{if } z > \bar{x}_1 \text{ and } z \neq R_j, \end{cases}$$

where $j = 1, \dots, m$. We have

$$F_{1+}'(R_i) = \begin{cases} H_+'(R_i) - \alpha c, & \text{if } R_i \leq \bar{x}_1, \\ H_+'(R_i) - \alpha c + \alpha \int_0^{R_i-\bar{x}_1} H'(R_i - b) \phi(b) db, & \text{if } R_i > \bar{x}_1 \end{cases}$$

and

$$F_{1-}'(R_i) = \begin{cases} H_-'(R_i) - \alpha c, & \text{if } R_i \leq \bar{x}_1, \\ H_-'(R_i) - \alpha c + \alpha \int_0^{R_i-\bar{x}_1} H'(R_i - b) \phi(b) db, & \text{if } R_i > \bar{x}_1, \end{cases}$$

in particular we see $F_{1-}'(R_i) \leq F_{1+}'(R_i)$ for all i ($1 \leq i \leq m$).

PROOF. Let $\bar{x}_1$ be an element in the interval $[R_{p-1}, R_p]$. If $z \leq \bar{x}_1$, our lemma is clear by Lemma 2.2. Therefore we may assume $\bar{x}_1 < z$. It also follows from Lemma 2.2 that we may only study a derivative of the function

$$\int_0^{z-\bar{x}_1} \{H(z-b) - H(\bar{x}_1)\} \phi(b) db.$$

For the sake of it we suppose that z is on the interval $[R_{i-1}, R_i]$. Then we see $p \leq i$ and have a sequence

$$R_{p-1} < \bar{x}_1 \leq R_p < \cdots < R_{k-1} < R_k < \cdots < R_{i-1} \leq z \leq R_i.$$

Put $g(z, b) = \{H(z-b) - H(\bar{x}_1)\} \phi(b)$ and let

$$K = \{(z, b) \mid R_{i-1} \leq z \leq R_i, 0 \leq b \leq z - \bar{x}_1\}.$$

Then the function $H(z-b)$ of two real variables z, b are continuous on the set K . Since $z - R_k \leq b \leq z - R_{k-1}$ if and only if $R_{k-1} \leq z - b \leq R_k$, we see that the function $H'(z-b)$ of two real variables z, b are continuous on the set $K - \cup_k L_k$, where $L_k = \{(z, b) \mid b = z - R_k\}$ ($p \leq k \leq i-1$). If $\phi(b)$ is continuous on $[0, z - \bar{x}_1]$ and $R_{i-1} < z < R_i$, then

$$\frac{d}{dz} \int_0^{z-\bar{x}_1} g(z, b) db = \int_0^{z-\bar{x}_1} \frac{\partial g}{\partial z}(z, b) db + g(z, z - \bar{x}_1) = \int_0^{z-\bar{x}_1} H'(z-b) \phi(b) db. \quad (2.9)$$

Similarly we obtain that if $z = R_{i-1}$, then

$$D_+ \int_0^{R_{i-1}-\bar{x}_1} g(R_{i-1}, b) db = \int_0^{R_{i-1}-\bar{x}_1} H'(R_{i-1}-b) \phi(b) db, \quad (2.10)$$

and that if $z = R_i$, then

$$D_- \int_0^{R_i-\bar{x}_1} g(R_i, b) db = \int_0^{R_i-\bar{x}_1} H'(R_i-b) \phi(b) db. \quad (2.11)$$

We have assumed that the function $\phi(b)$ is piecewise continuous, so suppose that $\phi(b)$ is discontinuous at $b = a_j$ ($1 \leq j \leq l$), where $a_1, a_2, \dots, a_l$ is an increasing sequence with $a_j \in [0, z - \bar{x}_1]$.

At first suppose that $z \in (R_{i-1}, R_i)$. In this case there is a positive number δ such that the following conditions:

1. We have $R_{i-1} \leq z - \delta, z + \delta \leq R_i$.
2. If $z - \delta \leq w < z$, then $w - R_k \neq a_j$ for all j, k .
3. If $z < w \leq z + \delta$, then $w - R_k \neq a_j$ for all j, k .

Let

$$K_{\delta-} = \{(w, b) \mid z - \delta \leq w \leq z, 0 \leq b \leq z - \bar{x}_1\},$$

$$K_{\delta+} = \{(w, b) \mid z \leq w \leq z + \delta, 0 \leq b \leq z - \bar{x}_1\}.$$

Considering the fact that the function $H'(z - b)\phi(b)$ of two real variables z, b are continuous at every point in the set $K_{\delta-}$ which is not on any lines $b = z - R_k$ ($p \leq k \leq i - 1$) and $b = a_j$ ($1 \leq j \leq l$), we are able to calculate a left-hand derivative by the method on $K_{\delta-}$ used in (2.11). Similarly by the same method on $K_{\delta+}$ used in (2.10), we may have a right-hand derivative at z , and these are shown to be equal. This implies that (2.9) holds in this case. There are remained cases where $z = R_{i-1}$ or $z = R_i$, and the equations (2.10) and (2.11) can be shown similarly in these cases.

The inequality

$$H'_-(R_i) \leq H'_+(R_i)$$

leads us to the relations

$$F'_{1-}(R_i) \leq F'_{1+}(R_i)$$

for all i ($1 \leq i \leq m$), therefore we complete our proof. $\square$

LEMMA 2.4. $F_1(z)$ has a second continuous derivative on $[R_{i-1}, R_i]$ for all i ($1 \leq i \leq m + 1$), given

$$F''_1(z) = \begin{cases} H''(z), & \text{if } z \leq \bar{x}_1 \text{ and } z \neq R_j, \\ H''(z) + \alpha \int_0^{z-\bar{x}_1} H''(z-b)\phi(b)db, & \text{if } z > \bar{x}_1 \text{ and } z \neq R_j, \end{cases}$$

where $j = 1, \dots, m$. We obtain

$$F''_{1+}(R_i) = \begin{cases} H''_+(R_i), & \text{if } R_i \leq \bar{x}_1, \\ H''_+(R_i) + \alpha \int_0^{R_i-\bar{x}_1} H''(R_i-b)\phi(b)db, & \text{if } R_i > \bar{x}_1 \end{cases}$$

and

$$F''_{1-}(R_i) = \begin{cases} H''_-(R_i), & \text{if } R_i \leq \bar{x}_1, \\ H''_-(R_i) + \alpha \int_0^{R_i-\bar{x}_1} H''(R_i-b)\phi(b)db, & \text{if } R_i > \bar{x}_1. \end{cases}$$

We have $F''_1(z) \geq 0$ on (R_{i-1}, R_i) for all i .

PROOF. As the method we used in Lemma 2.3 we can calculate a second derivative of $F_1(z)$. It also follows from the inequality $H''(z) \geq 0$ on (R_{i-1}, R_i) that $F_1(z)$ is a convex function on $[R_{i-1}, R_i]$ and hence our lemma is proved. $\square$

LEMMA 2.5. We have

$$\lim_{z \rightarrow -\infty} F'_1(z) < 0, \quad \lim_{z \rightarrow \infty} F'_1(z) > c.$$

PROOF. By (2.2) and Lemma 2.3 we see that

$$\lim_{z \rightarrow -\infty} F'_1(z) = \lim_{z \rightarrow -\infty} (H'(z) - \alpha c) < 0.$$

If $0 \leq b \leq z - \bar{x}_1$, then $z - b \geq \bar{x}_1$ and therefore $H'(z - b) \geq 0$. It thus follows that

$$\begin{aligned} \lim_{z \rightarrow \infty} F'_1(z) &= \lim_{z \rightarrow \infty} \left(H'(z) - \alpha c + \alpha \int_0^{z - \bar{x}_1} H'(z - b) \phi(b) db \right) \\ &= \lim_{z \rightarrow \infty} H'(z) - \alpha c + \alpha \int_0^{\infty} \lim_{z \rightarrow \infty} H'(z) \phi(b) db \\ &= \lim_{z \rightarrow \infty} H'(z) (1 + \alpha) - \alpha c > c. \end{aligned}$$

This completes the proof. $\square$

Now we have the following fundamental Theorem .

THEOREM 2.6. *Assume that $1 \leq k \leq N - 1$. Then there is a real number $\bar{x}_k$ such that $\bar{x}_k = \inf\{z \mid F'_{(k-1)+}(z) \geq 0\}$. We have the following statemens:*

1.

$$f_k(x) = \begin{cases} -cx + F_{k-1}(\bar{x}_k), & \text{if } x \leq \bar{x}_k, \\ -cx + F_{k-1}(x), & \text{if } x \geq \bar{x}_k. \end{cases}$$

In particular $f_k(x)$ is continuous on $\mathbf{R}$.

2.

$$F_k(z) = \begin{cases} H(z) - \alpha cz + \alpha \{F_{k-1}(\bar{x}_1) + cE(B)\}, & \text{if } z \leq \bar{x}_k, \\ H(z) - \alpha cz + \alpha \int_0^{z - \bar{x}_k} \{F_{k-1}(z - b) - F_{k-1}(\bar{x}_k)\} \phi(b) db \\ \quad + \alpha \{F_{k-1}(\bar{x}_k) + cE(B)\}, & \text{if } z > \bar{x}_k. \end{cases}$$

Furthermore $F_k(z)$ is continuous on $\mathbf{R}$.

3. $F_k(z)$ has a derivative on $[R_{i-1}, R_i]$ for all i ($1 \leq i \leq m + 1$), given by

$$F'_k(z) = \begin{cases} H'(z) - \alpha c, & \text{if } z \leq \bar{x}_k \text{ and } z \neq R_j, \\ H'(z) - \alpha c + \alpha \int_0^{z - \bar{x}_k} F'_{k-1}(z - b) \phi(b) db, & \text{if } z > \bar{x}_k \text{ and } z \neq R_j, \end{cases}$$

where $j = 1, \dots, m$. We have

$$F'_{k+}(R_i) = \begin{cases} H'_+(R_i) - \alpha c, & \text{if } R_i \leq \bar{x}_k, \\ H'_+(R_i) - \alpha c + \alpha \int_0^{R_i - \bar{x}_k} F'_{k-1}(R_i - b) \phi(b) db, & \text{if } R_i > \bar{x}_k \end{cases}$$

and

$$F'_{k-}(R_i) = \begin{cases} H'_-(R_i) - \alpha c, & \text{if } R_i \leq \bar{x}_k, \\ H'_-(R_i) - \alpha c + \alpha \int_0^{R_i - \bar{x}_k} F'_{k-1}(R_i - b) \phi(b) db, & \text{if } R_i > \bar{x}_k, \end{cases}$$

in particular we see $F'_{k-}(R_i) \leq F'_{k+}(R_i)$ for all i ($1 \leq i \leq m$)

4. $F_k(z)$ has a second continuous derivative on $[R_{i-1}, R_i]$ for all i ($1 \leq i \leq m+1$), given

$$F_k''(z) = \begin{cases} H''(z), & \text{if } z \leq \bar{x}_k \text{ and } z \neq R_j, \\ H''(z) + \alpha \int_0^{z-\bar{x}_k} F_{k-1}''(z-b)\phi(b)db, & \text{if } z > \bar{x}_k \text{ and } z \neq R_j, \end{cases}$$

where $j = 1, \dots, m$. We obtain

$$F_{k+}''(R_i) = \begin{cases} H_+''(R_i), & \text{if } R_i \leq \bar{x}_k, \\ H_+''(R_i) + \alpha \int_0^{R_i-\bar{x}_k} F_{k-1}''(R_i-b)\phi(b)db, & \text{if } R_i > \bar{x}_k \end{cases}$$

and

$$F_{k-}''(R_i) = \begin{cases} H_-''(R_i), & \text{if } R_i \leq \bar{x}_k, \\ H_-''(R_i) + \alpha \int_0^{R_i-\bar{x}_k} F_{k-1}''(R_i-b)\phi(b)db, & \text{if } R_i > \bar{x}_k. \end{cases}$$

We have $F_k''(z) \geq 0$ on (R_{i-1}, R_i) for all i .

5. $F_{k+}'(z)$ is a non-decreasing function on the set R of all real numbers.
6. We have

$$\lim_{z \rightarrow -\infty} F_k'(z) < 0, \quad \lim_{z \rightarrow \infty} F_k'(z) > c.$$

PROOF. If $k = 1$, our theorem follows from Lemma 2.1 through Lemma 2.5. Since we set $F_0(z) = H(z)$, the properties of the function $F_1(z)$ is gotten from those of $F_0(z)$. That is, we know by (2.6) and (2.7) that

$$\begin{aligned} f_2(x) &= \min_{z \geq x} \{-cx + F_1(z)\} \\ F_2(z) &= H(z) + \alpha \int_0^\infty f_2(z-b)\phi(b)db. \end{aligned}$$

We also see by Lemma 2.1 through Lemma 2.5 that the same assumptions which we suppose on the function $H(z)$ hold on $F_1(z)$. That is, $F_1(z)$ is continuous on $(-\infty, \infty)$ and has a second derivative on the interval $[R_{i-1}, R_i]$ for all i ($1 \leq i \leq m$). And also $F_1(z)$ is a convex function on $[R_{i-1}, R_i]$ for all i and we have

$$F_{1-}'(R_i) \leq F_{1+}'(R_i) \quad \text{for } i \ (1 \leq i \leq m)$$

and

$$\lim_{z \rightarrow -\infty} F_1'(z) < 0, \quad \lim_{z \rightarrow \infty} F_1'(z) > c.$$

Thus we can repeat the proof of lemmata to conclude that our theorem holds in the case $k = 2$. We are able to proceed to prove it in the case $k \geq 3$ and we complete our proof. $\square$

The following inequalities is valid to decide the sign of $F_k'(R_i)$ ($1 \leq k \leq N-1$, $1 \leq i \leq m$) (cf. Theorem 1 in Kodama and Sakaguchi, 2001b).

THEOREM 2.7. *We have*

$$\begin{aligned} H'(z) - \alpha c &\leq F'_k(z) \leq H'(z) - \alpha c + \alpha F'_{k-1}(z), \quad \text{if } z \geq \bar{x}_k \text{ and } z \neq R_i, \\ H'_+(R_i) - \alpha c &\leq F'_{k+}(R_i) \leq H'_+(R_i) - \alpha c + \alpha F'_{(k-1)+}(R_i), \quad \text{if } R_i \geq \bar{x}_k, \\ H'_-(R_i) - \alpha c &\leq F'_{k-}(R_i) \leq H'_-(R_i) - \alpha c + \alpha F'_{(k-1)-}(R_i), \quad \text{if } R_i \geq \bar{x}_k, \end{aligned}$$

for $1 \leq k \leq N-1$.

PROOF. Suppose that $\bar{x}_k \leq z$ and $z \neq R_i$. It follows from 3 in Theorem 2.6 that

$$F'_k(z) = H'(z) - \alpha c + \alpha \int_0^{z-\bar{x}_k} F'_{k-1}(z-b)\phi(b)db.$$

Note that $0 \leq b \leq z - \bar{x}_k$ if and only if $\bar{x}_k \leq z - b \leq z$. Since $F'_{k-1}(z)$ is a non-decreasing, we know that

$$0 \leq F'_{k-1}(\bar{x}_k) \leq F'_{k-1}(z-b) \leq F'_{k-1}(z) \quad \text{for } 0 \leq b \leq z - \bar{x}_k,$$

which implies

$$0 \leq \int_0^{z-\bar{x}_k} F'_{k-1}(z-b)\phi(b)db \leq \int_0^{z-\bar{x}_k} F'_{k-1}(z)\phi(b)db \leq F'_{k-1}(z).$$

Consequently we obtain

$$H'(z) - \alpha c \leq F'_k(z) \leq H'(z) - \alpha c + \alpha F'_{k-1}(z)$$

and, by 3 in Theorem 2.6, we also have

$$\begin{aligned} H'_+(R_i) - \alpha c &\leq F'_{k+}(R_i) \leq H'_+(R_i) - \alpha c + \alpha F'_{(k-1)+}(R_i), \\ H'_-(R_i) - \alpha c &\leq F'_{k-}(R_i) \leq H'_-(R_i) - \alpha c + \alpha F'_{(k-1)-}(R_i). \end{aligned}$$

We complete the proof. $\square$

3. Optimal policies

We wish to find the sequence $\bar{x}_1, \dots, \bar{x}_N$ to get optimal policies in the system of the probabilistic dynamic inventory problem. We studied in Kodama and Sakaguchi (2001a), Kodama and Sakaguchi (2001b) and Kodama and Sakaguchi (2001c) the interval $[R_{p-1}, R_p]$ which contains $\bar{x}_k$ under the assumption that the function $H'(z)$ is continuous. The interval may be found if we could seek the conditons $F'_{k-1}(R_{p-1}) < 0$ and $F'_{k-1}(R_p) \geq 0$. Although functions $F'_i(z)$ may be discontinuous in this paper, we have similar results considering the right derivatives $F'_{i+}(z)$ ($1 \leq i \leq N-1$) and we see them in the following theorem (cf. Theorem 2 in Kodama and Sakaguchi, 2001c).

THEOREM 3.1. *Let R be a real number. Then the following statements hold:*

1. *If $H'_+(R) \geq \alpha c$, then $F'_{k+}(R) \geq 0$ for all k $1 \leq k \leq N-1$.*
2. *If $1 \leq k \leq N-1$ and $F'_{(k-1)+}(R) \leq 0$, then $F'_{k+}(R) = H'_+(R) - \alpha c$.*

3. If $1 \leq k \leq N-1$, $H'_+(R) < \alpha c$ and $F'_{(k-1)+}(R) \leq 0$, then $F'_{k+}(R) < 0$.
4. If $H'_+(R) < 0$, then $F'_{k+}(R) < 0$ for all k $1 \leq k \leq N-1$.
5. If $1 \leq k \leq N-1$ and $H'_+(R) < \frac{\alpha c(1 + \alpha + \cdots + \alpha^{k-1})}{1 + \alpha + \cdots + \alpha^k}$, then $F'_{k+}(R) < 0$.
6. If $1 \leq k \leq N-1$ and $H'_+(R) = \frac{\alpha c(1 + \alpha + \cdots + \alpha^{k-1})}{1 + \alpha + \cdots + \alpha^k}$, then $F'_{k+}(R) \leq 0$.

PROOF. 1. If $R \leq \bar{x}_k$, then this is a direct consequence of 3 in Theorem 2.6. If $R > \bar{x}_k$, then it is also clear by Theorem 2.7.

2. In fact, if $F'_{(k-1)+}(R) < 0$, then $R < \bar{x}_k$, and hence $F'_{k+}(R) = H'_+(R) - \alpha c$ by 3 in Theorem 2.6. If $F'_{(k-1)+}(R) = 0$, then $\bar{x}_k \leq R$. It follows from Theorem 2.7 that

$$H'_+(R) - \alpha c \leq F'_{k+}(R) \leq H'_+(R) - \alpha c + \alpha F'_{(k-1)+}(R).$$

Substituting $F'_{(k-1)+}(R) = 0$ we have $F'_{k+}(R) = H'_+(R) - \alpha c$.

3. This easily follows from 2.
4. The statement 3 implies 4 immediately.
5. We see that $H'_+(R) < \alpha c$. Therefore by 3 we may assume that $F'_{i+}(R) > 0$ for all i $0 \leq i \leq k-1$. Hence it follows that $\bar{x}_{i+1} \leq R$. By Theorem 2.7 we have

$$F'_{(i+1)+}(R) \leq H'_+(R) - \alpha c + \alpha F'_{i+}(R) \quad \text{for all } i \text{ } 0 \leq i \leq k-1.$$

By these inequalities and our assumption we obtain

$$F'_{k+}(R) \leq H'_+(R)(1 + \alpha + \cdots + \alpha^k) - \alpha c(1 + \alpha + \cdots + \alpha^{k-1}) < 0.$$

6. We are also able to prove this by the same method of 5 that $F'_{k+}(R) \leq 0$, because

$$H'_+(R)(1 + \alpha + \cdots + \alpha^k) - \alpha c(1 + \alpha + \cdots + \alpha^{k-1}) = 0.$$

We complete our proof. □

COROLLARY 3.2. *Let p be an integer with $1 \leq p \leq m$. Then we have:*

1. If $H'_+(R_p) \geq \alpha c$, then $\bar{x}_k \leq R_p$ for all k $1 \leq k \leq N$.
2. If $H'_+(R_p) < 0$, then $R_p < \bar{x}_k$ for all k $1 \leq k \leq N$.
3. If $H'_+(R_p) < \frac{\alpha c(1 + \alpha + \cdots + \alpha^{l-1})}{1 + \alpha + \cdots + \alpha^l}$, then $R_p < \bar{x}_k$ for all k $l+1 \leq k \leq N$.

PROOF. These assertions are followed from Theorem 3.1, because $\bar{x}_k \leq R_p$ if and only if $F'_{(k-1)+}(R_p) \geq 0$, and the proof is complete. $\square$

For the purpose of calculating $\bar{x}_k$ we wish to get conditions that support us how to choose some functions $H_i(z)$ among the functions $H_1(z), \dots, H_{m+1}(z)$. Since there exist many complicated cases, we only consider simpler cases.

At first we know that

$$H'_+(R_1) \leq H'_+(R_2) \leq \dots \leq H'_+(R_m).$$

The corollary shows the following four particular cases:

Case A. If $H'_+(R_1) \geq \alpha c$, then $\bar{x}_k \leq R_1$ for all k $1 \leq k \leq N$.

Case B. If $H'_+(R_m) < 0$, then $R_m < \bar{x}_k$ for all k $1 \leq k \leq N$.

Case C. If $H'_+(R_{p-1}) < 0$ and $H'_+(R_p) \geq \alpha c$, then $R_{p-1} < \bar{x}_k \leq R_p$ for all k $1 \leq k \leq N$.

Case D_l. If $H'_+(R_{p-1}) < 0$, $F'_{(l-1)+}(R_p) \geq 0$ and $H'_+(R_m) < \frac{\alpha c(1 + \alpha + \dots + \alpha^{l-1})}{1 + \alpha + \dots + \alpha^l}$,

then $R_{p-1} < \bar{x}_k \leq R_p$ for all k $1 \leq k \leq l$ and $R_m < \bar{x}_k$ for all k $l+1 \leq k \leq N$.

When $m = 1$, the optimal policies are studied at the case D_l in Kodama (1998a).

Now we shall consider the probabilistic multi-period inventory model with zero delivery lag, backlogging of demand and linear purchasing cost $[c(y) = cy]$. Let $-cx + H(z)$ denote the expected one-period loss, given z is an amount on hand after an order is placed and let $f_k(x)$ denote the minimum total discount expected loss for period $1, 2, \dots, k$ where x is the initial stock level. From the principal of optimality, each $f_k(x)$ satisfies the functional equation (2.5) and (2.6). Then the *optimal policy* in our system of the probabilistic dynamic inventory problem is the following:

If $x < \bar{x}_k$, then order $(\bar{x}_k - x)$, otherwise do not order.

Now we study how to get the sequence $\bar{x}_1, \bar{x}_2, \dots, \bar{x}_N$ in simple cases. In general we have the following proposition (cf. Theorem 2.2 in Kodama, 1998a).

PROPOSITION 3.3. *We have*

$$\bar{x}_1 \leq \bar{x}_2 \leq \dots \leq \bar{x}_N.$$

PROOF. We prove $\bar{x}_p \leq \bar{x}_{p+1}$ by induction on p . By the definition of $\bar{x}_1$, $H'_+(z)$ is negative for $z < \bar{x}_1$ and therefore if $z < \bar{x}_1$, then $F'_{1+}(z) = H'_+(z) - \alpha c < 0$. This implies $\bar{x}_1 \leq \bar{x}_2$. Now we may assume $p \geq 2$. It is shown that $F'_{(p-1)+}(z) < 0$ for $z < \bar{x}_p$. Therefore if $z < \bar{x}_{p-1}$, then it follows from 3 in Theorem 2.6 that $F'_{(p-1)+}(z) = H'_+(z) - \alpha c < 0$.

By the inductive hypothesis we obtain $\bar{x}_{p-1} \leq \bar{x}_p$. Suppose that $\bar{x}_{p-1} \leq z < \bar{x}_p$. Then we have by 3 in Theorem 2.6

$$F'_{(p-1)+}(z) = H'_+(z) - \alpha c + \alpha \int_0^{z-\bar{x}_{p-1}} F'_{p-2}(z-b)\phi(b)db < 0. \quad (3.1)$$

If $0 \leq b \leq z - \bar{x}_{p-1}$ and $z - b \neq R_j$, then $\bar{x}_{p-1} \leq z - b \leq z < \bar{x}_p$, and so $F'_{p-2}(z-b) \geq 0$ because $F'_{p-2}(z)$ is not decreasing and $F'_{p-2}(\bar{x}_{p-1}) \geq 0$. This fact implies

$$\int_0^{z-\bar{x}_{p-1}} F'_{p-2}(z-b)\phi(b)db \geq 0,$$

and hence we obtain by (3.1) that $H'_+(z) - \alpha c < 0$ for $\bar{x}_{p-1} \leq z < \bar{x}_p$.

We have showed that $F'_{p+}(z) = H'_+(z) - \alpha c < 0$ for $z < \bar{x}_p$. Accordingly we conclude that $\bar{x}_p \leq \bar{x}_{p+1}$, and this proposition is proved. $\square$

The following proposition is obtained from 3 in Theorem 2.6.

PROPOSITION 3.4. (Section 5 in Kodama, 1998a) *We have for $z \geq \bar{x}_k$ with $z \neq R_i$ ($1 \leq i \leq m$)*

$$\begin{aligned}
 F'_k(z) = & H'(z) - \alpha c + \alpha \int_0^{z-\bar{x}_k} (H'(z-b_k) - \alpha c) \phi(b_k) db_k \\
 & + \alpha^2 \int_0^{z-\bar{x}_k} \int_0^{z-b_k-\bar{x}_{k-1}} (H'(z-b_k-b_{k-1}) - \alpha c) \phi(b_{k-1}) \phi(b_k) db_{k-1} db_k \\
 & + \alpha^3 \int_0^{z-\bar{x}_k} \int_0^{z-b_k-\bar{x}_{k-1}} \int_0^{z-b_k-b_{k-1}-\bar{x}_{k-2}} \\
 & \quad (H'(z-b_k-b_{k-1}-b_{k-2}) - \alpha c) \phi(b_{k-2}) \phi(b_{k-1}) \phi(b_k) db_{k-2} db_{k-1} db_k \\
 & + \dots \\
 & + \alpha^{k-1} \int_0^{z-\bar{x}_k} \int_0^{z-b_k-\bar{x}_{k-1}} \dots \int_0^{z-b_k-b_{k-1}-\dots-b_3-\bar{x}_2} \\
 & \quad (H'(z-b_k-b_{k-1}-\dots-b_2) - \alpha c) \phi(b_2) \phi(b_3) \dots \phi(b_k) db_2 db_3 \dots db_k \\
 & + \alpha^k \int_0^{z-\bar{x}_k} \int_0^{z-b_k-\bar{x}_{k-1}} \dots \int_0^{z-b_k-b_{k-1}-\dots-b_2-\bar{x}_1} \\
 & \quad H'(z-b_k-b_{k-1}-\dots-b_1) \phi(b_1) \phi(b_2) \dots \phi(b_k) db_1 db_2 \dots db_k.
 \end{aligned}$$

$\square$

Case A. Assume $H'_+(R_1) = H'_{2+}(R_1) \geq \alpha c$. Then it follows that $\bar{x}_k \leq R_1$ for all k ($1 \leq k \leq N$).

PROPOSITION 3.5. *Assume $H'_{1-}(R_1) < 0$ and $H'_{2+}(R_1) \geq \alpha c$. Then we have $\bar{x}_k = R_1$ for all k ($1 \leq k \leq N$).*

PROOF. Since $H'_{1-}(R_1) < 0$, $H'(z)$ is negative for $z < R_1$. Whence $\bar{x}_1 \geq R_1$, and we complete our proof by Proposition 3.3 and the condition in Case A. $\square$

PROPOSITION 3.6. *Suppose $H'_{2+}(R_1) \geq \alpha c$. If $\bar{x}_p = R_1$ for some p ($1 \leq p \leq N-1$), then $\bar{x}_k = R_1$ for all k with $p+1 \leq k \leq N$.*

PROOF. This is also clear by Proposition 3.3 and the condition in Case A. $\square$

Let $\bar{x}_{k+1} < R_1$. Then $\bar{x}_{k+1}$ is a solution of the equation $F'_k(z) = 0$, precisely it is the minimal solution of its equation. Because of $F'_0(z) = H'_1(z)$ for $z < R_1$, we may find $\bar{x}_1$ to solve the equation $H'_1(z) = 0$.

For $\bar{x}_1 \leq z < R_1$,

$$F'_1(z) = H'_1(z) - \alpha c + \alpha \int_0^{z-\bar{x}_1} H'_1(z-b_1) \phi(b_1) db_1.$$

Thus we can get $\bar{x}_2$ to solve the equation $F'_1(z) = 0$. It follows from Proposition 3.4 that the functions $F'_k(z)$ for $\bar{x}_k \leq z < R_1$ can be written by replacing $H'(z)$ with $H'_1(z)$, and we keep on this step.

Case B. Suppose $H'_+(R_m) < 0$. Then we have $R_m < \bar{x}_k$ for all k $1 \leq k \leq N$.

In this case there is a number $\bar{x}_1$ that is a solution of the equation $H'_{m+1}(z) = 0$ by the fact that $F'_0(z) = H'_{m+1}(z)$ for $z > R_m$. Replacing all the function $H'(z)$ in the equation of Proposition 3.4 with $H'_{m+1}(z)$, we are able to write $F'_k(z)$ for $z > R_m$ by using only the function $H'_{m+1}(z)$.

Case C. Assume that $H'_+(R_{p-1}) < 0$ and $H'_+(R_p) \geq \alpha c$. Then it follows that $R_{p-1} < \bar{x}_k \leq R_p$ for all k $1 \leq k \leq N$. Applying the properties in cases A and B to this case we know the following statements:

1. If $H'_-(R_p) < 0$, then $\bar{x}_k = R_p$ for all k ($1 \leq k \leq N$).
2. If $\bar{x}_k = R_p$ for some p ($1 \leq p \leq N-1$), then $\bar{x}_k = R_p$ for all k ($p+1 \leq k \leq N$).
3. If $\bar{x}_{k+1} < R_p$, then $R_{p-1} < \bar{x}_1 < R_p$. Hence we may find $\bar{x}_1$ to solve the equation $H'_p(z) = 0$. Replacing all the function $H'(z)$ in Proposition 3.4 with $H'_p(z)$, the functions $F'_k(z)$ ($R_{p-1} < z < R_p$) can be expressed in terms of the function $H'_p(z)$. For the sake of finding $\bar{x}_{k+1}$, it is enough to solve the equations $F'_i(z) = 0$ ($1 \leq i \leq k$) successively.

Case D_l. Suppose that $H'_+(R_{p-1}) < 0$, $F'_{(l-1)+}(R_p) \geq 0$ and $H'_+(R_m) < \frac{\alpha c(1 + \alpha + \dots + \alpha^{l-1})}{1 + \alpha + \dots + \alpha^l}$. Then we have $R_{p-1} < \bar{x}_k \leq R_p$ for all k $1 \leq k \leq l$ and $R_m < \bar{x}_k$ for all k $l+1 \leq k \leq N$.

In this case it is necessary to change the $H'_i(z)$ in order to express $F'_k(z)$. We write them explicitly. At first solve the equation $H'_p(z) = 0$ to get $\bar{x}_1$. Next for $2 \leq k \leq l-1$ and $\bar{x}_k \leq z < R_p$, we have by Proposition 3.4

$$\begin{aligned}
 F'_k(z) = & H'_p(z) - \alpha c + \alpha \int_0^{z-\bar{x}_k} (H'_p(z-b_k) - \alpha c) \phi(b_k) db_k \\
 & + \alpha^2 \int_0^{z-\bar{x}_k} \int_0^{z-b_k-\bar{x}_{k-1}} (H'_p(z-b_k-b_{k-1}) - \alpha c) \phi(b_{k-1}) \phi(b_k) db_{k-1} db_k \\
 & + \alpha^3 \int_0^{z-\bar{x}_k} \int_0^{z-b_k-\bar{x}_{k-1}} \int_0^{z-b_k-b_{k-1}-\bar{x}_{k-2}} \\
 & \quad (H'_p(z-b_k-b_{k-1}-b_{k-2}) - \alpha c) \phi(b_{k-2}) \phi(b_{k-1}) \phi(b_k) db_{k-2} db_{k-1} db_k \\
 & + \dots \\
 & + \alpha^{k-1} \int_0^{z-\bar{x}_k} \int_0^{z-b_k-\bar{x}_{k-1}} \dots \int_0^{z-b_k-b_{k-1}-\dots-b_3-\bar{x}_2} \\
 & \quad (H'_p(z-b_k-b_{k-1}-\dots-b_2) - \alpha c) \phi(b_2) \phi(b_3) \dots \phi(b_k) db_2 db_3 \dots db_k \\
 & + \alpha^k \int_0^{z-\bar{x}_k} \int_0^{z-b_k-\bar{x}_{k-1}} \dots \int_0^{z-b_k-b_{k-1}-\dots-b_2-\bar{x}_1} \\
 & \quad H'_p(z-b_k-b_{k-1}-\dots-b_1) \phi(b_1) \phi(b_2) \dots \phi(b_k) db_1 db_2 \dots db_k.
 \end{aligned}$$

We may continue to find a solution $\bar{x}_{k+1}$ of the equation $F'_k(z) = 0$, $\bar{x}_k \leq z < R_p$ for $2 \leq k \leq l-1$. If we could not find any solution, then $\bar{x}_{k+1} = R_p$.

Next we shall show $F'_l(z)$ by $H'_l(z)$, however it is complicated. For the simplicity we introduce the functions $h_l(z)$, $h_{l-1}(z)$, $h_{l-2}(z)$, $\dots$, $h_2(z)$, $h_1(z)$ for $z > R_m$ as follows.

$$\begin{aligned} h_{l-s}(z) &= \int_0^{z-\bar{x}_l} \int_0^{z-b_l-\bar{x}_{l-1}} \dots \int_0^{z-b_l-b_{l-1}-\dots-b_{l-s+1}-\bar{x}_{l-s}} \\ &\quad (H'(z-b_l-b_{l-1}-\dots-b_{l-s}) - \alpha c) \phi(b_{l-s}) \dots \phi(b_{l-1}) \phi(b_l) db_{l-s} \dots db_{l-1} db_l \\ &\quad (0 \leq s \leq l-2), \\ h_1(z) &= \int_0^{z-\bar{x}_l} \int_0^{z-b_l-\bar{x}_{l-1}} \dots \int_0^{z-b_l-b_{l-1}-\dots-b_2-\bar{x}_1} \\ &\quad H'(z-b_l-b_{l-1}-\dots-b_1) \phi(b_1) \phi(b_2) \dots \phi(b_l) db_1 db_2 \dots db_l. \end{aligned}$$

Then we obtain

$$\begin{aligned} h_l(z) &= \int_0^{z-\bar{x}_l} H'(z-b_l) \phi(b_l) db_l - \alpha c \int_0^{z-\bar{x}_l} \phi(b_l) db_l \\ &= \int_0^{z-R_m} H'(z-b_l) \phi(b_l) db_l + \sum_{t=p}^{m-1} \int_{z-R_{m+p-t}}^{z-R_{m+p-t-1}} H'(z-b_l) \phi(b_l) db_l \\ &\quad + \int_{z-R_p}^{z-\bar{x}_l} H'(z-b_l) \phi(b_l) db_l - \alpha c \int_0^{z-\bar{x}_l} \phi(b_l) db_l \\ &= \int_0^{z-R_m} H'_{m+1}(z-b_l) \phi(b_l) db_l + \sum_{t=p}^{m-1} \int_{z-R_{m+p-t}}^{z-R_{m+p-t-1}} H'_{m+p-t}(z-b_l) \phi(b_l) db_l \\ &\quad + \int_{z-R_p}^{z-\bar{x}_l} H'_p(z-b_l) \phi(b_l) db_l - \alpha c \int_0^{z-\bar{x}_l} \phi(b_l) db_l. \end{aligned}$$

Let $1 \leq s \leq l-1$. We set

$$\begin{aligned} K_{l-s} &= \left\{ (b_{l-s}, b_{l-s+1}, \dots, b_l) \mid 0 \leq b_{l-n} \leq z - \sum_{j=0}^{n-1} b_{l-j} - \bar{x}_{l-n}, 0 \leq n \leq s \right\}, \\ K_{l-sp} &= \left\{ (b_{l-s}, b_{l-s+1}, \dots, b_l) \mid z - R_p \leq \sum_{j=0}^s b_{l-s+j} \leq z - \bar{x}_{l-s}, \right. \\ &\quad \left. \sum_{j=0}^n b_{l-n+j} \leq z - \bar{x}_{l-n} \ (0 \leq n \leq s-1), \ b_{l-s+n} \geq 0 \ (0 \leq n \leq s) \right\}, \\ K_{l-si} &= \left\{ (b_{l-s}, b_{l-s+1}, \dots, b_l) \mid z - R_i \leq \sum_{j=0}^s b_{l-s+j} \leq z - R_{i-1}, \right. \\ &\quad \left. b_{l-s+n} \geq 0 \ (0 \leq n \leq s) \right\} \ (p+1 \leq i \leq m), \\ K_{l-sm+1} &= \left\{ (b_{l-s}, b_{l-s+1}, \dots, b_l) \mid \sum_{j=0}^s b_{l-s+j} \leq z - R_m, \ b_{l-s+n} \geq 0 \ (0 \leq n \leq s) \right\}. \end{aligned}$$

Then

$$K_{l-s} = K_{l-sp} \cup K_{l-sp+1} \cup \cdots \cup K_{l-sm} \cup K_{l-sm+1}.$$

Put

$$\begin{aligned} g(\mathbf{b}) &= g(b_{l-s}, b_{l-s+1}, \dots, b_l) \\ &= (H'(z - b_l - b_{l-1} - \cdots - b_{l-s}) - \alpha c) \phi(b_{l-s}) \phi(b_{l-s+1}) \cdots \phi(b_l) \\ g_t(\mathbf{b}) &= g_t(b_{l-s}, b_{l-s+1}, \dots, b_l) \\ &= (H'_t(z - b_l - b_{l-1} - \cdots - b_{l-s}) - \alpha c) \phi(b_{l-s}) \phi(b_{l-s+1}) \cdots \phi(b_l), \\ &\quad (p \leq t \leq m+1), \end{aligned}$$

where we make a minor change when $s = l - 1$. It is shown that

$$\begin{aligned} h_{l-s}(z) &= \int \int \cdots \int_{K_{l-s}} g(\mathbf{b}) db_{l-s} db_{l-s+1} \cdots db_l \\ &= \sum_{t=p}^{m+1} \int \int \cdots \int_{K_{l-st}} g(\mathbf{b}) db_{l-s} db_{l-s+1} \cdots db_l \\ &= \sum_{t=p}^{m+1} \int \int \cdots \int_{K_{l-st}} g_t(\mathbf{b}) db_{l-s} db_{l-s+1} \cdots db_l \end{aligned}$$

This yields us to express the function $F'_l(z)$ as follow

$$F'_l(z) = H'_{m+1}(z) - \alpha c + \sum_{i=0}^{l-1} \alpha^{i+1} h_{l-i}(z).$$

We need to keep on searching $\bar{x}_{l+2}$, $\bar{x}_{l+3}$, $\dots$, $\bar{x}_N$, and so it is necessary to write $F'_{l+1}(z)$, $F'_{l+2}(z)$, $\dots$, $F'_{N-1}(z)$. Unfortunately they demand more calucations than that of $F'_l(z)$ if we would search them in the general form.

We have discussed how to find x_N . However it is too difficult to deal with this inventory system even when it is rather easier case. We should develop the thoery to handle our dynamic inventory problem.

References

- Kabak, I.W. (1984). Partial Returns in the Single Period Inventory Model, *IE News*, **19**, 1-3.
- Kodama, M. (1986). Probabilistic Scheduling-Period-Order-Level Models with General Demand Pattern (in Japanese), *Society of Political Economy*, **51**, 35-44.
- Kodama, M. (1990a). Probabilistic Inventory Models with Partial Returns and Additional Orders (I) (in Japanese), *Society of Political Economy*, **55**, 31-48.
- Kodama, M. (1990b). Multi-Stage Inventory Models with Various Partial Returns and Additional Orders (in Japanese), *Society of Political Economy*, **56**, 1-26.
- Kodama, M. (1991a). Probabilistic Inventory Models with Partial Returns and Additional Orders (II) (in Japanese), *Society of Political Economy*, **56**, 277-293

- Kodama, M. (1991b). Optimal Policy for a Non-convex Expected Loss Functions(II) (in Japanese), *Society of Political Economy*, **57**, 175-198.
- Kodama, M. (1992). Optimal Policy for a Probabilistic System (I) (in Japanese), *Society of Political Economy*, **58**, 35-50.
- Kodama, M. (1993). Optimal Policy for a Probabilistic System (II) (in Japanese), *Society of Political Economy*, **59**, 17-27.
- Kodama, M. (1996). Some Probabilistic Inventory Problems with Various Demand Pattern, *Journal of Information & Optimization Science*, **17**, 17-48.
- Kodama, M. (1998a). Dynamic Inventory Models with Piecewise Cost Function (I) (in Japanese), *Journal of Economic Sciences*, **1**, 99-122.
- Kodama, M. (1998b). Dynamic Inventory Models with Piecewise Cost Function (II) (in Japanese), *Journal of Economic Sciences*, **2**, 33-60.
- Kodama, M. and Kitahara, T. (1983). A Study of Unified Inventory Model on Various Demand (in Japanese), *Society of Political Economy*, **47**, 49-72.
- Kodama, M. and Sakaguchi, M. (2001a). Some comments on the probabilistic dynamic inventory problems with piecewise cost functions, *Recent Developments in Operational Research*, Narosa New Delhi, 191-196.
- Kodama, M. and Sakaguchi, M. (2001b). Probabilistic analysis of dynamic inventory models with general demand pattern, *Journal of Information & Optimization Sciences*, **22**, 57-72.
- Kodama, M. and Sakaguchi, M. (2001c). Dynamic programming on the probabilistic inventory models with multiple piecewise cost functions, *Journal of Information & Optimization Sciences*, **22**, 419-440.
- Sorai, M., Arizono, I. and Ohta, H. (1986). A Solution of Single Period Inventory Model with Partial Returns and Additional Orders (in Japanese), *JIMA*, **37**, 100-105.

Received August 3, 2001

Revised February 28, 2002

Re-revised March 26, 2002