

Preparation of Anti-dinitrotoluene Polyclonal Antibody and Effect of the Hapten Spacer Length in Coating Antigen on Immunoassay Sensitivity

Nagatomo, Kazutaka

Laboratory of Food Analysis, Division of Food Biotechnology, Department of Bioscience and Biotechnology, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

Matsumoto, Kiyoshi

Laboratory of Food Analysis, Division of Food Biotechnology, Department of Bioscience and Biotechnology, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

Ishitobi, Sachiko

Laboratory of Food Analysis, Division of Food Biotechnology, Department of Bioscience and Biotechnology, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

Koga, Mao

Laboratory of Food Analysis, Division of Food Biotechnology, Department of Bioscience and Biotechnology, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

他

<https://doi.org/10.5109/14054>

出版情報：九州大学大学院農学研究院紀要. 54 (1), pp.165-171, 2009-02-27. Faculty of Agriculture, Kyushu University

バージョン：

権利関係：



Preparation of Anti-dinitrotoluene Polyclonal Antibody and Effect of the Hapten Spacer Length in Coating Antigen on Immunoassay Sensitivity

Kazutaka NAGATOMO¹, Kiyoshi MATSUMOTO*, Sachiko ISHITOBI¹,
Mao KOGA¹, Kiyoshi TOKO² and Norio MIURA³

Laboratory of Food Analysis, Division of Food Biotechnology, Department of
Bioscience and Biotechnology, Faculty of Agriculture,
Kyushu University, Fukuoka 812–8581, Japan

(Received October 28, 2008 and accepted December 5, 2008)

A polyclonal antibody against 2,4-dinitrotoluene (2,4-DNT) has been raised in rabbit, and the antibody was used to detect 2,4-DNT using an enzyme-linked immunosorbent assay (ELISA) method. A 2,4-dinitrophenyl-keyhole limpet hemocyanine (DNPh-KLH) conjugate was injected into a rabbit, and a polyclonal anti-DNPh-KLH antibody was realized after purification of the serum using protein G column. Aspects of the anti-DNPh-KLH antibody to various nitroaromatic compounds, such as cross-reactivities and avidity, were characterized. The temperature dependence of the avidity between the anti-DNPh-KLH antibody and 2,4-DNT was also evaluated. The quantification of 2,4-DNT was based on the principle of indirect competitive ELISA, in which the immunoreaction between the coating antigen-protein conjugates and the anti-DNPh-KLH antibody were inhibited in the presence of free 2,4-DNT in solution. The detection was performed using alkaline phosphatase-labeled anti-rabbit IgG with *p*-nitrophenylphosphate as a substrate. The addition of a mixture of free 2,4-DNT to the anti-DNPh-KLH antibody was found to decrease the absorbance at 405 nm due to the competitive effect of 2,4-DNT. The effect of the structure of the coating antigen-protein conjugate on the competition of free 2,4-DNT or coating antigen toward anti-DNPh-KLH antibody was also investigated. The immunoassay exhibited excellent sensitivity for the detection of 2,4-DNT in the concentration range of 1 ng mL⁻¹ to 10 µg mL⁻¹.

INTRODUCTION

Environmental pollution of soil, river water, and underground water by explosive-related compounds is a serious problem in the areas of buried landmines and/or landmine factories (Hilmi *et al.*, 1999). These compounds are suspected to have human toxicity, mutagenicity, and carcinogenicity (Kennel *et al.*, 2000; Sayama *et al.*, 1998; Tchounwou *et al.*, 2001; Banerjee *et al.*, 1999). Therefore, there is a pressing need to determine the concentrations and distributions of these compounds in the environment. The main component of landmines is 2,4,6-trinitrotoluene (TNT). TNT released into the environment is decomposed and converted to dinitrotoluenes (DNTs) and aminodinitrotoluenes (ADNTs), among other products, by microorganisms, light, and heat. These compounds have been measured by using HPLC or other instrumental analytical devices (Borch and Gerlach, 2004; Ahmad and Roberts; 1995). These instrumental analyses, however, require tedious pre-treatments such as extraction from samples (Halasz *et al.*, 2002). On the other hand, detection methods using an antigen-antibody reaction, such as enzyme-linked immunosorbent assay (ELISA) and fluorescent and chemiluminescent

immunosensors, are more useful and are able to detect ppb levels of explosive compounds (Goldman *et al.*, 2003; Green *et al.*, 2002; Wilson *et al.*, 2003).

We have been engaged in development of an antibody with high avidity to explosive-related compounds. In recent years, we have focused on the development of highly sensitive surface plasmon resonance (SPR) sensing methods for the detection of TNT using an antigen-antibody reaction (Sakai *et al.*, 2003; Shankaran *et al.*, 2004; Shankaran *et al.*, 2005a; Shankaran *et al.*, 2005b; Matsumoto *et al.*, 2005; Shankaran *et al.*, 2006). According to the reports of Jenkins *et al.* (2000, 2001), the concentrations of 2,4-DNT and other decomposed nitroaromatic compounds around and over buried landmines are higher than the concentration of TNT.

In the present study, a polyclonal antibody against 2,4-DNT was raised in a rabbit, and the antibody was applied to the detection of 2,4-DNT using ELISA. The aim was to prepare anti-DNPh-KLH antibody from a rabbit and to clarify the characteristics for the detection of 2,4-DNT based on the principle of indirect competitive ELISA. In the course of the detection of free 2,4-DNT, we recognized that the combination of the structure of the coating antigen-protein, free 2,4-DNT, and the anti-DNPh-KLH antibody are an important factor in sensitivity using an indirect competitive immunoassay. Therefore, the effect of the structures of the coating antigen-protein conjugates on the sensitivity of the detection of free 2,4-DNT was investigated in detail.

¹ Laboratory of Food Analysis, Division of Food Biotechnology, Department of Bioscience and Biotechnology, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

² Graduate School of Information Science and Electrical Engineering, Kyushu University, Fukuoka, 819–0395, Japan

³ Art, Science and Technology Center for Cooperative Research, Kyushu University, Kasuga, Fukuoka, 816–8580, Japan

* Corresponding author (E-mail: kmatsu@agr.kyushu-u.ac.jp)

MATERIALS AND METHODS

Materials

Bovine serum albumin (BSA), *p*-nitrophenyl phosphate disodium salt (*p*-NPP), sodium sulfate, 2,4,6-trinitrobenzenesulfonate sodium salt (TNBS), 1,6-diaminohexane, 6-aminohexanoic acid, *N*-hydroxysuccinimide (NHS), and gelatin were obtained from Nacalai Tesque, Inc. (Kyoto, Japan). Ovalbumin (OVA), keyhole limpet hemocyanin (KLH), alkaline phosphatase (ALP)-labeled anti-rabbit IgG, 1-ethyl-3-(3-dimethyl aminopropyl) carbodiimide (EDC), hexylamine, ethylamine, 2,4-dinitrotoluene (2,4-DNT), and *N*-(2,4-dinitrophenyl) glycine (2,4-DNPh-gly) were obtained from Sigma (St. Louis, MO, USA). 2,4,6-Trinitrophenol (TNPh-OH) and 1,3-dinitrobenzene were purchased from Kishida Chemical Co., Ltd. (Osaka, Japan). 2,6-Dinitrotoluene (2,6-DNT), *bis*(2-methoxyethyl) ether, 1,2-diaminoethane, 1,4-diaminobutane, 1,8-diaminooctane, 1,10-diaminodecane, and 1,12-diaminododecane were obtained from Wako Pure Chemicals Ind., Ltd. (Osaka, Japan). *N*-(2,4,6-trinitrophenyl) glycine (TNPh-gly) was obtained from Funakoshi Co. (Tokyo, Japan). 2,4-Dinitrobenzenesulfonate sodium salt (2,4-DNBS) was from Research Chemicals (Toronto, Canada). 2,4,6-Trinitrotoluene (TNT), 2-amino-4,6-dinitrotoluene (2-amino-4,6-DNT), and 4-amino-2,6-dinitrotoluene (4-amino-2,6-DNT) were purchased from Supelco (Bellefonte, PA, USA). Freund's complete adjuvant was obtained from Difco (Detroit, MI, USA). Rabbit (6 weeks old, female) was purchased from Kyoyu (Fukushima, Japan). The Hi Trap Protein G column and PD-10 column were purchased from Amersham Bioscience (Piscataway, NJ, USA). All other reagents were of analytical-reagent grade. All buffer solutions were prepared using water purified with a Milli-Q filter (Millipore, Bedford, MA, USA).

Apparatus

ELISA measurements were performed using 96-well immunoplates (NUNC, No. 446612, Roskilde, Denmark) and a microplate reader (Wallac 1420, Perkin Elmer Life Science Japan, Tokyo, Japan). Spectrophotometric measurements were performed using a Shimadzu Multi Spec 1500 (Kyoto, Japan).

Preparation of DNPh-KLH

2,4-DNBS (in 1 mg mL⁻¹ H₂O, 1 mL) was reacted with 1 mL of 0.48 mol L⁻¹ (M) NaHCO₃ solution (pH 8.5) containing 10 mg KLH for 12 h at 40 °C. After the reaction, the preparation was dialyzed against five changes of H₂O at 4 °C for 3 days, and then lyophilized to produce the DNPh-KLH conjugate. The molar ratios of the DNPh moieties to the protein were estimated using the molar adsorption coefficient of DNPh-NH- (1.74 × 10⁴ cm⁻¹ M⁻¹ at 360 nm).

Preparation of coating antigen-protein conjugates

DNPh-OVA: 2,4-DNBS (in 2 mg mL⁻¹ H₂O, 1 mL) was reacted with 1 mL of 25 mM borate buffer (pH 8.0) con-

taining 10 mg OVA for 15 h at 40 °C. After the reaction, the preparation was dialyzed against five changes of H₂O at room temperature for 2 days, and then lyophilized to produce the DNPh-OVA-1 conjugate. The other five conjugates were prepared by the same procedures as DNPh-OVA-1 except for the concentration of 2,4-DNBS (DNPh-OVA-2: 10 mg mL⁻¹; DNPh-OVA-3: 20 mg mL⁻¹; DNPh-OVA-4: 40 mg mL⁻¹; DNPh-OVA-5: 100 mg mL⁻¹; DNPh-OVA-6: 200 mg mL⁻¹).

DNPh-Cx-OVA: Six different DNPh-Cx-OVA conjugates with different length methylene chains (spacers) between the hapten and protein (OVA) were prepared. In a solution of 25 mM borate buffer (pH 8.0, 1 mL), 8.82 mg of OVA and 2.18 mg of NHS were dissolved, and then 52.9 mg of EDC was added to the solution and stirred over night at room temperature. 1,2-Diaminoethane (0.51 mg, liquid) was gently added to the reaction mixture and stirred over-night. The mixture was dialyzed against three changes of H₂O at room temperature and lyophilized to produce C2-OVA. After the lyophilized powder (2 mg) was dissolved in 1 mL of 4% NaHCO₃ solution, 1 mL of 2,4-DNBS solution (16 mg mL⁻¹) was added to the solution and stirred over-night at 40 °C. The reaction mixture was dialyzed against H₂O and lyophilized to produce DNPh-C2-OVA conjugate. Other conjugates were prepared using the same procedures except for the concentrations of diaminoalkanes and 2,4-DNBS (DNPh-C4-OVA: 0.78 mg of 1,4-diaminobutane in 200 μ L DMSO, 16 mg mL⁻¹ of 2,4-DNBS; DNPh-C6-OVA: 1.00 mg of 1,6-diaminohexane in 200 μ L DMSO, 16 mg mL⁻¹ of 2,4-DNBS; DNPh-C8-OVA: 1.24 mg of 1,8-diaminooctane in 200 μ L DMSO, 16 mg mL⁻¹ of 2,4-DNBS; DNPh-C10-OVA: 1.49 mg of diaminodecane in 200 μ L DMSO, 5 mg mL⁻¹ of 2,4-DNBS; DNPh-C12-OVA: 1.49 mg of 1,12-diaminododecane in 200 μ L DMSO, 7 mg mL⁻¹ of 2,4-DNBS).

DNPh-glycine-OVA: 25 mg of DNPh-glycine and 13 mg of NHS were dissolved in 1 mL of *bis*(2-methoxyethyl) ether. Fifty mg of sodium sulfate was added, and the mixture was cooled to 0 °C. To the mixture was added 34 mg of EDC, and the reaction mixture was allowed to warm to room temperature and stirred over-night. Ten mg of OVA were dissolved in 1 mL of 25 mM borate buffer (pH 8.0). At intervals of 30 min, three aliquots of the NHS-ester solution (each 62 μ L) were added slowly to the OVA solution under intense stirring. After reacting 2 h, the mixture was passed through a PD 10 gel filtration column, and then the eluted fraction was lyophilized to produce DNPh-glycine-OVA conjugate (DNPh-gly-OVA).

Synthesis of TNPh-derivatives

TNPh-derivatives were synthesized according to our previous report (Sakai *et al.*, 2003).

Immunization

A rabbit was immunized with DNPh-KLH conjugate according to the following procedure. Conjugate dissolved in PBS (0.6 mg mL⁻¹) was emulsified with an equal volume of Freund's complete adjuvant. On days 0, 14,

28, 42 and 56, 1 mL of the prepared mixture was intracutaneously injected into the rabbit. On days 0, 35, 49 and 63, the rabbit was bled from an ear vein, and the antiserum was collected by centrifugation of the blood sample. The antiserum was tested by direct enzyme-linked immunosorbent assay (direct-ELISA). Ninety-six-well immunoplates were coated with 100 μ L of DNPh-KLH conjugate (10 μ g mL⁻¹ in 50 mM carbonate buffer, pH 9.6) over-night at room temperature. The plates were then washed three times with PBS containing 0.05% Tween 20 (PBST), and treated with 150 μ L of 1% gelatin for 1 h at room temperature, and again washed three times with PBST and reacted with antiserum at eight different dilutions (1/1000 to 1/128000 in PBS), which were added to the wells (50 μ L to each well) and incubated for 2 h at room temperature. The plates were washed again three times with PBST, and a solution of ALP-labeled anti-rabbit IgG (2000-fold dilution in PBS) was added (100 μ L to each well) and incubated for 1 h at room temperature. The plates were washed again, and the substrate solution (2 mg mL⁻¹ of *p*-NPP in 50 mM carbonate buffer, pH 9.6, containing 1 mM MgCl₂ and 0.1 mM ZnCl₂) was added and incubated for 30 min at room temperature. Then absorbance was measured at 405 nm.

Preparation and purification of polyclonal anti-DNPh-KLH antibody

Whole blood of the rabbit was collected by cardiocentesis 7 days after the last injection. The purification was performed according to our previous paper (Sakai *et al.*, 2003) with a minor modification.

Indirect competitive ELISAs for 2,4-DNT

ELISAs for 2,4-DNT were performed as follows. Ninety-six-well immunoplates were coated with 100 μ L of DNPh-OVA conjugates (DNPh-OVA-1 to DNPh-OVA-6) or DNPh-Cx-OVA conjugates (5 μ g mL⁻¹ in 50 mM carbonate buffer, pH 9.8) over-night at room temperature. The following day, the plates were washed three times with PBST and treated with 150 μ L of 1% gelatin for 1 h at room temperature. The plates were washed three times with PBST and reacted with 100 μ L of the equivalent mixtures of anti-DNPh-KLH antibody (2 μ g mL⁻¹) and serially diluted DNPh-derivative antigen for 1 h at room temperature. The plates were washed three times with PBST, and then reacted with 100 μ L of ALP-labeled anti-rabbit IgG (2000-fold dilution in PBS) for 45 min at room temperature. After washing three times with PBST, 100 μ L of the substrate solution (2 mg mL⁻¹ of *p*-NPP in 50 mM carbonate buffer, pH 9.8, containing 1 mM MgCl₂ and 0.1 mM ZnCl₂) was added to each well and incubated for 30 min at room temperature. The absorbance at 405 nm was measured using a microplate reader.

Avidity of anti-DNPh-KLH antibody to various kinds of nitroaromatic compounds

The avidity of anti-DNPh-KLH antibody to nitroaromatic compounds was investigated by indirect competitive ELISAs using the DNPh-gly-OVA conjugate and

DNPh-C8-OVA as a coating antigen-protein conjugate. The ELISA procedures were the same as described in the section of Indirect competitive ELISAs for 2,4-DNT. The IC₅₀ was defined as the concentration of added DNPh derivative that yields 50% inhibition compared with no inhibition (100%). Molar cross-reactivities were related to 2,4-DNT (100); namely, all molar cross-reactivities were determined in relation to the 2,4-DNT standard inhibition curve. The molar cross-reactivity was calculated using the IC₅₀ of each derivative according to the following equation (Weiler and Zenk, 1976):

$$CR = (IC_{50}^*/IC_{50}) \times 100$$

where CR is molar cross-reactivity [%], IC₅₀^{*} is the IC₅₀ of the DNT standard [M], and IC₅₀ is the IC₅₀ of derivatives or related compounds [M].

RESULTS AND DISCUSSION

Immunization efficacy

The results of the immunization of the rabbit with DNPh-KLH conjugate are shown in Fig. 1. The antisera titers of the immunized rabbit increased, especially after booster immunization. The whole blood of the rabbit was collected at day 63. The concentration of the anti-DNPh-KLH antibody after Protein G treatment was estimated to be about 5 mg mL⁻¹ of serum, standardized as human IgG.

Temperature dependence of avidity between anti-DNPh-KLH antibody and DNT

Temperature dependencies of the avidity between the anti-DNPh-KLH antibody and 2,4-DNT was investigated by using indirect competitive-ELISA. DNPh-OVA (0.01 μ g mL⁻¹) was immobilized on the surface of the wells, and an antibody concentration of 10 μ g mL⁻¹ was used. Competitive reactions among free 2,4-DNT, bound DNPh-OVA, and anti-DNPh-KLH antibody were performed for 1 h. The results are shown in Fig. 2. The sen-




Fig. 1. Time course of antigen-specific antibody concentration in sera from a rabbit immunized with DNPh-KLH conjugate. The data in the figure are shown using 16,000-fold diluted sera with PBS.




Fig. 2. Dependence on temperature of the avidity between anti-DNPh-KLH antibody and 2,4-DNT measured by indirect-ELISAs. The concentrations of DNPh-OVA (coating antigen-protein conjugate) and anti-DNPh-KLH antibody were $0.01 \mu\text{g mL}^{-1}$ and $10 \mu\text{g mL}^{-1}$, respectively.

sitivity, or detection limit, of these analyses was defined as the concentration of antigen at which the percentage bound was depressed to 85%. The concentrations for 15% inhibition were almost the same at different temperatures, but there was a tendency toward higher sensitivity at lower temperatures. At lower temperatures, the non-specific binding of the antibody will be depressed, but simultaneously, the signal intensities will be weakened. In this case, differences in sensitivity between the temperatures were hard to distinguish, probably because of the lower signal intensities and avidity. At higher temperatures, the sensitivity observed was lower, but even so, the antibody was usable up to 50°C .

Optimization of number of DNPh-groups introduced on coating antigen-protein conjugates

To elucidate the effect of DNPh-group on the coating antigen-protein conjugate on the sensitivity, six DNPh-OVA conjugates having DNPh-groups of 0.4–5.3 mol per mol of OVA were examined. The OVA concentrations of the DNPh-groups were: DNPh-OVA-1 (0.4 mol per mol OVA), DNPh-OVA-2 (1.3 mol per mol OVA), DNPh-OVA-3 (2.7 mol per mol OVA), DNPh-OVA-4 (3.0 mol per mol OVA), DNPh-OVA-5 (5.0 mol per mol OVA), and DNPh-OVA-6 (5.3 mol per mol OVA). The DNPh-groups introduced on OVA were increased depending on the increased concentration of 2,4-DNBS, and the number was leveled off at more than 5% of 2,4-DNBS (final concentration). The results of indirect competitive ELISA using these coating antigen-protein conjugates are shown in Fig. 3. DNPh-OVA-5 and -6, having DNPh-group concentrations higher than 5 mol per mol OVA, did not show the association of free 2,4-DNT with antibody (no inhibition, random data), and others (DNPh-OVA-1 to -4) showed a decreased signal, depending on the concentration of free 2,4-DNT. From Fig. 3,




Fig. 3. Profiles of indirect competitive ELISA for 2,4-DNT using various DNPh-groups on the coating antigen-protein conjugates. Symbols are as follows: \square , DNPh-OVA-1; \blacklozenge , DNPh-OVA-2; $+$, DNPh-OVA-3; \diamond , DNPh-OVA-4; \bullet , DNPh-OVA-5; \blacktriangle , DNPh-OVA-6.

the preferable coating antigen-protein conjugates were judged to be DNPh-OVA-1 to -3, which have DNPh-groups at concentrations between 0.4 and 2.7 mol per mol OVA.

Optimization of spacer length introduced on coating DNPh-Cx-OVA conjugates

The spacer length on the coating antigen-protein conjugates also affects the sensitivity of 2,4-DNT detection when using indirect-ELISA. DNPh-Cx-OVA conjugates with different length methylene chains (C2–C12) were synthesized. In this case, the number of DNPh-groups on the coating antigen-protein conjugates was restricted to 1–2 mol per mol OVA, judging from the previous results (Fig. 3). The results of indirect competitive ELISA are shown in Fig. 4. A decrease of signals due to the association between free 2,4-DNT and antibody was observed for all DNPh-Cx-OVA conjugates. The sensitivity evaluated as IC_{50} was improved with increasing spacer length up to C10. The preferable coating antigen-protein conjugates were judged to be DNPh-C8-OVA or DNPh-C10-OVA. The improvement of the sensitivity was due to the following reasons: 1) the antibody could bind to the DNPh-moiety without the interference of a bulky protein-moiety of the coating antigen-protein conjugate, and 2) the antibody could recognize free 2,4-DNT and the DNPh-moiety equally because of the relatively long spacer. In general, recognition of a competitor by an antibody is reduced in heterologous systems. A small concentration of analyte, thus, will be sufficient to shift the equilibrium conditions against formation of the antibody-analyte immune complex, thereby inhibiting association of the antibody to the competitor and leading to very sensitive assays (Marco *et al.*, 1995). On the other hand, Kim *et al.* (2003) reported that heterology in the spacer length of the coating antigen had




Fig. 4. Profiles of indirect competitive ELISA for 2,4-DNT using various lengths of spacers on the coating antigen-protein conjugates. Symbols are as follows: \square , DNPh-OVA; \blacklozenge , DNPh-C2-OVA; \times , DNPh-C4-OVA; \triangle , DNPh-C6-OVA; \blacksquare , DNPh-C8-OVA; \bullet , DNPh-C10-OVA; \circ , DNPh-C12-OVA.

no significant effect on the sensitivity of ELISA, but heterology in the spacer structure of the coating antigen produced a remarkable improvement in the sensitivity of ELISA. Although the spacer structure was not examined in our experiment, the spacer length of the coating antigen-protein conjugate also affected the sensitivity of

ELISA in our case.

Avidity of anti-DNPh antibody to various kinds of nitroaromatic compounds

The avidity of the raised antibody to nitroaromatic compounds was evaluated by indirect competitive ELISAs. Two coating antigen-protein conjugates were used in this experiment: DNPh-C8-OVA and DNPh-gly-OVA. Midpoints (IC_{50}) and molar cross-reactivities are listed in Table 1. On using DNPh-C8-OVA, 2,6-DNT and 4-amino-2,6-DNT showed cross-reactivity of less than 1% or no association, when the molar cross-reactivity of 2,4-DNT was set as 100%. These results show that the anti-DNPh-KLH antibody recognized the nitro group of the 4-position of the benzene ring. 2-Amino-4,6-DNT showed about 3% molar cross-reactivity due to the amino group at the 2-position. TNPh-derivatives showed about 4–5% molar cross-reactivity or no association except to TNPh-amine (TNA). The anti-DNPh-KLH antibody seemed to discriminate between dinitrophenyl- and trinitrophenyl-compounds. On the other hand, 2,4-DNPh-glycine and TNPh-amine (TNA) showed high molar cross-reactivity, especially to 2,4-DNPh-glycine (>900%). These results indicate that the anti-DNPh-KLH antibody recognizes these compounds as an antigen. These phenomena occur because the dinitrophenyl moiety of the immunogen (2,4-DNPh-KLH conjugate) binds to KLH *via* an NH group and the antibody also recognizes the NH moiety of the 1-position of the benzene ring. 1,3-DNB showed molar cross-reactivity of about 5% because of the meta position of nitro groups.

Table 1. Avidity of anti-DNPh-KLH antibody to the nitroaromatic compounds

Nitroaromatic compounds	M.W.	Coating antigen-protein DNPh-C8-OVA		Coating antigen-protein DNPh-gly-OVA	
		IC_{50} (M)	Molar cross-reactivity (%)	IC_{50} (M)	Molar cross-reactivity (%)
2, 4-dinitrotoluene (2, 4-DNT)	182	1.4×10^{-7}	100	4.9×10^{-7}	100
2, 6-dinitrotoluene (2, 6-DNT)	182	3.4×10^{-5}	0.4	5.5×10^{-5}	0.9
2-amino-4, 6-dinitrotoluene (2-ADNT)	197	5.1×10^{-6}	2.7	2.5×10^{-6}	19.5
4-amino-2, 6-dinitrotoluene (4-ADNT)	197	N.I.*	–	N.I.*	–
2, 4-dinitrophenylglycine (2,4-DNP-gly)	241	1.5×10^{-8}	933	2.9×10^{-8}	1700
1,3-dinitrobenzene (1,3-DNB)	168	3.0×10^{-6}	4.7	2.7×10^{-6}	18.1
Trinitrophenol (TNP)	227	3.1×10^{-6}	4.5	1.3×10^{-6}	37.7
2, 4, 6-trinitroaniline (TNA)	228	1.3×10^{-7}	107.7	1.3×10^{-7}	375
N-(2, 4, 6-trinitrophenyl)-hexylamine (TNP-ha)	312	N.I.*	–	2.9×10^{-5}	1.7
N-(2, 4, 6-trinitrophenyl)-ethylamine (TNP-ea)	284	3.2×10^{-6}	4.4	7.8×10^{-7}	63.3
N-(2, 4, 6-trinitrophenyl)-glycine (TNP-gly)	286	3.5×10^{-6}	4.0	7.0×10^{-7}	70.7
Trinitrotoluene (TNT)	227	N.I.*	–	1.7×10^{-6}	28.1

N.I.*; not inhibited. Concentration-dependent inhibition was not observed in the range of the analyte concentrations studied (1.0×10^{-10} – 1.0×10^{-5} g mL⁻¹).

As shown in Table 1, the molar cross-reactivities and IC_{50} s showed different profiles when DNPh-gly-OVA was used as a coating antigen-protein conjugate. The selectivity for 2,4-DNT of DNPh-C8-OVA was much better than that of DNPh-gly-OVA. Optimization of a coating antigen-protein conjugate is an important factor for measurement by indirect competitive ELISA.

Detection of 2,4-DNT by indirect competitive ELISA

Figure 5 shows the standard curves of inhibition by 2,4-DNT using the anti-DNPh-KLH antibody and DNPh-C8-OVA as a coating antigen-protein conjugate in indirect competitive ELISA. As shown in Fig. 5, 2,4-DNT was detected at the concentration of 1 ng mL^{-1} (more than 15% decrease). Dynamic ranges, defined by the analyte concentrations that inhibited maximum signals by 15% and 80%, comprised a concentration range between 1 ng mL^{-1} and $10 \mu\text{g mL}^{-1}$.




Fig. 5. Standard curve of 2,4-DNT in indirect competitive ELISA. The error bars show SD of five measurements. The concentrations of coating antigen-protein conjugate (DNPh-C8-OVA) and the antibody (anti-DNPh-KLH antibody) were $0.5 \mu\text{g mL}^{-1}$ and $0.1 \mu\text{g mL}^{-1}$, respectively.

Evaluation of affinity constant between free 2,4-DNT and anti-DNPh-KLH antibody

Determination of an affinity constant, K_A , or its reciprocal, the dissociation constant, K_D , is frequently useful in the study of antigen-antibody interactions. Evaluation of K_D , however, is a difficult process because of the different values obtained by the method used for analysis. A convenient method developed by Seligman is widely used for calculating the K_D -value using indirect competitive ELISAs (Seligman, 1994). The basic equation is:

$$a_0/f = K_D/(1-f) + i_0$$

Here, f is the square root of $(A_0 - A_i)/A_0$ (A_i and A_0 are the absorbance with and without a competing antigen, respectively), and a_0 and i_0 stand for the initial concentration of the competing antigen (free analyte) and the initial antibody concentration, respectively. We evaluated the K_D -value with the inhibition curve of 2,4-DNT

obtained in indirect competitive ELISA using Seligman's method. First, we evaluated the K_D from the data shown in Fig. 3, which used DNPh-OVA-1 to -4 as coating antigen-protein conjugates. As expected from the data shown in Fig. 3, the dissociation constants between free 2,4-DNT and antibody were different, depending on the number of DNPh-groups on the coating antigen-protein conjugates. The evaluated values were 0.4×10^{-5} – $1.4 \times 10^{-5} \text{ M}$ ($K_A = 0.7 \times 10^5$ – $2.5 \times 10^5 \text{ M}^{-1}$). The K_D value was also evaluated by the data from Fig. 4, which used DNPh-C8-OVA as a coating antigen-protein conjugate. The calculated value was $0.2 \times 10^{-5} \text{ M}$ ($K_A = 5.0 \times 10^5 \text{ M}^{-1}$). This means that the association of the antibody with the coating antigen-protein conjugate was reduced and the association of the antibody with free 2,4-DNT was increased, improving the sensitivity of detection.

CONCLUSIONS

A polyclonal antibody against 2,4-DNT was successfully raised in a rabbit. The antibody was applied to detect 2,4-DNT by using indirect competitive ELISA. The measuring temperature scarcely affected on the detection sensitivity. The optimum coating antigen-protein conjugate on the microtiter plate was dinitrophenyl-C8-ovalbumin (DNPh-C8-OVA). The association constant between free 2,4-DNT and anti-DNPh-KLH antibody was $5.0 \times 10^5 \text{ M}^{-1}$, when DNPh-C8-OVA was used as a coating antigen-protein conjugate. The immunoassay exhibited excellent sensitivity for the detection of 2,4-DNT in the concentration range of 1 ng mL^{-1} to $10 \mu\text{g mL}^{-1}$. The combination of the anti-DNPh-KLH antibody and DNPh-C8-OVA as a coating antigen-protein conjugate is useful for the detection of 2,4-DNT.

ACKNOWLEDGEMENTS

This work was partly supported by a Grand-in-Aid from JST (Japan Science and Technology Corporation).

REFERENCES

- Ahmad F., and D. J. Roberts 1995 Use of narrow-bore high-performance liquid chromatography-diode array detection for the analysis of intermediates of the biological degradation of 2,4,6-trinitrotoluene. *J. Chromatogr. A*, **693**: 167–175
- Banerjee H. N., M. Verma, L. H. Hou, M. Ashraf, and S. K. Dutta 1999 Cytotoxicity of TNT and its metabolites. *Yale J. Biol. Med.*, **72**: 1–4
- Borch T., and R. Gerlach 2004 Use of reversed-phase high-performance liquid chromatography-diode array detection for complete separation of 2,4,6-trinitrotoluene metabolites and EPA Method 8330 explosives: influence of temperature and ion-pair reagent. *J. Chromatogr. A*, **1022**: 83–94
- Goldman E. R., A. Hayhurst, B. M. Lingerfelt, B. L. Iverson, G. Georgiou, and G. P. Anderson 2003 2,4,6-Trinitrotoluene detection using recombinant antibodies. *J. Environ. Monit.*, **5**: 380–383
- Green T. M., P. T. Charles, and G. P. Anderson 2002 Detection of 2,4,6-trinitrotoluene in sea water using a reversed-displacement immunosensor. *Anal. Biochem.*, **310**: 36–41
- Halasz A., C. Groom, E. Zhou, L. Paquet, C. Beaulier, S. Deschamps, A. Corriveau, S. Thiboutot, G. Ampleman, C. Dubois, and J.

- Hawari 2002 Detection of explosives and their degradation products in soil environments. *J. Chromatogr. A*, **963**: 411–418
- Hilmi A., J. H. T. Luong, and A. L. Nguyen 1999 Determination of explosives in soil and ground water by liquid chromatography–amperometric detection. *J. Chromatogr. A*, **844**: 97–110
- Jenkins T. F., M. E. Walsh, P. H. Miyares, J. A. Kopczyuski, T. A. Ranney, V. George, J. C. Pennington, and T. E. Berry Jr., 2000 “Cold Regions Research and Engineering Laboratory, Technical Report ERDC TR-00-5, August 2000”
- Jenkins T. F., D. C. Leggett, P. H. Miyares, H. E. Walsh, T. T. Ranney, J. H. Cragin, and V. George 2001 Chemical signature of TNT-filled landmines. *Talanta*, **54**: 501–503
- Kennel S. J., L. J. Foote, M. Morris, A. A. Vass, and W. H. Griest 2000 Mutation analyses of series of TNT-related compounds using the CHO–hprt assay. *J. Appl. Toxicol.*, **20**: 441–448
- Kim Y. J., Y. A. Cho, H.-S. Lee, Y. T. Lee, S. T. Gee, and B. D. Hammock 2003 Immunoassay of organophosphorous pesticides and effect of heterology in hapten spacer arm length on immunoassay sensitivity. *Anal. Chim. Acta*, **475**: 85–96
- Marco M.-P., S. Gee, and B. D. Hammock 1995 Immunochemical techniques for environmental analysis II. Antibody production and immunoassay developments. *Trends Anal. Chem.*, **14**: 415–425
- Matsumoto K., A. Torimaru, S. Ishitobi, T. Sakai, H. Ishikawa, K. Toko, N. Miura, and T. Imato 2005 Preparation and characterization of a polyclonal antibody from rabbit for detection of trinitrotoluene by a surface plasmon resonance biosensor. *Talanta*, **68**: 305–311
- Sakai T., A. Torimaru, K. Shinahara, N. Miura, T. Imato, K. Toko, and K. Matsumoto 2003 Preparation of a Polyclonal Antibody and a Bioassay for Nitroaromatic Compounds by an Enzyme-Linked Immunosorbent Assay Technique and a Surface Plasmon Resonance Immunosensor. *Sens. Mater.*, **15**: 439–452
- Sayama M., M. Mori, M. Shoji, S. Uda, M. Kakikawa, T. Kondo, and K. Kodaira 1998 Mutagenicities of 2,4- and 2,6-dinitrotoluenes and their reduced products in *Salmonella typhimurium* nitroreductase- and O-acetyltransferase- overproducing Ames test strains. *Mutat. Res.*, **420**: 27–32
- Seligman S. T. 1994 Influence of solid-phase antigen in competition enzyme-linked immunosorbent assays (ELISAs) on calculated antigen–antibody dissociation constants. *J. Immunol. Methods*, **168**: 101–110
- Shankaran D. R., K. V. Gobi, K. Matsumoto, T. Imato, K. Toko, and N. Miura 2004 Highly sensitive surface plasmon resonance immunosensor for parts-per-trillion level detection of 2,4,6-trinitrophenol. *Sens. Act. B*, **100**: 450–454
- Shankaran D. R., K. V. Gobi, T. Sakai, K. Matsumoto, K. Toko, and N. Miura 2005a Surface plasmon resonance immunosensor for highly sensitive detection of 2,4,6-trinitrotoluene. *Biosens. Bioelectron.*, **20**: 1750–1756
- Shankaran D. R., K. V. Gobi, T. Sakai, K. Matsumoto, T. Imato, K. Toko, and N. Miura 2005b A novel surface plasmon resonance immunosensor for 2,4,6-trinitrotoluene (TNT) based on indirect competitive immunoreaction: a promising approach for on-site landmine detection. *IEEE Sens. J.*, **5**: 616–621
- Shankaran D. R., K. Matsumoto, K. Toko, and N. Miura 2006 Development and comparison of two immunoassays for the detection of 2,4,6-trinitrotoluene (TNT) based on surface plasmon resonance. *Sens. Act. B*, **114**: 71–79
- Tchounwou P. B., B. A. Wilson, A. B. Ishaque, and J. Schneider 2001 Transcriptional activation of stress genes and cytotoxicity in human liver carcinoma cells (HepG2) exposed to 2,4,6-trinitrotoluene, 2,4-dinitrotoluene, and 2,6-dinitrotoluene. *Environ. Toxicol.*, **16**: 209–216
- Weiler E. W., and M. H. Zenk 1976 Radioimmunoassay for the determination of digoxin and related compounds in *Digitalis lanata*. *Phytochem.*, **15**: 1537–1545
- Wilson R., C. Clavering, and A. Hutchinson 2003 Electrochemiluminescence enzyme immunoassay for TNT. *Analyst*, **128**: 480–485