Relevance of Totally Laparoscopic Gastrectomy for Patients with Advanced Gastric Cancer

Saeki, Hiroshi Department of Surgery and Science, Graduate School of Medical Sciences, Kyushu University

Oki, Eiji Department of Surgery and Science, Graduate School of Medical Sciences, Kyushu University

Tsuda, Yasuo Department of Surgery and Science, Graduate School of Medical Sciences, Kyushu University

Ando, Koji Department of Surgery and Science, Graduate School of Medical Sciences, Kyushu University

他

https://doi.org/10.15017/1430773

出版情報:福岡醫學雑誌. 104 (11), pp.405-412, 2013-11-25. 福岡医学会 バージョン: 権利関係:

405

Original Article

Relevance of Totally Laparoscopic Gastrectomy for Patients with Advanced Gastric Cancer

Hiroshi Saeki¹), Eiji Oki¹), Yasuo Tsuda¹), Koji Ando¹), Yukiharu Hiyoshi¹⁾³), Shuhei Itoh¹), Masaru Morita¹), Tetsuo Ikeda¹), Keishi Sugimachi²), Yo-ichi Yamashita¹), Toru Ikegami¹), Hideaki Uchiyama¹), Tomoharu Yoshizumi¹), Yuji Soejima¹), Hirofumi Kawanaka¹), Koshi Mimori²), Masayuki Watanabe³) and Yoshihiko Maehara¹)

¹⁾Department of Surgery and Science, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan

²⁾Department of Surgery, Kyushu University Beppu Hospital, Beppu, Japan.

³⁾ Department of Gastroenterological Surgery, Graduate School of Life Sciences, Kumamoto University, Kumamoto, Japan

Abstract

Purpose : Although the use of laparoscopic gastrectomy for gastric cancer has been widespread, it has remained controversial whether it can be applied for the patients with advanced gastric cancer. The aim of this study was to clarify the safety and usefulness of totally laparoscopic gastrectomy for patients with advanced gastric cancer.

Patients and Methods : Totally laparoscopic gastrectomy was applied for a total of 38 patients with pStage IB-III advanced gastric cancer at our institute. The surgical and long-term results were analyzed in those patients.

Results : Twenty-seven patents underwent distal gastrectomy and 11 patients underwent total gastrectomy. The mean number of dissected lymph nodes was 41 (range, 16–87). The mean length of the operation and amount of blood loss was 324 min and 123 ml, respectively. Two cases of postoperative bleeding were noted, while neither anastomosis-related complications nor in-hospital death was observed. The follow-up period after surgery was 8–72 months. Postoperative recurrence was observed in 6 patients (peritoneal dissemination : 3 patients, pleural dissemination : 1 patient, liver metastasis : 1 patient, ovarian metastasis : 1 patient). The overall survival rates at 1, 3 and 5 years were 94.7%, 76.3% and 76.3%, respectively.

Conclusion : Totally laparoscopic gastrectomy is safe and can lead to satisfactory long-term outcomes in cases of advanced gastric cancer. Prospective controlled studies are warranted to confirm our findings.

Key words : Laparoscopic gastrectomy · Advanced gastric cancer · Less invasive surgery · Complication · Prognosis

Introduction

Although the annual incidence of mortality from gastric cancer has been decreasing yearly worldwide, gastric cancer is still the second leading cause of cancer-related death¹⁾²⁾. Chemotherapy and molecular targeting therapy for gastric cancer have recently made progress, however radical gastrectomy with regional lymph node dissection remains the only potentially

Corresponding Author : Hiroshi Saeki, MD

Department of Surgery and Science, Graduate School of Medical Sciences, Kyushu University

Address : 3-1-1 Maidashi, Fukuoka 812-8582, Japan

Tel: + 81-92-642-5466 Fax: + 81-92-642-5482

E-mail : h-saeki@surg2. med. kyushu-u. ac. jp

curative treatment available for advanced gastric cancer.

Laparoscopic gastrectomy has undergone rapid development and gained popularity in the past 20 years since the first reported case³⁾. The surgical techniques and technologies associated with laparoscopic surgeries have markedly advanced^{4)~7)}. Compared to traditional open gastrectomy, previous studies have reported that laparoscopic gastrectomy contributes to a faster recovery, better cosmetic outcomes and a better postoperative quality of life^{8)~10)}.

Laparoscopic surgery is currently being regarded as a treatment of choice for early gastric cancer, while its application for advanced gastric cancer has not achieved universal acceptance because of its uncertain oncological safety, particularly from the standpoint of sufficient lymphadenectomy¹¹⁾¹². Until now, there have been insufficient long-term follow-up results regarding the oncological adequacy of laparoscopic surgery for advanced gastric cancer.

We herein report the surgical and long-term clinical results of laparoscopic gastrectomy for patients with advanced gastric cancer. The aim of this study was to clarify the safety and usefulness of this operation for the patients with advanced gastric cancer.

Patients and Methods

Patients

Three hundred and twenty-eight patients with gastric cancer were surgically treated in the Department of Surgery and Science, Graduate School of Medical Sciences, Kyushu University, from 2007 to 2012. One hundred and seventy-one of these patients (52%) underwent totally laparos-copic gastrectomy, defined as both the resection and anastomosis being performed intracorporeally with a laparoscopic technique⁵⁾⁷⁾¹³⁾¹⁴⁾. The subjects included this study were 38 patients who underwent totally laparoscopic gastrectomy with pathological Stage IB-IIIC (potentially resectable) advanced gastric cancer, after excluding 125

Stage IA and 8 Stage IV patients. We investigated the background clinical data, surgical results, postoperative course and long-term prognosis of these 38 patients.

Surgical procedures

The distal or total gastrectomy and lymphadenectomy were performed according to the guidelines of the Japanese Gastric Cancer Association¹⁵⁾. The patient was placed under general anesthesia in the lithotomy position. A balloon blunt tip trocar was inserted in the umbilical region using a cut-down method and a laparoscope was inserted through the trocar. The liver was pulled up using a silicone disc by a phi-shaped technique as previously reported⁶⁾. All procedures were performed using two trocars for the operator and two trocars for the assistant. For distal gastrectomy, reconstruction was performed by the Billroth I method as the first choice, or by the Roux-en Y method. Billroth I reconstruction was performed with a delta anastomosis¹⁶⁾ or a book binding technique⁴⁾ and Roux-en Y reconstruction was also performed using a linear stapler¹⁷⁾. The resected specimen was pulled through the extended wound at the umbilicus. For total gastrectomy, Roux-en Y reconstruction with a functional end-to-end anastomosis using a linear stapler was performed as the first choice or otherwise was performed with a transorally inserted anvil delivery system¹⁸⁾. The resected specimen was pulled through the extended wound at the umbilicus for a functional end-to-end anastomosis, while it was pulled through the left upper abdomen when using a transorally inserted anvil delivery system. All anastomoses were completed intracorporeally without extending the wound for the anastomotic procedure.

Staging of the tumor and statistical analysis

The staging of the tumor was based on the Japanese Classification of Gastric Carcinoma¹⁵⁾ and the depth of invasion and lymph node metastasis were defined by the pathological

findings. The survival curve was plotted according to the Kaplan-Meier method. The data were analyzed using the StatView software package (Abacus Concepts, Inc., Berkeley, California).

Results

The background data and surgical results of 38 patients who underwent totally laparoscopic gastrectomy for advanced gastric cancer are shown in Table 1. Among the 38 patients, 18 patients (47%) had pStage III disease. D2 or D1 + lymph node dissection was applied for all patients except for 1 patient who underwent D1 lymph node dissection because he had simultaneous

far-advanced lung cancer. Twenty-seven patients underwent distal gastrectomy and 11 patients received total gastrectomy. The mean number of dissected lymph nodes was 41 (range, 16-87). The mean length of the operation and amount of blood loss was 324 min (range, 200-760 min) and 123 ml (range, 4-400 ml), respectively.

The postoperative clinical course, morbidity and mortality after totally laparoscopic gastrectomy for advanced gastric cancer are shown in Table 2. The mean day of resuming oral intake was postoperative day 5 (POD5) (range, POD2-16). The mean hospital stay after the operation was 13 days (range, 8-28 days). Two

 Table 1
 The background data and surgical results

Tuble 1 The background data and burglear results	
Factor	Cases (n=38)
Male/Female	26/12
Mean age (range)	66 (35-95)
Pathological Stage	
IB	6 (16)
IIA	9 (24)
IIB	5 (13)
IIIA	8 (21)
IIIB	5 (13)
IIIC	5 (13)
Operation	
Distal gastrectomy	27 (71)
Total gastrectomy	11 (29)
Lymph node dissection	
D1	1 (3)
D1 +	10 (26)
D2	27 (71)
Mean number of dissected lymph nodes (range)	41 (16-87)
Mean length of operation, min (range)	324 (200-760)
Mean amount of blood loss, ml (range)	123 (4-400)

The data in parentheses are percentages unless stated otherwise.

Table 2 The postoperative clinical course, morbidity and mo	ortality
---	----------

Factor	Cases (n=38)
Mean day of resuming oral intake, postoperative day (range)	5 (2-16)
Mean hospital stay after operation, days (range)	13 (8-28)
Overall morbidity	3 (8)
Anastomotic leakage	0 (0)
Anastomotic stenosis	0 (0)
Pulmonary complication	0 (0)
Cardiac complication	0 (0)
Bleeding	2 (5)
Pancreatic fistula	1 (3)
Mortality	0 (0)

The values in parentheses are percentages unless stated otherwise.



Fig. 1 The survival curve of the 38 patients with advanced gastric cancer who underwent totally laparoscopic gastrectomy. The 1, 3 and 5-year overall survival rates were 94.7%, 76.3% and 76.3%, respectively.

cases of postoperative bleeding were noted, however, no anastomosis-related complications, other severe complications or in-hospital death was observed.

The follow-up period after surgery was 8-72 months. Postoperative recurrence was observed in 6 patients (pStage IIA : peritoneal dissemination in 1 patient, pStage IIIA : peritoneal dissemination in 1 patient, pStage IIIB : liver metastasis in 1 patient, ovarian metastasis in 1 patient, pStage IIIC : peritoneal dissemination in 1 patient, the overall survival rates at 1, 3 and 5 years were 94.7%, 76.3% and 76.3%, respectively.

Discussion

Laparoscopic gastrectomy for early gastric cancer has still been recognized as an investigational treatment in daily practice, not as a standard operation, since the benefits of the procedure have been evaluated only by limited comparative studies, and randomized control trials to assess its usefulness in early gastric cancer are still $\text{ongoing}^{19)-21}$. Needless to say, the indications for its application, particularly for advanced gastric cancer, have not yet been determined. However, some clinical studies, including the current study, have emphasized that

laparoscopic gastrectomy is technically feasible for advanced gastric cancer and can yield sufficient short and long-term oncological outcomes compared with conventional open gastrectomy¹²⁾²²⁾.

Although this was a retrospective study performed at a single center, the amount of blood loss seemed to be advantageous, while the operation tended to be longer compared with the conventional operation. A reduction of the blood loss is a consistent finding in previous studies comparing laparoscopic and open operations for advanced gastric cancer²³⁾. Regarding the length of the operation, laparoscopic gastrectomy is generally more time-consuming than open gastrectomy²³⁾. However, various modified laparoscopic techniques could help to simplify the procedure and shorten the operation $^{6)18)24)}$. We expect that with the increased proficiency in the laparoscopic technique and continuous improvements of the equipment, the time required for laparoscopic gastrectomy will become shorter.

The laparoscopic approach for gastric surgery is expected to improve the postoperative recovery, morbidity and mortality for patients in comparison to open surgery. Totally laparoscopic gastrectomies are associated with less surgical trauma, less pain, a more rapid return to gastrointestinal function, a shorter hospital stay and a decreased risk of impaired respiratory function $^{25)26)}$. In the current study, the results concerning the day of resuming oral intake and the hospital stay after the operation were considered to be moderate compared with other studies²³⁾. However, the results for these factors largely depend on institutional principles, as well as the patient's postoperative clinical course. It is suggested that these factors can be further improved by the application of a critical pathway for laparoscopic gastrectomy, which may help to reduce the postoperative hospital stay and hospitalization cost²⁷⁾. Based on a systematic review including the randomized and observational trials limited solely to advanced gastric

408

cancer²⁸⁾, the overall postoperative morbidity associated with laparoscopy-assisted gastrectomy was similar to that associated with open gastrectomy. With regard to the postoperative complications in the current study, no anastomosis-related complications, other severe complications or in-hospital death, was observed. The surgical safety of totally laparoscopic gastrectomy for advanced gastric cancer should be investigated in larger scale studies.

Cancer recurrence and the long-term prognosis are critical outcomes for evaluating surgical interventions in oncological therapy. The postoperative cancer recurrence and the long-term survival rate in laparoscopic gastrectomy for advanced gastric cancer were previously reported to be similar to those in open gastrectomy by a meta-analysis²³⁾. Regarding the recurrence pattern, the concerns about the dissemination of gastric cancer due to insufflated gas from the pneumoperitoneum and port site or wound metastasis has been emphasized. Although port-site recurrence was reported in some studies $^{29)30)}\!\!,$ there have also been reports of wound metastasis in open gastrectomy, thus suggesting that such metastasis was not a peculiar event to laparoscopic gastrectomy³⁰⁾. Laparoscopic gastrectomy has been reported to not increase the risk of perigastric lymph node recurrence compared to open gastrectomy³¹⁾. The results of this study did not demonstrate any peculiar recurrence pattern related to the laparoscopic gastrectomy. With regard to the long-term prognosis, the results of laparoscopic gastrectomy were also reported to be comparable to those of open gastrectomy $^{31)-33)}$. In this study, the number of dissected lymph nodes was not inferior to that of open gastrectomy reported in previous study³⁴⁾. This might have positively contributed to the satisfactory long-term survival results in this study.

In conclusion, totally laparoscopic gastrectomy is safe and can contribute to satisfactory long-term outcomes in cases of advanced gastric cancer. Well-designed prospective and controlled studies are thus warranted to validate our findings.

Acknowledgments

We thank Brian Quinn for assisting with the preparation of the manuscript.

References

- Bertuccio P, Chatenoud L, Levi F, Praud D, Ferlay J, Negri E, Malvezzi M and La Vecchia C : Recent patterns in gastric cancer : a global overview. Int J Cancer. 125 : 666–673, 2009.
- Yako-Suketomo H and Katanoda K : Comparison of time trends in stomach cancer mortality (1990-2006) in the world, from the WHO mortality database. Jpn J Clin Oncol. 39 : 622-623, 2009.
- Kitano S, Iso Y, Moriyama M and Sugimachi K : Laparoscopy-assisted Billroth I gastrectomy. Surg Laparosc Endosc. 4 : 146–148, 1994.
- 4) Ikeda T, Kawano H, Hisamatsu Y, Ando K, Saeki H, Oki E, Ohga T, Kakeji Y, Tsujitani S, Kohnoe S and Maehara Y : Progression from laparoscopic-assisted to totally laparoscopic distal gastrectomy : comparison of circular stapler (i-DST) and linear stapler (BBT) for intracorporeal anastomosis. Surg Endosc. 27 : 325-332, 2013.
- 5) Oki E, Sakaguchi Y, Ohgaki K, Saeki H, Chinen Y, Minami K, Sakamoto Y, Toh Y, Kusumoto T and Maehara Y : Feasibility of delta-shaped anastomoses in totally laparoscopic distal gastrectomy. Eur Surg Res. 47 : 205–210, 2011.
- 6) Saeki H, Oki E, Kawano H, Ando K, Ida S, Kimura Y, Morita M, Kusumoto T, Ikeda T and Maehara Y : Newly developed liver-retraction method for laparoscopic gastric surgery using a silicone disc : the phi-shaped technique. J Am Coll Surg. 216 : e43-46, 2013.
- 7) Sakaguchi Y, Ikeda O, Ohgaki K, Oki E, Chinen Y, Sakamoto Y, Minami K, Toh Y and Okamura T : Totally Laparoscopic Gastrectomy for Gastric Cancer Associated with Recklinghausen's Disease. Diagn Ther Endosc. 2010 : 682401, 2010.
- 8) Yasunaga H, Horiguchi H, Kuwabara K, Matsuda S, Fushimi K, Hashimoto H and Ayanian JZ : Outcomes after laparoscopic or open distal gastrectomy for early-stage gastric cancer : a propensity-matched analysis. Ann Surg. 257 : 640-646, 2013.

- 9) Eom BW, Kim YW, Lee SE, Ryu KW, Lee JH, Yoon HM, Cho SJ, Kook MC and Kim SJ : Survival and surgical outcomes after laparoscopy-assisted total gastrectomy for gastric cancer : case-control study. Surg Endosc. 26 : 3273-3281, 2012.
- 10) Kim YW, Baik YH, Yun YH, Nam BH, Kim DH, Choi IJ and Bae JM : Improved quality of life outcomes after laparoscopy-assisted distal gastrectomy for early gastric cancer : results of a prospective randomized clinical trial. Ann Surg. 248 : 721-727, 2008.
- 11) Miura S, Kodera Y, Fujiwara M, Ito S, Mochizuki Y, Yamamura Y, Hibi K, Ito K, Akiyama S and Nakao A : Laparoscopy-assisted distal gastrectomy with systemic lymph node dissection : a critical reappraisal from the viewpoint of lymph node retrieval. J Am Coll Surg. 198 : 933–938, 2004.
- 12) Bo T, Peiwu Y, Feng Q, Yongliang Z, Yan S, Yingxue H and Huaxing L : Laparoscopy-assisted vs. open total gastrectomy for advanced gastric cancer : long-term outcomes and technical aspects of a case-control study. J Gastrointest Surg. 17 : 1202–1208, 2013.
- 13) Oki E, Sakaguchi Y, Ohgaki K, Minami K, Yasuo S, Akimoto T, Toh Y, Kusumoto T, Okamura T and Maehara Y : Surgical complications and the risk factors of totally laparoscopic distal gastrectomy. Surg Laparosc Endosc Percutan Tech. 21 : 146–150, 2011.
- 14) Oki E, Sakaguchi Y, Ohgaki K, Saeki H, Chinen Y, Minami K, Sakamoto Y, Toh Y, Kusumoto T, Okamura T and Maehara Y : The impact of obesity on the use of a totally laparoscopic distal gastrectomy in patients with gastric cancer. J Gastric Cancer. 12 : 108–112, 2012.
- Association JGC, ed. Japanese Classification of Gastric Carcinoma. 13th ed. Tokyo : Kanehara ; 1999.
- 16) Kanaya S, Gomi T, Momoi H, Tamaki N, Isobe H, Katayama T, Wada Y and Ohtoshi M : Delta-shaped anastomosis in totally laparoscopic Billroth I gastrectomy : new technique of intraabdominal gastroduodenostomy. J Am Coll Surg. 195 : 284-287, 2002.
- 17) Takaori K, Nomura E, Mabuchi H, Lee SW, Agui T, Miyamoto Y, Iwamoto M, Watanabe H and Tanigawa N : A secure technique of intracorporeal Roux-Y reconstruction after laparoscopic distal gastrectomy. Am J Surg. 189 : 178–183, 2005.
- 18) Liao GQ, Ou XW, Liu SQ, Zhang SR and Huang

W : Laparoscopy-assisted total gastrectomy with trans-orally inserted anvil (OrVil) : a single institution experience. World J Gastroenterol. 19 : 755-760, 2013.

- 19) Kodera Y, Fujiwara M, Ohashi N, Nakayama G, Koike M, Morita S and Nakao A : Laparoscopic surgery for gastric cancer : a collective review with meta-analysis of randomized trials. J Am Coll Surg. 211 : 677-686, 2010.
- 20) Japanese gastric cancer treatment guidelines 2010 (ver. 3). Gastric Cancer. 14 : 113–123, 2011.
- 21) NCCN Guidelines Version 2.2011 Gastric Cancer. <u>http://www.nccn.org/professionals/physic-</u> ian_gls/f_guidelines.asp.
- 22) Lee JH, Son SY, Lee CM, Ahn SH, Park do J and Kim HH : Morbidity and mortality after laparoscopic gastrectomy for advanced gastric cancer : results of a phase II clinical trial. Surg Endosc. 27 : 2877–2885, 2013.
- 23) Chen K, Xu XW, Mou YP, Pan Y, Zhou YC, Zhang RC and Wu D : Systematic review and meta-analysis of laparoscopic and open gastrectomy for advanced gastric cancer. World J Surg Oncol. 11 : 182, 2013.
- 24) Kim HS, Kim BS, Lee IS, Lee S, Yook JH and Kim BS : Comparison of totally laparoscopic total gastrectomy and open total gastrectomy for gastric cancer. J Laparoendosc Adv Surg Tech A. 23 : 323–331, 2013.
- 25) Mingoli A, Sgarzini G, Binda B, Brachini G, Belardi V, Huscher CG, Di Paola M and Ponzano C : Totally laparoscopic approach for treatment of early and advanced gastric cancer. J Am Coll Surg. 204 : 187–188, 2007.
- 26) Ikeda O, Sakaguchi Y, Aoki Y, Harimoto N, Taomoto J, Masuda T, Ohga T, Adachi E, Toh Y, Okamura T and Baba H : Advantages of totally laparoscopic distal gastrectomy over laparoscopically assisted distal gastrectomy for gastric cancer. Surg Endosc. 23 : 2374–2379, 2009.
- 27) Kim JW, Kim WS, Cheong JH, Hyung WJ, Choi SH and Noh SH : Safety and efficacy of fast-track surgery in laparoscopic distal gastrectomy for gastric cancer : a randomized clinical trial. World J Surg. 36 : 2879–2887, 2012.
- 28) Ye LY, Liu DR, Li C, Li XW, Huang LN, Ye S, Zheng YX and Chen L : Systematic review of laparoscopy-assisted versus open gastrectomy for advanced gastric cancer. J Zhejiang Univ Sci B. 14 : 468–478, 2013.
- 29) Hwang SI, Kim HO, Yoo CH, Shin JH and Son BH : Laparoscopic-assisted distal gastrectomy versus open distal gastrectomy for advanced

gastric cancer. Surg Endosc. 23 : 1252–1258, 2009.

- 30) Zhao Y, Yu P, Hao Y, Qian F, Tang B, Shi Y, Luo H and Zhang Y : Comparison of outcomes for laparoscopically assisted and open radical distal gastrectomy with lymphadenectomy for advanced gastric cancer. Surg Endosc. 25 : 2960–2966, 2011.
- 31) Park do J, Han SU, Hyung WJ, Kim MC, Kim W, Ryu SY, Ryu SW, Song KY, Lee HJ, Cho GS and Kim HH : Long-term outcomes after laparoscopy-assisted gastrectomy for advanced gastric cancer : a large-scale multicenter retrospective study. Surg Endosc. 26 : 1548–1553, 2012.
- 32) Park CH, Song KY and Kim SN : Treatment results for gastric cancer surgery : 12 years' experience at a single institute in Korea. Eur J Surg Oncol. 34 : 36–41, 2008.

- 33) Sato H, Shimada M, Kurita N, Iwata T, Nishioka M, Morimoto S, Yoshikawa K, Miyatani T, Goto M, Kashihara H and Takasu C : Comparison of long-term prognosis of laparoscopy-assisted gastrectomy and conventional open gastrectomy with special reference to D2 lymph node dissection. Surg Endosc. 26 : 2240-2246, 2012.
- 34) Martinez-Ramos D, Miralles-Tena JM, Cuesta MA, Escrig-Sos J, Van der Peet D, Hoashi JS and Salvador-Sanchis JL : Laparoscopy versus open surgery for advanced and resectable gastric cancer : a meta-analysis. Rev Esp Enferm Dig. 103 : 133-141, 2011.

(Received for publication October 4, 2013)

(和文抄録)

進行胃癌に対する完全鏡視下胃切除術の妥当性

1)九州大学 消化器・総合外科 2)九州大学病院 別府病院外科 3)熊本大学 消化器外科

佐伯浩司¹⁾, 沖 英次¹⁾, 津田康雄¹⁾, 安藤幸滋¹⁾, 日吉幸晴¹⁾³⁾, 伊藤修平¹⁾, 森田 勝¹⁾, 池田哲夫¹⁾, 杉町圭史²⁾, 山下洋市¹⁾, 池上 徹¹⁾, 内山秀昭¹⁾, 吉住朋晴¹⁾, 副島雄二¹⁾, 川中博文¹⁾, 三森功士²⁾, 渡邊雅之³⁾, 前原喜彦¹⁾

【目的】胃癌に対する腹腔鏡下手術は広く普及したが,進行胃癌に対する適応の是非に関しては議 論の余地がある.本研究の目的は,進行胃癌に対する完全腹腔鏡下胃切除術の有用性を明らかにす ることである.

【対象・方法】当科において完全腹腔鏡下胃切除術が施行された IB 期-III 期進行胃癌症例 38 例の 手術成績と予後を検討した.

【結果】幽門側胃切除27例,胃全摘11例が施行された.郭清リンパ節個数は平均41個(16-87個) であった.平均出血量123ml,平均手術時間324分,術後合併症として出血を2例に認めたが,吻 合部関連合併症および在院死は認めなかった.観察期間は8-72ヶ月で,再発は6例に認めた(腹膜 播種3例,胸膜播種1例,卵巣転移1例,肝転移1例).全生存率は1年94.7%,3年・5年76.3% であった.

【結語】進行胃癌に対する腹腔鏡下胃切除術は安全に施行可能であり,予後も良好である.これらの結果を検証するための,前向き比較試験が必要である.