The Popular Condensation Problem under Matroid Constraints

Kamiyama, Naoyuki
Institute of Mathematics for Industry, Kyushu University

<http://hdl.handle.net/2324/1434328>

出版情報 : Combinatorial optimization and applications : 8th International Conference, COCOA 2014, Wailea, Maui, HI, USA, December 19-21, 2014, Proceedings, pp.713-728, 2014-11-13.
Springer International Publishing
バージョン :
権利関係 :

MI Preprint Series

**Mathematics for Industry
Kyushu University**

The Popular Condensation Problem under Matroid Constraints

Naoyuki Kamiyama

MI 2014-4

(Received March 4, 2014)

Institute of Mathematics for Industry
Graduate School of Mathematics
Kyushu University
Fukuoka, JAPAN

The Popular Condensation Problem under Matroid Constraints

Naoyuki Kamiyama*

Institute of Mathematics for Industry, Kyushu University
kamiyama@imi.kyushu-u.ac.jp

Abstract

The popular matching problem introduced by Abraham, Irving, Kavitha, and Mehlhorn is one of assignment problems in strategic situations. It is known that a given instance of this problem may admit no popular matching. For coping with such instances, Wu, Lin, Wang, and Chao introduced the popular condensation problem whose goal is to transform a given instance so that it has a popular matching by deleting a minimum number of agents. In this paper, we consider a matroid generalization of the popular condensation problem, and give a polynomial-time algorithm for this problem.

1 Introduction

In this paper, we consider a problem of assigning applicants to posts in strategic situations. Such problems occur, e.g., when a school assigns students to lectures or a firm assigns workers to tasks. For such strategic assignment problems, several solution concepts have been introduced. For example, Gärdenfors [5] introduced the concept of popularity. Intuitively speaking, if a matching M is popular, then there exists no other matching N such that more applicants prefer N to M than prefer M to N . Using the concept of popularity, Abraham, Irving, Kavitha, and Mehlhorn [1] introduced the popular matching problem, and presented a linear-time algorithm for this problem. Several extensions of the popular matching problem have been investigated. For example, Manlove and Sng [10] considered a many-to-one variant of the popular matching problem, Mestre [11] considered a weighted variant, and Sng and Manlove [14] considered a weighted many-to-one variant. Furthermore, Kamiyama [6] introduced a matroid generalization of the popular matching problem, and gave a polynomial-time algorithm for this problem. This matroid generalization can represent a many-to-one variant of the popular matching problem presented by Manlove and Sng [10] and the popular matching with laminar capacity constraints (see [6] for details).

Unfortunately, it is known [1] that a given instance of the popular matching problem may admit no popular matching. For coping with such instances, several alternative solutions were presented. For example, Kavitha and Nasre [7] considered the problem of copying several posts so that a given instance admits a popular matching. Furthermore, Kavitha, Nasre, and Nimbhorkar [8] considered the problem of augmenting several posts with minimum costs. These problems have been shown to be hard in general. Wu, Lin, Wang, and Chao [15] considered the problem of transforming the set of agents so that a given instance admits a popular matching. More precisely, they introduced the popular condensation problem whose goal is to transform a given instance so that it has a popular matching by deleting a minimum number of agents,

*This work is partly supported by KAKENHI(25730006).

and gave a polynomial-time algorithm for this problem. In this paper, we consider a matroid generalization of the popular condensation problem (i.e., the popular condensation problem in the matroid setting presented by Kamiyama [6]), and give a polynomial-time algorithm for this problem. Our algorithm can be regarded as a matroid generalization of the algorithm presented by Wu, Lin, Wang, and Chao [15].

The rest of this paper is organized as follows. In Section 2, we give the formal definition of our problem. In Section 3, we review a characterization of the existence of a popular matching under matroid constraints presented by Kamiyama [6]. In Section 4, we give our algorithm, and prove its correctness.

2 Preliminaries

We denote by $\mathbb{Z}_+$ the set of non-negative integers. For each subset X and each element x , we define $X + x := X \cup \{x\}$ and $X - x := X \setminus \{x\}$.

An ordered pair $\mathcal{M} = (U, \mathcal{I})$ is called a *matroid*, if U is a finite set and $\mathcal{I}$ is a nonempty family of subsets of U satisfying the following conditions.

(I1) If $I \in \mathcal{I}$ and $J \subseteq I$, then $J \in \mathcal{I}$.

(I2) If $I, J \in \mathcal{I}$ and $|I| < |J|$, then there exists an element u in $J \setminus I$ with $I + u \in \mathcal{I}$.

2.1 Problem formulation

Throughout this paper, we are given a finite simple bipartite graph $G = (V, E)$ in which V is partitioned into two subsets A and P , and each edge in E connects a vertex in A and a vertex in P . We call a vertex in A an *applicant*, and a vertex in P a *post*. We denote by (a, p) the edge in E between an applicant a in A and a post p in P . For each vertex v in V and each subset M of E , we define $M(v)$ as the set of edges in M incident to v . Furthermore, for each subset X of A and each subset M of E , we write $M(X)$ instead of $\cup_{a \in X} M(a)$.

In addition, we are given an injective function $\pi: E \rightarrow \mathbb{Z}_+$. That is, $\pi(e) \neq \pi(e')$ for every distinct edges e, e' in E . Intuitively speaking, π represents preference lists of applicants. For each applicant a in A and each edges e, e' in $E(a)$, if $\pi(e) > \pi(e')$, then a prefers e to e' . Since π is injective, it represents “strict” preference lists of applicants. Without loss of generality, we assume that for each applicant a in A , there exists a post p_a in P such that $E(p_a) = \{(a, p_a)\}$ and

$$\forall e \in E(a) - (a, p_a): \pi(e) > \pi((a, p_a)).$$

Furthermore, for each post p in P , we are given a matroid $\mathcal{M}_p = (E(p), \mathcal{I}_p)$. We assume that for each applicant a in A , $\{(a, p_a)\} \in \mathcal{I}_{p_a}$. Furthermore, we assume that for each applicant a in A , there exists a post p in $P - p_a$ such that $(a, p) \in E$ and $\{(a, p)\} \in \mathcal{I}_p$.

For each subset X of A , a subset M of E is called a *matching with respect to X* , if it satisfies the following two conditions.

- For every applicant a in A ,

$$|M(a)| = \begin{cases} 1 & \text{if } a \in X \\ 0 & \text{if } a \notin X. \end{cases}$$

- For every post p in P , we have $M(p) \in \mathcal{I}_p$.

For each subset X of A , each matching M with respect to X , and each applicant a in X , we denote by $\mu_M(a)$ the unique edge in $M(a)$. Let M, N be matchings with respect to some subset X of A . We denote by $\text{pre}_M(N)$ the number of applicants a in X with

$$\pi(\mu_N(a)) > \pi(\mu_M(a)),$$

i.e., $\text{pre}_M(N)$ represents the number of applicants that prefer N to M .

A matching M with respect to a subset X of A is called a *popular matching with respect to X* , if

$$\text{pre}_N(M) \geq \text{pre}_M(N)$$

for every matching N with respect to X . That is, there exists no other matching N with respect to X such that more applicants in A prefer N to M than prefer M to N . A subset X of A is called a *popular condensation*, if there exists a popular matching with respect to X . Notice that for every applicant a in A , $\{a\}$ is a popular condensation. The goal of the *popular condensation problem under matroid constraints* is to find a maximum-size popular condensation.

2.2 Matroids

In this subsection, we give properties of matroids that will be used in the sequel.

Let $\mathcal{M} = (U, \mathcal{I})$ be a matroid. A subset I in $\mathcal{I}$ is called an *independent set in $\mathcal{M}$* . For each subset X of U , a subset B of X is called a *base of X in $\mathcal{M}$* , if B is an inclusion-wise maximal subset of X that is an independent set in $\mathcal{M}$. We call a base of U in $\mathcal{M}$ a *base in $\mathcal{M}$* . It follows from the condition **(I2)** that for each subset X of U , every two bases of X in $\mathcal{M}$ have the same size, which is called the *rank of X in $\mathcal{M}$* and denoted by $r_{\mathcal{M}}(X)$. It is known [12, Lemma 1.3.1] that

$$\forall X, Y \subseteq U: r_{\mathcal{M}}(X) + r_{\mathcal{M}}(Y) \geq r_{\mathcal{M}}(X \cup Y) + r_{\mathcal{M}}(X \cap Y). \quad (1)$$

It follows from (1) that for every subsets X, Y, Z of U such that $X \subseteq Y$ and $Z \cap Y = \emptyset$,

$$r_{\mathcal{M}}(Y \cup Z) - r_{\mathcal{M}}(Y) \leq r_{\mathcal{M}}(X \cup Z) - r_{\mathcal{M}}(X). \quad (2)$$

Furthermore, it follows from (1) and the non-negativity of $r_{\mathcal{M}}(\cdot)$ that

$$\forall X, Y \subseteq U: r_{\mathcal{M}}(X \cup Y) \leq r_{\mathcal{M}}(X) + r_{\mathcal{M}}(Y). \quad (3)$$

Let S be a subset of U . Define

$$\begin{aligned} \mathcal{I}|S &:= \{X \subseteq S \mid X \in \mathcal{I}\}, \\ \mathcal{M}|S &:= (S, \mathcal{I}|S). \end{aligned}$$

It is not difficult to see that $\mathcal{M}|S$ is a matroid and $r_{\mathcal{M}|S}(X) = r_{\mathcal{M}}(X)$ for every subset X of S . Define a function $r': 2^{U \setminus S} \rightarrow \mathbb{Z}_+$ by

$$r'(X) := r_{\mathcal{M}}(X \cup S) - r_{\mathcal{M}}(S).$$

In addition, we define

$$\begin{aligned} \mathcal{I}/S &:= \{X \subseteq U \setminus S \mid r'(X) = |X|\}, \\ \mathcal{M}/S &:= (U \setminus S, \mathcal{I}/S). \end{aligned}$$

It is known [12, Proposition 3.1.6] that $\mathcal{M}/S$ is a matroid and $r_{\mathcal{M}/S}(X) = r'(X)$ for each subset X of $U \setminus S$.

Let $\mathcal{M}_1 = (U_1, \mathcal{I}_1)$ and $\mathcal{M}_2 = (U_2, \mathcal{I}_2)$ be matroids with $U_1 \cap U_2 = \emptyset$. Define

$$\begin{aligned}\mathcal{I}_1 \oplus \mathcal{I}_2 &:= \{X \subseteq U_1 \cup U_2 \mid X \cap U_1 \in \mathcal{I}_1, X \cap U_2 \in \mathcal{I}_2\}, \\ \mathcal{M}_1 \oplus \mathcal{M}_2 &:= (U_1 \cup U_2, \mathcal{I}_1 \oplus \mathcal{I}_2).\end{aligned}$$

It is known [12, Proposition 4.2.12] that $\mathcal{M}_1 \oplus \mathcal{M}_2$ is a matroid and

$$\forall X \subseteq U_1 \cup U_2: r_{\mathcal{M}_1 \oplus \mathcal{M}_2}(X) = r_{\mathcal{M}_1}(X \cap U_1) + r_{\mathcal{M}_2}(X \cap U_2). \quad (4)$$

Let $\mathcal{M}_1 = (U, \mathcal{I}_1)$ and $\mathcal{M}_2 = (U, \mathcal{I}_2)$ be matroids on the same ground set U . A subset I of U is called a *common independent set* of $\mathcal{M}_1$ and $\mathcal{M}_2$, if $I \in \mathcal{I}_1 \cap \mathcal{I}_2$. It is known [2, 9] that we can find a maximum-size common independent set of $\mathcal{M}_1$ and $\mathcal{M}_2$ in $O(n^3\gamma)$ time, where $n := |U|$ and γ is the time required to check whether $I - u + u'$ and $I + u'$ belong to $\mathcal{I}_1$ (or $\mathcal{I}_2$) for each independent set I in $\mathcal{M}_1$ (or $\mathcal{M}_2$) and each elements u in I and u' in $U \setminus I$ (see also [13, Section 41.2]). Furthermore, the following characterization of the size of a maximum-size common independent set is known.

Theorem 1 (Edmonds [3]). *For each matroids $\mathcal{M}_1 = (U, \mathcal{I}_1)$ and $\mathcal{M}_2 = (U, \mathcal{I}_2)$, the size of a maximum-size common independent set of $\mathcal{M}_1$ and $\mathcal{M}_2$ is equal to*

$$\min_{X \subseteq U} (r_{\mathcal{M}_1}(X) + r_{\mathcal{M}_2}(U \setminus X)).$$

The following lemmas will be used in the sequel.

Lemma 2 (see, e.g., [6]). *Let $\mathcal{M} = (U, \mathcal{I})$, S , and B be a matroid, a subset of U , and a base in $\mathcal{M}|S$, respectively. Then, for every subset X of $U \setminus S$, X is an independent set in $\mathcal{M}/S$ if and only if $X \cup B$ is an independent set in $\mathcal{M}$.*

Lemma 3. *Let $\mathcal{M} = (U, \mathcal{I})$ be a matroid. For each subsets X, Y, Z of U such that $X \subseteq Y$ and $Z \cap Y = \emptyset$, we have*

$$r_{\mathcal{M}/X}(Z) - r_{\mathcal{M}/Y}(Z) \leq r_{\mathcal{M}}(Y) - r_{\mathcal{M}}(X).$$

Proof. It follows from the definition of $\mathcal{M}/X$ and $\mathcal{M}/Y$ that

$$\begin{aligned}r_{\mathcal{M}/X}(Z) &= r_{\mathcal{M}}(Z \cup X) - r_{\mathcal{M}}(X), \\ r_{\mathcal{M}/Y}(Z) &= r_{\mathcal{M}}(Z \cup Y) - r_{\mathcal{M}}(Y).\end{aligned}$$

This lemma follows from this and the monotonicity of $r_{\mathcal{M}}(\cdot)$. □

3 Characterization

For each applicant a in A , we define the *f-edge* $f(a)$ of a as the unique element in

$$\arg \max \{ \pi((a, p)) \mid (a, p) \in E(a), \{(a, p)\} \in \mathcal{I}_p \}.$$

For each subset X of A and each post p in P , we denote by $\Gamma_{X,p}$ the set of edges (a, p) in $E(p)$ such that $a \in X$ and $(a, p) = f(a)$. For each subset X of A and each applicant a in X , we define the *s-edge* $s_X(a)$ of a as the unique edge in

$$\arg \max \{ \pi((a, p)) \mid (a, p) \in E(a) - f(a), \{(a, p)\} \in \mathcal{I}_p / \Gamma_{X,p} \}.$$

Notice that $s_X(a)$ is well-defined because there exists the post p_a . For each subset X of A , we define the *reduced edge set* Π_X by

$$\Pi_X := \{f(a), s_X(a) \mid a \in X\}.$$

For each subset X of A , we define a matroid $\mathcal{A}_X = (\Pi_X, \mathcal{I}_X)$ by

$$\mathcal{I}_X := \{M \subseteq \Pi_X \mid \forall a \in X: |M(a)| \leq 1\}.$$

For each subset X of A and each post p in P , we define

$$\mathcal{P}_{X,p} := (\mathcal{M}_p|_{\Gamma_{X,p}} \oplus \mathcal{M}_p/\Gamma_{X,p})|_{\Pi_X(p)}.$$

Furthermore, for each subset X of A , we define

$$\mathcal{P}_X := \bigoplus_{p \in P} \mathcal{P}_{X,p}.$$

For simplicity, we define $s(\cdot) := s_A(\cdot)$, $\Gamma_p := \Gamma_{A,p}$, $\Pi := \Pi_A$, $\mathcal{A} := \mathcal{A}_A$, and $\mathcal{P} := \mathcal{P}_A$.

The following characterization of a popular condensation plays an important role. Precisely speaking, Kamiyama [6] proved Theorem 4 in the case of $X = A$, but Theorem 4 for a general subset X of A can be proved in the same way.

Theorem 4 (Kamiyama [6]). *For each subset X of A , X is a popular condensation if and only if there exists a common independent set M of $\mathcal{A}_X$ and $\mathcal{P}_X$ with $|M| = |X|$.*

The following lemmas will be used in the sequel.

Lemma 5. *There exists a subset D of A such that*

$$|D| + r_{\mathcal{P}}(\Pi(A \setminus D)) = \min_{F \subseteq \Pi} (r_{\mathcal{A}}(F) + r_{\mathcal{P}}(\Pi \setminus F)). \quad (5)$$

Proof. Let F be a minimizer of the right-hand side of (5). If $\Pi(X) = F$ for some subset X of A , then the proof is done. Let X be the set of applicants a in A such that there exists an edge e in F with $e \in \Pi(a)$, and assume that there exists an edge e' in $\Pi(X)$ with $e' \notin F$. Clearly, we have

$$\begin{aligned} r_{\mathcal{A}}(F + e') &= r_{\mathcal{A}}(F) \quad (= |X|), \\ r_{\mathcal{P}}(\Pi \setminus (F + e')) &\leq r_{\mathcal{P}}(\Pi \setminus F). \end{aligned}$$

This implies that there exists a subset D of A satisfying (5). □

Lemma 6. *For each subset X of A , if there exists a common independent set M of $\mathcal{A}_X$ and $\mathcal{P}_X$ with $|M| = |X|$, then*

$$\forall Y \subseteq X: |Y| \leq r_{\mathcal{P}_X}(\Pi_X(Y)).$$

Proof. It follows from Theorem 1 that

$$\forall F \subseteq \Pi_X: |X| \leq r_{\mathcal{A}_X}(F) + r_{\mathcal{P}_X}(\Pi_X \setminus F). \quad (6)$$

Let Y be a subset of X . It follows from (6) with $F = \Pi_X(X \setminus Y)$ that

$$\begin{aligned} |X| &\leq r_{\mathcal{A}_X}(\Pi_X(X \setminus Y)) + r_{\mathcal{P}_X}(\Pi_X \setminus \Pi_X(X \setminus Y)) \\ &= |X \setminus Y| + r_{\mathcal{P}_X}(\Pi_X(Y)), \end{aligned}$$

which completes the proof. □

4 Algorithm

Our algorithm **PCuMC** for the popular condensation problem under matroid constraints can be described as follows.

Algorithm PCuMC

Step 1. Find a maximum-size common independent set M of $\mathcal{A}$ and $\mathcal{P}$.

Step 2. Output the set Δ of applicants a in A with $M(a) \neq \emptyset$.

End of Algorithm

From now on, we prove the correctness of the algorithm **PCuMC**. Let M be the maximum-size common independent set found in **Step 1** of the algorithm **PCuMC**, and we denote by Δ the output of the algorithm of **PCuMC**.

We first prove that Δ is a popular condensation.

Lemma 7. $\Pi_\Delta = \Pi(\Delta)$.

Proof. It suffices to prove that for every applicant a in Δ , we have $s_\Delta(a) = s(a)$. For proving this, we prove that for every post p , there exists a base B in $\mathcal{M}_p|\Gamma_p$ with $B \subseteq \Gamma_{\Delta,p}$. If there exists such a base B , then B is also a base in $\mathcal{M}_p|\Gamma_{\Delta,p}$. Thus, the above statement follows from Lemma 2.

Let p be a post in P , and assume that any base in $\mathcal{M}_p|\Gamma_p$ is not a subset of $\Gamma_{\Delta,p}$. Since $M \cap \Gamma_p$ is an independent set in $\mathcal{M}_p|\Gamma_p$, there exists a base B in $\mathcal{M}_p|\Gamma_p$ with $M \cap \Gamma_p \subseteq B$. The above assumption implies that there exists an edge $(a, p) \in B$ and $a \notin \Delta$. It follows from the definition of **Step 2** that $M(a) = \emptyset$, which implies that $M + (a, p)$ is an independent set of $\mathcal{A}$. Furthermore, it follows from the condition **(I1)** that $(M \cap \Gamma_p) + (a, p)$ is an independent set in $\mathcal{M}_p|\Gamma_p$, and thus $M + (a, p)$ is an independent set in $\mathcal{P}$. Since $(a, p) \notin M$, these facts contradict the fact that M is a maximum-size common independent set of $\mathcal{A}$ and $\mathcal{P}$. This completes the proof. $\square$

Lemma 8. Δ is a popular condensation.

Proof. It follows from Theorem 4 that if M is a common independent set of $\mathcal{A}_\Delta$ and $\mathcal{P}_\Delta$, then the proof is done. It follows from Lemma 7 that M is a subset of Π_Δ . Furthermore, M is clearly an independent set of $\mathcal{A}_\Delta$. What remains is to prove that M is an independent set of $\mathcal{P}_\Delta$.

Let p be a post of P . For proving that $M(p)$ is an independent set in $\mathcal{P}_{\Delta,p}$, we first prove that $M \cap \Gamma_{\Delta,p}$ is an independent set in $\mathcal{M}_p|\Gamma_{\Delta,p}$. It follows from the definition of **Step 2** that

$$M \cap \Gamma_{\Delta,p} = M \cap \Gamma_p. \quad (7)$$

Furthermore, $M \cap \Gamma_p$ is an independent set in $\mathcal{M}_p$. These facts implies that $M \cap \Gamma_{\Delta,p}$ is an independent set in $\mathcal{M}_p|\Gamma_{\Delta,p}$.

Next we prove that $M(p) \setminus \Gamma_{\Delta,p}$ is an independent set in $\mathcal{M}_p/\Gamma_{\Delta,p}$. It follows from (7) that $M(p) \setminus \Gamma_{\Delta,p} = M(p) \setminus \Gamma_p$. Moreover, in the same way as in the proof of Lemma 7, we can prove that there exists a base B in $\mathcal{M}_p|\Gamma_p$ with $B \subseteq \Gamma_{\Delta,p}$. Since $\Gamma_{\Delta,p}$ is a subset of Γ_p , B is also a base in $\mathcal{M}_p|\Gamma_{\Delta,p}$. Thus, since $M(p) \setminus \Gamma_p$ is an independent set in $\mathcal{M}_p/\Gamma_p$, it follows from these facts and Lemma 2 that $M(p) \setminus \Gamma_{\Delta,p}$ is an independent set in $\mathcal{M}_p/\Gamma_{\Delta,p}$. $\square$

Next we prove that Δ is a maximum-size popular condensation. Let D be a subset of A satisfying (5) in Lemma 5, and define $Q := A \setminus D$.

Lemma 9. $|A \setminus \Delta| = |Q| - r_{\mathcal{P}}(\Pi(Q))$.

Proof. It follows from $|\Delta| = |M|$ and Theorem 1 that

$$|\Delta| = |M| = |D| + r_{\mathcal{P}}(\Pi(A \setminus D)) = |A| - |Q| + r_{\mathcal{P}}(\Pi(Q)),$$

which completes the proof. $\square$

Lemma 10. For every popular condensation Ω , we have $|A \setminus \Omega| \geq |Q| - r_{\mathcal{P}}(\Pi(Q))$.

Proof. Define $\Omega_0 := A \setminus \Omega$, $\Omega_1 := Q \cap \Omega_0$, and $\Omega_2 := \Omega_0 \setminus \Omega_1$. Notice that $Q \setminus \Omega_1$ is a subset of Ω . Thus, it follows from Theorem 4 and Lemma 6 that

$$|Q \setminus \Omega_1| \leq r_{\mathcal{P}_\Omega}(\Pi_\Omega(Q \setminus \Omega_1)).$$

It follows from this that if

$$r_{\mathcal{P}_\Omega}(\Pi_\Omega(Q \setminus \Omega_1)) \leq r_{\mathcal{P}}(\Pi(Q)) + |\Omega_2|, \quad (8)$$

then the proof is done because $|A \setminus \Omega| = |\Omega_1| + |\Omega_2|$.

Let p be a post in P . Define F_1 as the set of edges (a, p) in Γ_p such that $a \in Q$ or $a \in \Omega_2$. In addition, define F_2 as the set of edges (a, p) in $\Pi(p)$ such that $a \in Q$ and $(a, p) = s(a)$. Notice that

$$F_1 \cup F_2 = (\Pi(Q) \cap \Pi(p)) \cup \{(a, p) \in \Gamma_p \mid a \in \Omega_2\}.$$

It follows from this and (3) that

$$\begin{aligned} r_{\mathcal{P}_p}(F_1 \cup F_2) &= r_{\mathcal{P}_p}((\Pi(Q) \cap \Pi(p)) \cup \{(a, p) \in \Gamma_p \mid a \in \Omega_2\}) \\ &\leq r_{\mathcal{P}_p}(\Pi(Q) \cap \Pi(p)) + r_{\mathcal{P}_p}(\{(a, p) \in \Gamma_p \mid a \in \Omega_2\}) \\ &\leq r_{\mathcal{P}_p}(\Pi(Q) \cap \Pi(p)) + |\{(a, p) \in \Gamma_p \mid a \in \Omega_2\}| \\ &= r_{\mathcal{P}_p}(\Pi(Q) \cap \Pi(p)) + |\{a \in \Omega_2 \mid f(a) \in \Pi(p)\}|. \end{aligned} \quad (9)$$

Define F'_1 as the set of edges (a, p) in $\Gamma_{\Omega, p}$ with $a \in Q \setminus \Omega_1$. In addition, define F'_2 as the set of edges (a, p) in $\Pi_\Omega(p)$ such that $a \in Q \setminus \Omega_1$ and $(a, p) = s_\Omega(a)$. Notice that

$$F'_1 \cup F'_2 = \Pi_\Omega(Q \setminus \Omega_1) \cap \Pi_\Omega(p). \quad (10)$$

It follows from (4), (9), and (10) that if

$$r_{\mathcal{P}_{\Omega, p}}(F'_1 \cup F'_2) \leq r_{\mathcal{P}_p}(F_1 \cup F_2), \quad (11)$$

then (8) follows and the proof is done because

$$\begin{aligned} r_{\mathcal{P}_\Omega}(\Pi_\Omega(Q \setminus \Omega_1)) &= \sum_{p \in P} r_{\mathcal{P}_{\Omega, p}}(\Pi_\Omega(Q \setminus \Omega_1) \cap \Pi_\Omega(p)) \\ &= \sum_{p \in P} r_{\mathcal{P}_{\Omega, p}}(F'_1 \cup F'_2) \\ &\leq \sum_{p \in P} r_{\mathcal{P}_p}(F_1 \cup F_2) \\ &\leq \sum_{p \in P} r_{\mathcal{P}_p}(\Pi(Q) \cap \Pi(p)) + \sum_{p \in P} |\{a \in \Omega_2 \mid f(a) \in \Pi(p)\}| \\ &= r_{\mathcal{P}}(\Pi(Q)) + |\Omega_2|. \end{aligned}$$

It follows from (4) that

$$\begin{aligned} r_{\mathcal{P}_p}(F_1 \cup F_2) &= r_{\mathcal{M}_p}(F_1) + r_{\mathcal{M}_p/\Gamma_p}(F_2), \\ r_{\mathcal{P}_{\Omega,p}}(F'_1 \cup F'_2) &= r_{\mathcal{M}_p}(F'_1) + r_{\mathcal{M}_p/\Gamma_{\Omega,p}}(F'_2). \end{aligned} \quad (12)$$

For every edge (a, p) in $F'_2 \setminus F_2$, we have

$$\pi(f(a)) > \pi((a, p)) > \pi(s(a)).$$

This implies that $(a, p) \notin \mathcal{I}_p/\Gamma_p$ for every edge (a, p) in $F'_2 \setminus F_2$. Thus, we have

$$r_{\mathcal{M}_p/\Gamma_p}(F_2) = r_{\mathcal{M}_p/\Gamma_p}(F_2 \cup F'_2). \quad (13)$$

Furthermore, it follows from the monotonicity of $r_{\mathcal{M}_p/\Gamma_{\Omega,p}}(\cdot)$ that

$$r_{\mathcal{M}_p/\Gamma_{\Omega,p}}(F'_2) \leq r_{\mathcal{M}_p/\Gamma_{\Omega,p}}(F_2 \cup F'_2). \quad (14)$$

It follows from $\Gamma_{\Omega,p} \subseteq \Gamma_p$ and Lemma 3 that

$$r_{\mathcal{M}_p/\Gamma_{\Omega,p}}(F_2 \cup F'_2) - r_{\mathcal{M}_p/\Gamma_p}(F_2 \cup F'_2) \leq r_{\mathcal{M}_p}(\Gamma_p) - r_{\mathcal{M}_p}(\Gamma_{\Omega,p}). \quad (15)$$

Since $\Gamma_p \setminus \Gamma_{\Omega,p} = F_1 \setminus F'_1$ and $F'_1 \subseteq \Gamma_{\Omega,p}$, it follows from (2) that

$$r_{\mathcal{M}_p}(\Gamma_p) - r_{\mathcal{M}_p}(\Gamma_{\Omega,p}) \leq r_{\mathcal{M}_p}(F_1) - r_{\mathcal{M}_p}(F'_1). \quad (16)$$

It follows from (13), (14), (15), and (16) that

$$\begin{aligned} r_{\mathcal{M}_p/\Gamma_{\Omega,p}}(F'_2) - r_{\mathcal{M}_p/\Gamma_p}(F_2) &\leq r_{\mathcal{M}_p/\Gamma_{\Omega,p}}(F_2 \cup F'_2) - r_{\mathcal{M}_p/\Gamma_p}(F_2 \cup F'_2) \\ &\leq r_{\mathcal{M}_p}(\Gamma_p) - r_{\mathcal{M}_p}(\Gamma_{\Omega,p}) \\ &\leq r_{\mathcal{M}_p}(F_1) - r_{\mathcal{M}_p}(F'_1). \end{aligned} \quad (17)$$

It follows from (12) and (17) that

$$\begin{aligned} r_{\mathcal{P}_{\Omega,p}}(F'_1 \cup F'_2) - r_{\mathcal{P}_p}(F_1 \cup F_2) &= r_{\mathcal{M}_p}(F'_1) - r_{\mathcal{M}_p}(F_1) + r_{\mathcal{M}_p/\Gamma_{\Omega,p}}(F'_2) - r_{\mathcal{M}_p/\Gamma_p}(F_2) \\ &\leq r_{\mathcal{M}_p}(F'_1) - r_{\mathcal{M}_p}(F_1) + r_{\mathcal{M}_p}(F_1) - r_{\mathcal{M}_p}(F'_1) \\ &= 0. \end{aligned}$$

This implies (11), which completes the proof. $\square$

Theorem 11. *The algorithm **PCuMC** correctly solves the popular condensation problem under matroid constraints.*

Proof. This theorem follows from Lemmas 8, 9 and 10. $\square$

Here we analyze the time complexity of the algorithm **PMuMC**. Define $m := |E|$, and we assume that for each post p in P , each independent set I in $\mathcal{M}_p$, and each edges e in I and e' in $E(p) \setminus I$, we can check in $O(\gamma)$ time whether $I - e + e'$ and $I + e'$ belong to $\mathcal{I}_p$. It follows from Lemma 2 that once we find a base in $\mathcal{M}_p/\Gamma_p$, for each independent set I in $\mathcal{M}_p/\Gamma_p$ and each edges e in I and e' in $E(p) \setminus (I \cup \Gamma_p)$, we can check in $O(\gamma)$ time whether $I - e + e'$ and $I + e'$ belong to $\mathcal{I}_p/\Gamma_p$. Thus, for each independent set I in $\mathcal{P}$ and each edges e in I and e' in $E \setminus I$, we can check in $O(\gamma)$ time whether $I - e + e'$ and $I + e'$ are independent sets in $\mathcal{P}$. This implies that the time complexity of the algorithm **PCuMC** is $O(m^3\gamma)$.

Next we consider a weighted variant of the popular condensation problem under matroid constraints. More precisely, in this problem, we are given a weight function $w: A \rightarrow \mathbb{Z}_+$. The goal is to find a maximum-size popular condensation X maximizing $\sum_{a \in X} w(a)$. This problem can be solved as follows. For each edge (a, p) in Π , we define the weight of (a, p) as $w(a)$. This weighted variant can be solved by finding a maximum-size common independent set of $\mathcal{A}$ and $\mathcal{P}$ with maximum-weight at **Step 1** of the algorithm **PCuMC**. It is known [4] that this problem can be solved in $O(m^3\gamma)$ time (see also [13, Section 41.3]). Thus, this weighted variant can be solved in $O(m^3\gamma)$ time.

References

- [1] D. J. Abraham, R. W. Irving, T. Kavitha, and K. Mehlhorn. Popular matchings. *SIAM Journal on Computing*, 37(4):1030–1045, 2007.
- [2] M. Aigner and T. A. Dowling. Matching theory for combinatorial geometries. *Transactions of the American Mathematical Society*, 158(1):231–245, 1971.
- [3] J. Edmonds. Submodular functions, matroids, and certain polyhedra. In R. Guy, H. Hanani, N. Sauer, and J. Schönheim, editors, *Combinatorial Structures and Their Applications*, pages 69–87. Gordon and Breach, 1970.
- [4] A. Frank. A weighted matroid intersection algorithm. *Journal of Algorithms*, 2(4):328–336, 1981.
- [5] P. Gärdenfors. Match making: Assignments based on bilateral preferences. *Behavioral Science*, 20(3):166–173, 1975.
- [6] N. Kamiyama. Popular matchings under matroid constraints. MI Preprint Series 2014-1, Kyushu University.
- [7] T. Kavitha and M. Nasre. Popular matchings with variable item copies. *Theoretical Computer Science*, 412(12):1263–1274, 2011.
- [8] T. Kavitha, M. Nasre, and P. Nimbhorkar. Popularity at minimum cost. In *Proceedings of the 21st International Symposium on Algorithms and Computation*, volume 6506 of *Lecture Notes in Computer Science*, pages 145–156, 2010.
- [9] E. L. Lawler. Matroid intersection algorithms. *Mathematical Programming*, 9(1):31–56, 1975.
- [10] D. F. Manlove and C. T. S. Sng. Popular matchings in the capacitated house allocation problem. In *Proceedings of the 14th Annual European Symposium on Algorithms*, volume 4168 of *Lecture Notes in Computer Science*, pages 492–503, 2006.
- [11] J. Mestre. Weighted popular matchings. In *Proceedings of the 33rd International Colloquium on Automata, Languages and Programming*, volume 4051 of *Lecture Notes in Computer Science*, pages 715–726, 2006.
- [12] J. G. Oxley. *Matroid theory*. Oxford University Press, 2nd edition, 2011.
- [13] A. Schrijver. *Combinatorial Optimization: Polyhedra and Efficiency*. Springer, 2003.

- [14] C. T. S. Sng and D. F. Manlove. Popular matchings in the weighted capacitated house allocation problem. *Journal of Discrete Algorithms*, 8(2):102–116, 2010.
- [15] Y.-W. Wu, W.-Y. Lin, H.-L. Wang, and K.-M. Chao. An optimal algorithm for the popular condensation problem. In *Proceedings of the 24th International Workshop on Combinatorial Algorithms*, volume 8288 of *Lecture Notes in Computer Science*, pages 412–422, 2013.

List of MI Preprint Series, Kyushu University

The Global COE Program

Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p-adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials
- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials

- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings
- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric τ -functions of the q-Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications

- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain
- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection
- MI2009-26 Manabu YOSHIDA
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance

- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA
Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI
Finite element computation for scattering problems of micro-hologram using DtN map
- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO
Abstract collision systems on groups

- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA
An algebraic approach to underdetermined experiments
- MI2010-10 Kei HIROSE & Sadanori KONISHI
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE
Lagrangian approach to weakly nonlinear stability of an elliptical flow
- MI2010-21 Hiroki MASUDA
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiro FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra

- MI2010-25 Toshimitsu TAKAESU
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI
Spin-spin correlation functions of the q -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and spike detection via Gaussian basis expansions
- MI2010-32 Nobutaka NAKAZONO
Hypergeometric τ functions of the q -Painlevé systems of type $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA
CAP representations of inner forms of $Sp(4)$ with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER
Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide FUKUMOTO & Alexander B. SAMOKHIN
Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI
Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI
Anderson-Darling test and the Malliavin calculus
- MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling via Compressed Sensing

- MI2011-5 Hiroshi INOUE
Implications in Compressed Sensing and the Restricted Isometry Property
- MI2011-6 Daeju KIM & Sadanori KONISHI
Predictive information criterion for nonlinear regression model based on basis expansion methods
- MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI
Group variable selection via relevance vector machine
- MI2011-8 Jan BREZINA & Yoshiyuki KAGEI
Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow
Group variable selection via relevance vector machine
- MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC
Recursive structures in the multispecies TASEP
- MI2011-10 Kazunori YASUTAKE
On projective space bundle with nef normalized tautological line bundle
- MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA
An explicit formula for the discrete power function associated with circle patterns of Schramm type
- MI2011-12 Yoshiyuki KAGEI
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow
- MI2011-13 Vladimír CHALUPECKÝ & Adrian MUNTEAN
Semi-discrete finite difference multiscale scheme for a concrete corrosion model: approximation estimates and convergence
- MI2011-14 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves
- MI2011-15 Hiroshi INOUE
A generalization of restricted isometry property and applications to compressed sensing
- MI2011-16 Yu KAWAKAMI
A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space
- MI2011-17 Naoyuki KAMIYAMA
Matroid intersection with priority constraints
- MI2012-1 Kazufumi KIMOTO & Masato WAKAYAMA
Spectrum of non-commutative harmonic oscillators and residual modular forms
- MI2012-2 Hiroki MASUDA
Mighty convergence of the Gaussian quasi-likelihood random fields for ergodic Levy driven SDE observed at high frequency

- MI2012-3 Hiroshi INOUE
A Weak RIP of theory of compressed sensing and LASSO
- MI2012-4 Yasuhide FUKUMOTO & Youich MIE
Hamiltonian bifurcation theory for a rotating flow subject to elliptic straining field
- MI2012-5 Yu KAWAKAMI
On the maximal number of exceptional values of Gauss maps for various classes of surfaces
- MI2012-6 Marcio GAMEIRO, Yasuaki HIRAOKA, Shunsuke IZUMI, Miroslav KRAMAR, Konstantin MISCHAIKOW & Vidit NANDA
Topological Measurement of Protein Compressibility via Persistence Diagrams
- MI2012-7 Nobutaka NAKAZONO & Seiji NISHIOKA
Solutions to a q -analog of Painlevé III equation of type $D_7^{(1)}$
- MI2012-8 Naoyuki KAMIYAMA
A new approach to the Pareto stable matching problem
- MI2012-9 Jan BREZINA & Yoshiyuki KAGEI
Spectral properties of the linearized compressible Navier-Stokes equation around time-periodic parallel flow
- MI2012-10 Jan BREZINA
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a time-periodic parallel flow
- MI2012-11 Daeju KIM, Shuichi KAWANO & Yoshiyuki NINOMIYA
Adaptive basis expansion via the extended fused lasso
- MI2012-12 Masato WAKAYAMA
On simplicity of the lowest eigenvalue of non-commutative harmonic oscillators
- MI2012-13 Masatoshi OKITA
On the convergence rates for the compressible Navier- Stokes equations with potential force
- MI2013-1 Abuduwaili PAERHATI & Yasuhide FUKUMOTO
A Counter-example to Thomson-Tait-Chetayev's Theorem
- MI2013-2 Yasuhide FUKUMOTO & Hirofumi SAKUMA
A unified view of topological invariants of barotropic and baroclinic fluids and their application to formal stability analysis of three-dimensional ideal gas flows
- MI2013-3 Hiroki MASUDA
Asymptotics for functionals of self-normalized residuals of discretely observed stochastic processes
- MI2013-4 Naoyuki KAMIYAMA
On Counting Output Patterns of Logic Circuits
- MI2013-5 Hiroshi INOUE
RIPless Theory for Compressed Sensing

- MI2013-6 Hiroshi INOUE
Improved bounds on Restricted isometry for compressed sensing
- MI2013-7 Hidetoshi MATSUI
Variable and boundary selection for functional data via multiclass logistic regression modeling
- MI2013-8 Hidetoshi MATSUI
Variable selection for varying coefficient models with the sparse regularization
- MI2013-9 Naoyuki KAMIYAMA
Packing Arborescences in Acyclic Temporal Networks
- MI2013-10 Masato WAKAYAMA
Equivalence between the eigenvalue problem of non-commutative harmonic oscillators and existence of holomorphic solutions of Heun's differential equations, eigenstates degeneration, and Rabi's model
- MI2013-11 Masatoshi OKITA
Optimal decay rate for strong solutions in critical spaces to the compressible Navier-Stokes equations
- MI2013-12 Shuichi KAWANO, Ibuki HOSHINA, Kazuki MATSUDA & Sadanori KONISHI
Predictive model selection criteria for Bayesian lasso
- MI2013-13 Hayato CHIBA
The First Painleve Equation on the Weighted Projective Space
- MI2013-14 Hidetoshi MATSUI
Variable selection for functional linear models with functional predictors and a functional response
- MI2013-15 Naoyuki KAMIYAMA
The Fault-Tolerant Facility Location Problem with Submodular Penalties
- MI2013-16 Hidetoshi MATSUI
Selection of classification boundaries using the logistic regression
- MI2014-1 Naoyuki KAMIYAMA
Popular Matchings under Matroid Constraints
- MI2014-2 Yasuhide FUKUMOTO & Youichi MIE
Lagrangian approach to weakly nonlinear interaction of Kelvin waves and a symmetry-breaking bifurcation of a rotating flow
- MI2014-3 Reika AOYAMA
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Parallel flow in a cylindrical domain
- MI2014-4 Naoyuki KAMIYAMA
The Popular Condensation Problem under Matroid Constraints