The Effectiveness on Water, Sediment Quality and Shrimp Production (*Penaeus monodon*) at Ponds Treated with *Streptomyces* sp. A1 Probiotic in Thua Thien Hue- Viet Nam

Chau, Ngo Thi Tuong
Ha Noi University of Science, Viet Nam National University

Matsumoto, Masaru
Institute of Tropical Agriculture, Kyushu University

Miyajima, Ikuo
Institute of Tropical Agriculture, Kyushu University

<https://doi.org/10.5109/1434377>

出版情報 : 九州大学大学院農学研究院紀要. 59 (1), pp.33-38, 2014-02-28. Faculty of Agriculture, Kyushu University

バージョン :

権利関係 :

The Effectiveness on Water, Sediment Quality and Shrimp Production (*Penaeus monodon*) at Ponds Treated with *Streptomyces* sp. A1 Probiotic in Thua Thien Hue– Viet Nam

Ngo Thi Tuong CHAU^{1*}, Masaru MATSUMOTO² and Ikuo MIYAJIMA²

Institute of Tropical Agriculture, Kyushu University, Fukuoka 812–8581, Japan

(Received October 28, 2013 and accepted November 11, 2013)

The use of probiotics has increasingly become popular in shrimp culture not only for disease prevention and nutrition improvement but also because of an increasing demand for environment–friendly aquaculture. However, in Viet Nam, most of probiotics were imported with unknown origin so that they can pose a threat of spreading pathogenic or genetically modified microorganisms, resulting in the broken safe biology. Also, the beneficial effect of using such microbial products in aquaculture is still debatable as their efficacy is yet unclear. The objectives of this paper is to assess the effectiveness on water, sediment quality and shrimp production in ponds treated with a *Streptomyces* sp. A1 probiotic in Phu Loc district, Thua Thien Hue province (Viet Nam). Results showed that the probiotic could significantly increase pH, salinity, dissolved oxygen and reduce biochemical oxygen demand, chemical oxygen demand, concentrations of $\text{NH}_4\text{-N}$, $\text{NO}_2\text{-N}$, $\text{NO}_3\text{-N}$, presumptive *Vibrio* count and total coliforms of shrimp pond water. Simultaneously, the probiotic was likely to maintain pH, moisture, concentrations of total organic carbon, total nitrogen, total phosphorus and total sulfur suitable for shrimp pond sediment, especially reduce total carbon. Moreover, an average of 364.38 kg shrimp/ha was obtained in treated ponds with a feed conversion ratio of 1.10, per day growth of 0.21 g and survival rate of 85.43% compared with 240 kg shrimp/ha, 1.67, 0.13 and 84.97, respectively, in control ponds. This indicates that the addition of the probiotic had a noticeable influence on water and sediment quality of shrimp ponds and shrimp production. The finding of this study will facilitate to apply effectively a novel and safe indigenous probiotic into shrimp aquaculture, thereby improving shrimp aquaculture environment, increasing shrimp production, limiting to use antibiotics and chemicals, ensuring safe products and reaching for the sustainable shrimp aquaculture development of Thua Thien Hue in particular and Viet Nam in general.

Key words: *Penaeus monodon*, probiotic, shrimp pond sediment, shrimp pond water, *Streptomyces* sp. A1

INTRODUCTION

Tam Giang – Cau Hai is largest and most typical lagoon of Vietnam, with area of water surface is about 22,000 ha, or 48.2% of total area of all coastal lagoons of Vietnam. With this advantage, Thua Thien Hue province (Viet Nam) has many favorable conditions for development of shrimp culture (IMOLA, 2005). In fact, remarkable increase in shrimp culture helps to restructure the local rural economy in a positive direction (Tuan, 2010). However, a rapid, large–scale and often unplanned increase in brackish water shrimp culture ponds in recent years have resulted in shrimp epidemic diseases (Thua Thien Hue Department of Fisheries, 2010).

The shrimp diseases have mainly accounted for polluted pond environment. The addition of substantial amounts of antibiotics is still the choice for control the disease. Yet the abuse of antibiotics has resulted in the development of multi–resistant bacterial strains and antibiotic residues in shrimp products (Tendencia *et al.*, 2001; Molina *et al.*, 2002; Jayasree *et al.*, 2006; Nakayama *et al.*, 2006). Given the world–wide trade in aquaculture

products, health problems related to antibiotic use in aquaculture are not limited to producing countries, but are also relevant to importing countries. Actually, there were many events of returning consignments to Viet Nam from Japan for not maintaining the prescribed standards on residual antibiotics of the products. Therefore, to make the shrimp aquaculture industry more sustainable, alternative strategies to control infections are urgently needed. One of the most successful alternative methods has been reported to be the use of probiotics as biological control agents in shrimp aquaculture (Verschuere *et al.*, 2000; Vine *et al.*, 2006; Tom *et al.*, 2007). However, most such probiotics in Viet Nam are either imports or domestic products with unknown sources (Ministry of Fisheries, 2005), resulting in a threat of spreading pathogenic and genetically modified strains. Meanwhile, a actinomycetes strain *Streptomyces* sp. A1, which we isolated from shrimp culture pond sediments in Thua Thien Hue (Viet Nam), showed the activity against pathogenic strains *Vibrio* spp. through production of inhibitory compounds and siderophore, the ability to produce extracellular enzymes to decompose organic compounds, the formation of heat– and desiccation–resistant spores and the stability in preservation. Besides, *Streptomyces* sp. A1 can be mass–cultured, harvested, and fortified to feed (Chau *et al.*, 2011).

This paper gives an account of the effectiveness on water, sediment quality and shrimp production (*Penaeus*

¹ Ha Noi University of Science, Viet Nam National University, 334 Nguyen Trai, Thanh Xuan, Ha Noi, Viet Nam

² Institute of Tropical Agriculture, Kyushu University, Fukuoka 812–8581 Japan

* Corresponding author: (E–mail: ngotuongchau@hus.edu.vn)

monodon) at ponds treated with *Streptomyces* sp. A1 probiotic in Phu Loc district, Thua Thien Hue province (Viet Nam). The results will facilitate to offer the effective application of actinomycetes for improving the shrimp aquaculture in Thua Thien Hue and Viet Nam.

MATERIALS AND METHODS

Preparation of *Streptomyces* sp. A1 probiotic

Streptomyces sp. A1 was cultivated in optimal starch casein broth (SCB) sterilized for 15 min at 121°C in 250 mL Erlenmeyer flasks, on an incubated shaker (Jeiotech SI-600R, Korea) at 150 rpm, 30°C for 2 days. Then, this suspension were transferred to sterilized SCB in a 10-L fermentor (Infors Labfors HT, Switzerland) with a ratio of 1: 10 (v/v) for 3 days. After that, the fermentor was left to stand for 30 min to allow the vegetative biomass (micro-colonies) of the *Streptomyces* sp. A1 to settle. The biomass was harvested using sterile filter papers and a vacuum filter, washed with sterile distilled water at least three times, dried at 35°C for 30 min, packaged and stored at 4°C.

Treated design

Tiger shrimp (*Penaeus monodon*) were cultured in high tide ponds in Phu Loc district, Thua Thien Hue province, Viet Nam (16°16'20.39"N, 107°52'29.77"E), during from Jun 28th to September 6th 2012, following the local farmer's experiences. The larvae of *P. monodon* (stage PL 9) were put into the ponds with each area of 1000 m² and stocking density of 30 shrimp/m². Shrimps were fed commercial feed with amounts equal to 1% of body weight divided into three times per day. After a month, with a lower density of 20 shrimp/m², *Streptomyces* sp. A1 probiotic was added directly with a density of 10⁵ CFU per liter into water of the treated pond at intervals of 10 days during shrimp culture period. The ponds without adding the probiotic served as control ones. There were 6 phases to collect water and sediment samples referred as I to VI.

Water analysis

Temperature, conductivity, turbidity, salinity, pH, dissolved oxygen (DO) and total dissolve solid (TDS) of water samples were measured in the field using the Horiba U-52 (Japan). Water samples were mixtures collected near the surface of the center and sides of each pond by filling a sterile hand-held bottles. Water samples were stored in pre-sterilized plastic bottles and chilled immediately with ice and brought back aseptically to the laboratory for analysis. The concentrations of total suspended solid (TSS), inorganic nitrogen compounds (NH₄-N, NO₂-N, and NO₃-N), soluble reactive phosphorus (SRP), chemical oxygen demand (COD), biochemical oxygen demand (BOD) and presumptive *Vibrio* count on thiosulfate citrate bile salt sucrose agar (TCBS) and total coliforms in water samples were measured according to the procedures of APHA (2002).

Sediment analysis

Sediment samples were mixtures collected randomly on the surface of pond bottom (0–10 cm). The samples were stored in Pyrex bottles and transported aseptically to the laboratory and processed immediately for analyses. The pH (H₂O) and pH (KCl) were measured by the glass electrode method using a Hach-C535 (Hach, USA). Organic carbon (TOC) and total carbon (TC) were determined after dry combustion (ISO 10694: 1995). Total nitrogen (TN) was determined by modified Kjeldahl method (ISO 11261:1995). Total phosphorus (TP) was determined by Ascorbic acid method (10 TCN 373–99). Total sulfur (TS) was determined by dry combustion (ISO 15178:2000).

Growth analysis

The growth parameters were calculated by using the following formulae (Felix and Sudharsan, 2004)

$$\text{Per day growth (g)} = \frac{\text{Mean weight gain (g)}}{\text{Number of days}}$$

$$\text{Food conversion ratio (FCR)} = \frac{\text{Total feed given (g)}}{\text{Wet weight gain (g)}}$$

Survival (%)

$$= \frac{\text{Number of shrimps survived at the end of the experiment}}{\text{Number of shrimps stocked at the start of the experiment}} \times 100$$

Final production (kg/ha)

$$= \frac{\text{Total shrimp weight harvested (kg)}}{\text{Area of pond (ha)}}$$

Statistical analysis

Data were analysed using one-way ANOVA (SAS, version 6.03) to find significant difference on various parameters between treated and control ponds. A significance level of $P < 0.05$ was used.

RESULTS

Proliferation of *Streptomyces* sp. A1

A large amount of actinomycete biomass (17.0 g/l) was obtained, meeting the requirement for further experiments.

Water quality

The results of water quality analysis were shown in Tables 1 and 2. Water temperatures were nearly the same in the treated ponds at sampling phases, and there was no considerable difference in water temperature between treated and control ponds. pH values of the water sam-

Table 1. Water quality in the treated ponds at various phases (from I to VI)

Parameters	I	II	III	IV	V	VI	Average
Temperature (°C)	29.2	30.5	29.0	29.0	31.0	32.0	29.9
pH	7.0	6.8	6.9	6.8	6.8	6.9	6.9
Salinity (‰)	12.9	15.0	16.7	19.9	23.1	15.4	17.2
Conductivity (mS/cm)	21.6	24.8	27.2	32.0	36.6	25.4	27.9
TDS (g/l)	13.4	15.4	16.9	19.5	22.3	15.8	17.2
DO (mg/l)	19.8	19.1	12.7	11.2	7.8	12.2	13.8
Turbidity (NTU)	13.0	50.0	111	81.2	84.8	136	79.3
TSS (mg/l)	6	28	51	38	36	58	36
BOD ₅ (mg/l)	8	3	4	6	6	12	7
COD (mg/l)	83	85	75	83	31	69	71
NH ₄ ⁺ -N (mg/l)	ND	ND	ND	ND	ND	ND	ND
NO ₂ ⁻ -N (mg/l)	ND	0.01	0.02	ND	0.00	ND	0.01
NO ₃ ⁻ -N (mg/l)	ND	0.11	0.05	0.14	0.12	0.02	0.09
SRP (mg/l)	0.1	0.2	0.1	0.2	0.4	ND	0.2
Presumptive <i>Vibrio</i> (10 ⁴ CFU/ml)	NT	NT	4.2	2.6	35	6	8.0
Total coliform (10 ³ CFU/100 ml)	0.17	0.7	1.75	4.8	10.8	46	10.7

ND: non-detected; NT: non-tested

Table 2. Water quality in the control ponds at various phases (from I to VI)

Parameters	I	II	III	IV	V	VI	Average
Temperature (°C)	29.2	30.5	28.9	29.9	30.9	31.7	30.2
pH	6.5	6.7	6.7	6.9	6.8	7.1	6.8
Salinity (‰)	13.1	14.4	16.2	19.0	21.8	14.2	16.5
Conductivity (mS/cm)	19.0	24.4	26.4	30.6	34.7	23.5	26.4
TDS (g/l)	13.5	15.1	16.4	18.7	21.1	14.6	16.6
DO (mg/l)	7.3	7.4	6.2	6.9	12.4	16.4	9.4
Turbidity (NTU)	23	91.5	114	157	129	43.7	93
TSS (mg/l)	13	58	87	85	69	60	62
BOD ₅ (mg/l)	8	5	2	6	6	18	7.5
COD (mg/l)	81	87	74	79	37	71	72
NH ₄ ⁺ -N (mg/l)	ND	0.02	ND	0.01	0.01	0.20	0.06
NO ₂ ⁻ -N (mg/l)	ND	0.01	0.02	0.03	ND	ND	0.02
NO ₃ ⁻ -N (mg/l)	0.01	0.06	0.01	ND	0.08	1.94	0.35
SRP (mg/l)	0.1	0.1	0.2	0.2	0.1	ND	0.1
Presumptive <i>Vibrio</i> (10 ⁴ CFU/ml)	NT	NT	15	23	188	39	44.2
Total coliform (10 ³ CFU/100 ml)	0.14	0.65	1.75	3.4	10.8	54	11.79

ND: non-detected; NT: non-tested

ples were not considerably changed during shrimp culture period. The water salinity of all ponds increased linearly from I to V phases. COD of water samples in the treated and control ponds were remarkably changed through survey phases with average values of 71 and 71.5 mg/l, respectively. Also, average BOD of water samples in the treated ponds (7.0 mg/l) was lower than that in control ponds (7.5 mg/l). The concentrations of NH₄⁺-N in all water samples collected from the treated ponds were did not detected whereas those from the control

ponds ranged from 0.01 to 0.2 mg/l with average value of 0.06 mg/l. Following the same trend with BOD and COD, the average value of NO₃⁻-N of water samples in treated ponds (0.09 mg/l) was lower than that in control ponds (0.35 mg/l). According to Boyd (1990), the suitable concentrations of NO₃⁻-N for shrimp aquaculture ranged from 0.1 to 10 mg/l, then concentration of NO₃⁻-N detected in all water samples in this study appeared to be reasonable. However, the average concentration of RSP in water samples of treated ponds (0.2 mg/l) was higher than that

of control ponds (0.1 mg/l). Total coliform of water samples in ponds increased linearly through phases in respond to increased accumulation of shrimp feces during shrimp growth. The average total coliform in treated ponds was lower than that in control ponds. Especially, presumptive *Vibrio* count on TCBS agar was significantly reduced in treated ponds. In comparison to the National Technical Regulation on Coastal Water Quality (QCVN10: 2008/BTNMT) and National Standard on Coastal Water Quality (TCVN 5943–1995) or the procedure for intensive culture of tiger shrimp (28 TCN 171: 2001), the average temperature, pH, salinity and $\text{NO}_2\text{-N}$ of water samples in all ponds generally were within the limits of the coastal water used for aquaculture purposes. However, COD was over the limit of the coastal water used for aquaculture purposes in QCVN10:2008/BTNMT. Also, average total coliform was over the limit of the coastal water used for aquaculture purposes (10^3 CFU/100 ml) in QCVN10:2008/BTNMT and TCVN 5943–1995.

Sediment quality

Sediment samples were collected from ponds and their properties were determined (Tables 3 and 4). Over the study periods, in the treated ponds, sediment pH (H_2O) averaged 7.39 in a range of 7.2 and 7.61, and pH (KCl) averaged 7.22 in a range of 7.15 to 7.3. It is likely to suitable because shrimp pond bottom soils always need to have a pH of 7.5 or above to encourage decomposition of organic matter (Boyd, 2000). The high pH reflects the effect of the applications of liming materials

into dry shrimp pond bottom and brackishwater buffers the soil so that acidic conditions usually do not develop in the surface layer of soil (Boyd, 2000). The sediment moisture in treated ponds significantly ranged from 32.11 to 70.22% and averaged 51.5%. It may be due to a long spell of heavy rain occurred in this time. The optimum moisture content was about 20% for organic matter composition (Boyd, 2000). The average total carbon (TC) concentration in sediment samples of treated ponds (4.64%) was lower than that of control ponds (4.91%). Conversely, the average total organic carbon (TOC) concentration in sediment samples of treated ponds (2.78%) was higher than that of control ponds (2.57%). However, these concentrations match with suggestion that total organic carbon (TOC) concentrations ranged from 0.18% to 7.20% in many samples of the upper soil layer of shrimp ponds in some countries (Boyd, 1995). Total nitrogen (TN) concentration in sediment samples of treated ponds averaged 0.17% (ranging from 0.13 to 0.22 %) and of control ones averaged 0.18% (ranging from 0.15 to 0.24%) were not significantly different. Also, total phosphorus (TP) concentration in sediment samples of treated pond averaged 0.08% and of control ponds averaged 0.07% and there was no significant effect of the treatment by probiotic supplement on TP concentration. Whereas, the average total sulfur (TS) concentration in sediment samples of treated ponds (0.27%) was higher than that of control ponds (0.24%).

Table 3. Sediment quality in the treated ponds at various phases (from I to VI)

Parameters	I	II	III	IV	V	VI	Average
pH (H_2O)	7.45	7.20	7.38	7.32	7.38	7.61	7.39
pH (KCl)	7.22	7.15	7.19	7.23	7.21	7.30	7.22
Moisture (%)	50.72	46.03	32.11	60.47	70.22	50.63	51.50
TC (%)	5.10	4.18	4.36	4.87	4.58	4.75	4.64
TOC (%)	3.02	2.70	2.86	2.70	2.38	3.02	2.78
TN (%)	0.17	0.13	0.16	0.18	0.18	0.22	0.17
TS (%)	0.37	0.26	0.30	0.16	0.25	0.28	0.27
TP (%)	0.08	0.07	0.08	0.08	0.07	0.07	0.08

Table 4. Sediment quality in the control ponds at various phases (from I to VI)

Parameters	I	II	III	IV	V	VI	Average
pH (H_2O)	7.47	7.39	7.36	7.38	7.53	7.61	7.46
pH (KCl)	7.22	7.25	7.24	7.16	7.15	7.39	7.24
Moisture (%)	34.80	52.75	37.00	59.29	54.28	56.56	49.31
TC (%)	4.96	4.64	4.56	5.21	5.59	4.47	4.91
TOC (%)	2.86	2.54	2.70	2.54	2.70	2.07	2.57
TN (%)	0.16	0.16	0.16	0.15	0.18	0.24	0.18
TS (%)	0.18	0.21	0.29	0.22	0.26	0.27	0.24
TP (%)	0.10	0.05	0.05	0.09	0.07	0.07	0.07

Table 5. Growth parameters in treated and control ponds

Treatment	Per day growth (g)	FCR	Survival rate (%)	Production (kg/ha)
Treated ponds	0.21	1.10	85.43	364.38
Control ponds	0.13	1.67	84.97	240

Shrimp production

Shrimp from ponds treated with probiotic had significantly higher survival rate (85.43%), per day growth (0.21 g) and final production (365.38 kg/ha) with feed conversion rate (FCR, 1.67%) in comparison with those of control ponds ($P < 0.05$) (Table 5). This indicates that the addition of the probiotic had a noticeable influence on shrimp production.

DISCUSSION

Microbial processes are intricately intertwined with water, sediment quality and production in shrimp ponds. Uneaten feed, shrimp feces, remains of shrimp settle to pond bottoms where microbial decomposition of this settled organic matter may deplete dissolved oxygen (DO) concentrations. Resultant anaerobic decomposition depresses the redox potential and high concentration of nitrite, hydrogen sulfide and other reduced substances may result. In extensive ponds, shrimp obtain their nutrient by eating natural food organisms growing on the pond soil. In other words, excessive concentrations of organic matter in sediments may cause anaerobic conditions at the bottom which harm benthos and shrimp. Besides, the proliferation of opportunistically pathogenic *Vibrio* spp. commonly present in brackishwater and its sediments is considered as the major causation of the shrimp mortality in water rich in organic nutrients.

Despite the source of several novel antibiotics, marine actinobacteria has been given no attention for use as a probiotic in aquaculture. However, it was recommended that marine actinomycetes are promising candidates to be utilized in marine aquaculture (You *et al.*, 2005, 2007). You *et al.* (2005) described the potential of actinomycetes against shrimp pathogenic *Vibrio* spp. and recommended marine actinomycetes as potential probiotic strains due to their ability to degrade macromolecules, such as starch and protein, in culture pond water; the production of antimicrobial agents; and the formation of heat- and desiccation-resistant spores. More recently, there were a few studies on the possible use of marine actinomycetes in disease prevention against aquatic pathogens. Surajit *et al.* (2006) reported a preliminary study on the use of *Streptomyces* on the growth of black tiger shrimp. Kumar *et al.* (2006) extracted the antibiotic product from marine actinomycetes and incorporated it into feed to observe the *in vivo* effect on white spot syndrome virus in black tiger shrimp. You *et al.* (2007) recommended the use of actinomycetes to prevent the disease caused by *Vibrio* spp. All of this research indicated the importance of marine actinobacteria in aquaculture.

The development of suitable probiotics requires

empirical and fundamental experimental researches. It may be mentioned that a good number of probiotics, of foreign origin were tested in some hatcheries and culture ponds and was not found to be effective. This observation demands to discover new probiotics from the indigenous origin to be effective in local environment (Shafiqur *et al.*, 2009).

Given this circumstances, the addition of an indigenous actinomycetes strain, *Streptomyces* sp. A1, with the high activity against pathogenic strains *Vibrio* spp. through production of inhibitory compounds, the ability to produce extracellular enzymes to decompose organic compounds, the formation of heat- and desiccation-resistant spores and the stability in preservation mass-cultured, harvested, and fortified to feed appear to be an appropriate choice for shrimp aquaculture in Thua Thien Hue. In fact, findings of this study showed that the *Streptomyces* sp. A1 probiotic can be used to treat tiger shrimp (*Penaeus monodon*) ponds with ability to increase pH, salinity, DO and reduce BOD, COD, $\text{NH}_4\text{-N}$, $\text{NO}_2\text{-N}$, $\text{NO}_3\text{-N}$ and total coliforms of shrimp pond water. Especially, presumptive *Vibrio* count on thiosulfate citrate bile salt sucrose (TCBS) agar was significantly reduced in treated ponds compared to control ponds. Also, the probiotic was likely to maintain suitable quality of shrimp pond sediment and improve shrimp production. The results may offer the potential application of actinomycetes for improving the sustainable shrimp aquaculture in Thua Thien Hue and Viet Nam.

ACKNOWLEDGEMENT

This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number **106.03–2011.59**.

REFERENCES

- APHA 2002 Standard methods for the examination of water and wastewater, 20th ed., Press Inc Chapman & Hall, New York, p. 457
- Boyd, C. E. 1990 Water quality in ponds for aquaculture, Ala. Agar. Exp. Sta., Auburn Univer. Ala., p. 462
- Boyd, C. E. 2000 Chemical and physical properties of shrimp pond bottom soil in Ecuador. *J. World Aquacult. Soc.*, **31**(3): 358–375
- Chau, N. T. T., N. X. Hieu, P. T. N. Lan, M. Matsumoto and I. Miyajima. 2011 Identification and characterization of actinomycetes antagonistic to pathogenic *Vibrio* spp. isolated from shrimp culture pond sediments in Thua Thien Hue–Viet Nam. *J. Fac. Agr., Kyushu Univ.*, **56**(1), 15–22
- Felix, N. and M. Sudharsan 2004 Effect of glycine betaine, a feed attractant affecting growth and feed conversion of juvenile

- freshwater prawn *Macrobrachium rosenbergii*. *J. Aquacult. Nutr.*, **10**: 193–197
- IMOLA 2005 The proceedings of the National Workshop on Lagoons in Thua Thien Hue. Department of Science and Technology of Thua Thien Hue Province, The International Union Conservation of Nature and Natural Resources (IUCN) Office in Vietnam and Integrated Management of Lagoon Activities (IMOLA) Project, pp. 8–9 (in Vietnamese)
- Jayasree, L., P. Janakiram and R. Madhavi 2006 Characterization of *Vibrio* spp. associated with diseased shrimp from culture ponds of Andhra Pradesh (India). *J. World Aquacult. Soc.*, **37**(4): 523–532
- Kumar, S. S., R. Philip and C. T. Achuthankutty 2006 Antiviral property of marine actinomycetes against white spot syndrome virus in penaeid shrimps. *Curr. Sci.*, **91**: 807–811
- Ministry of Fisheries 2005 Sustainable development of shrimp farming– Status, opportunities and challenges for Vietnam. Thematic Information (in Vietnamese).
- Molina, A. A., G. A. Garcia, G. A. Abreu, M. C. Bolan, A. Roque and G. B. Gomez 2002 Plasmid profiling and antibiotic resistance of *Vibrio* strains isolated from cultured penaeid shrimp. *FEMS Microbiol. Lett.*, **213**(1): 7–12
- Nakayama, T., E. Ito, N. Nomura and M. Matsumura 2006 Comparison of *Vibrio harveyi* strains isolated from shrimp farms and from culture collection in terms of toxicity and antibiotic resistance. *FEMS Microbiol. Lett.*, **258**(2): 194–199
- QCVN 10 : 2008/BTNMT Vietnam National technical regulation on coastal water quality (in Vietnamese)
- Shafiqur, R., S. N. Khan, M. N. Naser, and M. M. Karim 2009 Application of probiotic bacteria: a novel approach towards ensuring food safety in shrimp aquaculture. *Journal of Bangladesh Academy of Sciences*, **33**(1): 139–144
- Surajit D., P. S. Lyla and S. A. Khan 2006 Application of *Streptomyces* as a probiotic in the laboratory culture of *Penaeus monodon* (Fabricius). *The Israeli Journal of Aquaculture– Bamidgeh* **58**(3): 198–204
- TCVN 5943–95 Water quality – Vietnam Coastal water quality standard (in Vietnamese)
- Tendencia, E. A., M. R. and De La Pena 2001 Antibiotic resistance of bacteria from shrimp ponds. *Aquaculture* **195**: 193–204
- Thua Thien Hue Department of Fisheries 2010 Report on disease situation by the Department of Fisheries in Thua Thien Hue province (in Vietnamese)
- Tom, D., B. Nico, S. Patrick, V. Willy and B. Peter 2007 Alternatives to antibiotics to control bacterial infections: luminescent vibriosis in aquaculture as an example. *Trends Biotechnol.*, **25**(10): 472–479
- Tuan, N. 2010 Thua Thien Hue on its way to be a centrally– governed city. Thuan Hoa Publishing House, Hue, Vietnam, pp. 69–70.
- Verschuere, L., G. Rombaut, P. Sorgeloos and W. Verstraete 2000 Probiotic bacteria as biological control agents in aquaculture. *Microbiol. Mol. Biol. R.*, **64**(4): 655–671
- Vine, N. G., W. D. Leukes and H. Kaiser 2006 Probiotics in marine larviculture. *FEMS Microbiol. Rev.*, **30**(3): 404–427
- You, J., L. X. Cao, G. F. Liu, S. N. Zhou, H. M. Tan and Y. C. Lin 2005 Isolation and characterization of actinomycetes antagonistic to pathogenic *Vibrio* spp. from nearshore marine sediments. *World. J. Microb. Biot.*, **21**(5): 679–682
- You, J., X. Xue, L. Cao, X. Lu, J. Wang, L. Zhang and S. Zhou 2007 Inhibition of *Vibrio* biofilm formation by a marine actinomycete strain A66. *Appl. Microbiol. Biotechnol.*, **76**(5): 1137–1144
- 28 TCN 171 : 2001 The procedure for intensive culture of Tiger shrimp in Viet Nam (in Vietnamese)