九州大学学術情報リポジトリ Kyushu University Institutional Repository

Monodromies of splitting families for degenerations of Riemann surfaces

奥田, 喬之

https://doi.org/10.15017/1441045

出版情報:九州大学,2013,博士(数理学),課程博士 バージョン: 権利関係:全文ファイル公表済

Monodromies of splitting families for degenerations of Riemann surfaces

Takayuki OKUDA

Abstract

When we study degenerations of Riemann surfaces from a topological viewpoint, the topological monodromies play a very important role. In this paper, as an analogy, we introduce the concept of "topological monodromies of splitting families" for degenerations of Riemann surfaces, and their "monodromy sets". We show that the monodromy sets of barking families associated with tame simple crusts act as a pseudo-periodic homeomorphism of negative twist on each irreducible component of the main fibers. As an application of our results, we show an interesting example of two splitting families for one degeneration that have different topological monodromies, although they give the same splitting.

1 Introduction

A degeneration of Riemann surfaces is a family of complex curves over an open disk in \mathbb{C} such that the central fiber is singular and the other fibers are all smooth complex curves. When we classify degenerations of Riemann surfaces from a topological viewpoint, the topological monodromies play a very important role.

Earle-Sipe [ES] and Shiga-Tanigawa [ST] showed that the topological monodromy of a degeneration is always represented by a pseudo-periodic homeomorphism of negative twist¹. The converse of this result was proved by

¹This terminology is used in [MM].

Matsumoto and Montesinos [MM]. Namely, given a pseudo-periodic homeomorphism f of negative twist, they constructed a degeneration with singular fiber whose monodromy homeomorphism coincides with f up to conjugacy.

We are interested in "splittings of singular fibers of degenerations". For a degeneration of Riemann surfaces, we say that its singular fiber splits into several singular fibers if there exists a complex 1-parameter family of families of complex curves such that the family of complex curves over the origin coincides with the given degeneration and the other families have at least two singular fibers. Such a complex 1-parameter family of families of complex curves is called a *splitting family* for the degeneration of Riemann surfaces. In this paper, as an analogy of topological monodromies of degenerations of Riemann surfaces, we introduce the concept of "topological monodromies of splitting families" for degenerations of Riemann surfaces, and their "monodromy sets". See Section 3 for the precise definitions.

As the main theorem of this paper, we show that the monodromy sets of *barking families* associated with "tame" simple crusts act as a pseudoperiodic homeomorphism of negative twist on each irreducible component of the main fibers. See Theorem 6.2 for the more precise statement. A barking family is a splitting family for degenerations of Riemann surfaces which was is introduced by Takamura in [Ta3]. If the singular fiber of the given degeneration has a subdivisor satisfying certain conditions, then such a subdivisor is called a *simple crust*, and we have an associated barking family. In a barking family, the original singular fiber X_0 is deformed to a simpler one in such a way that a part of X_0 looks "barked" off from X_0 . See Figure 4 in Section 5.

In Section 11, as an application of our results, we show an interesting example of two splitting families for one degeneration that have different topological monodromies, although they give the same splitting (that is, the types of the singular fibers appearing in respective splitting families coincide). This example indicates that the topological monodromy of splitting families plays a very important role when we classify the "topologically distinct" splitting families. Since a splitting family is a 2-parameter degenerating family of Riemann surfaces, the topological monodromies are also expected to be useful in studying fibered complex 3-dimensional manifolds.

2 Preliminaries

Let $\pi: M \to \Delta$ be a family of complex curves of genus $g \ge 1$ over an open disk Δ in \mathbb{C} centered at the origin, that is, a proper surjective holomorphic map from a smooth complex surface M to Δ such that all but finitely many fibers are smooth complex curves of genus g. We call such a π a *degeneration* of Riemann surfaces of genus g if the fiber $X_0 := \pi^{-1}(0)$ over the origin is singular and the other fibers $X_s := \pi^{-1}(s), s \ne 0$, are all smooth.

Two degenerations $\pi_i : M_i \to \Delta$ (i = 1, 2) are topologically equivalent if there exist two orientation preserving homeomorphisms $H : M_1 \to M_2$ and $h : \Delta \to \Delta$ such that h(0) = 0 and the following diagram commutes:

$$\begin{array}{cccc} M_1 & \stackrel{H}{\longrightarrow} & M_2 \\ \pi_1 & & & & \downarrow \pi_2 \\ \Delta & \stackrel{h}{\longrightarrow} & \Delta. \end{array}$$

On the topological classification of degenerations, the following is known.

Theorem 2.1 (Matsumoto-Montesinos [MM]). The topological equivalence classes of minimal degenerations of Riemann surfaces of genus $g \ge 2$ are in bijective correspondence with the conjugacy classes in MCG_g represented by pseudo-periodic homeomorphisms of negative type, via topological monodromy, where MCG_g denotes the mapping class group of an oriented closed real surface of genus g.

For a given degeneration $\pi : M \to \Delta$, we define a splitting family as follows. Let \mathcal{M} be a complex 3-dimensional manifold and set $\Delta^{\dagger} :=$ $\{t \in \mathbb{C} : |t| < \varepsilon\}$, an open disk with sufficiently small radius $\varepsilon > 0$. Consider a proper flat surjective holomorphic map $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$ such that the composition $\operatorname{pr}_2 \circ \Psi : \mathcal{M} \to \Delta^{\dagger}$ with the second projection $\operatorname{pr}_2 : \Delta \times \Delta^{\dagger} \to \Delta^{\dagger}$ is a submersion. For each $t \in \Delta^{\dagger}$, set $\Delta_t := \Delta \times \{t\}$, $M_t := \Psi^{-1}(\Delta_t)$ and $\pi_t := \Psi|_{M_t} : M_t \to \Delta_t$. Note that M_t is a smooth complex surface, and $\pi_t : M_t \to \Delta_t$ is a family of complex curves over Δ_t . Suppose that $\pi_0 : M_0 \to \Delta_0$ coincides with $\pi : M \to \Delta$. Then we call $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$ a *deformation family* for the degeneration $\pi : M \to \Delta$ and each $\pi_t : M_t \to \Delta_t$, $t \in \Delta^{\dagger} \setminus \{0\}$, a *deformation* of $\pi : M \to \Delta$.



Figure 1: The singular fiber of a degeneration of Riemann surfaces of genus two splits into some singular fibers.

Let Sing Ψ be the set of singular points of Ψ , and set $\mathcal{D} := \Psi(\text{Sing }\Psi)$, the singular value locus of Ψ , which is also called the *discriminant* of Ψ . From the assumption that Δ^{\dagger} is sufficiently small, it follows that \mathcal{D} is a plane curve in $\Delta \times \Delta^{\dagger}$ with at most one singularity at (0, 0).

In particular, $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$ is called a *splitting family* if for some integer $N \geq 2$, every deformation $\pi_t : M_t \to \Delta_t$ of the degeneration $\pi : M \to \Delta$ is a family of complex curves with N singular fibers. Set $X_{s,t} := \Psi^{-1}(s,t) (= \pi_t^{-1}(s))$ for each $(s,t) \in \Delta \times \Delta^{\dagger}$. For the deformation $\pi_t : M_t \to \Delta_t$ for a fixed $t \in \Delta^{\dagger} \setminus \{0\}$, denote the singular values of π_t by s_1, s_2, \ldots, s_N . Note that the singular values s_1, s_2, \ldots, s_N themselves depend on t, while the topological types of the singular fibers $X_{s_1,t}, X_{s_2,t}, \ldots, X_{s_N,t}$ over them do not. Then we say that the singular fiber X_0 splits into the singular fibers $X_{s_1,t}, X_{s_2,t}, \ldots, X_{s_N,t}$, and we write

$$X_0 \longrightarrow X_{s_1,t} + X_{s_2,t} + \dots + X_{s_N,t}.$$

See Figure 1.

3 Monodromies of splitting families

In this section, we introduce the new concept of "topological monodromies of splitting families."

Let $\pi : M \to \Delta$ be a degeneration of Riemann surfaces of genus $g \geq 1$ and $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$ be a splitting family for the degeneration $\pi : M \to \Delta$. Recall that the discriminant \mathcal{D} of Ψ is a plane curve in $\Delta \times \Delta^{\dagger}$ with at most one singularity at (0,0). Suppose that each deformation $\pi_t : M_t \to \Delta_t$ of the degeneration $\pi : M \to \Delta$ has N singular fibers, then the natural projection $\operatorname{pr}_2 : \mathcal{D} \setminus \{(0,0)\} \to \Delta^{\dagger} \setminus \{0\}$ is an N-fold covering map.

We first take a point $t_0 \in \Delta^{\dagger} \setminus \{0\}$, which will be fixed. Note that $\pi_{t_0} : M_{t_0} \to \Delta_{t_0}$ is a family of complex curves with at least two singular fibers. Let D_{t_0} denote its singular value locus, that is, $D_{t_0} = \mathcal{D} \cap \Delta_{t_0}$. Before proceeding, we define the mapping class group of $\pi_{t_0} : M_{t_0} \to \Delta_{t_0}$ as follows. Let $F : M_{t_0} \to M_{t_0}$ and $f : \Delta_{t_0} \to \Delta_{t_0}$ be orientation preserving homeomorphisms that make the diagram

$$\begin{array}{cccc} M_{t_0} & \xrightarrow{F} & M_{t_0} \\ \pi_{t_0} & & & \downarrow \pi_{t_0} \\ \Delta_{t_0} & \xrightarrow{f} & \Delta_{t_0} \end{array}$$

commutative. Clearly $f(D_{t_0}) = D_{t_0}$. Then we call the pair (F, f) a topological automorphism of $\pi_{t_0} : M_{t_0} \to \Delta_{t_0}$. Denote by \mathcal{H} the group of topological automorphisms of π_{t_0} , that is,

$$\mathcal{H} := \left\{ (F, f) \in \operatorname{Homeo}^+(M_{t_0}) \times \operatorname{Homeo}^+(\Delta_{t_0}) : f \circ \pi_{t_0} = \pi_{t_0} \circ F \right\},\$$

where Homeo⁺(M_{t_0}) (resp. Homeo⁺(Δ_{t_0})) is the group of orientation preserving homeomorphisms of M_{t_0} (resp. Δ_{t_0}). The group \mathcal{H} naturally has the structure of a topological group with respect to the compact open topology. Now we define the (*fiber preserving*) mapping class group MCG(π_{t_0}) of π_{t_0} as the group

$$\operatorname{MCG}(\pi_{t_0}) := \pi_0(\mathcal{H}).$$

In other words, $MCG(\pi_{t_0})$ is the group of isotopy classes of topological automorphisms in \mathcal{H} .

Take a smooth simple closed curve γ in $\Delta^{\dagger} \setminus \{0\}$ with base point t_0 that goes once around the origin in the counterclockwise direction. Then $\Delta \times \gamma$ is an open solid torus. Setting $L := \mathcal{D} \cap (\Delta \times \gamma)$, we see that L is a closed braid in the open solid torus $\Delta \times \gamma$. In fact, the natural projection $\operatorname{pr}_2 : L \to \gamma$ is an unramified N-fold covering map.

Note that $\Psi^{-1}(\Delta \times \gamma) \ (= (\mathrm{pr}_2 \circ \Psi)^{-1}(\gamma))$ is a smooth real 5-dimensional manifold. Now we consider the diagram

$$\Psi^{-1}(\Delta \times \gamma) \xrightarrow{\Psi} \Delta \times \gamma \xrightarrow{\operatorname{pr}_2} \gamma$$

of smooth real manifolds. For the closed braid $L \subset \Delta \times \gamma$, we set $W := \Psi^{-1}(L)$, which is nothing but the union of all singular fibers over $\Delta \times \gamma$. We see that there exists a Thom stratification² $(\mathcal{S}, \mathcal{S}')$ for the smooth map $\Psi : \Psi^{-1}(\Delta \times \gamma) \to \Delta \times \gamma$ such that (i) $\operatorname{pr}_2 : \Delta \times \gamma \to \gamma$ maps each stratum of \mathcal{S}' into γ submersively, and that (ii) for any strata $V \in \mathcal{S}$ and $K \in \mathcal{S}'$, the restrictions $\Psi : \overline{V} \cap \Psi^{-1}(\Delta \times \gamma) \to \Delta \times \gamma$ and $\operatorname{pr}_2 : \overline{K} \cap (\Delta \times \gamma) \to \gamma$ are proper. Then, by Thom's second isotopy lemma [GWPL], the stratified map $\Psi : \Psi^{-1}(\Delta \times \gamma) \to \Delta \times \gamma$ is topologically locally trivial over γ . By pasting these trivializations along the simple closed curve γ , we obtain two orientation preserving homeomorphisms $F : M_{t_0} \to M_{t_0}$ and $f : \Delta_{t_0} \to \Delta_{t_0}$ such that F (resp. f) maps $M_{t_0} \cap W$ (resp. D_{t_0}) to itself homeomorphically and that the following diagram is commutative:

$$\begin{array}{cccc} M_{t_0} & \xrightarrow{F} & M_{t_0} \\ \pi_{t_0} & & & \downarrow \pi_{t_0} \\ \Delta_{t_0} & \xrightarrow{f} & \Delta_{t_0}. \end{array}$$

Thus the pair (F, f) is a topological automorphism of $\pi_{t_0} : M_{t_0} \to \Delta_{t_0}$ and it is uniquely determined up to isotopy. We call the isotopy class [F, f] in $MCG(\pi_{t_0})$ represented by (F, f) the topological monodromy of $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$. See Figure 2.

Let us consider the restriction of F to the singular fibers. Since $f(D_{t_0}) = D_{t_0}$, a singular fiber is mapped to some singular fiber (possibly to itself).

 $^{^2 {\}rm For}$ the explanation of the Thom stratifications and Thom's second isotopy lemma, see Chapter 2 in [GWPL].



Figure 2: The topological monodromy of a splitting family of a degeneration of Riemann surfaces.

Recall that $L = \mathcal{D} \cap (\Delta \times \gamma)$ is a closed braid in the open solid torus $\Delta \times \gamma$. Let K_1, K_2, \ldots, K_c be the connected components of L, where c is a positive integer. For each $i \ (i = 1, 2, \ldots, c)$, the intersection $\Delta_{t_0} \cap K_i$ is contained in D_{t_0} , and consequently it consists of singular values of π_{t_0} , say,

$$\Delta_{t_0} \cap K_i = \left\{ s_1^{(i)}, s_2^{(i)}, \dots, s_{l_i}^{(i)} \right\} \subset D_{t_0}.$$

Then f cyclically permutes $s_1^{(i)}, s_2^{(i)}, \ldots, s_{l_i}^{(i)}$, while F cyclically permutes the corresponding singular fibers $X_{s_1^{(i)}, t_0}, X_{s_2^{(i)}, t_0}, \ldots, X_{s_{l_i}^{(i)}, t_0}$. Let \mathbb{X}_i denote the disjoint union of the singular fibers over $\Delta_{t_0} \cap K_i$:

$$\mathbb{X}_i := X_{s_1^{(i)}, t_0} \sqcup X_{s_2^{(i)}, t_0} \sqcup \cdots \sqcup X_{s_{l_i}^{(i)}, t_0}.$$

We call the union \mathbb{X}_i the *tassel*³ over K_i . Consider the restriction

$$F_i := F\Big|_{\mathbb{X}_i} : \mathbb{X}_i \to \mathbb{X}_i$$

³This terminology is introduced only for barking families by Takamura [Ta3].

of F to the tassel \mathbb{X}_i . Clearly for each $j = 1, 2, \ldots, l_i$, we have $F_i^{l_i}(X_{s_j^{(i)}, t_0}) = X_{s_j^{(i)}, t_0}$. We call the (c+1)-tuple $(F_1, F_2, \ldots, F_c; f)$ a monodromy set of the splitting family $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$. A monodromy set is uniquely determined up to isotopy.

4 Linear degenerations

This section reviews the concept of a linear degeneration, which is a representative of an equivalence class of degenerations.

Let $\pi: M \to \Delta$ be a degeneration of complex curves of genus $g \ge 1$ with singular fiber $X_0 = \sum_i m_i \Theta_i$, where each Θ_i is an irreducible component of X_0 with multiplicity m_i . Denote by X_0^{red} the underlying reduced curve of X_0 , that is, $X_0^{\text{red}} := \sum_i \Theta_i$. We say that the singular fiber X_0 (or more precisely, its underlying reduced curve X_0^{red}) has at most simple normal crossings if (i) every singularity of X_0^{red} is a node and (ii) none of the irreducible components Θ_i intersects itself (and therefore, each Θ_i is smooth). It is known that an arbitrary degeneration of Riemann surfaces, by successive blowing-ups, can be arranged so that its singular fiber has at most simple normal crossings.

In what follows, we assume that the singular fibers of any given degenerations have at most simple normal crossings.

Let $\Theta_1, \Theta_2, \ldots, \Theta_\lambda$ ($\lambda \ge 1$) be Riemann spheres contained in X_0 as irreducible components that satisfy the following conditions.

- Θ_i intersects Θ_{i-1} and Θ_{i+1} at exactly one point, respectively, and does not intersect other irreducible components of X_0 , $i = 2, 3, ..., \lambda 1$.
- Θ_1 intersects at most one irreducible component other than Θ_2 , and if it exists, say Θ_0 , then Θ_1 intersects Θ_0 at exactly one point.
- Θ_{λ} intersects at most one irreducible component other than $\Theta_{\lambda-1}$, and if it exists, say $\Theta_{\lambda+1}$, then Θ_{λ} intersects $\Theta_{\lambda+1}$ at exactly one point.

Let m_i is the multiplicity of Θ_i in X_0 $(i = 1, 2, ..., \lambda)$. Then the divisor

$$\mathbf{Ch} := m_1 \Theta_1 + m_2 \Theta_2 + \dots + m_\lambda \Theta_\lambda$$



Figure 3: This singular fiber has three cores, four branches and two trunks. The numbers stand for the multiplicity of each irreducible component and each intersection point is a node.

is called a *chain of Riemann spheres*. In what follows, we assume that, when we express a chain of Riemann spheres in this form, the Riemann spheres are arranged in this order. If Θ_1 (resp. Θ_{λ}) intersects an irreducible component Θ_0 (resp. $\Theta_{\lambda+1}$), then let m_0 (resp. $m_{\lambda+1}$) denote the multiplicity of Θ_0 (resp. $\Theta_{\lambda+1}$), and otherwise set $m_0 := 0$ (resp. $m_{\lambda+1} := 0$). Then we call the sequence of nonnegative integers $m_0, m_1, \ldots, m_{\lambda+1}$ the *multiplicity sequence* associated with the chain **Ch**.

An irreducible component of the singular fiber X_0 is called a *core* if it intersects the other irreducible components at at least three points or its genus is positive. A *branch* is a chain of Riemann spheres attached with a core on one end, while a *trunk* is a chain of Riemann spheres attached with cores on both ends. The singular fiber X_0 consists of cores, branches and trunks. See Figure 3. We say that X_0 is a *stellar* singular fiber if X_0 consists of exactly one core and some branches emanating from the core. Otherwise X_0 is said to be *constellar*. It is known that a constellar singular fiber is obtained from stellar singular fibers by "Matsumoto-Montesinos bonding" — Matsumoto-Montesinos bonding yields a trunk from two branches (see [Ta3] for details).

For an irreducible component Θ_i of X_0 , we denote by N_i the normal bundle of Θ_i in M. Let $\{p_1, p_2, \ldots, p_h\}$ be the set of the intersection points on Θ_i with the other irreducible components of X_0 and m_j be the multiplicity of the irreducible component intersecting Θ_i at p_j (j = 1, 2, ..., h). Note that

$$r_i := \frac{\sum_{j=1}^h m_j}{m_i}$$

is a positive integer. In fact, the self-intersection number of Θ_i in M is equal to $-r_i$, which follows from the adjunction formula. Then there exists a holomorphic section σ_i of the line bundle $N_i^{\otimes (-m_i)}$ over Θ_i such that

$$\operatorname{div}(\sigma_i) = \sum_{j=1}^h m_j p_j,$$

where div(σ_i) denotes the divisor defined by σ_i . Here σ_i has a zero of order m_j at p_j . Note that σ_i is uniquely determined up to multiplication by a constant. We call σ_i the standard section of the line bundle $N_i^{\otimes (-m_i)}$ over Θ_i .

For each *i*, take an open covering $\Theta_i = \bigcup_{\alpha} U_{\alpha}$ such that $U_{\alpha} \times \mathbb{C}$ is a local trivialization of the normal bundle N_i . We denote by $(z_{\alpha}, \zeta_{\alpha})$ coordinates of $U_{\alpha} \times \mathbb{C}$. Now define the holomorphic functions $\pi_{i,\alpha} : U_{\alpha} \times \mathbb{C} \to \mathbb{C}$ by

$$\pi_{i,\alpha}(z_{\alpha},\zeta_{\alpha}):=\sigma_{i,\alpha}(z_{\alpha})\zeta_{\alpha}^{m_{i}}$$

where $\sigma_{i,\alpha}$ is the local expression of σ_i on U_{α} . Then we see that the set $\{\pi_{i,\alpha}\}_{\alpha}$ of holomorphic functions defines a global holomorphic function $\pi_i : N_i \to \mathbb{C}$.

Definition 4.1. A degeneration $\pi : M \to \Delta$ is said to be *linear* if for every irreducible component Θ_i of its singular fiber X_0 ,

- (i) a tubular neighborhood $N(\Theta_i)$ of Θ_i in M is biholomorphic to a tubular neighborhood of the zero section of the normal bundle N_i , and
- (ii) under the identification by the biholomorphic map of (i), the following conditions are satisfied:
 - (a) the restriction $\pi|_{N(\Theta_i)}$ coincides with the holomorphic function π_i defined above, and
 - (b) if Θ_i intersects Θ_j at a point $p, j \neq i$, then there exist local trivializations $U_{\alpha} \times \mathbb{C}$ of N_i and $U_{\beta} \times \mathbb{C}$ of N_j around p such

that neighborhoods of p in $N(\Theta_i)$ and in $N(\Theta_j)$ are identified by plumbing, $(z_{\alpha}, \zeta_{\alpha}) = (\zeta_{\beta}, z_{\beta})$, and π is locally expressed as

$$\pi\big|_{N(\Theta_i)}(z_\alpha,\zeta_\alpha)=z_\alpha^{m_j}\zeta_\alpha^{m_i},\quad \pi\big|_{N(\Theta_j)}(z_\beta,\zeta_\beta)=z_\beta^{m_i}\zeta_\beta^{m_j},$$

where $(z_{\alpha}, \zeta_{\alpha}) \in U_{\alpha} \times \mathbb{C}$ and $(z_{\beta}, \zeta_{\beta}) \in U_{\beta} \times \mathbb{C}$.

In a linear degeneration, tubular neighborhoods of the branches and the trunks can be constructed explicitly:

Lemma 4.2. Let $m_0, m_1, \ldots, m_{\lambda+1}$ ($\lambda \ge 1$) be nonnegative integers such that

• $m_0, m_1, \ldots, m_{\lambda}$ are positive integers, and

•
$$r_i := \frac{m_{i-1} + m_{i+1}}{m_i}$$
 is a positive integer $(i = 1, 2, \dots, \lambda)$.

Then there exist a smooth complex surface T and a linear degeneration π : $T \rightarrow \Delta$ with the singular fiber

$$X_0 = m_0 V_0 + m_1 \Theta_1 + m_2 \Theta_2 + \dots + m_\lambda \Theta_\lambda + m_{\lambda+1} U_{\lambda+1},$$

where V_0 and $U_{\lambda+1}$ are copies of \mathbb{C} , $m_1\Theta_1 + m_2\Theta_2 + \cdots + m_{\lambda}\Theta_{\lambda}$ is a chain of Riemann spheres, and V_0 (resp. $U_{\lambda+1}$) intersects Θ_1 (resp. Θ_{λ}) at exactly one point.

Proof. We take λ copies $\Theta_1, \Theta_2, \ldots, \Theta_{\lambda}$ of $\mathbb{C}P^1$. For each $i = 1, 2, \ldots, \lambda$, let $\Theta_i = U_i \cup V_i$ be an open covering by two copies U_i, V_i of \mathbb{C} with coordinates $w_i \in U_i \setminus \{0\}$ and $z_i \in V_i \setminus \{0\}$ satisfying $z_i = 1/w_i$. Then we obtain a line bundle N_i over Θ_i of degree $-r_i$ from $U_i \times \mathbb{C}$ and $V_i \times \mathbb{C}$ by identifying $(z_i, \zeta_i) \in (V_i \setminus \{0\}) \times \mathbb{C}$ with $(w_i, \eta_i) \in (U_i \setminus \{0\}) \times \mathbb{C}$ via

$$g_i: z_i = \frac{1}{w_i}, \quad \zeta_i = w_i^{r_i} \eta_i.$$

Now patch N_i and N_{i+1} by plumbing, $(\zeta_i, z_i) = (w_{i+1}, \eta_{i+1})$, for each $i = 1, 2, \ldots, \lambda - 1$, then we obtain a smooth complex surface \widehat{T} .

Let us define the holomorphic functions $\pi_i: N_i \to \mathbb{C}$ by

$$\pi_i = \begin{cases} w_i^{m_{i-1}} \eta_i^{m_i}, & \text{on } U_i \times \mathbb{C}, \\ z_i^{m_{i+1}} \zeta_i^{m_i}, & \text{on } V_i \times \mathbb{C}. \end{cases}$$

The holomorphic functions $\{\pi_i\}$ together define a holomorphic function π : $\widehat{T} \to \mathbb{C}$ and the central fiber is

$$\pi^{-1}(0) = m_0 V_0 + m_1 \Theta_1 + m_2 \Theta_2 + \dots + m_\lambda \Theta_\lambda + m_{\lambda+1} U_{\lambda+1},$$

where $V_0 := \{0\} \times \mathbb{C} \subset U_1 \times \mathbb{C}$ and $U_{\lambda+1} := \{0\} \times \mathbb{C} \subset V_\lambda \times \mathbb{C}$. Thus, setting $T := \pi^{-1}(\Delta)$ for an open disk Δ in \mathbb{C} centered at the origin, the restriction $\pi : T \to \Delta$ of the holomorphic function $\pi : \widehat{T} \to \mathbb{C}$ is the desired linear degeneration.

Remark 4.3. To be precise, since $\pi : T \to \Delta$ obtained in Lemma 4.2 is not proper, it is not a degeneration. However, it can be identified with the restriction of some degeneration to a tubular neighborhood T of a chain $m_1\Theta_1 + m_2\Theta_2 + \cdots + m_\lambda\Theta_\lambda$ contained in the singular fiber.

5 Tame simple crusts and barking families

Let us review Takamura's theory of barking families. For a degeneration, Takamura defined a simple crust as a subdivisor of its singular fiber that satisfies certain conditions, and constructed a splitting family associated to each such simple crust. A splitting family constructed by his method is called a barking family. For details see [Ta3]. In this paper, we consider only simple crusts that satisfy some additional conditions and call them *tame* simple crusts. See Definition 5.4.

Let $\pi : M \to \Delta$ be a linear degeneration of Riemann surfaces with the singular fiber $X_0 = \sum_i m_i \Theta_i$. Let Y be an effective subdivisor of $X_0 = \sum_i m_i \Theta_i$. We express Y as

$$Y = \sum_{i} n_i \Theta_i,$$

where n_i is a nonnegative integer less than or equal to m_i . We define the *underlying reduced curve* of Y as $Y^{\text{red}} := \sum_i \Theta_i$, where the sum runs over all i with $n_i \ge 1$. Namely, an irreducible component Θ_i of X_0 is contained in Y^{red} if and only if $n_i \ge 1$. Let $\text{Core}(X_0)$ denote the set of all cores of X_0 and Core(Y) denote the set of the cores of X_0 that are contained in Y^{red} . We first assume that

- Y (or, more precisely, Y^{red}) is connected, and
- at least one irreducible component of Y is a core of X (or equivalently, $\operatorname{Core}(Y) \neq \emptyset$).

Let **Br** be a branch of X_0 attached with a core Θ_0 , and express it as

$$\mathbf{Br} = m_1 \Theta_1 + m_2 \Theta_2 + \dots + m_\lambda \Theta_\lambda,$$

where Θ_1 is attached with the core Θ_0 . Namely, denoting by m_0 the multiplicity of Θ_0 , the branch **Br** is a chain associated with the multiplicity sequence $m_0, m_1, m_2, \ldots, m_\lambda, m_{\lambda+1} := 0$. For each $i = 1, 2, \ldots, \lambda$, we set

$$r_i := \frac{m_{i-1} + m_{i+1}}{m_i},$$

which is a positive integer. Recall that the self-intersection number of Θ_i in M is equal to $-r_i$. Let n_i be the multiplicity of Θ_i in $Y, i = 0, 1, ..., \lambda$. Now set

$$\mathbf{br} := n_1 \Theta_1 + n_2 \Theta_2 + \dots + n_\lambda \Theta_\lambda.$$

From the above assumptions for Y, if $n_i = 0$ for some i, then $n_{i'} = 0$ for any $i' \ge i$. We thus may express as

$$\mathbf{br} = \emptyset$$
, or $\mathbf{br} = n_1 \Theta_1 + n_2 \Theta_2 + \dots + n_{\nu} \Theta_{\nu}$,

where ν is the least positive integer with $n_{\nu} \neq 0$ among $1, 2, ..., \lambda$. By convention, we set $\nu := 0$ if $\mathbf{br} = \emptyset$. We call \mathbf{br} a *subbranch* of \mathbf{Br} if one of the following conditions is satisfied.

• $\nu = 0$ or 1.

•
$$\nu \ge 2$$
, and $r_i = \frac{n_{i-1} + n_{i+1}}{n_i}$ for each $i = 1, 2, \dots, \nu - 1$.

Set $\overline{n}_{\nu+1} := r_{\nu}n_{\nu} - n_{\nu-1}$. If $\nu = 0$ (that is, $\mathbf{br} = \emptyset$), then we set $\overline{n}_{\nu+1} := 0$. **Definition 5.1.** Let l be a positive integer.

(A) A subbranch **br** of **Br** is of type A_l if $ln_i \leq m_i$ for each $i = 0, 1, ..., \nu$, and $\overline{n}_{\nu+1} \leq 0$.

- (B) A subbranch **br** of **Br** is of type B_l if $ln_i \leq m_i$ for each $i = 0, 1, ..., \nu$, $n_{\nu} = 1$ and $m_{\nu} = l$.
- (C) A subbranch **br** of **Br** is of type C_l if $ln_i \leq m_i$ for each $i = 0, 1, ..., \nu$, $n_{\nu} = \overline{n}_{\nu+1}$ and $m_{\nu} m_{\nu+1}$ divides l.

Now let \mathbf{Tk} be a trunk of X_0 and express it as

$$\mathbf{T}\mathbf{k} = m_1\Theta_1 + m_2\Theta_2 + \dots + m_\lambda\Theta_\lambda.$$

Let Θ_0 (resp. $\Theta_{\lambda+1}$) be the core intersecting Θ_1 (resp. Θ_{λ}) and let m_0 (resp. $m_{\lambda+1}$) denote its multiplicity. Then the trunk **Tk** is a chain of Riemann sphere associated with the multiplicity sequence $m_0, m_1, m_2, \ldots, m_{\lambda}, m_{\lambda+1}$. Recall that, for each $i = 1, 2, \ldots, \lambda$, the self-intersection number of Θ_i in M is equal to $-r_i$, where

$$r_i := \frac{m_{i-1} + m_{i+1}}{m_i}$$

Let n_i be the multiplicity of Θ_i in Y, $i = 0, 1, \ldots, \lambda + 1$. Now set

$$\mathbf{t}\mathbf{k} := n_1\Theta_1 + n_2\Theta_2 + \dots + n_\lambda\Theta_\lambda.$$

Since Y is connected and $\operatorname{Core}(Y) \neq \emptyset$, either n_0 or $n_{\lambda+1}$ or both must be positive.

Definition 5.2. Let l be a positive integer. We call **tk** a *tame subtrunk* of **Tk** with *barking multiplicity* l if the following condition is satisfied.

• $0 < ln_i \le m_i$ and $r_i = \frac{n_{i-1} + n_{i+1}}{n_i}$ for each $i = 1, 2, \dots, \lambda$.

We next consider the cores of X_0 . Let Θ_0 be a core of X_0 and let N_0 denote the normal bundle of Θ_0 in M. Recall that there exists a holomorphic section σ_0 of the line bundle $N_0^{\otimes (-m_0)}$ over Θ_0 such that

$$\operatorname{div}(\sigma_0) = \sum_{j=1}^h m_j p_j,$$

where p_j are the points at which Θ_0 intersects the other irreducible components of X_0 and m_j are the corresponding multiplicities. Let n_0 denote the multiplicity of Θ_0 in Y. Now suppose that there exists a meromorphic section τ of the line bundle $N_0^{\otimes n_0}$ over Θ_0 such that

$$\operatorname{div}(\tau) = -\sum_{j=1}^{h} n_j p_j + D$$

for some nonnegative divisor $D = \sum_{j=h+1}^{h'} a_j p_j$ on Θ_0 , where $p_1, p_2, \ldots, p_{h'}$ are all distinct points of Θ_0 . Then we call the meromorphic section τ a core section over Θ_0 for Y. Note that τ is not uniquely determined by Y. It follows that $r_0 := (\sum_{j=1}^h m_j)/m_0$ is a positive integer, while $r'_0 := (\sum_{j=1}^h n_j)/n_0$ is not necessarily an integer. Furthermore, we have the following (see [Ta3] Section 3.4).

Lemma 5.3. Suppose that the core Θ_0 is a Riemann sphere. Then Y has a core section τ over the core Θ_0 if and only if $r_0 \leq r'_0$. Moreover, τ has no zero, that is, D = 0, exactly when $r_0 = r'_0$.

Now we define tame simple crusts. Recall that $\operatorname{Core}(X_0)$ denotes the set of all cores of X_0 and $\operatorname{Core}(Y)$ denotes the set of the cores of X_0 that are contained in Y^{red} . Set $\operatorname{Adja}(Y) := \{ \Theta \in \operatorname{Core}(X_0) \setminus \operatorname{Core}(Y) : \Theta \cap Y^{\operatorname{red}} \neq \emptyset \}$, whose elements are said to be *adjacent* to Y.

Definition 5.4. Let Y be a connected subdivisor of X_0 such that $\text{Core}(Y) \neq \emptyset$, and let l be a positive integer. We call Y a *tame simple crust* of X_0 with *barking multiplicity* l if the following conditions are satisfied.

- $ln_0 \leq m_0$ for each core $\Theta_0 \in \text{Core}(Y)$, where m_0 and n_0 are the multiplicities of Θ_0 in X_0 and Y, respectively.
- The subdivisor **br** of each branch **Br** of X_0 for Y is a subbranch of type A_l , B_l or C_l .
- The subdivisor \mathbf{tk} of each trunk \mathbf{Tk} of X_0 for Y is a tame subtrunk with barking multiplicity l.
- For each core $\Theta_0 \in \text{Core}(Y) \cup \text{Adja}(Y)$, there exists a core section over Θ_0 for Y which has no zeros.



Figure 4: In a barking family, the singular fiber is deformed to the main fiber in such a way that the simple crust looks "barked" off from the original singular fiber.

In fact, Takamura [Ta3] defined simple crusts — subdivisors of X_0 satisfying more general conditions. (Note that tame simple crusts which we defined above are simple crusts in the sense of [Ta3].) He constructed a deformation family of the given degeneration $\pi : M \to \Delta$ associated with the simple crust Y and its barking multiplicity l. We call a deformation family obtained by his method a barking family. In a barking family, the original singular fiber X_0 is deformed to a simpler singular fiber in such a way that the subdivisor Y looks "barked" off from X_0 as depicted in Figure 4. The resulting singular fiber appears over the origin of Δ_t , so we denote it by $X_{0,t}$ and call it the main fiber. For each irreducible component Θ of the main fiber, exactly one of the following phenomena occurs as $t \to 0$.

- (1) The irreducible component Θ approaches to a union of irreducible components of X_0 which contains Y^{red} .
- (2) The irreducible component Θ approaches to one irreducible component of X_0 .

The irreducible component Θ of $X_{0,t}$ is called a *barked component* if (1) occurs, and a *stable component* if (2) occurs.

In a barking family, there necessarily appear not only the main fiber but also other singular fibers over some points away from the origin of Δ_t , which are called subordinate fibers. Under the deformation, the topological type of the singular fiber over the origin changes, so the local monodromy around it also changes. On the other hand, the global monodromies before and after the deformation — that is, the two monodromies each of which is induced by a loop in Δ (resp. Δ_t) parallel and closed to its boundary $\partial\Delta$ (resp. $\partial\Delta_t$) coincide. We then deduce that there should necessarily appear other singular fibers with nontrivial monodromies, since we see that the monodromies before and after the deformation are distinct. Thus every barking family turns out to be a splitting family. Therefore, we have the following.

Theorem 5.5 (Takamura [Ta3]). Let $\pi : M \to \Delta$ be a linear degeneration with the singular fiber X_0 . If X_0 has a simple crust Y, then $\pi : M \to \Delta$ admits a splitting family $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$.

Remark 5.6. In this paper, for a degeneration which is *not* necessarily relatively minimal, a splitting family is defined in such a way that each deformation has at least two singular fibers. Thus some singular fibers of a deformation in a splitting family may possibly become smooth fibers by blowing-downs. Such singular fibers are said to be *fake*.

6 Monodromy sets of barking families

Let $\pi : M \to \Delta$ be a linear degeneration of Riemann surfaces of genus $g \ge 1$ and $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$ be a barking family for $\pi : M \to \Delta$ associated with a tame simple crust Y. Denote by \mathcal{D} the discriminant of Ψ , that is, $\mathcal{D} := \Psi(\operatorname{Sing} \Psi).$

Let us consider the deformation $\pi_{t_0} : M_{t_0} \to \Delta_{t_0}$ of $\pi : M \to \Delta$ for a fixed point $t_0 \in \Delta^{\dagger} \setminus \{0\}$. Take a smooth simple closed curve γ in $\Delta^{\dagger} \setminus \{0\}$ with base point t_0 that goes once around the origin in the counterclockwise direction: then $L := \mathcal{D} \cap (\Delta \times \gamma)$ is a closed braid in the open solid torus $\Delta \times \gamma$, and we obtain the topological monodromy $[F, f] \in \text{MCG}(\pi_{t_0})$ of the barking family $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$. Let K_1, K_2, \ldots, K_c be the knot components of L, and \mathbb{X}_j be the tassel over $K_j, j = 1, 2, \ldots, c$. Then we obtain the monodromy set $(F_1, F_2, \ldots, F_c; f)$ of Ψ . See Section 3 for details. Since the main fiber of the deformation $\pi_t : M_t \to \Delta_t$ for any $t \in \gamma$ lies over the origin of Δ_t , the core curve $\{0\} \times \gamma$ in the open solid torus $\Delta \times \gamma$ is nothing but the knot component over which the main fibers lie. In what follows, we denote the knot component $\{0\} \times \gamma$ by K_1 , and we call the tassel X_1 over K_1 the main tassel. Clearly $X_1 = X_{0,t_0}$. On the other hand, we call the other tassels X_2, X_3, \ldots, X_c subordinate tassels. Each subordinate fiber of the deformation $\pi_{t_0} : M_{t_0} \to \Delta_{t_0}$ is contained in one of the subordinate tassels.

By using results on determination of subordinate fibers, we see that each irreducible component of \mathcal{D} is the hypersurface

$$\{(s,t)\in\Delta\times\Delta^{\dagger}: s^n=\kappa t^m\},\$$

for some relatively prime positive integers m, n (arising from the multiplicities of X_0 and Y) and some complex number κ . See [O] for instance. Then the intersection of the irreducible component and $\Delta \times \gamma$ is one of the knot components of L, and moreover we see that it is an (m, n)-torus knot in the open solid torus $\Delta \times \gamma$ (more precisely, when the solid torus is embedded in the standard way in a 3-sphere). Hence we have the following.

Proposition 6.1. Let K_1 be the knot component of L over which the main tassel lies, and let K_2, K_3, \ldots, K_c be the knot components of L over which the subordinate tassels lie. Then, we have the following.

- The knot component K_1 is a trivial closed braid in $\Delta \times \gamma$.
- The knot component K_j (j = 2, 3, ..., c) is a torus knot in $\Delta \times \gamma$.

We consider the monodromy homeomorphism $F_1 : X_{0,t_0} \to X_{0,t_0}$ on the main fiber, where $\mathbb{X}_1 = X_{0,t_0}$. We will show that F_1 acts as a pseudo-periodic homeomorphism of negative twist on each irreducible component of the main fiber. To be more precise, we have the following, which is a summary of Propositions 7.2, 9.2 and 9.3.

Theorem 6.2. Let $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$ be a barking family for a linear degeneration $\pi : \mathcal{M} \to \Delta$ associated with a tame simple crust Y. Let us consider the deformation $\pi_{t_0} : M_{t_0} \to \Delta_{t_0}$ of $\pi : \mathcal{M} \to \Delta$ for a fixed $t_0 \in \Delta^{\dagger}$.

We denote by $(F_1, F_2, \ldots, F_c; f)$ the monodromy set of Ψ , and let F_1 be the monodromy homeomorphism on the main fiber X_{0,t_0} . Let $\Theta_1, \Theta_2, \ldots, \Theta_a$ be the stable components of X_{0,t_0} and let $\Xi_1, \Xi_2, \ldots, \Xi_b$ be the barked components of X_{0,t_0} . Then we have the following.

- (1) For each i = 1, 2, ..., a, we have $F_1(\Theta_i) = \Theta_i$, and the restriction $F_1|_{\Theta_i} : \Theta_i \to \Theta_i$ is isotopic to the identity map of Θ_i .
- (2) If b = 1, then we have F₁(Ξ₁) = Ξ₁, and the isotopy class of the restriction F₁|_{Ξ₁} : Ξ₁ → Ξ₁ is conjugate to the topological monodromy of a degeneration of Riemann surfaces whose singular fiber is the enlargement⁴ of the tame simple crust Y. In particular, F₁|_{Ξ₁} : Ξ₁ → Ξ₁ is isotopic to a pseudo-periodic homeomorphism of negative twist.
- (3) If $b \ge 2$, then F_1 cyclically permutes $\Xi_1, \Xi_2, \ldots, \Xi_b$, and the restriction

$$F_1\big|_{\Xi_1\cup\Xi_2\cup\cdots\cup\Xi_b}:\Xi_1\cup\Xi_2\cup\cdots\cup\Xi_b\to\Xi_1\cup\Xi_2\cup\cdots\cup\Xi_b$$

coincides with the monodromy homeomorphism of a degeneration of disjoint unions of b Riemann surfaces whose singular fiber is the enlargement of the tame simple crust Y, up to isotopy and conjugacy. In particular, for each k = 1, 2, ..., b, $F_1^b|_{\Xi_k} : \Xi_k \to \Xi_k$ is isotopic to a pseudo-periodic homeomorphism of negative twist.

7 Degenerations of stable components

Let $\pi: M \to \Delta$ be a linear degeneration of Riemann surfaces of genus $g \geq 1$ and $\Psi: \mathcal{M} \to \Delta \times \Delta^{\dagger}$ be a barking family for $\pi: M \to \Delta$ associated with a tame simple crust Y with barking multiplicity l. For a base point $t_0 \in \Delta^{\dagger} \setminus \{0\}$, we denote the monodromy set of $\Psi: \mathcal{M} \to \Delta \times \Delta^{\dagger}$ by $(F_1, F_2, \ldots, F_c; f)$, where $F_j: \mathbb{X}_j \to \mathbb{X}_j$ is the monodromy homeomorphism on the j-th tassel $\mathbb{X}_j, j = 1, 2, \ldots, c$, and $f: \Delta_{t_0} \to \Delta_{t_0}$ is the associated homeomorphism on the open disk Δ_{t_0} over the base point t_0 . We assume that \mathbb{X}_1 is the main tassel (so $\mathbb{X}_1 = X_{0,t_0}$).

 $^{^{4}}$ See Section 8.

In this section, we investigate the restriction of $F_1 : X_{0,t_0} \to X_{0,t_0}$ to a stable component Θ of X_{0,t_0} . For this purpose, we construct a degeneration such that the stable component Θ coincides with a smooth fiber of it and that its monodromy homeomorphism is isotopic to $F_1|_{\Theta}$.

For each $s \in \Delta$, we set $\Delta_s^{\dagger} := \{s\} \times \Delta^{\dagger}$ and $M_s^{\dagger} := \Psi^{-1}(\Delta_s^{\dagger})$. Then we obtain the map

$$\pi_s^{\dagger} := \Psi \big|_{M^{\dagger}} : M_s^{\dagger} \to \Delta_s^{\dagger}.$$

It might be plausible that π_s^{\dagger} is a family of complex curves if we regard the deformation parameter t of Ψ as a degeneration parameter. However, it is not the case unless M_s^{\dagger} is a smooth complex surface. Note that, for the case s = 0, the central fiber $(\pi_0^{\dagger})^{-1}(0)$ of $\pi_0^{\dagger} : M_0^{\dagger} \to \Delta_0^{\dagger}$, coincides with the singular fiber X_0 of the original degeneration $\pi : M \to \Delta$, while a general fiber $(\pi_0^{\dagger})^{-1}(t), t \neq 0$, coincides with the main fiber $X_{0,t}$ of the deformation $\pi_t : M_t \to \Delta_t$.

Now we consider the restriction of $\pi_0^{\dagger} : M_0^{\dagger} \to \Delta_0^{\dagger}$ to a certain smooth complex surface, which is a degeneration of Riemann surfaces. Let Θ be a stable component of the main fiber X_{0,t_0} of the deformation $\pi_{t_0} : M_{t_0} \to \Delta_{t_0}$. Recall that, as $t_0 \to 0$, the component Θ approaches to some irreducible component of the singular fiber X_0 of $\pi : M \to \Delta$, say Θ_0 . Let N_0 be the normal bundle of Θ_0 in M, with coordinates (z, ζ) , where z is the base coordinate and ζ is the fiber coordinate. We have the following (see [Ta3] Section 16.2).

Lemma 7.1. The complex 3-dimensional manifold \mathcal{M} is locally expressed near the core Θ_0 as the hypersurface

$$\left\{ (z,\zeta,s,t) \in N_0 \times \Delta \times \Delta^{\dagger} : \sigma(z)\zeta^{m_0-ln_0} \left(\zeta^{n_0} + t\tau(z)\right)^l - s = 0 \right\},\$$

where m_0 and n_0 are the multiplicities of Θ_0 in X_0 and Y, respectively, σ is the standard section of $N_0^{\otimes (-m_0)}$ and τ is a core section of $N_0^{\otimes n_0}$ for Y. Furthermore, $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$ locally coincides with the restriction of the projection $N_0 \times \Delta \times \Delta^{\dagger} \to \Delta \times \Delta^{\dagger}$ to the hypersurface \mathcal{M} .

Note that the degeneration $\pi : M \to \Delta$ (that is, $\pi_0 : M_0 \to \Delta_0$) corresponds to the restriction of the projection $N_0 \times \Delta \times \{0\} \to \Delta \times \{0\}$ to the

hypersurface given by

$$\sigma(z)\zeta^{m_0} - s = 0, \quad \text{in } N_0 \times \Delta \times \{0\},$$

while $\pi_0^{\dagger} : M_0^{\dagger} \to \Delta_0^{\dagger}$ corresponds to the restriction of the projection $N_0 \times \{0\} \times \Delta^{\dagger} \to \{0\} \times \Delta^{\dagger}$ to the hypersurface given by

$$\sigma(z)\zeta^{m_0-ln_0} \left(\zeta^{n_0} + t\tau(z)\right)^l = 0, \quad \text{in } N_0 \times \{0\} \times \Delta^{\dagger}.$$

Now consider the hypersurface given by

$$\zeta = 0, \quad \text{in } N_0 \times \{0\} \times \Delta^{\dagger},$$

which is contained in M_0^{\dagger} . This hypersurface is nothing but $\Theta_0 \times \Delta_0^{\dagger}$, and the restriction of $\pi_0^{\dagger}: M_0^{\dagger} \to \Delta_0^{\dagger}$ to $\Theta_0 \times \Delta_0^{\dagger}$ coincides with the trivial degeneration $\pi_0^{\dagger}: \Theta_0 \times \Delta_0^{\dagger} \to \Delta_0^{\dagger}$ of Riemann surfaces. Note that the fiber $(\pi_0^{\dagger})^{-1}(t_0)$ over $t_0 \in \Delta_0^{\dagger}$ coincides with the stable component Θ . Since the restriction of F_1 to Θ coincides with the monodromy homeomorphism of this trivial degeneration, $F_1|_{\Theta}$ is isotopic to the identity map of Θ . Thus we have the following.

Proposition 7.2. Let $\Theta_1, \Theta_2, \ldots, \Theta_a$ be the stable components of the main fiber X_{0,t_0} of the deformation $\pi_{t_0} : M_{t_0} \to \Delta_{t_0}$. Then, for each $i = 1, 2, \ldots, a$, we have

$$F_1(\Theta_i) = \Theta_i,$$

and the restriction $F_1|_{\Theta_i} : \Theta_i \to \Theta_i$ is isotopic to the identity map of Θ_i .

Remark 7.3. In fact, Proposition 7.2 holds for barking families associated with simple crusts (not necessarily tame simple crusts).

8 Enlargements of tame simple crusts

Let $\pi: M \to \Delta$ be a linear degeneration of Riemann surfaces of genus $g \ge 1$ and $\Psi: \mathcal{M} \to \Delta \times \Delta^{\dagger}$ be a barking family for $\pi: \mathcal{M} \to \Delta$ associated with a tame simple crust Y with barking multiplicity l.

In this section, we study the restriction of the monodromy homeomorphism $F_1: X_{0,t_0} \to X_{0,t_0}$ to a barked component Ξ of the main fiber X_{0,t_0} . Recall that, as $t_0 \to 0$, the component Ξ approaches to a certain union of irreducible components of X_0 which contains Y^{red} . Unfortunately, unlike the case of stable components in Section 7, we cannot always construct a degeneration of Riemann surfaces such that Ξ coincides with a smooth fiber of it and that the degeneration itself can be identified with a certain restriction of $\pi_0^{\dagger} : M_0^{\dagger} \to \Delta_0^{\dagger}$. However, we can construct a degeneration of Riemann surfaces such that its restriction to the complement of a thin subset coincides with a certain restriction of π_0^{\dagger} , and that its singular fiber is identified with an "enlargement" of Y.

We introduce the concept of "enlargements" of tame simple crusts as follows. First let us define the enlargements of subbranches. Let **Br** be a branch of the singular fiber X_0 of the degeneration $\pi : M \to \Delta$ and **br** be a subbranch of **Br** for the tame simple crust Y. Express them as

$$\begin{cases} \mathbf{Br} = m_1 \Theta_1 + m_2 \Theta_2 + \dots + m_\lambda \Theta_\lambda, \text{ and} \\ \mathbf{br} = n_1 \Theta_1 + n_2 \Theta_2 + \dots + n_\nu \Theta_\nu, \text{ or } \emptyset, \end{cases}$$

where $0 \leq \nu \leq \lambda$, and $0 \leq n_i \leq m_i$, $i = 1, 2, \dots, \nu$.

First suppose that **br** is a subbranch of type A_l . Then, from the definition, it follows that $\overline{n}_{\nu+1}$ (= $r_{\nu}n_{\nu} - n_{\nu-1}$) may possibly be a negative integer. We define the decreasing sequence of nonnegative integers

$$n_{\nu} > \widetilde{n}_{\nu+1} > \widetilde{n}_{\nu+2} > \dots > \widetilde{n}_{\mu} > \widetilde{n}_{\mu+1} = 0$$

by the Euclidean division algorithm of negative type: namely, we choose the integers so that

$$\begin{cases} \widetilde{r}_{\nu} := \frac{n_{\nu-1} + \widetilde{n}_{\nu+1}}{n_{\nu}}, \ \widetilde{r}_{\nu+1} := \frac{n_{\nu} + \widetilde{n}_{\nu+2}}{\widetilde{n}_{\nu+1}}, \text{ and} \\ \widetilde{r}_i := \frac{\widetilde{n}_{i-1} + \widetilde{n}_{i+1}}{\widetilde{n}_i} \ (i = \nu + 2, \nu + 3, \dots, \mu) \end{cases}$$

are integers greater than or equal to 2. If $\nu = 0$ (that is, $\mathbf{br} = \emptyset$), then we set $\tilde{n}_{\nu+1} := 0$. We now consider the sequence

$$n_0, n_1, n_2, \ldots, n_{\nu}, \tilde{n}_{\nu+1}, \tilde{n}_{\nu+2}, \ldots, \tilde{n}_{\mu}, \tilde{n}_{\mu+1} = 0.$$

By Lemma 4.2, there exists a linear degeneration $\pi: \widetilde{T} \to \Delta$ with the singular fiber

$$n_0V_0 + n_1\Theta_1 + n_2\Theta_2 + \dots + n_{\nu-1}\Theta_{\nu-1} + n_{\nu}\widetilde{\Theta}_{\nu} + \widetilde{n}_{\nu+1}\widetilde{\Theta}_{\nu+1} + \dots + \widetilde{n}_{\mu}\widetilde{\Theta}_{\mu},$$

where V_0 is a copy of \mathbb{C} . We call the chain $\widetilde{\mathbf{br}} := n_1 \Theta_1 + n_2 \Theta_2 + \cdots + \widetilde{n}_{\mu} \widetilde{\Theta}_{\mu}$ the *enlargement* of the subbranch **br**.

Suppose that **br** is of type B_l or C_l . we then consider the sequence

$$n_0, n_1, n_2, \ldots, n_{\nu}, \widetilde{n}_{\nu+1} := 0.$$

Note that $\tilde{r}_{\nu} := (n_{\nu-1} + \tilde{n}_{\nu+1})/n_{\nu} = n_{\nu-1}/n_{\nu}$ is a positive integer. In fact, if **br** is of type B_l , then $\tilde{r}_{\nu} = n_{\nu-1}$. On the other hand, if **br** is of type C_l , then $\tilde{r}_{\nu} = r_{\nu} - 1$. By Lemma 4.2, there exists a linear degeneration $\pi : \tilde{T} \to \Delta$ with the singular fiber

$$n_0V_0 + n_1\Theta_1 + n_2\Theta_2 + \dots + n_{\nu-1}\Theta_{\nu-1} + n_{\nu}\widetilde{\Theta}_{\nu}$$

where V_0 is a copy of \mathbb{C} . We call the chain $\widetilde{\mathbf{br}} := n_1 \Theta_1 + n_2 \Theta_2 + \cdots + n_{\nu} \widetilde{\Theta}_{\nu}$ the *enlargement* of the subbranch **br**. By convention, we set $\mu := \nu$.

Remark 8.1. If a subbranch **br** is of both type A_l and B_l , then the enlargements of **br** defined above by the two methods coincide. In this case, the length ν of **br** is equal to λ . Note that subbranches of type A_l are not of type C_l .

Let \mathbf{Tk} be a trunk of the singular fiber X_0 of the degeneration $\pi : M \to \Delta$ and \mathbf{tk} be a subtrunk of \mathbf{Tk} for the tame simple crust Y. Express them as

$$\begin{cases} \mathbf{T}\mathbf{k} = m_1\Theta_1 + m_2\Theta_2 + \dots + m_\lambda\Theta_\lambda \text{ and} \\ \mathbf{t}\mathbf{k} = n_1\Theta_1 + n_2\Theta_2 + \dots + n_\lambda\Theta_\lambda. \end{cases}$$

We define the *enlargement* $\widetilde{\mathbf{tk}}$ of \mathbf{tk} as \mathbf{tk} itself, that is, we set $\widetilde{\mathbf{tk}} := \mathbf{tk}$. In fact, by Lemma 4.2, there exists a linear degeneration $\pi : \widetilde{T} \to \Delta$ with the singular fiber

$$n_0V_0 + n_1\Theta_1 + n_2\Theta_2 + \dots + n_\lambda\Theta_\lambda + n_{\lambda+1}U_{\lambda+1},$$

where V_0 and $U_{\lambda+1}$ are copies of \mathbb{C} .

Recall that \mathcal{M} is expressed near the core Θ_0 as the hypersurface given by

$$\sigma(z)\zeta^{m_0-ln_0}\left(\zeta^{n_0}+t\tau(z)\right)^l-s=0,\quad\text{in }N_0\times\Delta\times\Delta^\dagger,$$

and that Ψ coincides with the restriction of the projection map $N_0 \times \Delta \times \Delta^{\dagger} \rightarrow \Delta \times \Delta^{\dagger}$ to the hypersurface \mathcal{M} . Here σ is the standard section of the line bundle $N_0^{\otimes (-m_0)}$ satisfying

$$\operatorname{div}(\sigma) = \sum_{j=1}^{h} m_j p_j,$$

where p_1, p_2, \ldots, p_h are the intersection points on Θ_0 with the other irreducible components of X_0 . On the other hand, τ is a core section of the line bundle $N_0^{\otimes n_0}$ for Y satisfying

$$\operatorname{div}(\tau) = -\sum_{j=1}^{h} n_j p_j,$$

Note that, from the definition of tame simple crusts, the core section τ has no zeros. Then $\tau^{-1} (= 1/\tau)$ is a holomorphic section of $N_0^{\otimes (-n_0)}$ which has a zero of order n_j at p_j , j = 1, 2, ..., h, and $\tau^{-1}(z)\zeta^{n_0}$ defines a holomorphic function on N_0 . Now consider the hypersurface W_0 in $N_0 \times \Delta_0^{\dagger}$ defined by

$$\tau^{-1}(z)\zeta^{n_0} + t = 0,$$

where $\Delta_0^{\dagger} := \{0\} \times \Delta^{\dagger}$. Then the restriction of the projection map $N_0 \times \Delta_0^{\dagger} \to \Delta_0^{\dagger}$ to the hypersurface W_0 is a degeneration of punctured Riemann surfaces whose singular fiber is

$$n_0\Theta_0 + \sum_{j=1}^h n_j U_j,$$

where U_j is the fiber of N_0 over the point p_j (j = 1, 2, ..., h), that is, $U_j = \{(p_j, \zeta) \in N_0\}$.

Now we define the enlargement of the tame simple crust Y. Express Y as

$$Y := \sum_{i} n_i \Theta_i + \sum_{j} \mathbf{br}^{(j)} + \sum_{k} \mathbf{tk}^{(k)},$$

where Θ_i , $\mathbf{br}^{(j)}$ and $\mathbf{tk}^{(k)}$ are the cores, the subbranches and the subtrunks of Y, respectively. For each $\mathbf{br}^{(j)}$ (resp. $\mathbf{tk}^{(k)}$), let $\widetilde{\mathbf{br}}^{(j)}$ (resp. $\widetilde{\mathbf{tk}}^{(k)}$) be its enlargement defined as above. By the same argument as that for linear degenerations, we patch the tubular neighborhoods of the cores Θ_i and the enlargements $\widetilde{\mathbf{br}}^{(j)}$ and $\widetilde{\mathbf{tk}}^{(k)}$ to obtain a smooth complex surface \widetilde{M} . From the construction, it is easy to see that the holomorphic map $\widetilde{\pi} : \widetilde{M} \to \Delta^{\dagger}$ defined by the degeneration maps of these neighborhoods is a degeneration with the singular fiber

$$\widetilde{Y} := \sum_{i} n_i \Theta_i + \sum_{j} \widetilde{\mathbf{br}}^{(j)} + \sum_{k} \widetilde{\mathbf{tk}}^{(k)}.$$

We call \widetilde{Y} the *enlargement* of Y.

9 Constructing degenerations of barked components

Recall that the restriction $\pi_0^{\dagger}: M_0^{\dagger} \to \Delta_0^{\dagger}$ of $\Psi: \mathcal{M} \to \Delta \times \Delta^{\dagger}$ to the preimage $M_0^{\dagger} := \Psi^{-1}(\Delta_0^{\dagger})$ of $\Delta_0^{\dagger} := \{0\} \times \Delta^{\dagger}$ is not a family of complex curves, but that the central fiber $(\pi_0^{\dagger})^{-1}(0)$ coincides with the singular fiber X_0 of the original degeneration $\pi: \mathcal{M} \to \Delta$ and the general fiber $(\pi_0^{\dagger})^{-1}(t), t \neq 0$, coincides with the main fiber $X_{0,t}$ of the deformation $\pi_t: \mathcal{M}_t \to \Delta_t$.

We will show that the restriction of $\widetilde{\pi} : M \to \Delta^{\dagger}$ to the complement of a thin subset of \widetilde{M} coincides with a certain restriction of $\pi_0^{\dagger} : M_0^{\dagger} \to \Delta_0^{\dagger}$.

Lemma 9.1. For the enlargement

$$\widetilde{\mathbf{br}}^{(j)} = n_1 \Theta_1 + n_2 \Theta_2 + \dots + n_{\nu-1} \Theta_{\nu-1} + n_\nu \widetilde{\Theta}_\nu + \widetilde{n}_{\nu+1} \widetilde{\Theta}_{\nu+1} + \dots + \widetilde{n}_\mu \widetilde{\Theta}_\mu$$

of each subbranch $\mathbf{br}^{(j)}$ of Y, set $E^{(j)} := \widetilde{\Theta}_{\nu+1} \cup \widetilde{\Theta}_{\nu+2} \cup \cdots \cup \widetilde{\Theta}_{\mu} \cup \widetilde{U}_{\mu}$, where \widetilde{U}_{μ} is the fiber over the end⁵ of $\widetilde{\mathbf{br}}^{(j)}$ of the normal bundle \widetilde{N}_{μ} of $\widetilde{\Theta}_{\mu}$. Then

$$\widetilde{M}^{\times} := \widetilde{M} \setminus \bigcup_{j} E^{(j)}$$

⁵A point on $\widetilde{\Theta}_{\mu}$ away form the attachment point with $\widetilde{\Theta}_{\mu-1}$.

is naturally contained in M_0^{\dagger} . Moreover, the restriction of $\widetilde{\pi} : \widetilde{M} \to \Delta^{\dagger}$ to \widetilde{M}^{\times} can be identified with a certain restriction of $\pi_0^{\dagger} : M_0^{\dagger} \to \Delta_0^{\dagger}$.

Proof. The degeneration $\widetilde{\pi} : \widetilde{M} \to \Delta^{\dagger}$ is locally expressed near a core Θ_i as the restriction of the projection map $N_i \times \Delta_0^{\dagger} \to \Delta_0^{\dagger}$ to the hypersurface W_i in $N_i \times \Delta_0^{\dagger}$ defined by

$$\tau^{-1}(z)\zeta^{n_i} + t = 0,$$

while $\pi_0^{\dagger}: M_0^{\dagger} \to \Delta_0^{\dagger}$ is locally expressed as the restriction of the projection map of $N_i \times \Delta_0^{\dagger}$ to the hypersurface given by

$$\sigma(z)\tau^{l}(z)\zeta^{m_{i}-ln_{i}}\left(\tau^{-1}(z)\zeta^{n_{i}}+t\right)^{l}=0.$$

Thus \widetilde{M} is contained in M_0^{\dagger} near the core Θ_i , and the restrictions of $\widetilde{\pi} : \widetilde{M} \to \Delta^{\dagger}$ and $\pi_0^{\dagger} : M_0^{\dagger} \to \Delta_0^{\dagger}$. coincide.

We next consider the enlargement $\widetilde{\mathbf{br}}^{(j)}$ of each subbranch $\mathbf{br}^{(j)}$ of Y. Let Θ_i $(i = 1, 2, ..., \nu - 1)$ be the Riemann sphere contained in $\widetilde{\mathbf{br}}^{(j)}$ (or $\mathbf{br}^{(j)}$) as the irreducible component. Let $\Theta_i = U_i \cup V_i$ be an open covering by two copies U_i, V_i of \mathbb{C} with coordinates $w_i \in U_i \setminus \{0\}$ and $z_i \in V_i \setminus \{0\}$ satisfying $z_i = 1/w_i$. Then we obtain a line bundle N_i over Θ_i of degree $-r_i$ from $U_i \times \mathbb{C}$ and $V_i \times \mathbb{C}$ by identifying $(z_i, \zeta_i) \in (V_i \setminus \{0\}) \times \mathbb{C}$ with $(w_i, \eta_i) \in (U_i \setminus \{0\}) \times \mathbb{C}$ via

$$g_i: z_i = \frac{1}{w_i}, \quad \zeta_i = w_i^{r_i} \eta_i,$$

where $r_i = (n_{i-1} + n_{i+1})/n_i = (m_{i-1} + m_{i+1})/m_i$.

The degeneration $\widetilde{\pi} : \widetilde{M} \to \Delta^{\dagger}$ is locally expressed near Θ_i as the restriction of the projection map $N_i \times \Delta_0^{\dagger} \to \Delta_0^{\dagger}$ to the hypersurface H_i in $N_i \times \Delta_0^{\dagger}$ defined by

$$\begin{cases} w_i^{n_{i-1}} \eta_i^{n_i} + t = 0, & (w_i, \eta_i, t) \in U_i \times \mathbb{C} \times \Delta_0^{\dagger}, \\ z_i^{n_{i+1}} \zeta_i^{n_i} + t = 0, & (z_i, \zeta_i, t) \in V_i \times \mathbb{C} \times \Delta_0^{\dagger}. \end{cases}$$

On the other hand, $\pi_0^{\dagger} : M_0^{\dagger} \to \Delta_0^{\dagger}$ is locally expressed as the restriction of the projection map of $N_i \times \Delta_0^{\dagger}$ to the hypersurface given by

$$\begin{cases} w_i^{m_{i-1}-ln_{i-1}} \eta_i^{m_i-ln_i} (w_i^{n_{i-1}} \eta_i^{n_i} + t)^l = 0, & (w_i, \eta_i, t) \in U_i \times \mathbb{C} \times \Delta_0^{\dagger}, \\ z_i^{m_{i+1}-ln_{i+1}} \zeta_i^{m_i-ln_i} (z_i^{n_{i+1}} \zeta_i^{n_i} + t)^l = 0, & (z_i, \zeta_i, t) \in V_i \times \mathbb{C} \times \Delta_0^{\dagger}. \end{cases}$$

Thus \widetilde{M} is contained in M_0^{\dagger} near $\widetilde{\mathbf{br}}^{(j)} \setminus \bigcup_j E^{(j)}$, and the restrictions of $\widetilde{\pi}$: $\widetilde{M} \to \Delta^{\dagger}$ and $\pi_0^{\dagger} : M_0^{\dagger} \to \Delta_0^{\dagger}$. coincide.

By the same argument as that for $\widetilde{\mathbf{br}}^{(j)}$, we see that \widetilde{M} is contained in M_0^{\dagger} near the subtrunk $\mathbf{tk}^{(k)} \ (= \widetilde{\mathbf{tk}}^{(k)})$ and the restrictions of $\widetilde{\pi} : \widetilde{M} \to \Delta^{\dagger}$ and $\pi_0^{\dagger} : M_0^{\dagger} \to \Delta_0^{\dagger}$. coincide. This completes the proof of the assertion. \Box

Under the identification of $\tilde{\pi} : \widetilde{M}^{\times} \to \Delta^{\dagger}$ with a certain restriction of $\pi_0^{\dagger} : M_0^{\dagger} \to \Delta_0^{\dagger}$ in Lemma 9.1, the central fiber of $\tilde{\pi} : \widetilde{M}^{\times} \to \Delta^{\dagger}$ corresponds to the singular curve obtained by puncturing the simple crust Y at the end of each subbranch, while a general fiber over $t_0 \in \Delta^{\dagger} \setminus \{0\}$ is the disjoint union of punctured Riemann surfaces obtained from the barked components of the main fiber X_{0,t_0} of the deformation $\pi_{t_0} : M_{t_0} \to \Delta_{t_0}$.

Suppose that X_{0,t_0} has exactly one barked component, and denote it by Ξ . Then the restriction of the monodromy homeomorphism F_1 of the barking family to the punctured barked component (that is, the Riemann surface obtained by puncturing the barked component Ξ at the attachment points with other irreducible components) forms a self-homeomorphism, and it coincides with the restriction of the monodromy homeomorphism of the degeneration of Riemann surfaces $\tilde{\pi} : \widetilde{M} \to \Delta^{\dagger}$ to the punctured general fiber of $\tilde{\pi} : \widetilde{M}^{\times} \to \Delta^{\dagger}$ up to isotopy. Since a self-homeomorphism of a punctured real surface uniquely induces a self-homeomorphism of its compactification, the monodromy homeomorphism of the degeneration of Riemann surfaces $\tilde{\pi} : \widetilde{M} \to \Delta^{\dagger}$ is isotopic to $F_1|_{\Xi} : \Xi \to \Xi$. Hence, we have the following.

Proposition 9.2. We denote by $(F_1, F_2, \ldots, F_c; f)$ the monodromy set of Ψ . Suppose that the main fiber X_{0,t_0} of the deformation $\pi_{t_0} : M_{t_0} \to \Delta_{t_0}$ has exactly one barked component, say Ξ . Then we have $F_1(\Xi) = \Xi$, and the isotopy class of $F_1|_{\Xi} : \Xi \to \Xi$ is conjugate to the topological monodromy of a degeneration of Riemann surfaces whose singular fiber is the enlargement of the tame simple crust Y. In particular, $F_1|_{\Xi} : \Xi \to \Xi$ is isotopic to a pseudo-periodic homeomorphism of negative twist.

Now, let us consider the general case: let $\Xi_1, \Xi_2, \ldots, \Xi_b$ be the barked components of X_{0,t_0} . Note that $\widetilde{\pi} : \widetilde{M} \to \Delta^{\dagger}$ is not necessarily a degeneration of *Riemann surfaces* but is a degeneration of *disjoint unions of Riemann* surfaces (see Remark 9.4 below). In other words, each general fiber is a disjoint union of of Riemann surfaces. In this case, the fiber $\tilde{\pi}^{-1}(t_0)$ over $t_0 \in \Delta^{\dagger}$ consists of $\Xi_1, \Xi_2, \ldots, \Xi_b$. By an argument similar to the above, the monodromy homeomorphism of $\tilde{\pi} : \widetilde{M} \to \Delta^{\dagger}$ is isotopic to

$$F_1\big|_{\Xi_1\cup\Xi_2\cup\cdots\cup\Xi_b}:\Xi_1\cup\Xi_2\cup\cdots\cup\Xi_b\to\Xi_1\cup\Xi_2\cup\cdots\cup\Xi_b$$

Hence, we have the following.

Proposition 9.3. We denote by $(F_1, F_2, \ldots, F_c; f)$ the monodromy set of Ψ . Let $\Xi_1, \Xi_2, \ldots, \Xi_b$ be the barked components of the main fiber X_{0,t_0} of the deformation $\pi_{t_0} : M_{t_0} \to \Delta_{t_0}$. Then F_1 cyclically permutes $\Xi_1, \Xi_2, \ldots, \Xi_b$, and

$$F_1\big|_{\Xi_1\cup\Xi_2\cup\cdots\cup\Xi_b}:\Xi_1\cup\Xi_2\cup\cdots\cup\Xi_b\to\Xi_1\cup\Xi_2\cup\cdots\cup\Xi_b$$

coincides with the monodromy homeomorphism of a degeneration of disjoint unions of b Riemann surfaces whose singular fiber is the enlargement of the tame simple crust Y, up to isotopy and conjugacy. In particular, for each k = $1, 2, \ldots, b, F_1^b|_{\Xi_k} : \Xi_k \to \Xi_k$ is isotopic to a pseudo-periodic homeomorphism of negative twist.

Remark 9.4. Given a degeneration $\pi : M \to \Delta$ of Riemann surfaces of genus g with singular fiber $X_0 = \sum_i m_i \Theta_i$, for a positive integer $b \ge 2$, the composition $\rho \circ \pi : M \to \Delta$ with the holomorphic map $\rho : \Delta \ni s \mapsto s^b \in \Delta$ is a degeneration of disjoint unions of b Riemann surfaces of genus g with singular fiber $bX_0 = \sum_i (m_i b) \Theta_i$. Furthermore, its monodromy homeomorphism F cyclically permutes the b Riemann surfaces, and the restriction of F^b to one of the Riemann surfaces coincides with the monodromy homeomorphism of $\pi : M \to \Delta$ up to isotopy and conjugacy.

10 Singular fibers appearing in splitting families

In this section, we state some lemmas which help us to determinate the topological types of singular fibers appearing in splitting families. The proofs of the lemmas can be found in [O].

The first lemma is for general splitting families. For a singular fiber X, we denote by $\mathcal{E}(X)$ the *Euler contribution* of X, that is, we set $\mathcal{E}(X) := e(X) - 2(1 - g)$, where e(X) is the topological Euler characteristic of the underlying reduced curve of X.

Lemma 10.1. Let $\pi : M \to \Delta$ be a degeneration of Riemann surfaces of genus $g \ge 1$ with the singular fiber X_0 and let $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$ be a splitting family of $\pi : M \to \Delta$ such that X_0 splits into singular fibers X_1, X_2, \ldots, X_N ($N \ge 2$). Then, we have the following.

(1) $\mathcal{E}(X_i) \ge 0$ for each $i = 0, 1, \dots, N$. (2) $\mathcal{E}(X_0) = \sum_{i=1}^N \mathcal{E}(X_i)$.

Now let $\pi : M \to \Delta$ be a linear degeneration of Riemann surfaces with singular fiber X_0 . Suppose that X_0 has a simple crust Y with barking multiplicity l. Then, we have a barking family $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$ of $\pi : \mathcal{M} \to \Delta$ associated with Y.

Lemma 10.2. Every subordinate fiber X appearing in $\Psi : \mathcal{M} \to \Delta \times \Delta^{\dagger}$ is a reduced curve at most with A-singularities⁶. In particular, $\mathcal{E}(X) \ge 1$, where the equality holds exactly when X is a Lefschetz fiber.

Let **br** be a subbranch of a branch **Br** of X_0 for Y. Express them as

$$\begin{cases} \mathbf{Br} = m_1 \Theta_1 + m_2 \Theta_2 + \dots + m_\lambda \Theta_\lambda, \text{ and} \\ \mathbf{br} = n_1 \Theta_1 + n_2 \Theta_2 + \dots + n_\nu \Theta_\nu, \text{ or } \emptyset, \end{cases}$$

where⁷ $0 \leq \nu \leq \lambda$, and $0 \leq ln_i \leq m_i$, $i = 1, 2, ..., \nu$. Let Θ_0 be the core intersecting the component Θ_1 and let m_0 (resp. n_0) be the multiplicity of Θ_0 in X_0 (resp. Y). We say that **br** is *proportional* if $\nu = \lambda$ and

$$\frac{n_0}{m_0} = \frac{n_1}{m_1} = \dots = \frac{n_\lambda}{m_\lambda}.$$

⁶An A-singularity is a singularity analytically equivalent to $y^2 = x^{e+1}$ for some positive integer e.

⁷We set $\nu := 0$ if $\mathbf{br} = \emptyset$. See the paragraph above Definition 5.1.

Now we assume that the singular fiber X_0 is stellar, that is, X_0 consists of exactly one core Θ_0 and some branches emanating from Θ_0 . The following two lemmas give us the number of the subordinate fibers and that of their singularities.

Lemma 10.3. Suppose that (i) the core Θ_0 is a Riemann sphere, (ii) X_0 has three branches, (iii) the core section τ for Y has no zero, and (iv) Y has no proportional subbranches. Let m_0 (resp. n_0) denote the multiplicity of the core Θ_0 in X_0 (resp. Y). Then we have the following.

- (a) Each deformation of the degeneration has exactly \bar{n}_0 subordinate fibers.
- (b) Each subordinate fiber in (a) has c singularities.

Here $c := \text{gcd}(m_0, n_0)$, the greatest common divisor of m_0 and n_0 , and $\bar{n}_0 := n_0/c$.

Lemma 10.4. Suppose that (i) the core Θ_0 is a Riemann sphere, (ii) X_0 has three branches, (iii) the core section τ for Y has no zero, and (iv) Y has a proportional subbranch $\mathbf{br} = n_1\Theta_1 + n_2\Theta_2 + \cdots + n_\lambda\Theta_\lambda$ of a branch $\mathbf{Br} = m_1\Theta_1 + m_2\Theta_2 + \cdots + m_\lambda\Theta_\lambda$ of X_0 . Then no other subbranches are proportional, and moreover we have the following.

- (a) Each deformation of the degeneration has exactly \bar{n}_{λ} subordinate fibers.
- (b) Each subordinate fiber in (a) has c singularities.

Here $c := \text{gcd}(m_{\lambda}, n_{\lambda})$, the greatest common divisor of m_{λ} and n_{λ} , and $\bar{n}_{\lambda} := n_{\lambda}/c$.

11 Barking families giving the same splitting

As an application of our results, we show an interesting example of two splitting families for one degeneration which give the same splitting (that is, the topological types of the singular fibers appearing in the two splitting families coincide) and which have, nevertheless, the different topological monodromies. This example indicates that the topological monodromies of splitting families play a very important role when we classify "topologically distinct" splitting families.

Let us consider the linear degeneration $\pi: M \to \Delta$ of Riemann surfaces of genus two whose singular fiber is stellar and is of the form

$$X_0 = 10\Theta_0 + \sum_{j=1}^3 \mathbf{Br}^{(j)},$$

where the core Θ_0 is a Riemann sphere and the three branches, attached to Θ_0 , are as follows:

$$\begin{cases} \mathbf{Br}^{(1)} = 5\Theta_1^{(1)}, \\ \mathbf{Br}^{(2)} = 4\Theta_1^{(2)} + 2\Theta_2^{(2)}, \\ \mathbf{Br}^{(3)} = 1\Theta_1^{(3)}. \end{cases}$$

Here the core Θ_0 intersects $\Theta_1^{(1)}$, $\Theta_1^{(2)}$, and $\Theta_1^{(3)}$.

Barking 1. We first define the connected subdivisor Y_1 as

$$Y_1 = 6\Theta_0 + \sum_{j=1}^{3} \mathbf{br}_1^{(j)}, \quad \text{where} \quad \begin{cases} \mathbf{br}_1^{(1)} = 3\Theta_1^{(1)}, \\ \mathbf{br}_1^{(2)} = 2\Theta_1^{(2)}, \\ \mathbf{br}_1^{(3)} = 1\Theta_1^{(3)}. \end{cases}$$

See Figure 5. Lemma 5.3 ensures that Y_1 has a core section over Θ_0 which has no zeros, and

- $\mathbf{br}_1^{(1)}$ is a subbranch of $\mathbf{Br}_1^{(1)}$ of type A_1 ,
- $\mathbf{br}_1^{(2)}$ is a subbranch of $\mathbf{Br}_1^{(2)}$ of type A_1 ,
- $\mathbf{br}_1^{(3)}$ is a subbranch of $\mathbf{Br}_1^{(3)}$ of type B_1 .

Therefore Y_1 is a tame simple crust of X_0 with barking multiplicity 1 and induces a barking family $\Psi_1 : \mathcal{M}_1 \to \Delta \times \Delta^{\dagger}$, in which the singular fiber X_0 is deformed to the main fiber X'_0 as depicted in Figure 5. Then the set of the subordinate fibers in each deformation of the degeneration $\pi : \mathcal{M} \to \Delta$ for the barking family $\Psi_1 : \mathcal{M}_1 \to \Delta \times \Delta^{\dagger}$ consists of three Lefschetz fibers. In fact, since $\mathbf{br}^{(1)}$ is proportional and Y_1 satisfies the assumptions of Lemma



Figure 5: The barking family associated with the tame simple crust Y_1 with barking multiplicity 1.

10.4, we see that there are exactly three subordinate fibers and each of them has exactly one singularity. On the other hand, we have

$$\mathcal{E}(X_0) = 8 \quad \text{and} \quad \mathcal{E}(X'_0) = 5,$$

where $\mathcal{E}(X)$ denotes the Euler contribution⁸ of a singular fiber X. Thus, by Lemma 10.1, the sum of the Euler contributions of the subordinate fibers is equal to three. Furthermore, from Lemma 10.2, we see that the Euler contribution of each of the three subordinate fibers is equal to one, and that the three subordinate fibers are all Lefschetz fibers.

Barking 2. We next define the connected subdivisor Y_2 as

$$Y_2 = 3\Theta_0 + \sum_{j=1}^{3} \mathbf{br}_2^{(j)}, \quad \text{where} \quad \begin{cases} \mathbf{br}_2^{(1)} = 1\Theta_1^{(1)}, \\ \mathbf{br}_2^{(2)} = 1\Theta_1^{(2)}, \\ \mathbf{br}_2^{(3)} = 1\Theta_1^{(3)}. \end{cases}$$

See Figure 6. Lemma 5.3 ensures that Y_2 has a core section over Θ_0 which has no zeros, and

⁸That is, $\mathcal{E}(X) := e(X) - 2(1 - g)$, where e(X) is the topological Euler characteristic of the underlying reduced curve of X.



Figure 6: The barking family associated with the tame simple crust Y_2 with barking multiplicity 1.

- $\mathbf{br}_2^{(1)}$ is a subbranch of $\mathbf{Br}_2^{(1)}$ of type A_1 ,
- $\mathbf{br}_2^{(2)}$ is a subbranch of $\mathbf{Br}_2^{(2)}$ of type A_1 ,
- $\mathbf{br}_2^{(3)}$ is a subbranch of $\mathbf{Br}_2^{(3)}$ of type B_1 .

Therefore Y_2 is a tame simple crust of X_0 with barking multiplicity 1 and induces a barking family $\Psi_2 : \mathcal{M}_2 \to \Delta \times \Delta^{\dagger}$, in which the singular fiber X_0 is deformed to the main fiber X_0'' as depicted in Figure 6. Then the set of the subordinate fibers in each deformation of the degeneration $\pi : M \to \Delta$ for the barking family $\Psi_2 : \mathcal{M}_2 \to \Delta \times \Delta^{\dagger}$ consists of three Lefschetz fibers. In fact, since Y_2 satisfies the assumptions of Lemma 10.3, we see that there are exactly three subordinate fibers and each of them has exactly one singularity. On the other hand, we have

$$\mathcal{E}(X_0) = 8 \quad \text{and} \quad \mathcal{E}(X_0'') = 5.$$

Thus by the same argument as that for Y_1 , we see that the three subordinate fibers are all Lefschetz fibers.

Comparison. The main fibers X'_0 and X''_0 appearing in the deformations of the barking families $\Psi_1 : \mathcal{M}_1 \to \Delta \times \Delta^{\dagger}$ and $\Psi_2 : \mathcal{M}_2 \to \Delta \times \Delta^{\dagger}$, respectively, apparently have different topological types. However, both of them turn into Lefschetz fibers, by successive blowing-downs. To be more precise, recall that every (-1)-curve in a complex surface M is preserved under an arbitrary deformation of M by Kodaira's stability theorem [K]. Namely, there exists an analytic family of (-1)-curves in \mathcal{M}_i for each i = 1, 2. Furthermore, by [FN], we can blow down them simultaneously and then the resulting family is a splitting family of the degeneration obtained from $\pi: M \to \Delta$ by a blowing-down. Repeating this process four times, we obtain a splitting family $\overline{\Psi}_i: \overline{\mathcal{M}}_i \to \Delta \times \Delta^{\dagger}$ of the relatively minimal degeneration $\overline{\pi}: \overline{M} \to \Delta$ (i = 1, 2). In the splitting family $\overline{\Psi}_1$ (resp. $\overline{\Psi}_2$), the singular fiber \overline{X}'_0 (resp. \overline{X}''_0), which is obtained by blowing down the main fiber X'_0 (resp. X''_0), is a Lefschetz fiber. Hence, both $\overline{\Psi}_1$ and $\overline{\Psi}_2$ split the singular fiber of the minimal degeneration $\overline{\pi}: \overline{M} \to \Delta$ into four Lefschetz fibers. In particular, they give the same splitting.

Now we investigate the monodromy set of these splitting families. Note that the Lefschetz fiber $\overline{X'_0}$ is obtained from the unique barked component Ξ of the main fiber X'_0 by identifying the two attachment points on it in the above blowing-down process. Since the complement of the family of (-1)-curves is preserved under the simultaneous blowing-downs, the monodromy homeomorphism F' on the singular fiber $\overline{X'_0}$ of the splitting family $\overline{\Psi}_1: \overline{\mathcal{M}}_1 \to \Delta \times \Delta^{\dagger}$ is induced from that on the the barked component Ξ of the main fiber X'_0 . From Theorem 6.2, we see that the monodromy homeomorphism on the the barked component Ξ corresponds to the topological monodromy of the degeneration whose singular fiber is the enlargement Y_1 of Y_1 . Here, the multiplicity of the core Θ_0 in Y_1 is six, and so is that in Y_1 . Since the topological monodromy of a degeneration with stellar singular fiber whose core has multiplicity m is periodic of order m, the monodromy homeomorphism F' corresponds to a periodic mapping classes of order six. Similarly, the monodromy homeomorphism F'' of the singular fiber $\overline{X_0''}$ corresponds to the topological monodromy of the degeneration whose singular fiber is the enlargement \widetilde{Y}_2 of Y_2 , that is, a periodic mapping class of order three. Thus, the monodromy homeomorphisms F' and F'' are distinct up to isotopy and conjugacy. On the other hand, since the three subordinate fibers in the respective barking family form one subordinate tassel, the monodromy homeomorphism F' does not correspond to the monodromy homeomorphism of the subordinate tassel of $\overline{\Psi}_2$. Thus the monodromy set of the splitting families $\overline{\Psi}_1$ and $\overline{\Psi}_2$ are distinct up to isotopy and conjugacy. Hence, we have the following.

Proposition 11.1. There exist two splitting families for one degeneration that have different topological monodromies, although they give the same splitting.

Acknowledgements

The author would like to express his deep gratitude to his supervisor Osamu Saeki for helpful suggestions and warm encouragement. The author would also like to thank Shigeru Takamura for insightful comments.

References

- [ES] C.J. Earle and P.L. Sipe, Families of Riemann surfaces over the punctured disk, Pacific J. Math. 150 (1991), 79–96.
- [FN] A. Fujiki and S. Nakano, Supplement to "On the inverse of monoidal transformation", Publ. Res. Inst. Math. Sci. 7 (1971/72), 637–644.
- [GWPL] C.G. Gibson, K. Wirthmüller, A.A. du Plessis and E.J.N. Looijenga, *Topological Stability of Smooth Mappings*, Lecture Notes in Math., Vol. 552, Springer-Verlag, 1976.
- [K] K. Kodaira, On stability of compact submanifolds of complex manifolds, Amer. J. Math. 85 (1963), 79–94.
- [MM] Y. Matsumoto and J.M. Montesinos-Amilibia, Pseudo-periodic Maps and Degeneration of Riemann Surfaces, Lecture Notes in Math., Vol. 2030, Springer-Verlag, 2011.
- [O] T. Okuda, Singular fibers in barking families of degenerations of elliptic curves, preprint (arXiv:1205.1592 [math.GT]), to appear in "Singularities in Geometry and Topology 2011", Adv. Stud. Pure Math.

- [ST] H. Shiga and H. Tanigawa, On the Maskit coordinates of Teichmüller spaces and modular transformations, Kodai Math. J. 12 (1989), 437– 443.
- [Ta3] S. Takamura, Towards the Classification of Atoms of Degenerations, III — Splitting Deformations of Degenerations of Complex Curves, Lecture Notes in Math., Vol. 1886, Springer-Verlag, 2006.

Takayuki OKUDA

GRADUATE SCHOOL OF MATHEMATICS, KYUSHU UNIVERSITY, MOTOOKA 744, NISHI-KU, FUKUOKA 819-0395, JAPAN *E-mail address*: t-okuda@math.kyushu-u.ac.jp