

On the complex K-group of certain manifold

Fujino, Tsutomu

Department of Mathematics, Fukuoka Technological University

Ishikawa, Nobuhiro

Department of Mathematics, College of General Education, Kyushu University

Kamata, Masayoshi

Department of Mathematics, College of General Education, Kyushu University

<https://doi.org/10.15017/1448953>

出版情報：九州大学教養部数学雑誌. 9 (1), pp.1-6, 1973-11. 九州大学教養部数学教室
バージョン：
権利関係：

On the complex K -group of certain manifold

By

Tsutomu FUJINO, Nobuhiro ISHIKAWA
and Masayoshi KAMATA

(Received Jul. 28, 1973)

1. Let D_p be a dihedral group of order $2p$ which is generated by g of order p and t of order 2 such that $tgt = g^{-1}$. We consider an action of D_p over a product space $S^{2l+1} \times S^m \subset C^{l+1} \times R^{m+1}$ of spheres, given by

$$(1.1) \quad g^k t^j (z, x) = (\rho^k c^j(z), (-1)^j x),$$

where c is the conjugation and $\rho = \exp 2\pi i / p$. Denote by $D_p(l, m) = (S^{2l+1} \times S^m) / D_p$ the orbit space [3]. We are concerned with the complex K -group of $D_p(2k+1, 2n+1)$ where p is an odd prime.

Consider an action of Z_2 , a cyclic group of order 2, over the complex K -group $K(X)$, given by the conjugation automorphism $[\xi] = [\bar{\xi}]$. Let $K(X)^{Z_2}$ be the invariant subgroup of $K(X)$ under the involution. In this paper we obtain

THEOREM 1.1. *Suppose that p is an odd prime. Then, there exists an isomorphism*

$$\tilde{K}(D_p(2k+1, 2n+1)) \cong Z \oplus \tilde{K}(L^{2k+1}(p))^{Z_2} \oplus \tilde{K}(RP^{2n+1}),$$

where $L^{2k+1}(p)$ is a standard $(4k+3)$ -dimensional lens space and RP^{2n+1} is a $(2n+1)$ -dimensional real projective space.

2. Denote by $K_G(X)$ the equivariant K -group of G -space X . It is well-known that if the action of G is free then $K_G(X) \cong K(X/G)$. There is a canonical homomorphism from the representation ring $R(G)$ to $K_G(X)$ which maps a representation space M to a equivariant G -bundle $X \times M$. If X is free G -space then there is a homomorphism

$$\pi: R(G) \longrightarrow K_G(X) \cong K(X/G).$$

Let $\alpha: H \longrightarrow G$ be a homomorphism and $f: Y \longrightarrow X$ be an equivariant map from H -space Y to G -space X , that is, $f(h \cdot y) = \alpha(h) \cdot f(y)$. The equivariant map f induces the homomorphism

$$f^!: K_G(X) \longrightarrow K_H(Y) \quad [6].$$

We take a Z_2 -action on S^m by

$$t^l \cdot x = (-1)^l x, \quad t \in Z_2 \text{ a generator,}$$

a Z_p -action on S^{2l+1} by

$$g^l \cdot z = \rho^l z, \quad \rho = \exp(2\pi\sqrt{-1}/p), \quad g \in Z_p \text{ a generator}$$

and a D_p -action on $S^{2l+1} \times S^m$ given by (1.1) in section 1. We have these orbit spaces an m -dimensional real projective space RP^m , a $(2l+1)$ -dimensional lens space $L^l(p)$ and $D_p(l, m)$. There exist equivariant maps

$$\begin{aligned} i: S^{2l+1} &\longrightarrow S^{2l+1} \times S^m, \quad i(z) = (z, (1, 0, \dots, 0)), \\ j: S^m &\longrightarrow S^{2l+1} \times S^m, \quad j(x) = ((1, 0, \dots, 0), x) \end{aligned}$$

and

$$p: S^{2l+1} \times S^m \longrightarrow S^m, \quad p(z, x) = x$$

compatible with injections $\tilde{i}: Z_p \longrightarrow D_p$, $\tilde{j}: Z_2 \longrightarrow D_p$ and a projection $\tilde{p}: D_p \longrightarrow Z_2$ respectively. It follows immediately that

$$(2.1) \quad j^! p^! = 1.$$

Let H be a normal subgroup of a finite group G and A be a unitary representation of H . The induced representation A^G is defined as follows

$$(2.2) \quad A^G(g) = \begin{pmatrix} A(t_1 g t_1) & A(t_1 g t_2) & \cdots & A(t_1 g t_n) \\ A(t_2 g t_1) & A(t_2 g t_2) & \cdots & A(t_2 g t_n) \\ \vdots & \vdots & & \vdots \\ A(t_n g t_1) & A(t_n g t_2) & \cdots & A(t_n g t_n) \end{pmatrix}$$

where $G/H = \{\{t_1\}, \{t_2\}, \dots, \{t_n\}\}$ and $A(t_i g t_j) = 0$ if $t_i g t_j \notin H$.

Denote by L a standard representation space with $(\exp(2\pi\sqrt{-1}/p))$. For a standard Z_p -space S^{2l+1} with $S^{2l+1}/Z_p = L^l(p)$, we put

$$\xi_l = S^{2l+1} \times_{Z_p} L.$$

N. Mohammed [5] obtained that

$$K_{Z_p}(S^{2l+1}) \cong K(L^l(p)) \cong Z[\xi_l] / (\xi_l^p - 1, (\xi_l - 1)^{l+1}).$$

Hence, the homomorphism $\pi: R(Z_p) \longrightarrow K_{Z_p}(S^{2l+1})$ is surjective. Then, we define the homomorphism

$$i_*: K_{Z_p}(S^{2l+1}) \longrightarrow K_{D_p}(S^{2l+1} \times S^m)$$

by $i_*(S^{2l+1} \times M) = S^{2l+1} \times S^m \times M^{D_p}$, where M is representation space of Z_p and M^{D_p} is the induced representation space.

We consider a Z_2 -action over $K_{Z_p}(S^{2l+1})$ given by

$$t(S^{2l+1} \times M) = S^{2l+1} \times \bar{M},$$

where \bar{M} is a conjugate representation space of M and t is a generator of Z_2 .

Let $K_{Z_p}(S^{2l+1})^{Z_2}$ be the invariant subgroup under the Z_2 -action. Then, we have

PROPOSITION 2.1. For $\gamma \in K_{Z_p}(S^{2l+1})^{Z_2}$,

$$i!i_*(\gamma) = 2\gamma.$$

PROOF. Suppose that $\gamma = S^{2l+1} \times M \in K_{Z_p}(S^{2l+1})^{Z_2}$, that is, $\bar{M} = M$. Let A be the representation of M . Then, $i_*(\gamma) = S^{2l+1} \times S^m \times M^{D_p}$, where M^{D_p} is the induced representation space of M . The representation A^{D_p} of M^{D_p} is given as follows:

$$A^{D_p}(g) = \begin{pmatrix} A(g) & 0 \\ 0 & \bar{A}(g) \end{pmatrix}, \quad g \in Z_p.$$

Since $\bar{M} = M$ and $\bar{A} = A$,

$$i!i_*(\gamma) = 2\gamma. \quad \text{q. e. d.}$$

THEOREM 2.2. The homomorphism

$$\theta: \tilde{K}_{Z_p}(S^{2l+1})^{Z_2} \oplus \tilde{K}_{Z_2}(S^m) \longrightarrow \tilde{K}_{D_p}(S^{2l+1} \times S^m)$$

given by $\theta(\gamma, \nu) = i_*(\gamma) + p!(\nu)$ is injective.

PROOF. Suppose that $\theta(\gamma, \nu) = 0$. Since $i!p! = 0$, we have $i!i_*(\gamma) = 0$. Hence, from Proposition 2.1, it follows that $\gamma = 0$. On the other hand, from (2.1), we have $\nu = j!p!(\nu) = 0$. q. e. d.

3. The manifold $D_p(l, m)$ is homeomorphic to an orbit space $(L^l(p) \times S^m)/Z_2$, where a Z_2 -action on $L^l(p) \times S^m$ is given by

$$t^l([z], x) = ([c^l(z)], (-1)^l x),$$

where t is a generator of Z_2 . Denote by (C_i, D_i) a cell of $(L^l(p) \times S^m)/Z_2$ represented by a standard cell C_i of $L^l(p)$ and a standard cell D_i of S^m

and by (c^i, d^j) a dual cochain of (C_i, D_j) . The coboundary relations are given by

$$\begin{aligned}\partial(c^{2i+1}, d^j) &= p(c^{2(i+1)}, d^j) + \{(-1)^i + (-1)^j\}(c^{2i+1}, d^{j+1}), \\ \partial(c^{2i}, d^j) &= \{(-1)^i + (-1)^{j+1}\}(c^{2i}, d^{j+1}).\end{aligned}$$

Therefore, we have the following.

PROPOSITION 3.1. *The integral cohomology group $\tilde{H}^*(D_p(2k+1, 2n+1); Z)$ is a direct sum of the following groups:*

*free groups generated by (c^0, d^{2n+1}) , (c^{4k+3}, d^0) and (c^{4k+3}, d^{2n+1}) ,
torsion groups generated by (c^0, d^{2j}) and (c^{4k+3}, d^{2j}) whose orders are 2
and torsion groups generated by (c^{4i}, d^0) and (c^{4i}, d^{2n+1}) whose orders are p ,
where $1 \leq j \leq n$ and $1 \leq i \leq k$.*

Denote by Y the $(4k+2n+3)$ -skeleton of $D_p(2k+1, 2n+1)$. Then, we have

$$(3.1) \quad \tilde{H}^i(D_p(2k+1, 2n+1)/Y; Z) = \begin{cases} Z & i = 4k+2n+4, \\ 0 & \text{otherwise} \end{cases}$$

and

$$(3.2) \quad \tilde{H}^i(Y; Z) \sim \begin{cases} \tilde{H}^i(D_p(2k+1, 2n+1); Z) & i \leq 4k+2n+3, \\ 0 & \text{otherwise.} \end{cases}$$

PROPOSITION 3.2. *There exists a short exact sequence*

$$0 \rightarrow Z \rightarrow \tilde{K}(D_p(2k+1, 2n+1)) \rightarrow \tilde{K}(Y) \rightarrow 0.$$

PROOF. Consider the exact sequence of K -groups with respect to a pair $(D_p(2k+1, 2n+1), Y)$,

$$\begin{aligned}\cdots \rightarrow \tilde{K}^{-1}(D_p(2k+1, 2n+1)) &\xrightarrow{i^!} \tilde{K}^{-1}(Y) \rightarrow \tilde{K}(D_p(2k+1, 2n+1)/Y) \\ &\rightarrow \tilde{K}(D_p(2k+1, 2n+1)) \rightarrow \tilde{K}(Y) \rightarrow \tilde{K}^1(D_p(2k+1, 2n+1)/Y) \rightarrow \cdots\end{aligned}$$

Note that from (3.1) we have

$$\tilde{K}^i(D_p(2k+1, 2n+1)/Y) \cong \begin{cases} Z & \text{if } i \text{ is even,} \\ 0 & \text{if } i \text{ is odd.} \end{cases}$$

From the discussion of the Atiyah-Hirzebruch spectral sequence for $\tilde{K}^{-1}(X)$ with $E_2^{s,t}(X) \cong \tilde{H}^s(X; \tilde{K}^t(S^0))$ [2], $X = D_p(2k+1, 2n+1)$ or Y , we have the following,

the free part of $\tilde{K}^{-1}(X) \cong Z \oplus Z$.

Hence, it follows that $i^! : \tilde{K}^{-1}(D_p(2k+1, 2n+1)) \longrightarrow \tilde{K}^{-1}(Y)$ is isomorphic. q. e. d.

PROPOSITION 3.3. *The order of the group $\tilde{K}(Y) \leq p^k 2^n$.*

PROOF. The order of $E_{\infty}^{s,t}(Y)$ is not more than that of $E_2^{s,t}(Y)$, $s+t =$ even. From Proposition 3.1, the proposition follows.

PROOF OF THEOREM 1.1.

Denote by c and r the complexification and the real restriction. Put $\sigma_m = \xi_m - 1$, then

$$\sigma_m + \bar{\sigma}_m = c r \sigma_m.$$

In [4], it is proved that $r\sigma_m, (r\sigma_m)^2, \dots, (r\sigma_m)^{(p-1)/2}$ are linearly independent in $\widetilde{KO}(L^m(p))$ and if $m = s(p-1) + t$, $0 \leq t < p-1$ then

$$\text{the order of } (r\sigma_m)^t = \begin{cases} p^{s+1} & \text{if } 2t \leq t, \\ p^s & \text{if } 2t > t. \end{cases}$$

We note that if $m = 2k+1$, the complexification c is injective and $\sigma_m + \bar{\sigma}_m$ belongs to $\tilde{K}(L^m(p))^{Z_2}$. Therefore, we have that the order of $\tilde{K}(L^{2k+1}(p))^{Z_2} \geq p^k$. Since $\tilde{K}_{Z_2}(S^{2n+1}) \cong \tilde{K}(RP^{2n+1}) \cong Z_{2^n} [1]$,

$$\text{the order of } \tilde{K}(L^{2k+1}(p))^{Z_2} \oplus \tilde{K}(RP^{2n+1}) \geq p^k 2^n.$$

It follows from Theorem 2.2, Proposition 3.3 that the torsion part of $\tilde{K}(D_p(2k+1, 2n+1))$ is isomorphic to $\tilde{K}(L^{2k+1}(p))^{Z_2} \oplus \tilde{K}(RP^{2n+1})$. q. e. d.

Department of Mathematics,
Fukuoka Technological University,

Department of Mathematics,
College of General Education,
Kyushu University
and

Department of Mathematics,
College of General Education,
Kyushu University.

References

- [1] J. F. Adams, *Vector fields on spheres*, Ann. of Math., **75**(1962), 603-622.
- [2] M. F. Atiyah and F. Hirzebruch, *Vector bundles and homogeneous spaces*, Proc. Symp. Pure Math. vol. III, Amer. Math. Soc., Providence, R. I., 1961, 7-38.
- [3] M. Kamata and H. Mimami, *Bordism groups of dihedral groups*, J. Math. Soc. Japan, **25**(1973), 334-341.
- [4] T. Kambe, *The structure of K_Λ -ring of lens space and their application*, J. Math. Soc. Japan, **18**(1966), 135-146.
- [5] N. Mahammed, *A propos de la K -théorie des espaces lenticulaires*, C. R. Acad. Sci. Paris, Ser. A-B, **271**(1970), 639-642.
- [6] G. B. Segal, *Equivariant K -theory*, Inst. Hautes Études Sci, Publ. Math., **34**(1968), 129-155.