Spectroscopic and Crystallographic Studies on the Insertion Reaction of Aryl Isocyanides into the Bond between Palladium and Carbon, which Contribute to Understanding the trans-[Br2Ni(CNAr)2]-Catalyzed Ethylene Polymerization

Tsuchiya, Kazuhiro Department of Molecular and Material Sciences, Interdisciplinary Graduate School of Engineering Sciences, Kyushu University

Nagashima, Hideo Institute for Materials Chemistry and Engineering, Kyushu University | Department of Molecular and Material Sciences, Interdisciplinary Graduate School of Engineering Sciences, Kyushu University

https://doi.org/10.15017/14564

出版情報:九州大学大学院総合理工学報告.28(3),pp.335-342,2006-12.九州大学大学院総合理工学府 バージョン: 権利関係:

### Spectroscopic and Crystallographic Studies on the Insertion Reaction of Aryl Isocyanides into the Bond between Palladium and Carbon, which Contribute to Understanding the *trans*-[Br<sub>2</sub>Ni(CNAr)<sub>2</sub>]-Catalyzed Ethylene Polymerization

Kazuhiro TSUCHIYA\*1 and Hideo NAGASHIMA\*1,\*2,  $^{\dagger}$ 

\*E-mail of corresponding author: *nagasima@cm.kyushu-u.ac.jp* 

### (Received October 31, 2006)

The insertion of two sterically different isocyanides into a Pd-Me bond was investigated by spectroscopy and crystallography; this contribute to better understanding of ethylene polymerization catalyzed by trans-[Br<sub>2</sub>Ni(CNAr)<sub>2</sub>]. Thus, the reaction of MePdCl(COD) (COD = cyclooctadiene) with a small isocyanide (CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>) or a bulky isocyanide (CNC<sub>6</sub>H<sub>3</sub>·2,6-Ph<sub>2</sub>) demonstrated that isocyanide immediately inserted into the Pd-Me bond. Altough the  $C_6H_3$ -2,6-Ph<sub>2</sub> group is more bulky than the  $C_6H_3$ -2,6-Me<sub>2</sub> moiety, the anisotropic structure of ortho phenyl groups could provide enough space for the insertion by their Similarly, appropriate rotations. the molecular structure of trans-[ClPd{C(Me)=NC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub>}(CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub>)<sub>2</sub>] (**5b**) provided another example of appropriate rotations of phenyl rings, which ingeniously avoid the inter-ligand repulsions.

Key words: organometallic chemistry, palladium complex, aryl isocyanide, insertion reaction

of

### 1. Inroduction

Organometallic complexes have been actively investigated as the catalyst for ethylene polymerization since the discovery of the metallocene catalyst.<sup>1)</sup> Brookhart and co-workers found Ni complexes bearing  $\alpha$ -diimine ligands showing high catalytic activity for ethylene polymerization in 1995,<sup>2)</sup> opening the field of the "post-metallocene catalyst".<sup>3)</sup> We have recently found that the trans [Br<sub>2</sub>Ni(CNAr)<sub>2</sub>] (Ar = aryl group) showed considerably high activity for ethylene polymerization in the presence of methylaluminoxane (MAO) (Scheme 1).<sup>4)</sup> The catalytically active species must be molecule; this is supported by the catalytic activity depends on the structure of the isocyanide ligands. For instance, the Ni complex having CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> showed the activity 40 times higher than that of CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>.

Although the performance

1. Department of Molecular and Material Sciences

2. Institute for Materials Chemistry and Engineering







Fig. 2 The azanickellacyclopentene complex (I)

trans-[Br<sub>2</sub>Ni(CNAr)<sub>2</sub>] is unique as ล post-metallocene catalyst, the mechanism is totally unknown. From the analogy to well-investigated intermediates of the Brookhart's catalyst, which are cationic alkyl nickel species bearing coordinated ethylene as shown in Fig. 1, cationic species A may be generated from *trans*-[Br<sub>2</sub>Ni(CNAr)<sub>2</sub>] by way of mono-methylation and subsequent halogen abstraction by MAO (Scheme 2). However, this mechanism is problematic due to the trans-configuration of **A** between Me group and ethylene, which prevents insertion of ethylene between the Ni-Me bond. Possible insertion of the isocyanide ligand in A between the Ni-Me bond posed another problem for the mechanistic consideration, which could produce new nickel intermediates having а Ni-C(=NAr)Me moiety. It is really attractive to consider alternative mechanisms, in which intermediates having a Ni-C(=NAr)Me moiety or its analogues play an important role in the *trans*-[Br<sub>2</sub>Ni(CNAr)<sub>2</sub>]-catalyzed ethylene polymerization. In fact, a support of this possibility was available by the discovery that a nickel complex I, which can be formed by insertion of three molecules of CNAr to Ni-Me species generated from *trans*-[Br<sub>2</sub>Ni(CNAr)<sub>2</sub>] with MAO, actually behaves as an ethylene polymerization catalyst (Fig. 2).<sup>5)</sup>

We felt high probability of the mechanisms involving insertion of CNAr into Ni-Me species. However, a problem of this mechanism is the activity of ethylene polymerization depends on the structures of isocyanides, typically, trans-[Br<sub>2</sub>Ni(CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub>)<sub>2</sub>] >> trans-[Br<sub>2</sub>Ni(CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>)<sub>2</sub>]. As described in section 2.4 in detail, the molecular modeling study strongly suggests that there is a substantial difference in steric hindrance between CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> and CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>; the former is more bulky than the latter. A question is whether CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> may be too bulky for the smooth insertion into the Ni-Me bond; this provides difficulties in generating the catalytically active species. In order to solve this problem, we were interested in the model reaction using palladium as central metal to



**Fig. 3** The reaction of MePdCl(COD) with 2 or 3 equivalents of ArNC.

investigate facile the how insertion of CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> and CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub> into Pd-Me bond of in situ generated MePdCl(CNAr)<sub>2</sub>. Palladium is in the same triad as nickel, affording more stable methyl complexes. of High stability the palladium-methyl complexes are useful for the mechanistic studies of the *trans*-[Br<sub>2</sub>Ni(CNAr)<sub>2</sub>] catalyzed ethylene polymerization, in which insertion of CNAr to MeNiX(CNAr)2 is possibly involved as the mechanism.

### 2. Results and Dicussion

2.1 The spectroscopic study of the insertion reaction of  $CNC_6H_3\mathchar`-2,6\mathchar`-Me_2$  into a Pd-Me bond

Vincente and co-workers have recently the insertion reaction of reported CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub> to Pd-Me bond, which is the small isocyanide described above, into a Pd-(C<sub>6</sub>H<sub>4</sub>-2-NH<sub>2</sub>) bond.<sup>6)</sup> There was no report on the insertion reaction of CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub> into a Pd-Me bond, which is more important for the consideration of the mechanism of the MAO-assisted trans-[Br<sub>2</sub>Ni(CNAr)<sub>2</sub>]-catalyzed ethylene polymerization. As a related study, Anderson and co-workers reported spectroscopic studies on the insertion reaction of CN<sup>t</sup>Bu into a Pd-Me bond.<sup>7)</sup> This report has shown the reaction of MePdCl(COD) and three equivalents of CN<sup>t</sup>Bu to give the Pd complex having a structure similar to 5 shown in Fig. 3. In the related study to this Anderson's research, Uson and co-workers discussed the possibility

of intermediates similar to **3** and **4**.<sup>8)</sup>

We carried out the reaction of MePdCl(COD) with two or three equivalents of  $CNC_6H_3$ -2,6-Me<sub>2</sub> in  $CD_2Cl_2$ , and monitored the formed products by <sup>1</sup>H NMR spectroscopy. As shown in Fig. 3, the reaction of MePdCl(COD) and two equivalents of CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub> was expected to form 2a by the ligand exchanging of COD and CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>. If **2a** is unstable and insertion of CNAr rapidly takes place, 3a and 4a could be formed. Addition of another equivalent of CNC6H3-2,6-Me2 was likely to give 5a as a single stable product. The actual <sup>1</sup>H NMR spectra (the range of Me on aryl group) for these reactions are shown in Fig 4. The reaction of MePdCl(COD) with two equivalents of CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub> afforded significant amount of precipitation, whereas addition of additional  $CNC_6H_3$ -2,6-Me<sub>2</sub> resulted in dissolution of the precipitates. A <sup>1</sup>H resonance due to the Pd-Me group is generally observed around or higher than 1 ppm; however, no singlet was observed around 1 ppm on <sup>1</sup>H NMR spectra. The obtained addition of spectrum by three equivalents of CNC6H3-2,6-Me2 was unequivocally showed that the sole species assignable to 5a exist in the solution. Thus, the signal of Me group originated from Pd-Me was seen at 2.68 ppm, significantly downfield from 1 ppm, suggesting the insertion reaction of isocyanide to form the  $Me(Pd)C=N(C_6H_3-2,6-Me_2)$  moiety. The signals due to the coordinated CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub> were observed at 2.10 and 2.40 ppm, respectively. The integral ratio of these three singlets was 3:6:12, which is in accord with the structure of 5a. It is well known that the signals of aromatic protons are generally seen in 7-8 ppm. The signals of triplet (2H) and doublet (4H) at 7.29 and 7.16 ppm were thus assignable to para and meta protons of the coordinated  $CNC_6H_{\mathcal{F}}2$ , 6-Me<sub>2</sub>, respectively. In contrast, the multiplet (3H) at 6.88-6.95 ppm appeared unusually upfield as the aromatic protons. The upfield shift of aromatic protons is often observed, when the electron donating group such as nitrogen functional groups having a lone pair electron. Compared with the  $N\equiv C$ group of isocyanide (Ar-N $\equiv$ C $\rightarrow$ Pd) acting as an



 Table 1
 Representative IR absorptions of obtained mixtures under each condition

| condition              | 2eq. of CNAr |             | 3eq. of CNAr |
|------------------------|--------------|-------------|--------------|
|                        | solution     | precipitate |              |
| C≡N(cm <sup>-1</sup> ) | 2175         | 2189        | 2174         |
| C=N(cm <sup>-1</sup> ) | 1630-1660    | 1663        | 1666         |
|                        | 1590         | 1560        |              |

electron acceptor, the N=C group of the imidoyl (Ar-N=C(Me)-Pd) could behave as an electron donor, giving the upfield shift described above. Two IR absorptions were visible at 2174 and 1666 cm<sup>-1</sup> due to the stretching vibration of the coordinated isocyanide (C=N-Ar) and imidoyl group (C=N-Ar), respectively (Table 1). Thus all of the <sup>1</sup>H and IR data are consistent with the structure of **5a**.

The <sup>1</sup>H NMR spectrum of the reaction of two equivalents of CNC6H3-2,6-Me2 with MePdCl(COD) was complicated, suggesting the existence of more than two palladium species besides 5a. The IR spectrum also gave several absorptions including the peaks due to 5a. As described above, no <sup>1</sup>H resonance around 1 ppm indicates rapid insertion of CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub> into the Pd-Me bond. At present, we postulate the species formed by addition of two equivalents of CNC6H3-2,6-Me2 would be a mixture of 3a and 4a, which are presumably a mixture of cis and trans isomers. This hypothesis is supported by the IR analyses of the precipitates and the solid recovered from the solution. The precipitate had absorption at 1560 cm<sup>-1</sup> due to the bridging imidoyl group of 4a, whereas the absorptions of the sample obtained from the solution were mainly observed at the range of 1630-1660 cm<sup>-1</sup> which can be assignable to **3a** and **5a**. It is important that dual species existing in the reaction mixture of two equivalents of  $CNC_6H_3$ -2,6-Me<sub>2</sub> with MePdCl(COD) were converted to a single product **5a** by addition of another equivalent of  $CNC_6H_3$ -2,6-Me<sub>2</sub>. these results suggest that the spectral change is not inconsistent with the scheme shown in Fig. 3.

These spectroscopic results demonstrated that the small isocyanide,  $CNC_6H_3$ -2,6-Me<sub>2</sub>, quickly inserted into the Pd-Me bond of intermediary MePdCl( $CNC_6H_3$ -2,6-Me<sub>2</sub>)<sub>2</sub> to form the imidoyl structure. This provided a good basis to investigate the homologous reaction of more bulky  $CNC_6H_3$ -2,6-Ph<sub>2</sub> with MePdCl(COD).

# 2.2 The spectroscopic study of the insertion reaction of $CNC_6H_3\mathchar`-2,6\mathchar`-Ph_2$ into a Pd-Me bond

Similar pathways to Fig. 3 (Ar =C<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>) can be drawn for the reactions of MePdCl(COD) and two or three equivalents of  $CNC_6H_3-2, 6-Ph_2$  [Fig. 3 (Ar =  $C_6H_3-2, 6-Ph_2$ )]. The change of actual <sup>1</sup>H NMR spectra (the range of Me) for the reactions of CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> with MePdCl(COD) are shown in Fig. 5. In contrast to the above described reaction with the small isocyanide, the reaction with two equivalents of  $CNC_6H_3$ -2,6-Ph<sub>2</sub> kept homogeneous. The 1H NMR showed two singlets due to the Me groups at 0.98 and 1.62 ppm. A small singlet appeared at 1.15 ppm will be discussed later. On addition of another equivalent of CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub>, the singlet at 1.62 ppm disappeared, and all of the <sup>1</sup>H resonances, both methyl and aryl regions, became consistent with the structure of 5b. Thus, the signals observed at 7.61-7.40 ppm are due to all of the protons on aryl groups of the  $CNC_6H_{3}2$ , 6-Ph<sub>2</sub> ligand coordinated to the Pd center. The two triplets at 6.60 and 6.71 ppm were due to the protons on the para and meta position of  $C_6H_3$ -2,6-*Ph*<sub>2</sub> of the imidoyl group, respectively. The multiplets observed at 6.80-6.86 ppm and a doublet at 7.10 ppm were due to the protons of  $C_6H_{\mathcal{F}}2$ , 6-Ph<sub>2</sub> and ortho protons of  $C_6H_3$ -2,6-*Ph*<sub>2</sub> of the imidoyl group, respectively. The IR spectrum showed two



Fig. 5 <sup>1</sup>H NMR spectrum of the reaction of MePdCl(COD) with 2 (above) or 3 (below) equivalents of  $CNC_6H_3$ -2,6-Ph<sub>2</sub> in  $CD_2Cl_2$ 

**Table 2.** Representative IR absorptions of obtained mixtures under each condition

| condition              | 2eq. of CNAr | 3eq. of CNAr |
|------------------------|--------------|--------------|
| C≡N(cm <sup>-1</sup> ) | 2172         | 2175         |
| C=N(cm <sup>-1</sup> ) | 1653         | 1653         |
|                        | 1583         | 1585         |

absorptions at 2175 and 1653 cm<sup>-1</sup> due to the stretching vibration of the coordinated isocyanide (C $\equiv$ N-Ar) and imidoyl group (C=N-Ar), respectively, which is also consistent with the structure of **5b**.

A solution formed by the reaction of two equivalents of this bulky isocyanide with MePdCl(COD) apparently contained **5b** and another species showing a <sup>1</sup>H resonance at 1.62 ppm. The IR absorptions mainly appeared at 2172 and 1653 cm<sup>-1</sup>, which are assignable to the isocyanide ligand and non-bridging imidoyl groups. This suggests that the species showing the singlet at 1.62 ppm on the NMR could be **3b**. Evidence for the presence of **4b** may be a small absorption at 1583 cm<sup>-1</sup>, which may correspond to the singlet at 1.15 ppm on the <sup>1</sup>H NMR spectrum.

All of these results suggest that the insertion of isocyanide into the Pd-Me bond is also rapid even with the bulky isocyanide such as  $CNC_6H_{32}$ , 6-Ph<sub>2</sub>. For the explanation of the spectroscopic data described above consistently, existence of **2b** in solution must be excluded, which might be detectable in solution when the bulky isocyanide slowly undergoes the insertion between the Pd-Me bond. One may claim that the <sup>1</sup>H resonances due to the methyl groups appeared at 0.98-1.62 ppm as described above, which could be assignable to Pd-Me moieties. Since the methyl group of the imidoyl moiety in **5a** appeared significantly downfield from this region, we have to explain the unusual chemical shift of the imidoyl methyl group in **5b**. This should be due to the long range shielding effect induced by local magnetic field of the benzene ring, which can clearly be visible in the molecular structure of **5b** described in the next section.

# 2.3 The molecular structure of the complex **5b** formed by insertion of the bulky isocyanide, CNC<sub>6</sub>H<sub>3</sub>·2,6-Ph<sub>2</sub>

The described above spectroscopic studies indicated that even the bulky isocyanide,  $CNC_6H_3$ -2,6-Ph<sub>2</sub>, quickly insert into the Pd-Me bond. The insertion reaction proceeded so smoothly even at room temperature. However, considerations using the molecular model described later indicate that bulkiness of CNC6H3-2,6-Ph2 may provide sterically crowded very circumstances on the molecular structure of 5b. In fact, inter-ligand steric repulsions around the Pd centre looked enormous when we built the simple molecular model for 5b. In other words, the bulkiness of isocyanide may affect the thermodynamic stability of **5b**, though it does not give little effect on the kinetic formation of **5b**. The crystallography to determine the molecular structure of 5b was thus carried out.

The single crystals were grown from a  $CH_2Cl_2$ /hexane solution containing a mixture obtained by the reaction of MePdCl(COD) with 4 equivalents of  $CNC_6H_3$ -2,6-Ph<sub>2</sub>. There was no problem on the crystallographic data, and the molecular structure of **5b** was revealed to be just the same as we expected. The ORTEP drawing is illustrated in Fig. 6.

Arrangement of the four groups connecting to the Pd atom is square planer, and two  $CNC_6H_3$ ·2,6·Ph<sub>2</sub> ligands coordinated to the Pd centre are in trans configuration. The imidoyl group,  $C(Me)=NC_6H_3$ ·2,6·Ph<sub>2</sub>, exists at the opposite site of the chlorine atom. The two  $CNC_6H_3$ ·2,6·Ph<sub>2</sub> ligands are bonded to the Pd



Fig. 6 The molecular structure of 5b. Ellipsoids represent 50% probability; hydrogen atoms are omitted for clarity.

| Table 3 | Representative | bond lengths | and angles for <b>5b</b> |
|---------|----------------|--------------|--------------------------|
|---------|----------------|--------------|--------------------------|

| -           |          | =                | -        |  |
|-------------|----------|------------------|----------|--|
| Pd(1)-Cl(1) | 2.439(1) | Cl(1)-Pd(1)-C(1) | 175.7(1) |  |
| Pd(1)-C(1)  | 2.022(5) | Cl(1)-Pd(1)-C(4) | 96.7(1)  |  |
| Pd(1)-C(4)  | 1.990(5) | C(4)-Pd(1)-C(5)  | 173.0(2) |  |
| Pd(1)-C(5)  | 1.972(5) | C(1)-N(1)-C(3)   | 125.0(4) |  |
| N(1)-C(1)   | 1.277(6) | C(2)-C(1)-Pd(1)  | 116.6(3) |  |
| N(2)-C(4)   | 1.154(6) |                  |          |  |
| N(3)-C(5)   | 1.143(6) |                  |          |  |

 $Table \ 4 \quad {\rm Crystallographic \ data \ for \ 5b}$ 

| Crystal system                                 | monoclinic                             |
|--|--|
| Space group                                    | P2 <sub>1</sub> /c                     |
| a, (Å)   | 9.561(2)                               |
| b, (Å)   | 17.843(4)                              |
| c, (Å)   | 27.205(5)                              |
| β, (deg.)                                      | 94.7442(9)                             |
| V, (Å <sup>3</sup> )                           | 4632.8(16)                             |
| Z  | 4                                      |
| $\mu(MoK\alpha)$ (cm <sup>-1</sup> )           | 15.28                                  |
| No. of reflections measured                    | total: 27045, unique: 7638             |
|  | $(R_{int} = 0.069)$                    |
| Good ness of fit on $F^2$                      | 1.006                                  |
| Final R indices $[\lambda > 2\sigma(\lambda)]$ | $R_1 = 0.0530$                         |
| R indices (all data)                           | $wR_2 = 0.1490$                        |
| Largest diff. peak and hole                    | $2.54$ and -0.77 $e{\rm \AA}^{\cdot3}$ |
|  |  |

atom with bond lengths of 1.990(5) and 1.972(5)Å for Pd-C, whereas those of 1.154(6) and 1.143(6) Å for C=N. The C(Me)=NC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> group has a typical structure of the imidoyl group, of which N(1)-C(1) bond distance [1.277(6) Å] suggested the existence of a N=C double bond. whereas the angles of C(1)-N(1)-C(3)[125.0(4)]deg.] and C(2)-C(1)-Pd(1) [116.6(3) deg.] contribute to sp<sup>2</sup> hybridization of C(1)and N(1). The arrangement of the phenyl groups, which are located at the ortho-positions of three  $C_6H_3$ -2,6-Ph<sub>2</sub> moieties, was demonstrated to avoid the steric repulsions of the Ph rings ingeniously. This is achieved by rotation of the Ph planes in the CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> groups. Thus, the possible rotation to avoid the steric repulsions explain both the thermodynamic stability of **5b**; this explanation can be extended to interpretation of the facile insertion of  $CNC_6H_3$ -2,6-Ph<sub>2</sub> into the Pd-Me bond discussed above.

The crystallographic study of **5b** also gave a clear answer to the question why the <sup>1</sup>H resonance due to the imidoyl methyl group in 5b appeared at unusually higher field. In the molecular structure of **5b**, the methyl group is located at the face of two phenyl moieties of the  $CNC_6H_3$ -2,6-Ph<sub>2</sub> ligands. It is well known that the ring current of the aromatic ring gives local magnetic field leading to substantial change of the chemical shifts of protons nearby in the NMR spectroscopy(see Experimental section). The proton spatially located on the face of the phenyl ring generally gives significant upfield shifts by 1-2 ppm. The location of the methyl group is fitted just in this case, and its chemical shift (0.98 ppm) is explained by the long range shielding of the phenyl rings nearby.

## 2.4 Considerations on the steric effect of isocyanides

As described above, we started this research from our hypothesis that CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> may be sterically too bulky to promote insertion into the Pd-Me bond, which is facilely accomplished by smaller CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>. This hypothesis seemed to be reasonable, when we saw the structures of two isocyanides, CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> and CNC6H3-2,6-Me2, determined by semi-empirical MO (PM3) calculations, of which CPK model illustrations are shown in Fig. 7. It is well known that a methyl group is isotropic, and free rotation along with the C(Me)-C(aryl) bond in  $CNC_6H_3$ -2,6-Me<sub>2</sub> provides а spherical sterically exclusive region, of which diameter is ca. 5 Å. According to the estimation of the spherical sterically exclusive region, free rotation of the phenyl group affords a sphere of which diameter is ca. 8 Å. In this sense, CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> must be more bulky than CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>. However, a feature of phenyl group is its anisotropic structure, and planarity of the phenyl group would provide



Fig. 7 The space filling model of  $CNC_6H_3$ -2,6-Me<sub>2</sub> (left) and  $CNC_6H_3$ -2,6-Ph<sub>2</sub> (right), simulated by PM3 method



Fig. 8 The isocyanides;  $CNC_6H_3$ -2,6-Me<sub>2</sub> [a],  $CNC_6H_3$ -2,6-Ph<sub>2</sub> [b(edge)] and [b(face)]



Fig. 9 The skeleton model (upper) and space filling model (bottom) of complex **5a** (left), There were simulated by MOPAC based on the structure of *trans*-[Pd{C(= $NC_6H_3$ -2,6- $Me_2$ )C<sub>6</sub>H<sub>4</sub> $NH_2$ -2}I(CNC<sub>6</sub>H<sub>3</sub>-2,6- $Me_2$ )]. Those of the complex **5b** (right) were determined by crystallographic data

much smaller steric influence. This is typically seen in Fig. 8, in which the rotation gives two extreme configuration, "edge" and "face". If the rotation is restricted so as to minimize the steric repulsion among the ligands, the C<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> groups would not afford severe steric hindrance to the complex 5b. The molecular structure of 5a was shown in Fig. 9, was estimated from the crystallographically determined structure of trans- $[Pd{C(=NC_6H_3-2,6-Me_2)C_6H_4NH_2-2}]I(CN$  $C_6H_3$ -2,6-Me<sub>2</sub>)], of which the  $C_6H_4NH_2$ -2 moiety was replaced by the Me group. The steric circumstance around the palladium in 5a is not crowded, and there is enough space for the free rotation of the methyl groups in C<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub> moieties. In sharp contrast, the ortho-phenyl groups in **5b** are apparently large, and caused substantial steric repulsions

among the ligands bonded to the palladium atom. In other words, it is apparent that coordination of all of the ligands in 5b is accomplished by the fact that the motion of the phenyl groups should be severely restricted to minimize the inter-ligand steric interactions. This is accomplished by anisotropic structure of the phenyl groups which enable to alternate their steric influence by appropriate rotation. The importance of the anisotropic structure of the phenyl groups in the  $C_6H_3$ -2,6-Ph<sub>2</sub> groups can be expanded to the explanation for the facile insertion of CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub>. Although the phenyl group in  $CNC_6H_3$ -2,6-Ph<sub>2</sub> is more bulky than the methyl group in CNC<sub>6</sub>H<sub>3</sub>-2,6-Me<sub>2</sub>, the phenyl groups can produce enough space for the insertion into the Pd-Me bond by their appropriate rotations. In other words, success of the CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> ligand in the *trans*-[Br<sub>2</sub>Ni(CNAr)<sub>2</sub>]-catalyzed polymerization of ethylene could be attributed to the steric flexibility of CNC6H3-2,6-Ph2 derived from the anisotropic structure of the phenyl groups, which facilely produces the catalytically active species by the insertion reaction into the intermediary Ni-Me bond as well as giving the appropriate steric influence on the active species for the ethylene polymerization. Importance of appropriate steric influence of the ligand on the ethylene polymerization has been well established in the chemistry of the Brookhart's catalysts.

### Concluion

We reported in this paper a model study for elucidating the mechanisms of *trans*-[Br<sub>2</sub>Ni(CNAr)<sub>2</sub>]-catalyzed

polymerization of ethylene. We believe that insertion of the isocyanide ligand into a Ni-Me bond formed by pretreatment of MAO with *trans*-[Br<sub>2</sub>Ni(CNAr)<sub>2</sub>] plays an important role in catalysis. The spectroscopic as well as the crystallographic studies using palladium homologues as models for this insertion reaction solve a question whether the bulky  $CNC_6H_3$ -2,6-Ph<sub>2</sub> really inserts between the Ni-Me bond. Although complete elucidation of the polymerization mechanisms requires further investigation, the results described in





this paper supports a possible mechanism 10. Thus, shown in Fig. *trans* [MeBrNi(CNAr)<sub>2</sub>] species formed by the pretreatment of nickel catalyst with MAO is followed by the insertion of CNAr into the bond. The Ni-Me resulting imidoyl intermediate 6 reacts with ethylene to afford the intermediate 7. Replacement of the Br ligand by a methyl group is accomplished by methylation with MAO to form 8, and insertion of coordinated ethylene into the Ni-Me bond of 8 initiates the ethylene polymerization.

We are planning further elucidation of the polymerization mechanisms and the catalyst design to improve the catalytic activity and polymerization behavior. The present study suggests that model studies using palladium will help for these future plans.

### Acknowledgement

The authors are grateful to Mr. Masao Tanabiki and memberes of TOSOH Co. Ltd. for helpful discussions. Help for crystallographic studies by Dr. Yusuke Taisuke Sunada and Matsumoto is acknowledged.

#### Experimental section

General.: All experiments were carried out under an argon atmosphere. All solvents were distilled over  $CaH_2$  prior to use ( $CD_2Cl_2$  and  $CH_2Cl_2$ ). All NMR experiments were carried out using  $CD_2Cl_2$  solution placed in a 5 $\Phi$  NMR tube and degassed several times. The tube was sealed in flame, while the solution was kept in a liquefied nitrogen bath in vacuum. The <sup>1</sup>H NMR spectra were taken with a JEOL Lambda 400 spectrometer. Chemical shifts were recorded in ppm from the internal standard (<sup>1</sup>H: solvent). IR spectra were recorded in cm<sup>-1</sup> on a JASCO FT/IR-550 spectrometer.

## NMR identification of the species formed by the reaction of MePdCl(COD) with isocyanides:

(I) MePdCl(COD) with two or three equivalents of  $CNC_6H_3$ -2,6- $Me_2$ ; In a NMR tube were placed  $CNC_6H_3$ -2,6- $Me_2$ ; (2.7 mg, 20 µmol, or 4.1 mg, 30 µmol) and MePdCl(COD) (2.8 mg, 10 µmol), and then  $CD_2Cl_2$  (ca. 0.3 mL) was transferred to by vacuum.

Spectraldatafor[ClPd][C(Me)=NC6H3\*2,6·Me2](CNC6H3\*2,6·Me2)2] (5a); <sup>1</sup>HNMR (395MHz, CD2Cl2, r.t.);  $\delta7.61$  (d, 1H, Ph, J = 7.2Hz).7.59 (d, 1H, Ph, J = 7.2Hz), 7.50\*7.40 (m, Ph, 21H), 7.10(d, 4H, J = 7.7Hz), 6.86\*6.80 (m, 3H), 6.71 (t, 4H, Ph, J = 7.7Hz), 6.60 (t, 2H, Ph, J = 7.7Hz), 0.98 (s, Me, 3H). IR: $v_{C = N} 2175$ , 1652 cm<sup>-1</sup>

(II) MePdCl(COD) with two or three equivalents of  $CNC_6H_3$ -2,6-Ph<sub>2</sub>; The reaction of  $CNC_6H_3$ -2,6-Ph<sub>2</sub>; (7.7 mg, 30 µmol, or 12 mg, 45 µ mol) and MePdCl(COD) (4.0 mg, 15 µmol) was carried out in similar fashion.

Isolation and crystallographic study of ClPd[C(Me)=NC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub>](CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub>)<sub>2</sub> (5b); In CH<sub>2</sub>Cl<sub>2</sub> (10 mL), CNC<sub>6</sub>H<sub>3</sub>-2,6-Ph<sub>2</sub> (123 mg, 0.48 mmol) and MePdCl(COD) (34.4mg, 0.13 mmol) were dissolved at -78°C, and the mixuture was stirred for 2h at room temperature. The resulting solution was removed in vacuo, and the residue was washed with hexane. The desired complex 5b was formed as white powder. Recrystallization of this crude product from CH<sub>2</sub>Cl<sub>2</sub>-hexane gave **5b** as needle in 99% yield (119 mg). <sup>1</sup>H NMR (395MHz CD<sub>2</sub>Cl<sub>2</sub>, r.t.);  $\delta$ 7.61 (d, 1H, Ph, J = 7.2Hz). 7.59 (d, 1H, Ph, J=7.2Hz), 7.50-7.40 (m, Ph, 21H), 7.10 (d, 4H, Ph, 7.7Hz), 6.86-6.80 (m, Ph, 3H), 6.71 (t, 4H, Ph, J = 7.7Hz), 6.60 (t, 2H, Ph, J = 7.7Hz), 0.98 (s, Me, 3H). IR:  $\nu_{C\equiv N}$  2175, 1652 cm  $^{\text{-}1}$ 

Crystallographic studies; X-ray crystallography was performed on a Rigaku Saturn CCD area detector with graphite monochromated Mo K $\alpha$  radiation ( $\lambda = 0.71070$ Å). The data were collected at 293(1) K using  $\omega$  scans in the  $\theta$  range of  $-110.0^{\circ} \le \theta \le 70^{\circ}$ . Data were collected and processed using CrystalClear (Rigaku) on a Pentium computer. The data were corrected for Lorentz and polarization effects. The structure was solved by direct methods<sup>9)</sup> and expanded using Fourier techniques.<sup>10)</sup> The non-hydrogen atoms were refined anisotropically. Hydrogen atoms were refined using the riding model. The final cycle of full-matrix least squares refinement on  $F^2$  was based on 10380 observed reflections and 610 variable parameters. Neutral atom scattering factors were taken from Cromer and Waber.<sup>11)</sup> All calculations were performed using the CrystalStructure<sup>12),13)</sup> crystallographic software package. Details of final refinement are summarized in Table 4, and the numbering scheme employed is shown in Fig. 6.

### References

- (a) E. Heins, H. Hinck, W. Kaminsky, G. Oppermann, P. Raulinat and H. Sinn, Makromol. Chem., 134, 1 (1970). (b) W. Kaminsky, H. J. Vollmer, E. Heins and H. Sinn, Makromol. Chem., 175, 443 (1974). For reviews; (c) R. H. Grubbs and G. W. Coates, Acc. Chem. Res., 29, 85 (1996). (d) A. K. Rappé, W. M. Skiff and C. Casewit, J. Chem. Rev., 100, 1435 (2000).
- (a) L. K. Johnson, C. M. Killan and M. Brookhart, J. Am. Chem. Soc., 117, 6414 (1995). (b) D. P. Gate, S. A. Svejda, E. Onate, C. M. Killan, L. K. Johnson, P. S. White and M. Brookhart, Macromolecules, 33, 2320 (2000). (c) S. D. Ittel, L. K. Johnson and M. Brookhart, Chem. Rev., 100, 1169 (2000).
- For a review; (a) V. C. Gibson and S. K. Spitzmesser, Chem. Rev., 103, 283 (2003) and reference therein. Typical examples, see; (b) G. J. P. Britovesk, S. P. D. Baugh, O. Hoarau, V. C. Gibson, D. F. Wass, A. J. P. White and D. J. Williams, Inorg. Chim. Acta., 345, 279 (2003). (c) T. R. Youkin, E. F. Conner, J. I. Henderson, S. K. Friedrich, R. H. Grubbs and D. A. Bansleben, Science, 287, 460 (2000). (d) S. Y. Desjardins, K. J. Cavell, J. L. Hoare, B. W. Skelton, A. N. Sobolev, A. H. White and W. Keim, J. Organomet. Chem., 544, 163 (1997).
- M. Tanabiki, K. Tsuchiya, Y. Kumanomido, K. Matsubara, Y. Motoyama and H. Nagashima, Organometallics, 23, 3976 (2004).
- 5) unpublished data
- J. Vincente, J.-A. Abad, E. M-Viviente and P. G. Jones, Organometallics, 21, 4554 (2002).
- E. T. Ladipo and G. K. Anderson, Organometallics, 13, 303 (1994).
- R. Uson, J. Fornies, P. Espinet and E. Lalinde, J. Organomet. Chem., 254, 371 (1983). (b) R. Uson, J. Fornies, P. Espinet, E. Lalinde, P. G. Jones and G. M. Sheldrick, J. Organomet. Chem., 253, C47 (1983). (c) R. Uson, J. Fornies, P. Espinet, E. Lalinde, P. G. Jones and G. M. Sheldrick, J. Organomet. Chem., 288, 249 (1985). (d) R. Uson, J. Fornies, P. Espinet and E. Lalinde, J. Organomet. Chem., 349, 269 (1988).
- <u>SIR92</u>: A. Altomare, G. Cascarano, C. Giacovazzo, A. Guagliardi, M. Burla, G. Polidori and M. Camalli, J. Appl. Cryst., 27, 435 (1994).
- 10) <u>DIRDIF99</u>: P. T. Beurskens, G. Admiraal, G. Beurskens, W. P. Bosman, R. de Gelder, R. Israel and J. M. M. Smits, (1999). The DIRDIF-99 program system, Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands.
- (11) D. T. Cromer and J. T. Waber "International Tables for X-ray Crystallography", Vol. IV, The Kynoch Press, Birmingham, England, Table 2.2 A (1974).
- (12) <u>CrystalStructure 3.5.1</u>: Crystal Structure Analysis Package, Rigaku and Rigaku/MSC (2000-2003). 9009 New Trails Dr. The Woodlands TX 77381 USA.
- (13) CRYSTALS Issue 10: D. J. Watkin, C. K. Prout, J. R. Carruthers, and P. W. Betteridge, Chemical Crystallography Laboratory, Oxford, UK. (1996)