Using Cut-Column Pollination Method to Overcome Crossing Barriers in Phalaenopsis Sunrise Goldmour 'KHM637'

Chuang, Hua-Ting Laboratory of Floriculture, Department of HorticulturalScience, National Chiayi University

Huang, Kuang-Liang Laboratory of Floriculture, Department of HorticulturalScience, National Chiayi University

Shen, Rong-Show Laboratory of Floriculture, Department of HorticulturalScience, National Chiayi University

Miyajima, Ikuo Institute of Tropical Agriculture, Kyushu University

他

https://doi.org/10.5109/1467626

出版情報:九州大学大学院農学研究院紀要. 59(2), pp.265-271, 2014-08-29. Faculty of Agriculture, Kyushu University バージョン: 権利関係:

Using Cut–Column Pollination Method to Overcome Crossing Barriers in *Phalaenopsis* Sunrise Goldmour 'KHM637'

Hua–Ting CHUANG¹, Kuang–Liang HUANG¹, Rong–Show SHEN¹, Ikuo MIYAJIMA² and Shan–Te HSU¹*

Institute of Tropical Agriculture, Kyushu University, Fukuoka 812–8581, Japan (Received April 25, 2014 and accepted May 12, 2014)

Crossing barriers occur frequently when crossing between commercial Phalaenopsis cultivars with desirable horticultural traits are attempted. In this study, Phalaenopsis Sunrise Goldmour 'KHM637', a tetraploid with 2n=4x=76 chromosomes, known to be difficult in hybridization was chosen to examine crossing barriers and to explore the cut-column pollination method to overcome these barriers. P. Sunrise Goldmour 'KHM637' was either self-crossed, or used as the pollen or pistillate parent in reciprocal crossings with three tetraploid cultivars, including P. Brother Irene Irene 'Feng Fong', P. Sogo Yukidian 'V3', and P. Tai Lin Red Angel 'V31' that do not exhibit breeding barriers. Results showed that both stigma cavity closure and pollen tube growth did not occur when P. Sunrise Goldmour 'KHM637' was used as the pistillate parent. No offspring was obtained when using P. Sunrise Goldmour 'KHM637' as either pollen or pistillate parent. To explore performance of cut-column pollination method, P. Sunrise Goldmour 'KHM637' was used as the pistillate parent, after the removal of the upper part of column but with the stigma cavity intact, and then pollinated with pollinia of these three tetraploid cultivars and one species, P. violacea, respectively. The results showed that 22.2%-100.0% capsule sets were achieved from the four combinations, and offsprings were successfully obtained from three combinations. The authors suggested that the cut-column pollination method be applicable to other *Phalaenopsis* cultivars with similar crossing behaviors of *P*. Sunrise Goldmour 'KHM637' for overcoming the crossing barriers, and the applicability of the given method tended to be less limited to pollen parents without breeding barriers.

Key words: breeding barrier, maternal breeding barrier, reciprocal crossing, self–crossing, stigma cavity closure

INTRODUCTION

The Orchidaceae is one of the largest families of flowering plants with about 30,000 species classified into around 725 genera. The Phalaenopsis is a monopodial orchid in the Orchidaceae family. The genus Phalaenopsis comprising over 60 species belongs to the tribe Vandaeae under subfamily Vandoideae, which contains five subgenera, viz. subgenus Proboscidioides, subgenus Aphyllae, subgenus Parishianae, subgenus Polychilos, and subgenus Phalaenopsis (Christenson, 2001). Moth orchid is a common name for the genus Phalaenopsis. In commercial practice, plants of Phalaenopsis, Doritis and Doritaenopsis, an intergeneric hybrid between the former two genera, are collectively named as phalaenopsis orchids. At present, the Royal Horticultural Society (RHS) committee has decided to sink Doritis into Phalaenopsis; therefore Doritis and Doritaenopsis are now considered to be Phalaenopsis by RHS. Currently, more than 31,000 Phalaenopsis greges have been registered in RHS of UK (Royal

Horticultural Society, 2013). Phalaenopsis orchid has been a favorite consumer product due to its attractive color, graceful and long–lasting flowers. Nowadays, pot phalaenopsis has become one of the most important orchids in the trade.

Through years of research and development of new cultivars and promotion of the phalaenopsis industry, the mass production of high quality phalaenopsis orchids has led Taiwan to be known as "the kingdom of Phalaenopsis". Phalaenopsis orchids are mainly exported to the United States, Japan, European Union, Korea and China. In 2011, its gross export value reached \$98.5 million USD (Customs Administration, Ministry of Finance, R.O.C., 2012). In order to increase its competitiveness in facing the challenge from international competition, the industry has to continuously develop excellent new cultivars in addition to mass production of seedlings with the highest quality. For developing new cultivars through conventional breeding, failures in producing capsules with fertile seeds are frequently encountered in intergeneric, inter- and intra-specific crossings of *Phalaenopsis*. It is not clear which type of incompatibility and incongruity may cause the barriers in breeding of *Phalaenopsis* and whether these barriers are caused by genetic or external factors. Until now, there are no complete and detailed studies in addressing these issues.

In order to elucidate the basis for the reproductive barriers between specific crossing combinations for getting new cultivars, the genetic and physiological characteristics of some *Phalaenopsis* cultivars were examined

¹ Laboratory of Floriculture, Department of Horticultural Science, National Chiayi University, Chiayi 60004, Taiwan (R.O.C.)

² Institute of Tropical Agriculture, Kyushu University, Fukuoka 812–8581, Japan

^{*} Corresponding author: Department of Horticultural Science, National Chiayi University, No.300 Syuefu Rd., Chiayi City 60004, Taiwan (R.O.C.), E-mail: hsust@mail.ncyu.edu.tw, telephone: +88652717423, fax: +88652775472.

to identify the reproductive barriers between specific crossing combinations in our previous studies (Chuang et al., 2008; Hsu et al., 2010). These results indicated that crossing difficulties in those selected cultivars might be caused by odd chromosome set and some other factors of pollen or pistillate parents. The authors also demonstrated that the cut-column pollination method innovated in this study proved to overcome breeding barriers in Phalaenopsis Taipei Gold 'STM', a triploid with 2n=3x=57 chromosomes, known to have stigma closure problem after pollination and be difficult in hybridization (Chuang et al., 2013). P. Sunrise Goldmour 'KHM637', a tetraploid with 2n=4x=76 chromosomes, has also shown similar hybridization behaviors to P. Taipei Gold 'STM' (Chuang et al., 2008). Therefore, this study aimed to examine the efficacy of cut-column pollination method for overcoming the breeding barriers of P. Sunrise Goldmour 'KHM637', and to explore the applicable extensibility of the given method.

MATERIALS AND METHODS

Plant material and cultivation conditions

The mature clonal plants of *Phalaenopsis* Sunrise Goldmour 'KHM637' (SG), a tetraploid with 2n=4x=76chromosomes, known to have stigma closure problem after pollination and to be difficult in hybridization (Chuang *et al.*, 2008), *P*. Brother Irene Irene 'Feng Fong' (BII), *P*. Sogo Yukidian 'V3' (V3), *P*. Tai Lin Red Angel 'V31' (V31) and *P. violacea* were purchased or offered from several professional growers in Taiwan (Table 1) and grown in a greenhouse at Horticultural Technology Center, National Chiayi University, Chiayi, Taiwan, R.O.C.. For all experiments, plants were grown in a greenhouse at a temperature regime of 25/20°C day/night and a natural photoperiod ($23^{\circ}47$ 'N, $120^{\circ}48$ 'E) with a light intensity of 300 to $400 \,\mu$ mol·m⁻²·s⁻¹ during 1000 to 1200 h on the sunny days

Self- and reciprocal crossings

First, all four cultivars presented in Table 1 were self-crossed to check their crossing behaviors. Next, the cultivar of SG with difficulty in hybridization was reciprocal-crossed with each of the three normal cultivars, BII, V3 and V31. The occurrence of stigma cavity closure, *in vivo* pollen tube elongation, and capsule set were collected at three days after pollination (DAP), 7 DAP, and 60 DAP, respectively. The fertilities were evaluated by asymbiotic seed germinating abilities according to the method recommended by Lin *et al.* (2008). Two separate sets of experiments were conducted for investigating pollen tube development and fertility, respectively. Nine flowers randomly selected from three plants were examined in each self- and reciprocal crossings.

Microscopic observation of pollen tube growth

To monitor pollen tube behavior in different crossing combinations, columns were harvested evenly from nine flowers on three plants of each crossing seven days after pollination, fixed in FAA (formalin: acetic acid: 70% ethanol=1:1:18) for 24 h at room temperature, and softened in 8 N NaOH at 60°C for 1.5 h. The samples were then stained with a 0.1% aniline blue solution (Kho and Baër, 1968) at 4°C for 24 h, quashed onto slides in a drop of glycerin and observed under a fluorescent microscope (Zeiss Axioskop2, EM: BP 365 to 390 nm). A pollen grain showing tube length exceeding twice its own diameter was considered germinated.

Examining the efficacy of cut-column pollination method

The cultivar of SG with difficulty in hybridization, used as the pistillate parent, was treated with cut-column pollination method used cultivars of BII, V3 and V31, and *P. violacea* as the pollen parents without breeding barriers, for evaluating the efficacy of the given method on overcoming the breeding barriers of SG. The procedures of cut-column pollination method used in this study were conducted as follows. 1. The sepals and petals of pistillate parent were removed from the base. 2. The upper part of column was cut off, with the stigma cavity intact. 3. Pollinia of pollen parent were then placed onto stigma cavity of pistillate parent. 4. The stigma cavity was subsequently sealed with paraffin film (Fig. 1). The occurrence of capsule set was collected at 60 DAP. The fertility and offspring production were examined by asym-

Table 1. Crossing performance, chromosome number and source of the Phalaenopsis cultivars/species used in this study

Description	Cultivar/species name (abbreviation)	Chromosome number	Reference	Source of plant materials
Difficult in hybridization	Phalaenopsis Sunrise Goldmour 'KHM637' (SG)	2n=4x=76	Chuang <i>et al.</i> , 2008	I–Hsin Biotechnology Inc.
Normal in hybridization	P. Brother Irene Irene 'Feng Fong' (BII)	2n=4x=76	Hsu <i>et al.</i> , 2010	Star Orchids
	P. Sogo Yukidian 'V3' (V3)	2n=4x=76	Hsu et al., 2010	Taiwan Sugar Corporation
	P. Tai Lin Redangel 'V31' (V31)	2n=4x=76	Hsu et al., 2010	Star Orchids
	P. violacea	2n=2x=38	Kao <i>et al.</i> , 2001	NCYU ^z

^z NCYU: National Chiayi University.



Fig. 1. Schematic illustration of the cut-column pollination method used in this study to overcome breeding barriers in *Phalaenopsis* Sunrise Goldmour 'KHM637', which had stigma closure problem after pollination. The sepals and petals of intact flower (A) of pistillate parent were removed from the base (B). The upper part of column was cut off, with the stigma cavity intact (C). Pollinia of pollen parent were then placed onto stigma cavity of pistillate parent (D). The stigma cavity was subsequently sealed with paraffin film (E). biotic seed germinating abilities according to the method recommended by Lin *et al.* (2008). Four to nine crossings were evaluated in each of crossing combinations.

RESULTS

Crossing behaviors in self-pollination

In the self-crossings, the three normal cultivars, *Phalaenopsis* Brother Irene Irene 'Feng Fong' (BII), *P*. Sogo Yukidian 'V3' (V3), and *P*. Tai Lin Red Angel 'V31' (V31), all showed 100% of stigma cavity closure at three days after pollination (DAP), 100% of in vivo pollen tube elongation at 7 DAP, and 100% capsule set at 60 DAP, and finally produced over 900 offsprings (Table 2). However, the stigma cavity closure and pollen tube elongation did not occur, and none of offspring had been produced in the self-crossings of *P*. Sunrise Goldmour 'KHM637' (SG) with difficulties in hybridization (Table 2).

Crossing behaviors in reciprocal pollination

To further elucidate breeding barriers, the cultivar with difficulties in hybridization (SG) was reciprocally– crossed with each of the three normal cultivars (BII, V3 and V31). Both phenomena of stigma cavity closure and *in vivo* pollen tube elongation did not occur when SG was used as the pistillate parent (Table 3). When BII, V3 and V31 were used as pistillate parents, the phenom-

Cultivar name ^v	Stigma cavity closure (%) (3 DAP)	In vivo pollen tube elongation (%) (7 DAP)	Capsule set (%) (60 DAP)	Fertile capsule (%)	No. of offsprings
SG	0 b ^x	0 b	0 b	0 b	0
BII	100 a	100 a	100 a	100 a	>900
V3	100 a	100 a	100 a	100 a	>900
V31	100 a	100 a	100 a	100 a	>900

Table 2. Occurrence of stigma cavity closure, pollen tube elongation, capsule set, and fertility in self-crossings² of three normal *Phalaenopsis* cultivars and the cultivar of *P*. Sunrise Goldmour 'KHM637' with breeding barriers

 $^{\rm z}$ Two separate experiments were conducted to investigate pollen tube development and fertility. Nine flowers were examined in each self–crossing.

^y Abbreviations of the cultivar name as shown in Table 1. DAP, days after pollination.

 $^{\rm x}$ Mean separation within columns by LSD at 5% level.

Table 3. Occurrence of stigma cavity closure, pollen tube elongation, capsule set, and fertility in reciprocal-crossed
combinations^z of three normal *Phalaenopsis* cultivars and the cultivar of *P*. Sunrise Goldmour 'KHM637' with
breeding barriers

Combination ^y $(\stackrel{\circ}{+} \times \stackrel{\circ}{\triangleleft})$	Stigma cavity closure (%) (3 DAP)	In vivo pollen tube elongation (%) (7 DAP)	Capsule set (%) (60 DAP)	Fertile capsule (%)	No. of offsprings
SG×BII	0 b ^x	0 a	0	0	0
SG×V3	0 b	0 a	0	0	0
SG×V31	0 b	0 a	0	0	0
BII×SG	77.8 a	11.1 a	0	0	0
V3×SG	77.8 a	0 a	0	0	0
V31×SG	88.9 a	11.1 a	0	0	0

^z Two separate experiments were conducted to investigate pollen tube development and fertility. Nine flowers were examined in each reciprocal–crossing.

Abbreviations of the cultivar name as shown in Table 1. DAP, days after pollination.

 $^{\rm x}$ Mean separation within columns by LSD at 5% level.

ena of stigma cavity closure were observed at 3 DAP and reached 77.8%, 77.8% and 88.9%, respectively, and *in vivo* pollen tube elongation were partially observed at 7 DAP and reached 11.1%, 0% and 11.1%, respectively (Table 3). No offspring was obtained when using SG as either pollen or pistillate parent (Table 3).

Efficacy of cut-column pollination method

To explore performance of cut-column pollination method, the cultivar of SG with difficulty in hybridization was used as the pistillate parent, after the removal of the upper part of column but with the stigma cavity intact, and then pollinated with pollinia of three cultivars, BII, V3 and V31, and one species, *P. violacea*, respectively. The results showed that capsule sets collected at 60 DAP reached 22.2%, 77.8%, 100.0% and 75.0%, for these four

combinations used pollinia of BII, V3, V31 and P. violacea, respectively (Table 4). Capsule sets were achieved by cut-column pollination method in these four combinations (Fig. 2A, left). In contrast, the flower of SG without conducting procedures of cut-column pollination method wilted after pollination (Fig. 2A, right). Both the capsule fertility and offspring production were then examined using aseptic seed germination when mature capsule (Fig. 2B) was successfully obtained. The results showed that 55.6%-77.8% of fertile capsule formations were achieved except the combination of using pollinia from BII, and offsprings were successfully obtained from other three of these four combinations (Table 4). Some of these hybrid seedlings have been cultivated into healthy flowering plants (Fig. 2) for further selection and breeding.

 Table 4. Effects of cut-column pollination method on capsule set, fertility, and offspring production in *Phalaenopsis* Sunrise Goldmour 'KHM637', the cultivar with stigma closure problem controlled by pistillate parent itself, used as the pistillate parent

Combination ^z $(\stackrel{\circ}{+} \times \stackrel{\circ}{\mathcal{J}})$	No. of crossing	Capsule set % (60 DAP)	Fertile capsule (%)	No. of offsprings
SG×BII	9	22.2 c ^y	0 b	0
SG×V3	9	77.8 b	77.8 a	83
SG×V31	9	100.0 a	55.6 a	45
SG×P. violacea	4	75.0 b	75.0 a	149

^z Abbreviations of the cultivar name as shown in Table 1. DAP, days after pollination.

^y Mean separation within columns by LSD at 5% level.



Fig. 2. The performance of using cut–column pollination method to overcome breeding barriers in *Phalaenopsis* Sunrise Goldmour 'KHM637', which had stigma closure problem after pollination. The capsule was set following cut–column pollination method (A, left). The flower wilted at 16 days after pollination without cut–column (A, right). Mature capsule was successfully obtained (B). Seeds germinated under aseptic culture (C). The potted hybrid seedlings were obtained (D). The flower of a hybrid derived from crossing the cut–column, pistillate parent *P*. Sunrise Goldmour 'KHM637' with the pollen parent *P. violacea* (E). Scale bars, 1 cm.

DISCUSSION

Breeding barriers in producing hybrids by intergeneric, inter- and intraspecific hybridization in Phalaenopsis have been observed. However, the causes and mechanisms of such breeding barriers have not been thoroughly studied. Approaches to elucidate the basis of reproductive barriers between specific crossing combinations in *Phalaenopsis* have been conducted by our research team (Chuang et al., 2008; Chuang et al., 2013; Hsu et al., 2010). Six cultivars which are known to be difficult in hybridization, including Doritaenopsis Ever Spring Prince '#1' (ESP), D. Queen Beer 'KHM 159' (QB), D. Taida Salu 'KHM 101' (TS), P. Brother Girl 'B' (BG), P. Taipei Gold 'STM' (TG), and P. Sunrise Goldamour 'KHM637' (SG), and three cultivars with normal behaviors in crossing, including P. Brother Irene Irene 'Feng Fong' (BII), P. Sogo Yukidian 'V3' (V3), and P. Tai Lin Red Angel 'V31' (V31), were investigated for the mechanisms underlying the crossing barriers.

The chromosome numbers analyzed by root tip squash showed that the five of selected cultivars with difficulties in hybridization are triploids with 2n=3x=57 chromosomes, while SG and the other three normal cultivars are tetraploids with 2n=4x=76 chromosomes (Chuang *et al.*, 2008; Hsu *et al.*, 2010). These results suggested that the triploidy of these five selected cultivars might have contributed to their crossing difficulties.

The crossing results showed that the closure of stigma cavities occurred in TS and BG, while such stigma closure was never observed when ESP or QB was used as the pollen parent in crosses (Hsu *et al.*, 2010). Neither closure of the stigma cavities nor elongation of pollen tubes occurred when TG or SG was self–crossed, or was used as the pistillate parent during intraspecfic crossing (Chuang *et al.*, 2008). These results suggested that crossing difficulties in those selected cultivars might be caused by odd chromosome sets and some other factors of pollen or pistillate parents.

Stigma closure problem of *P*. Sunrise Goldmour 'KHM637' not limited to specific crossing combinations

P. Sunrise Goldmour 'KHM637' (SG) is known to have stigma closure problem after pollination and be difficult in hybridization (Chuang *et al.*, 2008). After pollination a series of physiological and morphological changes, such as pollen tube growth and fertilization to the seed set may develop. Usually the stigma cavity of *Phalaenopsis* flower would close 48 hours after pollination (Goh *et al.*, 1982; O'Neill *et al.*, 1993; Zhang and O'Neill, 1993). Use of fluorescence to monitor pollen tube behavior in the style and ovary provided a rapid and correct interpretation in studying self– and cross–incompatibility. Therefore, the phenomena of stigma cavity closure and pollen tube behavior in the column could be easily observed for evaluating the pollination process of each crossing combination.

In the self-crossings, as expected the three normal

cultivars, BII, V3 and V31, all showed stigma cavity closure and pollen tube elongation, and finally produced offsprings (Table 2). However, the stigma cavity closure and pollen tube elongation did not occur, and none of offspring had been produced in the self-crossings of SG (Table 2).

To further elucidate breeding barriers, SG was reciprocally-crossed with each of the three normal cultivars (BII, V3 and V31). Both phenomena of stigma cavity closure and *in vivo* pollen tube elongation did not occur when SG was used as the pistillate parent (Table 3). The failure to produce any offspring in self- and reciprocal crossings of SG might be caused by some uncertain maternal parent factors in itself and not limited to specific crossing combinations, since stigma cavity closure and pollen tube elongation did not occur when it was used as a pistillate parent, no matter what pollen parents used (Tables 2 and 3; Fig. 2A, right).

Stigma closure problem not the only reason accountable for failure to produce offspring in crossing of *P*. Sunrise Goldmour 'KHM637'

Several methods including the stigma cavity sealing with paraffin film, applying different quantity of pollinia, and auxins treatments have been conducted by us to attempt to overcome breeding barriers in P. Sunrise Goldmour 'KHM637' (SG) and P. Taipei Gold 'STM' (TG) (Chuang *et al.*, 2008). However, *in vivo* pollen tube elongation did not occur by sealing stigma cavity with paraffin film when SG or TG was self-crossed, or was used as the pistillate parent and pollinated with pollonia of V3 (Chuang *et al.*, 2008). Obviously the effect of stigma closure problem after pollination could not be corrected simply by sealing.

The postpollination development in orchid flowers is precisely and completely triggered by pollination. This postpollination developmental syndrome includes the induction and coordination of ovary and ovule development in preparation for fertilization. The development of the ovary as well as other aspects of the postpollination syndrome has an absolute requirement for the participatory action of auxin (Zhang and O'Neill, 1993). It has long been proposed that the causative agent in the postpollination response is auxin deposited on the stigma with the pollinia (Burg and Dijkman, 1967). Indeed, there are a number of reports indicated that auxin is a natural component of orchid pollen (Arditti, 1979; Stead, 1992).

In light of this association between auxin and postpollination response in orchids, the authors have also examined the effects of quantity of pollinia and auxin on the stigma cavity closure and *in vivo* pollen tube elongation in crossing combinations of using SG or TG as the pistillate parent and V3 as the pollen parent (Chuang *et al.*, 2008). However, both phenomena of stigma cavity closure and *in vivo* pollen tube elongation did not occur even when four pollinia were used in each pollination. The authors found that exogenous application of 2.9 mM indole–3–acetic acid (IAA) or naphthalene acetic acid (NAA) to the stigmatic surface of SG and TG results in occurrence of stigma cavity closure, however, not in *in* *vivo* pollen tube elongation, capsule set, and offspring production. These results indicated that stigma closure problem might not be the only reason accountable for failure to produce offspring in crossing of *P*. Sunrise Goldmour 'KHM637'.

Applicability of the cut–column pollination method less limited to pollen parents

Sexual barriers preventing intra- or interspecific hybridization have been distinguished into pre- and post-fertilization barriers. The nature of the barrier determines the method to be used to overcome the specific barrier. A range of techniques such as pollen heating, electrical stimulation, chemical treatment, bud pollination, mentor pollination, style grafting, cut-style and placental pollination have been developed to overcome pre-fertilization barriers for diverse crops (Janson, 1993; Janson *et al.*, 1993; Magdalita *et al.*, 1998; Ronald and Ascher, 1975; Shen and Liao, 1976; Van Tuyl *et al.*, 1988; Van Tuyl *et al.*, 1991; Vervaeke *et al.*, 2004; Visser, 1983).

The authors have developed a technique for overcoming breeding barriers in *Phalaenopsis* and named the technique as "cut-column pollination method" (Chuang *et al.*, 2013), since the technique was modified from the cut-style pollination using in lily (Janson *et al.*, 1993; Van Tuyl *et al.*, 1991) and a distinguishing feature of the Orchidaceae is the gynostemium or column, which is the fusion of the style, stigma, and stamens.

The authors have also demonstrated that this cutcolumn pollination method proved to overcome breeding barriers in *P*. Taipei Gold 'STM' (TG), known to have stigma closure problem after pollination and be difficult in hybridization (Chuang *et al.*, 2013). *P*. Sunrise Goldmour 'KHM637' (SG) has shown similar hybridization behavior to TG (Chuang *et al.*, 2008). In this study, the authors aimed to examine the efficacy of cut-column pollination method for overcoming the breeding barriers of SG, and to explore the applicable extensibility of the given method.

The cultivar of SG, used as the pistillate parent, was treated with cut-column pollination method and then pollinated with pollinia of three cultivars, BII, V3 and V31, and one species, P. violacea, respectively. The results showed that capsule sets were all achieved by cut-column pollination method in these four combinations, and both the capsule fertility and offspring production were mostly obtained except the combination of using pollinia from BII (Table 4). The similar situation was also found in combinations which cultivar of TG, used as the pistillate parent, was treated with cut-column pollination method and then pollinated with pollinia of three cultivars, BII, V3 and V31, and two species, P. violacea and P. fasciata, respectively (Chuang et al., 2013). These results suggested that the failure to produce any offspring in combination of SG and BII, and of TG and BII might be caused by similar uncertain interaction between partners of each combination.

In a comparison of the cut-style technique using in lily (Janson *et al.*, 1993; Van Tuyl *et al.*, 1991), it was shown that some pre-fertilization barriers in *Phalaenopsis* could be circumvented by using the cutcolumn pollination method. If the sterility was caused due to existence of pre-fertilization barriers in upper part of column, fertility might be restored by the cut-column pollination method. In this way pollinia bypassed column barriers which could inhibit pollen tube growth, without incidence of stigma cavity closure.

In conclusion, the cut-column pollination method might be applicable to other *Phalaenopsis* cultivars with similar crossing behaviors of *P*. Sunrise Goldmour 'KHM637' and *P*. Taipei Gold 'STM' for overcoming the crossing barriers, and the applicability of the given method tends to be less limited to pollen parents without breeding barriers.

REFERENCES

- Ardl
tti, J. 1979 Aspects of the physiology of orchids. Adv. Bot. Res.
 $\mathbf{7}:$ 421–655
- Burg, S. P. and M. J. Dijkman 1967 Ethylene and auxin participation in pollen induced fading of Vanda orchid blossoms. *Plant Physiol.* 42: 1648–1650
- Christenson, E. A. 2001 *Phalaenopsis*: A monograph. International *Phalaenopsis* Alliance, Timber Press, Inc. U.S.A.
- Chuang, H. T., K. L. Huang, and S. T. Hsu. 2013 Using cut–column pollination method to overcome breeding barriers in *Phalaenopsis* Taipei Gold 'STM'. J. Taiwan Soc. Hort. Sci. 59: 103–111
- Chuang, H. T., S. T. Hsu, and T. M. Shen. 2008 Breeding barriers in yellow *Phalaenopsis* orchids. J. Taiwan Soc. Hort. Sci. 54: 59–66
- Customs Administration. 2012 Trade statistics. Trade statistics search system. http://www.customs.gov.tw/StatisticWeb/News. aspx>
- Goh, C. J., M. S. Strauss and J. Arditti 1982 Flower induction and physiology in orchids, p. 213–241. In: J. Arditti (ed.). Orchid Biology: Reviews and Perspectives, II. Cornell Univ. Press, Ithaca, N.Y. U.S.A.
- Hsu, S. T., H. T. Chuang, and T. M. Shen 2010 Breeding barriers in red *Phalaenopsis* orchids. Acta Hort. 878: 145–152
- Janson, J. 1993 Placental pollination in *Lilium longiflorum* Thunb. *Plant Sci.* 90: 105–115
- Janson, J., M. C. Reinders, J. M. Van Tuyl and C. J. Keijzer 1993 Pollen tube growth in *Lilium longiflorum* following different pollination techniques and flower manipulations. *Acta Botanica Neerlandica* 42: 461–472
- Kao, Y. Y., S. B. Chang, T. Y. Lin, C. H. Hsieh, Y. H. Chen, W. H. Chen, and C. C. Chen. 2001 Differential accumulation of heterochromatin as a cause for karyotype variation in *Phalaenopsis* orchids. *Ann. Bot.* 87: 387–395
- Kho, Y. O. and J. Baër. 1968 Observing pollen tubes by means of fluorescence. *Euphytica* 17: 298–302
- Lin, H. J., H. T. Chuang, S. T. Hsu and T. M. Shen 2008 Effects of developmental temperature and capsule harvest time on the aseptic seed germination in *Phalaenopsis aphrodite* subsp. *formosana. Plant Nursery* 10: 59–71
- Magdalita, P. M., R. A. Drew, I. D. Godwin and S. W. Adkins 1998 An efficient interspecific hybridization protocol for *Carica* papaya L. × C. cauliflora Jacq. Aust. J. Exp. Agric. 38: 523–530
- O'Neill, S. D., J. A. Nadeau, X. S. Zhang, A. Q. Bui and A. H. Halevy 1993 Interorgan regulation of ethylene biosynthetic genes by pollination. *Plant Cell* 5: 419–432
- Ronald, W. G. and P. D. Ascher 1975 Effects of high temperature treatments on seed yield and self incompatibility in *Chrysanthemum. Euphytica* 24: 317–322
- Royal Horticultural Society. 2013 Plant registration. http://www.

rhs.org.uk/Plants/Plant-science/Plant-registration/Orchids>

- Shen, T. F. and C. I. Liao 1976 Factors affecting the degree of selfincompatibility and the breaking of self-incompatibility in Yehsen cabbage. *Sci. Dev.* **4**: 2298–2305
- Stead, A. D. 1992 Pollination-induced flower senescence: A review. *Plant Growth Regul.* **11**: 13–20
- Van Tuyl, J. M., M. P. Van Diën, M. G. M. Van Creij, T. C. M. Van Kleinwee, J. Franken and R. J. Bino 1991 Application of *in vitro* pollination, ovary culture, ovule culture and embryo rescue for overcoming incongruity barriers in interspecific *Lilium* crosses. *Plant Science* 74: 115–126

Van Tuyl, J. M., T. P. Straathof, R. J. Bino and A. A. M. Kwakkenbos

1988 Effect of three pollination methods on embryo development and seed set in intra- and interspecific crosses between seven *Lilium* species. *Sex Plant Reprod.* **1**: 119–123

- Vervaeke, I., R. Delen, J. Wouters, R. Deroose and M. P. de Proft 2004 Semi in vivo pollen tube growth of Aechmea fasciata. Plant Cell, Tissue Organ Cult. 76: 67–73
- Visser, T. 1983 The role of pioneer pollen in compatible and incompatible pollinations of apple and pear. *Acta Hortic.* **139**: 51–57
- Zhang, X. S. and S. D. O'Neill 1993 Ovary and gametophyte development are coordinately regulated by auxin and ethylene following pollination. *Plant Cell* 5: 403–418