Safety Evaluation and Antimutagenic Activity of Bamboo/Wood Vinegars Collected at Different Temperatures

Lin, Han Chien

Laboratory of Environment Functional Materials, Department of Wood Based Materials and Design,
College of Agriculture, National Chiayi University

Chen, Po-Kuang

Graduate Institute of Forest Products Science and Furniture Engineering, College of
Agriculture, National Chiayi University : Master

Lai, Ying-Jang

Department of Food Science, College of Science and Engineering, National Quemoy University

Wu, She-Ching

Department of Food Science, College of Life Sciences, National Chiayi University,

他

<https://doi.org/10.5109/1467647>

出版情報 : 九州大学大学院農学研究院紀要. 59 (2), pp.359-368, 2014-08-29. Faculty of
Agriculture, Kyushu University

バージョン :

権利関係 :

Safety Evaluation and Antimutagenic Activity of Bamboo/Wood Vinegars Collected at Different Temperatures

Han Chien LIN^{1*}, Po-Kuang CHEN², Ying-Jang LAI³, She-Ching WU⁴,
Gwo-Shyong HWANG⁵ and Noboru FUJIMOTO⁶

Laboratory of Wood Material Technology, Division of Sustainable Bioresources Science,
Department of Agro-environmental Sciences, Faculty of Agriculture,
Kyushu University, Fukuoka 812-8581, Japan

(Received April 30, 2014 and accepted May 12, 2014)

The biological action of bamboo/wood vinegars collected at different temperatures from the exit of chimney of earthen kiln was evaluated by *Salmonella* mutagenesis assay, as a safety evaluation (Ames test) and reverse mutation assay (antimutagenic activity). The compounds of the vinegars were analyzed using gas chromatography–mass spectroscopy analysis. The acid, phenol and ketone compounds of bamboo vinegars were 10.65–20.09%, 57.87–65.98% and 10.13–18.76%, and the compounds of wood vinegars were 4.27–14.51%, 50.23–65.03% and 12.93–25.26%, respectively. The vinegars' safety showed that neither cytotoxicity nor mutagenicity toward *Salmonella typhimurium* TA98 and TA100 with S9 mix (an external metabolic activation system) at the diluting percent content of vinegars were lower than 20.00% or less, and without the S9 mix was at 33.33% or less. The vinegars at a diluting percent content below 20.00% expressed a dose-dependent inhibitory effect against both 4-nitroquinoline-*N*-oxide (NQNO), a direct mutagen, and the mutagenicity of aflatoxin B₁ (AFB₁), an indirect mutagen which requires metabolic activation, in *Salmonella typhimurium* TA98 and TA100. The inhibition of the vinegars against NQNO and AFB₁ toward TA100 was better than those toward TA98. In addition, the main percent of phenol and ketone compounds in the vinegars showed cytotoxicity/mutagenicity and an antimutagenic effect against the strains mentioned above, which may partially account for the biological action of bamboo/wood vinegars.

Key words: Ames test, Antimutagenic Activity, Bamboo/Wood Vinegar, Gas Chromatography–Mass Spectroscopy, Safety Evaluation

INTRODUCTION

Both bamboo and wood vinegar, by-products of the bamboo or wood charcoal manufacturing, are brown-red transparent liquids collected during pyrolysis of bamboo or wood charcoal. Yatagai *et al.* (1988) reported that with the application of bamboo/wood vinegars, even the compounds are complex and different, and are mainly able to be divided into three main portions: acid, phenol and neutral compounds. The vinegars consist of 80–200 compounds: 32% organic acid, 40% phenolic compound, 3% aldehyde, 5% alkone compound, 5% alcohol compound, 4% ester compound and 5% others. When both are dehydrated there is usually 80% water (Ikimoto and Ikeshima,

2000; Nomura, 2004; Uchimura *et al.*, 2000; Tsai *et al.*, 2009). The organic compounds in bamboo/wood vinegars may have practical applications even when present in only trace quantities (Uchimura *et al.*, 2000; Lin *et al.*, 2006).

Bamboo vinegar is effective in improving soil, promoting crops and preventing worm growth, as well as reducing agricultural chemicals, compost odor and sterilization (Ikimoto and Ikeshima, 2000). Recently, bamboo vinegar products have been developed that are beneficial for promoting growth of plants to as a plant root growth promoter or a pH value adjuster of cultural media (Huang *et al.*, 2011; Lin *et al.*, 2011; Ho *et al.*, 2013). It is also effective when used against allergies (Hageta, 2004), in healthy drinks (Kobahasi, 2004; Nomura, 2004; Yoshie, 2004), as a virus/fungi/bacterial resistant (Kou, 2004; Lin *et al.*, 2006; Lin and Shiah, 2006; Lu *et al.*, 2007; Lin *et al.*, 2008; Chen *et al.*, 2010) and as an agent of antioxidation, especially for a resistant lipid oxidation effect (Tsai *et al.*, 2009). The compounds of wood vinegar are much more complex than those of bamboo vinegar (Chen *et al.*, 2006; Hwang *et al.*, 2008), and can repel termites and disease germs of plants, be fungi resistant, improve the soil, promote crops or prevent worm growth and reduce the use of agricultural chemicals, such as a natural insecticide or soil fungicide (Ishihara, 1996; Nishimiya *et al.*, 1998; Yatagai, 2002; Lu *et al.*, 2007; Lin *et al.*, 2009). As stated in the above references, the commercial production of bamboo/wood vinegar is being increased and highly valued for its various effective uses in Taiwan.

Horne and Paul (1996) reported that bamboo/wood

¹ Laboratory of Environment Functional Materials, Department of Wood Based Materials and Design, College of Agriculture, National Chiayi University, Chiayi, Taiwan, ROC.

² Master, Graduate Institute of Forest Products Science and Furniture Engineering, College of Agriculture, National Chiayi University, Chiayi, Taiwan, ROC.

³ Department of Food Science, College of Science and Engineering, National Quemoy University, Quemoy, Taiwan ROC.

⁴ Department of Food Science, College of Life Sciences, National Chiayi University, Chiayi, Taiwan, ROC.

⁵ Divisions of Forest Utilization, Taiwan Forestry Research Institute (TFRI), Taipei, Taiwan, ROC.

⁶ Laboratory of Wood Material Technology, Division of Sustainable Bioresources Science, Department of Agro-environmental Sciences, Faculty of Agriculture, Kyushu University, Japan.

* Corresponding author (E-mail: alexhlin@mail.ncyu.edu.tw)

vinegars collected at the exit of chimney of earthen or furnace kiln, when the carbonization temperature of bamboo/wood was raised to over 500°C, produced some carcinogenic polycyclic aromatic hydrocarbons (PAHs), such as naphthalene and phenanthrene (Namiki, 1990; Vasavada and Cornforth, 2006). The concentrations of these toxics increased with the increase in temperature as well. Even though the council of agriculture in Taiwan has submitted certified agricultural standards of forest products (2004) to prove that it is necessary for bamboo/wood vinegars to be collected at 80–140°C at an exit of chimney of earthen kiln and below 350°C for a furnace kiln, collected bamboo/wood vinegars from above this range of temperatures are necessary to evaluate the potential of mutagenic and carcinogenic agents, due to the fact that both of them are omnipresent in the human environment and seem impossible to completely eliminate. Ames *et al.* (1975) reported that for screening of environmental mutagens and carcinogens, the Ames test (Safety evaluation), a convenient method to evaluate mutagenic activities of these chemicals, has been developed, and McCann *et al.* (1975) and Shugimura *et al.* (1976) have suggested that the mutagenic activities of a number of chemicals correlate well with the carcinogenic activities.

To expect to be one and/or a reference of functional additives for bamboo/wood vinegars, the safety evaluation, including cytotoxicity and mutagenicity, of the vinegars using the bacterial mutation assay on *Salmonella typhimurium* TA98 and TA100 strains with and without an extrinsic metabolic activation system were evaluated. Moreover, because the antimutagenic properties have an array of prospective applications in human care, such as the increasing application in drinks, food antioxidation, etc., and have not been reported for the antimutagenic activities of bamboo/wood vinegars that have been made so far, the antimutagenic activity of the vinegars are investigated as well. These vinegars, collected at different temperatures from the exit of earthen kiln chimney, were also investigated for their basic compound by using gas chromatography–mass spectroscopy.

MATERIALS AND METHODS

Specimen preparation

Bamboo/wood vinegars

The vinegars were collected at different temperatures, ranging from 80°C to 150°C (Uchimura *et al.*, 2000), during the manufacturing process (pyrolysis) of charcoal from Moso bamboo (*Phyllostachys heterocycla* Milf) and branches/tree tops (BTW) of *Cryptomeria japonica* and were provided by the Division of Forest Utilization, TFRI Taipei, Taiwan (Hwang *et al.*, 2006; Hwang *et al.*, 2008). The bamboo and wood vinegars were based on the temperatures measured by a thermocouple at the exit of earthen kiln chimneys during bamboo/wood charcoal pyrolysis. The different groups of bamboo vinegar were collected at 80–150°C, and with categories over 80, 90–92, 99–102, 120–123 and 145–150°C. The various collected temperatures of wood vinegar were 80–159°C, and

with categories 80–90, 91–100, 101–120, 121–140 and 141–159°C. The basic properties of the bamboo/wood vinegars refer to (Lin *et al.*, 2008; Lin *et al.*, 2009).

Test strains

Salmonella typhimurium (*S. typhimurium*), including TA98 and TA100, was bought from the Bioresource Collection and Research Center, Food Industry Research and Development Institute.

Rat liver mixture

The rat liver mixture (S9 mix) (Organ Teknika Co., Switzerland) was prepared from Sprague–Dawley male rats treated with Aroclor 1254.

Mutagens

The mutagens are 4-nitroquinoline-*N*-oxide (NQNO) and aflatoxin B1 (AFB1). They were obtained from the Sigma Chemical Co. (Steinheim, Germany). All reagents used in the test were of analytical grade.

Experimental methods

Compounds analysis

Gas chromatography–mass spectroscopy (GC–MS) analysis was performed with a Varian CP–3800 chromatography instrument, combined with a Saturn–2000 mass spectrometer equipped with an electron ionization and quadrupole analyzer. The vinegars collected at different temperatures were diluted 10 times using acetone. The flow rate was set at 1.0 mL/min and the carrier gas was nitrogen. The size of the capillary column (DB–5MS, J&W Scientific, Folsom, CA, USA) was 30.0 m×250 µm i.d. and its film thickness was 0.25 µm. The temperatures of the ion source and interface were set at 280°C, and the mass scan range was from 40 to 550 amu. The electron energy was set at 70 eV. The oven temperature was programmed to hold at 40°C for 5 min and then, increased to 280°C at a rate of 12°C/min. The GC–MS chromatographic spectra of various bamboo/wood vinegar compounds were analyzed by the software of MS Workstation (Tsai *et al.*, 2009).

Ames test

Cytotoxicity

0.1 mL of bamboo/wood vinegars were diluted to a percent content of 50, 33.33, 25, 20, 13.33 and 10%. The specimens were put in the test tube and mixed with 0.1 mL phosphate buffer saline and 0.1 mL *S. typhimurium* TA98 and TA100, and cultured overnight in nutrient broth. If there was additional S9 mix, the aforesaid addition level was changed to 0.2 mL phosphate buffer saline, 0.1 mL *S. typhimurium* TA98 and TA100 and cultured overnight in nutrient broth and 0.5 mL S9 (with S9 mix) or zero S9 (without S9 mix). Afterwards, the test tube was pre-cultured at 37°C for 20 min. The mixed diluent was taken out, and then, 1 mL diluent was put in the plate, mixed with nutrient agar and shaken up. When the mixture solidified, the plate was placed in an incubator at 37°C for 48 h. The colony (bacterial) count was calculated; if the bacterial count of the test group (including with or without S9 mix) was greater than the bacterial count of the control group by 80% (the bacterial count rate, Survival; %), there was no toxicity (Ames and

McCann, 1975). The survival of cytotoxicity is calculated as follows:

$$\text{Survival (\%)} = \left(\frac{\text{the bacterial count of test group}}{\text{bacterial count of control group}} \right) \times 100$$

Mutagenicity

The mutagenicity is analyzed by using the method proposed by Maron and Ames (1983). The test vinegars for this mutagenicity test, with or without S9 mix, are the same as for the cytotoxicity test. 0.1 mL phosphate buffer, 0.1 mL *S. typhimurium* TA98 and TA100 cultured overnight in oxioid nutrient broth No.2 were put into the test tube, mixed with 0.5 mL phosphate buffer saline and cultured at 37°C for 20 min. The test mixture was mixed with 2 mL Molten Top Agar (including 0.05 mM L-histidine, 0.05 Mm Biotin and 0.09 M NaCl) uniformly; the nutrient agar then was poured into the plate. When the mixture solidified, the plate was put in the incubator at 37°C for 48 h, and then, the colony count was calculated. In addition, the phosphate buffer saline is only a pair of control groups (Blank). If the colony count of the TA98 and TA100 test groups is greater than that for the control group by more than two times, the specimen has mutagenicity. In other words, the Mutagenicity ratio (MR) = induced revertants per plate/spontaneous revertants per plate (Blank).

Antimutagenic activity

The test vinegars of the antimutagenic activity were assayed according to the Ames method (Maron and Ames, 1983). The mutagens that were diluted with dimethyl sulfoxide (DMSO) were NQNO (1 µg/plate for TA 98 and TA 100, respectively), a direct-acting mutagen; and AFB₁ (5 µg/plate for TA 98 and TA 100, respectively), which required S9 mix for metabolic activation. A mutagen (0.1 mL; contained 1 µg NQNO or 5 µg AFB₁) was added to the mixture of a strain (TA 98 or TA 100), and 0.1 mL of each test vinegar was added to the S9 mix for AFB₁ or to the phosphate buffer (0.1 mol/L, pH 7.4) for NQNO. The mutagenicity of each mutagen in the absence of an extract is defined as 100%. The inhibition (%) of mutagenicity of test vinegar is calculated as follows: Inhibition (%) = [1 - (Number of his⁺ revertants in the presence of the test vinegar - Number of spontaneous revertants) / (Number of his⁺ revertants in the absence of the test vinegar - Number of spontaneous revertants)] × 100

Statistical analysis

The test results of the cytotoxicity, mutagenicity and antimutagenicity are represented by an average value (standard deviation), and the control group and test group are compared by Duncan's analysis. If the *p* value is smaller than 0.05, meaning a significant difference between the test group and the control group, it is represented by different superscript upper case letters.

RESULTS AND DISCUSSION

Compounds of bamboo/wood vinegars

The identified compounds of bamboo/wood vinegars collected from different temperatures were analyzed

with GC-MS. The acid, phenol, ketone and other compounds were found in bamboo/wood vinegars. In general, the main fraction of bamboo/wood vinegars for an acid compound is acetic acid (Yatagai *et al.*, 1988). However, the retention time cannot be measured from the beginning point (zero) and is about 5 min, so that the acetic acid is not shown in the results of the study. In other words, the fraction percents of compounds from bamboo/wood vinegars are calculated only from the measured compounds, shown in Tables 1 and 2.

The acid, phenol, ketone and other compounds of bamboo vinegars were 10.65–20.09%, 57.87–65.98%, 10.13–18.76% and 9.66–15.30, respectively (Table 1). The acid compounds included butanoic acid, 2-methoxyethyl acetate, 4-hydroxy-butanoic acid and 4-hydroxy-3-methoxy-butanoic acid. The maximum fraction of acid compounds was the bamboo vinegar collected from 80–150°C. The phenol (5.93–16.60%), 2-methoxy-phenol (8.27–16.39%) and 4-ethyl-phenol (3.68–9.48%) were the main fractions of phenol compounds for bamboo vinegars. For ketone compounds, the 2-hydroxy-3-methyl-2-cyclopentenone-1-one, 2,3-dimethyl-2-cyclopentenone-1-one and maltol could be measured for bamboo vinegars collected at all temperatures. The maximum fraction of ketone compounds was the bamboo vinegar collected from 120–123°C.

For wood vinegars (Table 2), the acid, phenol, ketone and other compounds of wood vinegars were 4.27–14.51%, 50.23–65.03%, 12.93–25.26% and 9.57–19.92%, respectively. The main fractions of phenol compounds were phenol (3.36–11.30%), 2-methoxy-phenol (11.70–22.90%) and 2-methoxy-4-methyl-phenol (7.09–20.10%). Compared to both vinegars, the maximum fraction of ketone compounds for wood vinegar was also collected at temperatures from 120–123°C, and was higher than those for bamboo. Moreover, the range of the phenol compounds fraction for bamboo/wood vinegar was around 50–60%; among which, the maximum fractions of phenol compounds were phenol and 2-methoxy-phenol.

Cytotoxicity of bamboo/wood vinegars

According to the former results of compound analysis, phenol, ketone and other compounds are present in bamboo/wood vinegars. Some phenol compounds displayed antimicrobial activity (Yam *et al.*, 1997). If mutagenicity occurred in a treated material, the results of the antimutagenic assay would be affected and confused due to increased or decreased numbers of revertants of TA98 and TA100 (Duh and Yen, 1997). The phenol compounds of bamboo vinegars were 57.87–65.98%, and the compounds of wood vinegars were 50.23–65.03%. Therefore, the cytotoxicity of bamboo/wood vinegars must be determined before testing the mutagenicity and antimutagenicity of bamboo/wood vinegars. Cytotoxicity results for the bamboo/wood vinegars collected at temperatures of 90–92/105–109°C with the original vinegar (no diluting) and a range of diluting percent contents of 50.00, 33.33, 25.00, 20.00, 13.33 and 10.00% for *S. typhimurium* TA98 and TA100 without the S9 mix, as well as for with the S9 mix with diluting percent contents of 33.33,

Table 1. Various Compounds of bamboo vinegars collected from different temperatures by using Gas chromatography–mass spectroscopy

Identified compound	Collection temperature (°C)					
	80	90–92	99–102	120–123	145–150	80–150
Compound percent (%)						
Acid compound						
butanoic acid	1.91	1.77	1.37	1.79	2.23	4.58
2-methoxyethyl acetate	1.50	1.38	1.11	1.26	1.63	5.98
4-hydroxy–butanoic acid	8.98	9.02	8.17	6.84	7.89	8.11
4-hydroxy–3-methoxy–butanoic acid	–	–	–	1.44	0.84	1.42
Total	12.39	12.17	10.65	11.33	12.59	20.09
Phenol compound						
phenol	14.69	16.60	15.20	12.48	15.79	5.93
4-methyl–phenol	2.78	4.47	3.88	3.13	5.32	2.04
3-ethyl–phenol	4.60	6.44	6.06	8.35	8.81	6.69
2-methoxy–phenol	16.39	15.40	11.70	9.33	8.27	12.60
2,4-dimethyl–phenol	–	1.41	1.56	1.09	1.73	–
4-ethyl–phenol	8.26	9.48	8.60	6.64	6.98	3.68
2-methoxy–4-methyl–phenol	4.06	4.20	3.59	2.92	2.40	3.41
4-ethyl–2-methoxy–phenol	2.97	3.16	2.75	2.03	1.73	3.46
2,6-dimethoxy–phenol	5.77	4.82	9.69	11.90	8.76	20.20
Total	59.52	65.98	63.03	57.87	59.79	58.06
Ketone compound						
4-hydroxy–4-methyl–2-pentanone	3.07	2.80	2.28	2.14	5.73	–
1-hydroxy–3-methyl–2-butanone	–	–	1.66	1.46	1.51	–
1-methyl–2-propanone	–	–	0.86	0.94	–	–
3-methyl–2-cyclopentenone–1-one	1.19	–	0.90	2.31	2.22	1.38
2,5-dihydro–3,5-dimethyl–2-furanone	2.11	–	1.57	1.65	1.41	1.78
2-hydroxy–3-methyl–2-cyclopentenone–1-one	3.37	3.40	3.76	3.46	1.25	2.07
2,3-dimethyl–2-cyclopentenone–1-one	2.10	2.28	2.00	1.76	1.69	1.79
maltol	0.95	1.17	1.74	3.30	2.53	1.88
2-nonanone	–	1.05	–	0.89	1.02	–
1-(2,6-dihydroxy–4-methoxyphenyl)–ethanone	–	–	–	0.84	0.60	1.28
Total	12.79	10.70	14.77	18.76	17.96	10.13
Other compounds						
pyridine	9.36	6.46	5.86	5.02	4.32	5.87
4-methyl–1-penten–3-ol	4.92	3.65	3.08	2.35	–	–
3-propoxy–1-propene	–	–	–	1.17	1.72	1.96
tetrahydro–2,5-dimethyl–furan	–	–	1.30	1.50	1.66	–
2,6-dimethyl–heptane	1.02	1.04	1.31	2.00	1.96	3.89
Total	15.30	11.15	11.55	12.04	9.66	11.72

– : Trace amount (can not be measured)

25.00, 20.00, 13.33 and 10.00% are shown in Table 3.

The residual bacterial count of the control group (Blank) toward *S. typhimurium* TA98 and TA100 was 1838 and 1500. For bamboo vinegars collected at temperatures of 90–92°C, the residual bacterial count without S9 for various diluting percent contents was 0–1830 for TA98 and 87–1641 for TA100; for those with S9, it was 1605–2000 for TA98 and 1736–2249 for TA100. For wood vinegars at 105–109°C, the range of the residual

bacterial count without S9 was 607–2071 for TA98 and 1558–1917 for TA100, and with S9 was 1864–2242 for TA98 and 3181–3448 for TA100. Waleh *et al.* (1982) indicated that the residual bacteria rate of *S. typhimurium* must be over 80% of the control group to determine that the test group has no cytotoxicity for *S. typhimurium*. The residual bacteria rate (Survival, %) toward TA98 and TA100 for bamboo vinegar without the S9 mix at a diluting percent content 33.3 % or less, and with the

Table 2. Various Compounds of wood vinegars collected from different temperatures by using Gas chromatography–mass spectroscopy

Identified compound	Collection temperature (°C)					
	80–81	90–97	105–109	125–133	153–159	80–159
	Compound ratio (%)					
Acid compound						
butanoic acid	4.74	2.87	2.70	1.48	6.80	3.10
2-methoxyethyl acetate	3.96	3.29	2.66	1.27	2.43	4.36
4-hydroxy–butanoic acid	5.81	4.69	4.15	1.52	5.01	3.01
Total	14.51	10.85	9.51	4.27	14.24	10.47
Phenol compound						
phenol	8.12	8.70	9.48	10.88	11.30	3.36
4-methyl-phenol	2.94	3.60	4.46	4.81	4.36	1.60
3-ethyl-phenol	4.26	5.28	6.50	6.33	6.94	7.21
2-methoxy-phenol	18.20	13.39	12.40	14.98	11.70	22.90
2,4-dimethyl-phenol	–	1.75	2.05	1.29	1.98	–
4-ethyl-phenol	–	1.17	1.32	1.45	1.09	–
2-methoxy-4-methyl-phenol	13.80	11.29	10.30	7.09	9.83	20.10
4-ethyl-2-methoxy-phenol	4.97	4.02	3.77	3.15	3.90	7.50
2,6-dimethoxy-phenol	–	1.03	1.00	0.54	1.04	2.36
Total	52.29	50.23	51.28	50.55	52.11	65.03
Ketone compound						
4-hydroxy-4-methyl-2-pentanone	3.75	2.07	2.01	1.10	3.62	–
1-hydroxy-3-methyl-2-butanone	–	0.89	1.37	1.15	1.39	–
1-methyl-2-propanone	2.24	1.79	1.51	2.18	1.13	–
3-methyl-2-cyclopentenone-1-one	2.26	1.97	2.26	0.81	1.50	–
5-methoxy-2-pentanone	2.08	1.62	1.54	2.15	1.55	–
2,5-dihydro-3,5-dimethyl-2-furanone	2.26	2.36	2.08	3.12	1.44	2.21
2-hydroxy-3-methyl-2-cyclopentenone-1-one	6.98	6.45	6.57	5.70	5.50	5.21
2,3-dimethyl-2-cyclopentenone-1-one	2.08	1.52	1.51	1.87	1.63	2.83
maltol	1.98	3.58	4.09	5.00	3.10	2.68
2,9-decanedione	–	1.80	1.87	2.18	1.73	–
Total	23.63	24.05	24.81	25.26	22.59	12.93
Other compounds						
pyridine	3.14	3.88	3.22	4.57	2.70	3.43
4-methyl-1-penten-3-ol	4.25	3.55	2.70	4.68	–	–
2,5-dimethyl-furan	2.18	1.25	1.04	1.51	2.62	–
2-(1-methylethoxy)-1-propene	–	1.15	1.42	1.75	–	–
1-methoxy-cyclobutane	–	2.89	3.57	4.61	3.30	3.65
2,6-dimethyl-heptane	–	2.15	2.45	2.80	2.44	4.49
Total	9.57	14.87	14.40	19.92	11.06	11.57

– : Trace amount (can not be measured)

S9 mix at 25.00% or less was greater than 80%. The wood vinegars for survival without the S9 mix at a diluting percent content of 50.00% or less toward TA98 but for TA100 at 100% (original vinegar), and with the S9 mix at 33.33% or less toward TA98 and TA100 was higher than 80%.

For vinegars collected at all different temperatures, the Survival results are shown in Table 4. The Survival of all bamboo vinegars collected from 80–150, over 80,

90–92, 99–102, 120–123 and 145–150°C without the S9 mix at a diluting percent content of 33.33% or less were all higher than those for Blank by more than 80%. However, the Survival of the bamboo vinegar collected at temperatures of 99–102°C at a diluting percent content of 25% showed that the cytotoxicity toward *S. typhimurium* TA98 and TA100 with the S9 mix was lower than 80%, indicating “with toxicity”. For the collected temperatures of all wood vinegars at 80–159, 80–90,

for the mutagenicity test can be selected according to the results above.

Ames *et al.* (1975) reported that the number of spontaneous revertants induced by the specimen is less than for the control group by more than two times; the specimen has no mutagenicity. Table 5 shows the muta-

S9 mixture	Diluting percent content (%)	Bamboo vinegar (90–92°C)				Wood vinegar (105–109°C)			
		TA98	MR ¹⁾	TA100	MR	TA98	MR	TA100	MR
	Blank ²⁾	65 (4.93) ^{b3)}	1.00	132 (4.36) ^b	1.00	65 (4.93)b	1.00	132 (4.36) ^d	1.00
–S9	25.00	53 (8.19) ^a	0.81	129 (3.00) ^b	0.98	54 (5.29) ^a	0.83	98 (6.11) ^b	0.74
	20.00	42 (6.66) ^a	0.64	121 (4.51) ^{ab}	0.91	51 (1.73) ^a	0.78	120 (10.12)c	0.91
	13.33	53 (4.04) ^a	0.81	109 (5.69) ^a	0.83	50 (6.03) ^a	0.76	86 (7.81) ^{ab}	0.65
	10.00	41 (8.00) ^a	0.63	115 (10.26) ^a	0.87	51 (8.50) ^a	0.79	84 (2.31) ^a	0.63
	Blank	76 (0.58) ^b	1.00	140 (12.06) ^a	1.00	76 (0.58) ^a	1.00	140 (12.06) ^{ab}	1.00
S9	20.00	69 (4.51) ^{ab}	0.90	195 (10.97) ^b	1.39	81 (9.64) ^b	1.06	160 (3.51) ^{bc}	1.14
	13.33	73 (6.66) ^b	0.96	204 (13.23) ^b	1.45	92 (3.21) ^{bc}	1.20	188 (15.52) ^d	1.34
	10.00	74 (5.20) ^b	0.97	206 (3.06) ^b	1.47	82 (7.21) ^b	1.07	182 (15.82) ^{cd}	1.29

^{b)} Values (standard deviation) within a transverse with the different superscripts are significantly different ($p < 0.05$) by Duncan's multiple range tests

Specimens	Diluting percent content (%)	Collection temperature (°C)												
		80		90–92		99–102		120–123		145–150		80–150		
		TA98	TA100	TA98	TA100	TA98	TA100	TA98	TA100	TA98	TA100	TA98	TA100	
Bamboo vinegars	-S9	25.00	0.72	1.08	0.81	0.98	-	-	0.74	0.63	0.68	0.50	0.62	0.80
		20.00	0.68	1.02	0.64	0.91	0.65	0.90	0.81	0.62	0.59	0.54	0.67	0.76
		13.33	0.70	0.86	0.81	0.83	0.72	0.78	0.52	0.57	0.57	0.50	0.68	0.72
		10.00	0.61	0.93	0.63	0.87	0.75	1.00	0.69	0.77	0.69	0.56	0.73	0.60
	S9	20.00	1.10	1.47	0.90	1.39	0.88	1.57	1.18	1.19	1.32	1.32	1.16	0.83
		13.33	0.91	1.51	0.96	1.45	0.89	1.62	1.14	1.29	1.22	1.26	1.21	1.04
		10.00	1.04	1.42	0.97	1.47	0.96	1.66	1.00	1.28	1.20	1.40	1.35	1.05
		Wood vinegars	-S9	25.00	0.84	1.19	1.06	0.90	0.83	0.74	0.75	0.76	0.69	0.50
20.00	0.88			1.04	0.82	0.90	0.78	0.91	0.89	0.74	0.87	0.54	0.73	0.99
13.33	0.98			0.80	0.89	0.66	0.76	0.65	0.95	0.58	0.86	0.49	0.93	0.62
10.00	0.73			0.82	0.83	0.68	0.79	0.63	0.85	0.71	0.82	0.49	0.82	0.76
S9	20.00		1.07	1.27	0.99	1.14	1.06	1.14	1.07	1.22	1.22	1.32	1.06	1.15
	13.33		1.11	1.39	1.16	1.39	1.20	1.34	1.14	1.22	1.28	1.32	1.14	1.26
	10.00		1.09	1.25	1.08	1.26	1.07	1.29	1.06	1.26	1.11	1.34	1.15	1.28

genicity results for the bamboo/wood vinegars collected at temperatures of 90–92/105–109°C with a diluting percent content of 25.00, 20.00, 13.33 and 10.00% for *S. typhimurium* TA98 and TA100 without the S9 mix and

with the S9 for a diluting percent content of 20.00, 13.33 and 10.00%. The spontaneous revertants of Blank were 65 for TA98 and 132 for TA100, and of bamboo/wood vinegars without the S9 mix were from 41 to 54 for TA98

Table 7. Antimutagenicity of bamboo/wood vinegars collected at the temperatures of 90–92/105–109°C toward *Salmonella typhimurium* TA98 and TA100 without the S9 or with the S9 mix

Mutagens	Diluting percent content (%)	Bamboo vinegar (90–92°C)				Wood vinegar (105–109°C)			
		TA98	Inhibition ¹⁾ (%)	TA100	Inhibition (%)	TA98	Inhibition (%)	TA100	Inhibition (%)
	Blank ²⁾	1128 (93.15) ^{cd)}	0.00	1445 (75.80) ^c	0.00	1128 (93.15) ^b	0.00	1445 (75.8) ^e	0.00
NQNO (1 µg/plate)	25.00	807 (48.22) ^a	30.18	425 (52.79) ^a	77.65	906 (43.27) ^a	20.89	579 (61.13) ^b	65.92
	20.00	841 (71.59) ^a	27.04	446 (35.80) ^a	76.05	925 (70.27) ^a	19.07	620 (39.95) ^b	62.82
	13.33	891 (85.05) ^{ab}	22.33	594 (32.92) ^b	64.80	1058 (64.71) ^b	6.59	794 (57.58) ^c	49.57
	10.00	985 (64.26) ^b	13.49	616 (42.71) ^b	63.15	1084 (64.16) ^b	4.14	977 (61.78) ^d	35.65
Spontaneous revertants		65 (4.93)		132 (4.36)		65 (4.93)		132 (4.36)	
	Blank	1824 (57.86) ^d	0.00	2406 (86.00) ^c	0.00	1824 (57.86) ^c	0.00	2406 (86.00) ^d	0.00
AFB ₁ (5 µg/plate)	20.00	1114 (6.93) ^b	40.63	843 (25.48) ^b	68.97	1498 (83.21) ^{ab}	18.65	619 (21.39) ^a	78.89
	13.33	1292 (73.32) ^c	30.44	1012 (49.27) ^c	61.53	1510 (45.03) ^b	17.97	769 (45.71) ^b	72.27
	10.00	1319 (44.38) ^c	28.88	1202 (35.04) ^d	53.14	1587 (77.60) ^b	13.58	1051 (55.18) ^c	59.79
Spontaneous revertants		76 (0.58)		140 (12.06)		76 (0.58)		140 (12.06)	

¹⁾ Inhibition (%) = [1 – (Number of his⁺ revertants in the presence of the test vinegar – Number of spontaneous revertants) / (Number of his⁺ revertants in the absence of the test vinegar – Number of spontaneous revertants)] × 100

²⁾ Blank (the control group) was added without either bamboo or wood vinegars

³⁾ Values (standard deviation) within a transverse with the different superscripts are significantly different ($p < 0.05$) by Duncan's multiple range tests

Table 8. Inhibition of various bamboo/wood vinegars collected at different temperatures toward *Salmonella typhimurium* TA98, TA100 without the S9 or with the S9 mix

Specimens	Diluting percent content (%)	Collection temperature (°C)											
		80		90–92		99–102		120–123		145–150		80–150	
		TA98	TA100	TA98	TA100	TA98	TA100	TA98	TA100	TA98	TA100	TA98	TA100
Bamboo vinegars	25.00	25.03	70.14	30.18	77.65	–	–	27.85	74.45	2.76	81.77	22.96	68.97
	20.00	23.15	65.34	27.04	76.05	23.21	75.22	11.67	64.45	1.19	60.23	4.20	61.38
	13.33	12.80	60.64	22.33	64.80	20.58	58.91	4.08	61.55	1.00	40.71	0.75	57.31
	10.00	11.29	59.45	13.49	63.15	15.18	57.97	1.07	28.52	0.63	18.18	0.19	42.74
AFB ₁	20.00	24.38	62.94	40.63	68.97	37.80	66.35	20.03	75.24	11.63	73.06	20.14	74.86
	13.33	21.13	57.88	30.44	61.53	26.36	66.00	18.73	68.97	8.54	66.41	17.09	73.27
	10.00	18.31	37.43	28.88	53.14	24.26	54.88	13.20	59.88	0.92	60.50	7.44	58.44
Specimens	Diluting percent content (%)	80–81		90–97		105–109		125–133		153–159		80–159	
		TA98	TA100	TA98	TA100	TA98	TA100	TA98	TA100	TA98	TA100	TA98	TA100
Wood vinegar	25.00	16.88	43.37	19.39	41.26	20.89	65.92	22.02	52.77	5.02	67.19	31.43	54.95
	20.00	5.46	40.58	12.23	39.77	19.07	62.82	8.59	48.81	1.76	53.94	22.02	52.82
	13.33	4.89	32.86	8.72	37.07	6.59	49.57	1.69	33.42	–0.19	47.99	7.72	49.87
	10.00	2.32	32.10	5.40	35.63	4.14	35.65	0.75	32.96	–4.45	42.00	4.71	43.12
AFB ₁	20.00	18.96	61.88	20.71	71.68	18.65	78.89	16.94	69.18	7.13	70.56	25.06	71.15
	13.33	15.75	50.14	17.09	62.28	17.97	72.27	15.22	64.76	6.52	62.67	15.60	66.32
		11.67	12.03	8.93	61.88	13.58	59.79	14.99	59.06	3.62	59.35	11.56	64.53

Unit: %

and from 84 to 129 for TA100. With the S9 mix, the spontaneous revertants of bamboo/wood vinegars were 69–92 for TA98 and 160–206 for TA100. The results on the same Table 5 also show that the bamboo/wood vinegars in the test range did not exceed spontaneous revertants by more than two times for TA98 and TA100 without/with S9 mix; that is, the mutagenicity ratio (MR) was less than 2. The spontaneous revertants of the other bamboo/wood vinegars collected at various temperatures at the ranges of the diluting percent content were smaller than those of the control group by more than two times; that is, the MR was also less than 2 (results not shown in this paper). The bamboo/wood vinegars, therefore, have no mutagenicity toward *S. typhimurium* TA98 and TA100.

The MR of the bamboo/wood vinegars collected at all different temperatures toward *S. typhimurium* TA98 and TA100 without the S9 mix for a diluting percent content of 25.00, 20.00, 13.33 and 10.00% or with the S9 mix for a diluting percent content of 20.00, 13.33 and 10.00% is shown in Table 6. No matter what the vinegars were for all collected temperatures and *S. typhimurium* TA98 and TA100 without/with the S9 mix, the MR was less than 2. Neither mutagenicity nor toxicity was observed for bamboo/wood vinegars collected at various temperatures toward *S. typhimurium* TA 98 or TA 100 without/with the S9 mix. Hence, without the S9 mix a diluting percent content of 25.00 to 10.00% and with S9 mix at 20.00 to 10.00% a diluting percent content was selected for the antimutagenic assay.

Antimutagenicity of bamboo/wood vinegars

In the present study, NQNO and AFB₁ were used as direct mutagens requiring metabolic activation and indirect acting mutagen, respectively. Doses of mutagens, 1 µg for NQNO and 5 µg for AFB₁, were selected from a dose–response curve of a preliminary experiment (Yen *et al.*, 2001). The His⁺ revertants of strain are less than those of the control group, said that with antimutagenic activities. Meanwhile, the inhibitory effect of the specimen is expressed by inhibition (%), and the higher the inhibition, the more effective the antimutagenic activities (Maron and Ames, 1983). The inhibitory effects for one of the antimutagenicity results for the bamboo/wood vinegars collected at temperatures of 90–92/105–109°C with a diluting percent content of 25.00, 20.00, 13.33 and 10.00% for NQNO and at 20.00, 13.33 and 10.00% for AFB₁ are summarized in Table 7.

The His⁺ revertants of strain against the NQNO in Blank (without bamboo/wood vinegars) were 1128 for TA98, and 1445 for TA100, for AFB₁: they were 1824 for TA98, and 2406 for TA100. The spontaneous revertants without NQNO were 65 for TA98 and 132 for TA100, and without AFB₁ were 76 for TA98 and 140 for TA100. The His⁺ revertants of strain (inhibition) against the NQNO for bamboo vinegar at 90–92°C with different diluting percent contents were 807–985 (30.18–13.49%) for TA98 and 425–616 (77.65–63.15%) for TA100, for wood vinegar at 105–109°C they were 906–1084 (20.89–4.14%) for TA98 and 579–977 (65.92–35.65%) for TA100. For AFB₁, they were 1114–1319 (40.63–28.88%) for TA98 and 843–

1202 (68.97–53.14%) for TA100 from bamboo vinegar at 90–92°C with different diluting percents, from wood vinegar at 105–109°C they were 1498–1587 (18.65–13.58%) for TA98 and 619–1051 (78.89–59.79%) for TA100. The results also showed that the higher diluting percent content, the greater the inhibition; as well as, no matter what the vinegar was, the inhibition for TA100 was greater than that of TA98.

The inhibition of the bamboo/wood vinegars collected at all different temperatures against the NQNO for diluting percent contents of 25.00, 20.00, 13.33 and 10.00% or against the AFB₁ at 20.00, 13.33 and 10.00% is shown in Table 8. The inhibition of the bamboo vinegars to TA98 was 0.19–30.18% for NQNO and 0.92–40.63% for AFB₁; they were better than that of wood vinegar, –4.45–31.43% and 3.62–25.06%. For both vinegars with TA100 against NQNO (18.18–81.77%) and AFB₁ (12.03–78.89%), they were better than those for TA98. The antimutagenicity to NQNO was effective for bamboo/wood vinegars collected at various temperatures with a diluting percent content of 25.00% or less, and for AFB₁, it was also effective at a 20.00% or less diluting percent content. Furthermore, the bamboo/wood vinegars showed that the inhibitory effect on NQNO or AFB₁ toward TA100 was greater than that toward TA98. It is also indicated that the inhibition of the vinegars against AFB₁ toward TA98 and TA100 is better than that against NQNO.

In this study, the main percent of the phenol and ketone compounds in the bamboo/wood vinegars collected from various different temperatures was 50.23–65.98% (Table 1) and 10.13–25.26% (Table 2), resulting in no cytotoxicity (Table 4)/mutagenicity (Table 6) and with antimutagenic effect (Table 6) against strains. It is inferred that the phenol and ketone compounds may partially account for the biological action of bamboo/wood vinegars.

CONCLUSIONS

The compounds, safety evaluation and antimutagenic activity of bamboo/wood vinegars were investigated. The compounds of bamboo/wood vinegars include the acid, phenol, ketone and the other compounds. The main percent content of the phenol compounds for bamboo vinegars was phenol (5.93–16.60%) and 2-methoxyphenol (8.27–16.39%). For wood vinegars, the main compounds were 2-methoxyphenol (11.70–22.90%) and 2-methoxy-4-methylphenol (7.09–20.10%). The diluting percent content of vinegars was lower than 20.00% or less with the S9 mix and 33.33% or less without the S9 mix in cytotoxicity and mutagenicity toward *S. typhimurium* TA98 and TA100 because the rest of the bacterium at these percent contents was higher than 80% of the control, and the mutagenicity ratio was less than for the control group by more than two times. The diluting percent content of vinegars of 20.00% or less, expressed an amount-dependent inhibitory effect against both the mutagenicity of 4-nitroquinoline-N-oxide (NQNO) with 1 µg/plate and aflatoxin B₁ (AFB₁) with 5 µg/plate in *S. typhimurium* TA98 and TA100. It is suggested that

bamboo/wood vinegars with a diluting percent content to the least extent of 20.00% or less had no cytotoxicity and mutagenicity, and their antimutagenicity with NQNO and AFB₁ was effective.

ACKNOWLEDGEMENTS

The authors offer the sincere appreciation to the Forestry Bureau, Council of Agriculture Executive with No. 98-04.4-B-P-14 for financial support.

REFERENCES

- Ames, B. N., J. McCann and E. Yamasaki 1975 Methods for Detecting Carcinogens and Mutagens with the *Salmonella/Mammalian*-Microsome Mutagenicity Test. *Mutat. Res.*, **31**: 347-364
- Ames, B. N. and J. McCann 1976 Detection of Carcinogens as Mutagenes in the *Salmonella/Microsome* Test: Assay of 300 Chemicals. *Proc. Natl. Acad. Sci. U.S.A.*, **73**: 950-954
- Chen, C. H., K. T. Lu, L. W. Hung, C. T. Liu and C. H. Yen 2006 Manufacture of Wood Charcoals and Wood Vinegars from Floating Wood and Their Utilization on the Culture of Plants (I) Manufacture of Wood Charcoals and Wood Vinegars from Floating Softwood. *Quarterly Journal of Forest Research*, **28**(4): 87-98
- Chen, P. K., S. C. Wu, G. S. Hwang and H. C. Lin 2010 Investigation on Basic Properties and Antibacterial Activity of Bamboo/Wood Vinegars Collected from Different Temperature. *J. Agric For. (NCYU)*, **7**(2): 95-117
- Council of Agriculture 2004 *Certified Agricultural Standards of Forest Products*. COA, Executive Yuan, Taiwan ROC.
- Duh, P. D. and G. C. Yen 1997 Antioxidant activity of three herbal water extracts. *Food Chem.*, **60**: 639-645
- Hageta, Y. 2004 *Application of Wood/Bamboo Vinegar on the Cultivation of Grape*. Nobunkyou Publication, pp. 84-85.
- Ho, H. M., W. R. Kuo, Chia-Wen Peng, G. S. Hwang and H. C. Lin 2013 Investigation of Yield and Seedling Growth of Cabbage from Cultural Media with Moso Bamboo Charcoal. *J. Agric For. (NCYU)*, **10**(1): 43-60
- Horne, P. A. and T. W. Paul 1996 Influence of Temperature on the Products from the Flash Pyrolysis of Biomass. *Fuel*, **75**(9): 1051-1059
- Huang, Y. P., C. W. Peng, Y. H. Huang, T. R. Tsai, W. R. Kuo, and H. C. Lin 2011 Investigation of *Mikania micrantha* Charcoal Applied on Cultural Media. *Materials Science Forum*, **685**: 216-229
- Hwang, G. S., C. L. Ho, H. Y. Yu and Y. C. Su 2006 Bamboo Vinegar Collected during Charcoal Making with an Earthen Kiln and Its Basic Properties. *Taiwan J For Sci.*, **29**(1): 1-15
- Hwang, G. S., Y. J. Lin, C. L. Ho and H. Y. Yu 2008 Study on Charcoal Marking with Branches and Tops Wood of *Cryptomeria japonica* Using an Earthen Kiln Abstracts of The 2008 Annual meeting of The Chinese Forest Products Association, pp. 27-33
- Ikimoto, T. and O. Ikeshima 2000 *Usages of Bamboo Charcoal and Bamboo Vinegar*. Nobunkyou Publication, pp. 89-107
- Ishihara, S. 1996 Development of Wood-based Charcoal Materials, *Mokuzaigakkaishi*, **42**(8): 717-723
- Kobahasi, Y. 2004 *Reductive Effectives of Agricultural Chemical*. Nobunkyou Publication, pp. 146-149
- Kou, C. W. 2004 *Effects of Heating Rates on Fundamental Properties of Vinegar during Bamboo Carbonization and Its Application on Inhibit Plant Pathogenic Bacteria*, Master thesis of National Chung University, pp. 1-76
- Lin, H. C., T. Ohuchi, Y. Murase, T. C. Shiah, S. L. Shieh, P. C. Chiu, Y. M. Juan and S. C. Cheng 2006 Application of Bamboo Vinegar with Vacuum Process to Evaluate Fungi Resistance of Bamboo Materials. *Journal of the Faculty of Agriculture Kyushu University. Japan*, **51**(1): 5-11.
- Lin, H. C. and T. C. Shiah 2006 Evaluation of Fungi Resistance of Moso Bamboo Materials Using Bamboo Vinegar with Smoking Process. *Quart. Journal Forest Research of Taiwan. Taiwan ROC*, **28**(2): 51-66
- Lin, H. C., Y. Murase, T. C. Shiah, G. S. Hwang, P. K. Chen and W. L. Wu 2008 Application of Moso Bamboo Vinegar with Different Collection Temperatures to Evaluate Fungi Resistance of Moso Bamboo Materials. *Journal of the Faculty of Agriculture Kyushu University. Japan*, **53**(1): 107-113
- Lin, H. C., N. Fujimoto, G. S. Hwang, S. C. Wu, C. Y. Liu, P. J. Chao, W. L. Guo and Y. T. Sung 2009 Application of Wood Vinegars Prepared from Branches and Tree Tops of *Cryptomeria japonica* at Different Collection Temperatures to Evaluate their Fungi Resistance for Moso Bamboo Material. *Journal of the Faculty of Agriculture Kyushu University. Japan*, **54**(2): 457-462
- Lin, H. C., Y. H. Huang, T. R. Tsai, C. W. Peng, G. S. Hwang and W. R. Kuo 2011 Investigation of Promoting Growth of Tomato Plug Seedlings from Moso Bamboo Charcoal Cultural Media with Bamboo Vinegar. *J. Agric For. (NCYU)*, **8**(2): 58-72
- Lu, K. T., C. W. Kuo and C. T. Liu 2007 Inhibition Efficiency of a Mixed Solution of Bamboo Vinegar and Chitosan against *Ralstonia solanacearum*. *Taiwan J For Sci.*, **22**(3): 329 - 338
- Maron, D. M. and B. N. Ames 1983 Revised methods for the *Salmonella* mutagenicity Test. *Mutat Res.*, **113**: 173-215
- McCann, J., E. Choi, E. Yamasaki and B. N. Ames 1975 Detection of Carcinogens as Mutagens in the *Salmonella/Microsome* Test: Assay of 300 Chemicals. *Proc Natl Acad Sci U S A.*, **72**(12): 5135-5139
- Namiki, M. 1990 Antioxidants/Antimutagens in Foods. *Crit. Rev. Food Sci. Nutr.* **29**: 273-300
- Nishimiya, K., T. Hata, Y. Imamura and S. Ishihara 1998 Analysis of Chemical Structure of Wood Charcoal by X-ray Photoelectron Spectroscopy. *J. Wood Sci.*, **44**: 56-61
- Nomura, R. 2004. Healthy Effects of Bamboo Vinegar. *Nobunkyou Publication*, pp. 141-143
- Sugimura, T., S. Sato, M. Nagao, T. Yahagi, T. Matsushima T. and Y. Seino 1976 Overlapping of Carcinogens and Mutagens. *Fundam Cancer Prev Proc Int Symp Princess Takamatsu Cancer Res Fund 6th 1975*: pp. 191-215
- Tsai, C. T., S. C. Wu, K. J. Lin, G. S. Hwang and H. C. Lin 2009 Evaluation of Antioxidation Activity and Resist Lipid Oxidation Effect with Moso Bamboo Vinegar Collection From Different Temperature. *J. Agric For. (NCYU)*, **6**(1): 66-83
- Uchimura, T., H. Tanikai, and K. Hosoukawa 2000 *Issues of Bamboo Charcoal and Bamboo Vinegar*. Soumorisya Publication, pp. 138-168
- Vasavada, M. N. and D. P. Cornforth 2006 Evaluation of Antioxidant Effects of Raisin Paste in Cooked Ground Beef, Pork, and Chicken. *J Food Sci.*, **71**(4): 242-246
- Waleh, N. S., S. J. Rapport and K. E. Mortelmans 1982 Development of a toxicity test to be coupled to the Ames *Salmonella* assay and the method of construction of the required strains. *Mutat Res.*, **97**: 247-256
- Yam, T. S., S. Shah and J. M. T. Hamilton-Miller 1997 Microbiology Activity of Whole and Fractionated Crude Extracts of Tea (*Camellia sinensis*), and of Tea Components. *FEMS Microbiology Letters*, **152**: 169-174
- Yatagai, M., G. Unrinin and T. Orira 1988 By-products of Wood Carbonization IV. Components of Wood Vinegar. *Mokuzaigakkaishi*, **34**: 184-188
- Yatagai, M., M. Nishimoto, K. Hori, T. Ohira and A. Shibata 2002 Termiticidal Activity of Wood vinegar, its Components and Their Homologues. *J. Wood Sci.*, **48**: 338-342
- Yen G. C., P. D. Duh, Y. L. Hung 2001 Contributions of Major Components to the Antimutagenic Effect of Hsian-Tsao (*Mesona procumbens* Hemsl.), *Journal of Agricultural and Food Chemistry*. **49**: 5000-5004
- Yoshie, L. 2004 *Effects of Wood/Bamboo Vinegar on Cow Skin*. Nobunkyou Publication, pp. 132-133