

Elliptic curves and Fibonacci numbers arising from Lindenmayer system with Symbolic Computation

Yoshida, Hiroshi
Faculty of Mathematics, Kyushu University

Miwa, Yoshihiro
Graduate School of Comprehensive Human Sciences, University of Tsukuba

Kaneko, Masanobu
Faculty of Mathematics, Kyushu University

<https://hdl.handle.net/2324/14895>

出版情報 : 2011-03. Springer
バージョン :
権利関係 :

MI Preprint Series

Kyushu University
The Global COE Program
Math-for-Industry Education & Research Hub

Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations

H. Yoshida,
Y. Miwa & K. Masanobu

MI 2009-21

(Received May 22, 2009)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

Elliptic curves and Fibonacci numbers arising from Lindenmayer system with Symbolic Computation

Hiroshi Youshida ^{a,*}, Yoshihiro Miwa ^b, Masanobu Kaneko ^a

^a*Faculty of Mathematics, Kyushu University, Hakozaki 6-10-1, Higashi-ku, FUKUOKA 812-8581 Japan*

^b*Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tennoudai 1-1-1, Tsukuba-shi, IBARAKI 305-8575 Japan*

Abstract

Starting from an egg, the multicell becomes a set of cells comprising a variety of types to serve functions. This phenomenon brings us a bio-inspired Lindenmayer system. To investigate conditions for a variety of cell types, we have constructed a stochastic model over Lindenmayer systems. This model also considers interactive behaviors among cells, yielding complicated nonlinear polynomials. Using symbolic computation, we have derived the explicit relations between cell-type diversity and cell-type ratio constraint. These relations exhibit elliptic curve- and Fibonacci number-related patterns. This is the first example of elliptic curves to appear in the Lindenmayer context. A survey of the rational points and the quadratic irrational numbers on the derived curves has revealed Fibonacci-related periodic and quasiperiodic patterns. Further, we have found that in some region, there are only two elliptic curve-related periodic patterns.

PACS: 02.10; 82.40.C

Key words: Elliptic curves, Fibonacci numbers, Lindenmayer system, Symbolic computation

* Corresponding author. Faculty of Mathematics, Kyushu University, Hakozaki 6-10-1, Higashi-ku, FUKUOKA 812-8581 Japan. Tel/fax: +81-92-642-7396.
Email address: phiroschi@math.kyushu-u.ac.jp (Hiroshi Youshida).

1 Introduction

The multicell is thought to form according to specific rules, wherein the first cell type transforms into one different cell type or into two or more different cell types. Such cell-type transformation rules can be described by a tree diagram, termed a *cell lineage* diagram, whereas the developmental process of multicells is termed *cell differentiation* [8]. In 1968, such a cell differentiation process was modeled as rewriting rules by Lindenmayer [13, 14].

Here, we analyze a bio-inspired Lindenmayer system (L-system), where the rewriting rules are based on cell lineage diagrams. L-system models for developmental systems with cell lineages and for biological tissues such as blood vessels of the eye have been studied in past decades [10, 21, 22]. In these L-systems, the *interactions* among cells have been ignored because of the complexity or difficulty of the analysis. By contrast, in this paper, we present a symbolic computation, the quantifier elimination method, for the derivation of algebraic equations in an L-system with interactions (“IL-system” hereafter). Further, the stochastic aspects can be introduced into an L-system, termed a stochastic L-system [4, 5]. We studied the IL-system giving nonlinear compositions of extra cellular matrices in green algae [26]. Apart from L-system models, we studied compatible conditions for self-replication and high cell-type diversity by modeling the interaction of cells having chaotic dynamics [24, 25]. Chaotic dynamics yielded stochastic behavior. These previous studies let us know that interactive and stochastic behaviors play an important role in multicells.

In this paper, we extend our previous studies to a stochastic IL-system model (“sIL-system” hereafter) using algebraic equations between cell-type diversity and cell-type ratio constraint on the cell chain. To devise the basic cell differentiation rules, we consider three cell lineage diagrams: (i) the branching cell lineage $A \Rightarrow \{B, C\}$ without the cell-type order conservation rule, (ii) the linear diagram $A \Rightarrow B \Rightarrow C$ with the conservation rule, and (iii) the linear diagram without the rule. Using symbolic computation exhaustively, we have obtained five equations including *elliptic curves* which appears in the Lindenmayer context, for the first time.

Throughout this paper, we present the derived algebraic numbers or equations instead of approximate values. The reason is illustrated by a simple example. Consider the algebraic number $(1 + \sqrt{5})/2$ and its approximate value 1.618. The former is one root of $x^2 - x - 1 = 0$ in x , which can also be expressed as a continued

fraction: $[1; 1, 1, 1, \dots]$. The appearance of only 1 reveals the relationship between $(1 + \sqrt{5})/2$ and self-similarity (quasiperiodicity) of the corresponding sequence [11, 16]. By contrast, from the latter value, 1.618, we cannot submit such a strong proposal. This is why we made the extra effort to obtain the exact algebraic numbers or equations, to allow us to analyze the periodicity and quasiperiodicity of produced patterns.

2 Model: Bio-inspired Lindenmayer System

We study two basic subgraphs extracted from biological cell lineage diagrams: *branching* and *linear* cell differentiation rules of three cell types. This model is described by an sIL system for three cell types, A , B , and C .

We first define the following two basic cell differentiation graphs, *branching* $A \Rightarrow \{B, C\}$ and *linear* $A \Rightarrow B \Rightarrow C$. The proliferation of a type is denoted by $A \rightarrow AA$, $B \rightarrow BB$ or $C \rightarrow CC$, while the transition between types is denoted by $A \rightarrow B$, $A \rightarrow C$ (in the branching diagram), or $A \rightarrow B$, $B \rightarrow C$ (in the linear diagram). We next define the proliferation and transition rates (probabilities) of the branching diagram as follows:

$$A \rightarrow \begin{cases} AA & p_{1,1}, \\ B & p_{1,2}, \\ C & p_{1,3}, \\ A & 1 - p_{1,1} - p_{1,2} - p_{1,3}, \end{cases} \quad B \rightarrow \begin{cases} BB & p_{2,2}, \\ B & 1 - p_{2,2}, \end{cases} \quad C \rightarrow \begin{cases} CC & p_{3,3}, \\ C & 1 - p_{3,3} \end{cases} \quad (1)$$

with $0 < p_{i,j} < 1$ ($1 \leq i \leq j \leq 3$) and $p_{1,1} + p_{1,2} + p_{1,3} < 1$. The rule $A \rightarrow A$, $B \rightarrow B$, or $C \rightarrow C$ shows that the type does not change.

Likewise, the rates of the linear diagram are defined as:

$$A \rightarrow \begin{cases} AA & p_{1,1}, \\ B & p_{1,2}, \\ A & 1 - p_{1,1} - p_{1,2}, \end{cases} \quad B \rightarrow \begin{cases} BB & p_{2,2}, \\ C & p_{2,3}, \\ B & 1 - p_{2,2} - p_{2,3}, \end{cases} \quad C \rightarrow \begin{cases} CC & p_{3,3}, \\ C & 1 - p_{3,3} \end{cases} \quad (2)$$

with $0 < p_{i,j} < 1$ ($1 \leq i \leq j \leq 3$) and $p_{i,i} + p_{i,i+1} < 1$ ($1 \leq i \leq 2$). The one-dimensional cell chain becomes longer as these rules are applied. In addition

to the rewriting rules mentioned above, we adopt another rewriting rule, termed a *cell-type order conservation rule*:

$$AC \rightarrow ABC, CA \rightarrow CBA, \quad (3)$$

which guarantees the contiguity of cell types. When we adopt the above conservation rule, this rule is applied after each application of the branching or linear rewriting rules. This cell-type order conservation rule originates from the “intercalary regeneration” phenomenon in insect legs, as illustrated in Fig. 1. When portions of the legs with noncontiguous positional values are grafted together, new tissue is intercalated to fill the gap so that the noncontiguous positional values disappear [1, 6, 7]. Such a regeneration phenomenon occurs in our previous model of chaotic elements [24, 25] as well as in the actual wound repair in the early stages of the cell lineage diagram [2]. This is why we analyze the effect of the conservation rule. In this paper, we analyze the three diagrams: (i) a branching cell lineage diagram without the cell-type order conservation rule, (ii) a linear cell lineage diagram without the rule, and (iii) a linear diagram with the rule. We have not analyzed a branching diagram with the rule because we have not found a corresponding phenomenon in the actual cell-lineage diagrams.

Using these three models, we calculate the following composition in the cell chain:

$$AA, AB, BA, (AC, CA), BB, BC, CB, CC, \quad (4)$$

where (AC, CA) does not exist when the cell-type order conservation rule is adopted and applied during the elongation of a cell chain. In this paper, we adopt the above “two” contiguous composition instead of the “one” composition A, B, C because we do not aim to regard, for instance, $AAAAABBBBBCCCCC = A^5B^5C^5$ as a sequence with high cell-type diversity. In other words, we aim to evaluate the pattern-dependent diversity. Of course, we could adopt a more than two contiguous composition, but such composition gives an intractable calculation. For this reason, we have adopted the two contiguous composition (4).

Further, we assume the following constraint:

$$gN(AA) = N(BB) = N(CC) \wedge gN(AB) = N(BC), \quad (5)$$

where \wedge denotes “And,” g designates the parameter in this constraint, and $N(XY)$ denotes the number of the two contiguous cell types XY in the cell chain. $N(AB) = N(BA)$ and $N(BC) = N(CB)$ hold always because of the mirror symmetry of the

rewriting rules.

Under the constraint (5), we have calculated the maximum value of $N(AB)/N(AA)(= N(BC)/N(BB))$, which represents the ratio of junctions to nonjunctions between different cell types. Here, we have adopted this maximum as the measure of *cell-type diversity* in the sense that the appearance frequencies of two contiguous cell types become uniform as the ratio $N(AB)/N(AA)$ approaches 1 under the constraint (5). Notice that the large value of $N(AB)/N(AA)$ prevents us from regarding $A^n B^n C^n$ as a high-diversity pattern.

3 Method

3.1 Estimation of cell-type composition

We calculate the *growth matrix* of the two contiguous cell types based on [17, 23]. This matrix enables us to estimate its composition after n applications of the branching or linear rewriting rules (and the cell-type order conservation rule). In this paper, we have calculated three growth matrices for the branching cell lineage without the cell-type order conservation rule, and the linear diagram with and without the conservation rule, as shown in Table 1.

Let M be the growth matrix of the two contiguous cell types under some condition. Here, we start with AA ; that is, the axiom is AA , therefore, the composition at step n can be calculated by $(1, 0, 0, \dots)M^n$. We study the composition as n approaches infinity—in other words, for a sufficiently long cell chain. First, we derive the conditions for coexistence of the two contiguous cell types from the eigenvalues and the eigenvectors of the growth matrices. Second, under the constraint (5) and the coexistence condition, we estimate the composition using the growth matrix. Last, we derive the equation between the maximum value of $N(AB)/N(AA)(= N(BC)/N(BB))$ and the constraint parameter g in (5) using the quantifier elimination method.

3.2 Quantifier Elimination Method

In this paper, we investigate the maximum value of $N(AB)/N(AA)(= N(BC)/N(BB))$. For this purpose, we have used the quantifier elimination (QE) method. QE is one

of the main subjects in computer algebra [3, 15]. QE deals with first-order formulas comprising polynomial equations, inequalities, quantifiers (\exists, \forall), and boolean operators such as \wedge (And), \vee (Or), and \neg (Negation). QE computes an equivalent quantifier-free formula for a given first-order formula over the real field. For instance, for the input $\forall x(x^2 + bx + c > 0)$, QE outputs the equivalent quantifier-free formula $b^2 - 4c < 0$. QE obtains the relations between the unquantified variables that make the input formula true. We obtain the maximum value of an objective polynomial under some constraints by adding one extra variable, which is assigned to the objective polynomial [23]. For instance, to calculate the maximum value of $x/(p + y)$ under the constraints $x^2 + py^2 \leq 1 \wedge y \leq x^2$, we provide QE with the following input, adding an extra variable h :

$$\exists y \exists x (x^2 + py^2 \leq 1 \wedge y \leq x^2 \wedge h(p + y) == x).$$

For this formula, the QE output is $p < 0 \vee h^2 p^3 - p - h^2 < 0$, which indicates that the maximum value of $x/(p + y)$ is $\sqrt{p/(p^3 - 1)}$ when $p > 1$ and that this value does not exist ($+\infty$) when $p \leq 1$. Thus, even when parameters exist, we can obtain the maximum value; in other words, QE gives the equation between the maximum value and parameters. We can use QEPCAD-B or Mathematica Ver. 6/7 as QE-implemented software.

4 Results

4.1 Equations derived by QE

We first derive the coexistence condition of two contiguous cell types for the branching cell-lineage diagram. Analysis of the eigenvalues and their corresponding eigenvectors of the growth matrix reveals the coexistence condition: $2p_{1,1} - p_{1,2} - p_{1,3} > 0 \wedge 2p_{1,1} - p_{1,2} - p_{1,3} > 2p_{2,2} \wedge 2p_{1,1} - p_{1,2} - p_{1,3} > 2p_{3,3}$ ¹. Under this condition and the constraint (5), the following relations hold:

$$\frac{N(AB)}{N(AA)} = \frac{p_{1,2}(g(1 - p_{1,2} - p_{1,3}) - p_{1,3})}{2p_{1,3} + g(p_{1,2} + p_{1,3})},$$

$$p_{1,1} = \frac{-(1 - p_{1,2} - p_{1,3})(2p_{1,3} + g(p_{1,2} + p_{1,3}))}{2(p_{1,3} - g(1 - p_{1,2} - p_{1,3}))},$$

¹ See also [23, Section 5.2] for details.

$$\begin{aligned}
p_{2,2} &= (g(-2 + p_{1,2} + p_{1,3})(p_{1,3}(p_{1,2}^2 + 2p_{1,3}) + g(p_{1,2}^3 + p_{1,2}^2(-1 + p_{1,3}) + p_{1,2}(6 \\
&\quad p_{1,3} + p_{1,3}^2)))/(2(p_{1,3} + g(-1 + p_{1,2} + p_{1,3}))(g^2(p_{1,2}^2 + p_{1,2}(-2 + \\
&\quad p_{1,3}) - p_{1,3}) + p_{1,2}p_{1,3} + g(-1 + p_{1,2})(p_{1,2} + 2p_{1,3}))), \\
p_{3,3} &= (gp_{1,3}(-2 + p_{1,2} + p_{1,3})(p_{1,3}(2 + p_{1,3}) + g(p_{1,2} + p_{1,2}p_{1,3} + p_{1,3}^2)))/ \\
&\quad (2(p_{1,3} + g(-1 + p_{1,2} + p_{1,3}))(g^2(p_{1,2}^2 + p_{1,2}(-2 + p_{1,3}) - p_{1,3}) + p_{1,3}^2 \\
&\quad + gp_{1,3}(-3 + 2p_{1,2} + p_{1,3}))). \tag{7}
\end{aligned}$$

To obtain the maximum value of $N(AB)/N(AA)$ by QE, the following input is used:

$$\begin{aligned}
\exists p_{1,3} \exists p_{1,2} \left(\psi(p_{1,2}, p_{1,3}, g) \wedge h == N(AB)/N(AA) \wedge p_{1,2} > 0 \wedge \right. \\
\left. p_{1,3} > 0 \wedge p_{1,1} > 0 \wedge p_{2,2} > 0 \wedge p_{3,3} > 0 \wedge p_{1,1} + p_{1,2} + p_{1,3} < 1 \right),
\end{aligned}$$

where $\psi(p_{1,2}, p_{1,3}, g)$ is a formula derived by combining conjunctively all equations and inequalities appearing in the coexistence condition and the constraint (5). However, one cannot obtain the maximum if one provides the QE command of QEPCAD-B 1.48 or Mathematica 6.0.1 with the above formula as it is. This is because the computation requires too much memory and time. Therefore, we have transformed the above formula into the equivalent formulas in reduced form as follows.

Via factorization, we have derived the formulas:

$$\begin{aligned}
\exists p_{1,3} \exists p_{1,2} \left(0 < p_{1,2} \wedge 0 < p_{1,3} \wedge \right. \\
\left. -h(2p_{1,3} + g(p_{1,2} + p_{1,3})) == p_{1,2}(p_{1,3} + g(-1 + p_{1,2} + p_{1,3})) \wedge \right. \\
\left. e_1 < 0 \wedge e_2 < 0 \wedge e_3 < 0 \wedge e_4 < 0 \wedge e_5 < 0 \wedge e_6 < 0 \right)
\end{aligned}$$

with:

$$\begin{aligned}
e_1 &:= p_{1,3} + g(-1 + p_{1,2} + p_{1,3}), \\
e_2 &:= (g^2(p_{1,2}^2 + p_{1,2}(-2 + p_{1,3}) - p_{1,3}) + p_{1,2}p_{1,3} + g(-1 + p_{1,2})(p_{1,2} + 2p_{1,3})) \\
&\quad (p_{1,3}(p_{1,2}^2 + 2p_{1,3}) + g(p_{1,2}^3 + p_{1,2}^2(-1 + p_{1,3}) + p_{1,2}p_{1,3} + p_{1,3}^2)), \\
e_3 &:= (g^2(p_{1,2}^2 + p_{1,2}(-2 + p_{1,3}) - p_{1,3}) + p_{1,3}^2 + gp_{1,3}(-3 + 2p_{1,2} + p_{1,3})), \\
e_4 &:= (4p_{1,3} + g(-2 + 3p_{1,2} + 3p_{1,3})), \\
e_5 &:= g^2(p_{1,2}^2 + p_{1,2}(-2 + p_{1,3}) - p_{1,3}) + p_{1,2}p_{1,3} + g(-1 + p_{1,2})(p_{1,2} + 2p_{1,3}), \\
e_6 &:= g^2(p_{1,2}^2 + p_{1,2}(-2 + p_{1,3}) - p_{1,3}) + p_{1,3}^2 + gp_{1,3}(-3 + 2p_{1,2} + p_{1,3}),
\end{aligned}$$

under the physiological condition: $0 < h < 1, g > 0$. Mathematica 6.0.1 outputs: $0 < h < (\sqrt{8g + 5g^2} - g)/(2(2 + g))$. This shows that the equation between g and

the maximum value f of $N(AB)/N(AA)(= h)$ is $(g + 2)f^2 + gf - g = 0$.

Likewise, for the *linear* cell-lineage diagram without and with the cell-type order conservation rule, we have also used the QE method to derive the algebraic equations between f (the maximum of $N(AB)/N(AA)$) and g . The procedure for deriving these equations follows.

The eigenvalues of the growth matrix in the middle row of Table 1 are:

$$1 - p_{1,2}, 1 + 2p_{1,1} - p_{1,2}, (1 - p_{1,2})^2, 1 - p_{2,3}, 1 + 2p_{2,2} - p_{2,3}, \\ (1 - p_{1,2})(1 - p_{2,3}), (1 - p_{2,3})^2, 1 + 2p_{3,3}.$$

Therefore, the coexistence condition for the linear cell-lineage diagram without the rule is:

$$2p_{1,1} > p_{1,2} \wedge 2p_{1,1} - p_{1,2} > 2p_{2,2} - p_{2,3} \wedge 2p_{1,1} - p_{1,2} > 2p_{3,3}.$$

Under the constraint and the coexistence condition, we obtain the following relations:

$$N(AB)/N(AA) = \frac{(1 - p_{1,2})p_{1,2}}{2p_{1,1} + (1 - p_{1,2})p_{2,3}}, \\ p_{2,2} = (p_{1,1}(2p_{1,1} - p_{1,2} + p_{2,3})(p_{1,2}^2(-2p_{1,1} - (-1 + p_{1,2})(-2 + p_{2,3})))g \\ (4p_{1,1}^2 + -2p_{1,1}((-3 + p_{2,3})p_{2,3} + p_{1,2}(1 + p_{2,3})) + p_{2,3}(p_{1,2}^2 - (- \\ 2 + p_{2,3})p_{2,3} + p_{1,2}(-1 - 2p_{2,3} + p_{2,3}^2)))))/((-1 + p_{1,2})p_{1,2}(-4 \\ p_{1,1}^2 + (-1 + p_{1,2})p_{1,2}p_{2,3} - 2p_{1,1}(p_{1,2}(-1 + p_{2,3}) + p_{2,3}))2gp_{1,1}(\\ 4p_{1,1}^2 - 2p_{1,1}((-3 + p_{2,3})p_{2,3} + p_{1,2}(1 + p_{2,3})) + p_{2,3}(p_{1,2}^2 - (-2 + \\ p_{2,3})p_{2,3} + p_{1,2}(-1 - 2p_{2,3} + p_{2,3}^2))))), \\ p_{3,3} = ((1 + 2p_{1,1} - p_{1,2})(-1 + p_{1,2})p_{1,2}^2p_{2,3}^2 + gp_{1,1}(8p_{1,1}^3 - 4p_{1,1}^2((-2 + \\ p_{2,3})p_{2,3} + p_{1,2}(2 + p_{2,3})) + 2p_{1,1}(-p_{2,3}^2 + p_{1,2}^2(1 + 2p_{2,3}) + p_{1,2}p_{2,3} \\ (-3 + p_{2,3}^2)) + p_{2,3}(-p_{1,2}^3 + (-2 + p_{2,3})p_{2,3}^2 + p_{1,2}^2(1 + p_{2,3} - p_{2,3}^2) - \\ p_{1,2}p_{2,3}(1 - 3p_{2,3} + p_{2,3}^2)))))/((2p_{1,1} + p_{2,3} - p_{1,2}p_{2,3})(-(-1 + p_{1,2} \\)p_{1,2}p_{2,3} + 2gp_{1,1}(2p_{1,1} - p_{1,2} + 2p_{2,3} - p_{2,3}^2))).$$

To obtain the maximum value of $N(AB)/N(AA)$, we provided QE with the following input:

$$\exists p_{1,1} \exists p_{1,2} \exists p_{2,3} (\psi'(p_{1,1}, p_{1,2}, p_{2,3}, g) \wedge h == N(AB)/N(AA) \wedge 0 < p_{1,2} < 1 \wedge \\ 0 < p_{2,3} < 1 \wedge 0 < p_{1,1} \wedge 0 < p_{2,2} \wedge 0 < p_{3,3} < 1 \wedge p_{1,1} + p_{1,2} < 1 \wedge p_{2,2} + p_{2,3} < 1),$$

where $\psi'(p_{1,1}, p_{1,2}, p_{2,3}, g)$ is a formula derived by combining conjunctively all equations and inequalities appearing in the coexistence condition and the constraint. Again, the above formula was too complicated to analyze by the QE program (QEPCAD-B or Mathematica).

Therefore, we transform the formula in the following manner.

- From $h = N(AB)/N(AA)$,

$$p_{1,1} = \frac{(1 - p_{1,2})(p_{1,2} - hp_{2,3})}{2h} \quad (8)$$

holds. Using (8), we eliminate one variable, $p_{1,1}$.

- Next, via factorization, the inequalities with $p_{1,1}$ eliminated, can be transformed into the following formulas:

$$e'_1 < 0 \wedge e'_2 < 0 \wedge e'_3 > 0 \wedge e'_4 < 0 \wedge e'_5 < 0 \wedge e'_6 < 0 \wedge e'_7 < 0 \quad (9)$$

with:

$$\begin{aligned} e'_1 &:= -p_{1,2} + hp_{2,3}, \\ e'_2 &:= (1 - h - p_{1,2} + hp_{2,3})(-2h^2p_{1,2} + gp_{1,2} - hgp_{1,2} - hp_{1,2}^2 - gp_{1,2}^2 + hgp_{2,3} + 2h^2 \\ &\quad p_{1,2}p_{2,3} + hgp_{1,2}p_{2,3} - hgp_{2,3}^2)(-hp_{1,2}^2 + h^2p_{1,2}^2 - gp_{1,2}^2 + hgp_{1,2}^2 + hp_{1,2}^3 + gp_{1,2}^3 \\ &\quad + h^2p_{1,2}p_{2,3} - 2h^3p_{1,2}p_{2,3} - h^2gp_{1,2}p_{2,3} - 3h^2p_{1,2}^2p_{2,3} - 2hgp_{1,2}^2p_{2,3} + h^2gp_{2,3}^2 \\ &\quad + 2h^3p_{1,2}p_{2,3}^2 + hgp_{1,2}p_{2,3}^2 + h^2gp_{1,2}p_{2,3}^2 - h^2gp_{2,3}^3), \\ e'_3 &:= gp_{1,2}^3 - 2hgp_{1,2}^3 + h^2gp_{1,2}^3 - 2gp_{1,2}^4 + 2hgp_{1,2}^4 + gp_{1,2}^5 - 2hgp_{1,2}^2p_{2,3} + 3h^2gp_{1,2}^2 \\ &\quad p_{2,3} - h^3gp_{1,2}^2p_{2,3} + 5hgp_{1,2}^3p_{2,3} - 4h^2gp_{1,2}^3p_{2,3} - 3hgp_{1,2}^4p_{2,3} - 2h^4p_{1,2}p_{2,3}^2 \\ &\quad - h^2gp_{1,2}p_{2,3}^2 - h^3gp_{1,2}p_{2,3}^2 - 2h^3p_{1,2}^2p_{2,3}^2 - hgp_{1,2}^2p_{2,3}^2 - 3h^2gp_{1,2}^2p_{2,3}^2 + 2h^3 \\ &\quad gp_{1,2}^2p_{2,3}^2 + hgp_{1,2}^3p_{2,3}^2 + 3h^2gp_{1,2}^3p_{2,3}^2 + 2h^3gp_{2,3}^3 + 2h^4p_{1,2}p_{2,3}^3 + 3h^2gp_{1,2}p_{2,3}^3 \\ &\quad - 2h^2gp_{1,2}^2p_{2,3}^3 - h^3gp_{1,2}^2p_{2,3}^3 - 2h^3gp_{2,3}^4 + h^3gp_{1,2}p_{2,3}^4, \\ e'_4 &:= -2h^2p_{1,2}^2 + 2h^3p_{1,2}^2 - 2hgp_{1,2}^2 + 2h^2gp_{1,2}^2 + 2h^3p_{1,2}^3 + gp_{1,2}^3 + h^2gp_{1,2}^3 - hp_{1,2}^4 \\ &\quad + 3h^2p_{1,2}^4 - 2gp_{1,2}^4 + 2hgp_{1,2}^4 + hp_{1,2}^5 + gp_{1,2}^5 + 2h^3p_{1,2}p_{2,3} - 4h^4p_{1,2}p_{2,3} - 2h^3 \\ &\quad gp_{1,2}p_{2,3} + 2h^2p_{1,2}^2p_{2,3} - 6h^3p_{1,2}^2p_{2,3} - 2h^4p_{1,2}^2p_{2,3} + 2hgp_{1,2}^2p_{2,3} - 5h^2gp_{1,2}^2 \\ &\quad p_{2,3} - h^3gp_{1,2}^2p_{2,3} + h^2p_{1,2}^3p_{2,3} - 7h^3p_{1,2}^3p_{2,3} + hgp_{1,2}^3p_{2,3} - 4h^2gp_{1,2}^3p_{2,3} - 4 \\ &\quad h^2p_{1,2}^4p_{2,3} - 3hgp_{1,2}^4p_{2,3} + 2h^3gp_{2,3}^2 - 2h^3p_{1,2}p_{2,3}^2 + 8h^4p_{1,2}p_{2,3}^2 + h^2gp_{1,2}p_{2,3}^2 \\ &\quad + 5h^3gp_{1,2}p_{2,3}^2 + 4h^3p_{1,2}^2p_{2,3}^2 + 4h^4p_{1,2}^2p_{2,3}^2 - hgp_{1,2}^2p_{2,3}^2 + 5h^2gp_{1,2}^2p_{2,3}^2 + 2h^3 \\ &\quad gp_{1,2}^2p_{2,3}^2 + 5h^3p_{1,2}^3p_{2,3}^2 + hgp_{1,2}^3p_{2,3}^2 + 3h^2gp_{1,2}^3p_{2,3}^2 - 4h^3gp_{2,3}^3 - 4h^4p_{1,2}p_{2,3}^3 \\ &\quad - h^2gp_{1,2}p_{2,3}^3 - 4h^3gp_{1,2}p_{2,3}^3 - 2h^4p_{1,2}^2p_{2,3}^3 - 2h^2gp_{1,2}^2p_{2,3}^3 - h^3gp_{1,2}^2p_{2,3}^3 + 2 \\ &\quad h^3gp_{2,3}^4 + h^3gp_{1,2}p_{2,3}^4, \end{aligned}$$

$$\begin{aligned}
e'_5 &:= -1 + h + p_{1,2} - hp_{2,3}, \\
e'_6 &:= -hp_{1,2}^2 + h^2p_{1,2}^2 - gp_{1,2}^2 + hgp_{1,2}^2 + hp_{1,2}^3 + gp_{1,2}^3 + h^2p_{1,2}p_{2,3} - 2h^3p_{1,2}p_{2,3} - h^2 \\
&\quad gp_{1,2}p_{2,3} - 3h^2p_{1,2}^2p_{2,3} - 2hgp_{1,2}^2p_{2,3} + h^2gp_{2,3}^2 + 2h^3p_{1,2}p_{2,3}^2 + hgp_{1,2}p_{2,3}^2 + \\
&\quad h^2gp_{1,2}p_{2,3}^2 - h^2gp_{2,3}^3, \\
e'_7 &:= -gp_{1,2}^2 + hgp_{1,2}^2 + gp_{1,2}^3 - h^2p_{1,2}p_{2,3} - h^2gp_{1,2}p_{2,3} - 2hgp_{1,2}^2p_{2,3} + h^2gp_{2,3}^2 + \\
&\quad hgp_{1,2}p_{2,3}^2 + h^2gp_{1,2}p_{2,3}^2 - h^2gp_{2,3}^3.
\end{aligned}$$

Notice that in the above formulas, we show only those inequalities that maximize $N(AB)/N(AA)$; it is relatively easy to find these via polynomial factorization and confirmation by QE input for fixed values of g .

- Furthermore, the equality $gN(AB) = N(BC)$ in the constraint condition yields:

$$\begin{aligned}
0 = & -4gp_{1,1}^2p_{1,2} + 2gp_{1,1}p_{1,2}^2 + 4gp_{1,1}^2p_{1,2}^2 - 2gp_{1,1}p_{1,2}^3 + 4gp_{1,1}^2p_{2,3} - 2gp_{1,1}p_{1,2} \\
& p_{2,3} + p_{1,2}^2p_{2,3} + 2p_{1,1}p_{1,2}^2p_{2,3} + 2gp_{1,1}p_{1,2}^2p_{2,3} - 2p_{1,2}^3p_{2,3} - 2p_{1,1}p_{1,2}^3p_{2,3} + p_{1,2}^4 \\
& p_{2,3} + 2gp_{1,1}p_{2,3}^2 - 4gp_{1,1}^2p_{2,3}^2 - 2gp_{1,1}p_{1,2}p_{2,3}^2 - 2gp_{1,1}p_{2,3}^3 + 2gp_{1,1}p_{1,2}p_{2,3}^3.
\end{aligned}$$

Substituting (8) into the above equality yields:

$$g = \frac{hp_{1,2}p_{2,3}(-h - p_{1,2} + hp_{2,3})}{(-p_{1,2} + hp_{2,3})(p_{1,2} - hp_{1,2} - p_{1,2}^2 - p_{2,3} + hp_{1,2}p_{2,3} + p_{2,3}^2)}. \quad (10)$$

We therefore eliminate one more variable, g .

- By eliminating g from the equalities and inequalities and factorizing, we obtain the following reduced formulas suitable for QEPCAD-B:

$$\exists p_{2,3} \exists p_{1,2} (0 < h < 1 \wedge g > 0 \wedge 0 < p_{1,2} < 1 \wedge 0 < p_{2,3} < 1 \wedge e_{p0} == 0 \wedge e_{p1} < 0 \wedge e_{p2} < 0)$$

with:

$$\begin{aligned}
e_{p0} &:= hp_{1,2}p_{2,3}(-h - p_{1,2} + hp_{2,3}) - g(-p_{1,2} + hp_{2,3})(p_{1,2} - hp_{1,2} - p_{1,2}^2 \\
&\quad - p_{2,3} + hp_{1,2}p_{2,3} + p_{2,3}^2) \\
e_{p1} &:= 2hp_{1,2}^2 - 2h^2p_{1,2}^2 + p_{1,2}^3 - 3hp_{1,2}^3 - p_{1,2}^4 - 3hp_{1,2}p_{2,3} - h^2p_{1,2}p_{2,3} + \\
&\quad 2h^3p_{1,2}p_{2,3} - 2p_{1,2}^2p_{2,3} - hp_{1,2}^2p_{2,3} + 7h^2p_{1,2}^2p_{2,3} + p_{1,2}^3p_{2,3} + 4hp_{1,2}^3 \\
&\quad p_{2,3} + h^2p_{2,3}^2 + 5hp_{1,2}p_{2,3}^2 - 4h^3p_{1,2}p_{2,3}^2 + p_{1,2}^2p_{2,3}^2 - 2hp_{1,2}^2p_{2,3}^2 - 5h^2 \\
&\quad p_{1,2}^2p_{2,3}^2 - 2h^2p_{2,3}^3 - 2hp_{1,2}p_{2,3}^3 + h^2p_{1,2}p_{2,3}^3 + 2h^3p_{1,2}p_{2,3}^3 + h^2p_{2,3}^4, \\
e_{p2} &:= -p_{1,2} + hp_{1,2} + p_{1,2}^2 + hp_{2,3} + 2h^2p_{2,3} - hp_{1,2}p_{2,3} + hp_{2,3}^2.
\end{aligned}$$

- Last, because of the RAM limitation (up to 4 GB) of 32-bit software, we provided the above formulas for the QE program (we used QEPCAD-B Ver. 1.48 with measure-zero-error and +N670000000 option. It took from 4.5 to 12 hours

and 3.7 GB RAM with an Intel Xeon CPU 2.33 GHz processor), with g -space divided into $(0, 1/7]$, $(1/7, 1/4]$, $(1/4, 1)$, $[1, 2]$, $(2, \infty)$.

After all, with respect to the *linear* cell-lineage diagram without the rule, we obtained the following equivalent quantifier-free formulas as QE outputs:

$$\left\{ \begin{array}{ll} g \geq \text{root}_{-1} h^3 g^2 - 2h^2 g^2 - hg^2 + 2g^2 + 2h^4 g + h^3 g - 5h^2 g + 2h^4 & \text{For } (0, 1/7], \\ & (1/7, 1/4], (1/4, 1), \\ \left(\begin{array}{l} h^3 g^2 - 2h^2 g^2 - hg^2 + 2g^2 + 2h^4 g + h^3 g - 5h^2 g + 2h^4 \geq 0 \\ \wedge 6h^2 g + 3hg - 3g - 2h^2 - h \leq 0 \end{array} \right) & \text{For } [1, 2], \\ 6h^2 g + 3hg - 3g - 2h^2 - h \leq 0 & \text{For } (2, \infty), \end{array} \right. \quad (11)$$

where root_{-1} denotes the largest real root of the equation $h^3 g^2 - 2h^2 g^2 - hg^2 + 2g^2 + 2h^4 g + h^3 g - 5h^2 g + 2h^4 = 0$ in g . The equations for the linear diagram with the rule is the same as in [23]. The equations obtained between f and g are summarized in Table 2.

5 Discussion

So far, we have derived the algebraic equations between the cell-type diversity measure f and the parameter g under the constraint (5). We summarize in Table 2 and illustrate in Fig. 2 the equations between g and f in the three diagrams: the branching and linear cell lineage diagrams with and without the cell-type order conservation rule (3).

Let us look closely at the values of the natural number g that make f a rational number. The rational numbers of the maximum value of $N(AB)/N(AA)$ show that the pattern becomes periodic with respect to the two contiguous cell types as the cell chain becomes sufficiently long. We have searched for $g \in \mathbb{N}$ with corresponding rational f on the three curves (Table 2 and Fig. 2). Only on the curve of the branching diagram without the rule $(g+2)f^2 + gf - g = 0$ does there exist a pair that satisfies $\{(g, f) \mid g \in \mathbb{N}, f \in \mathbb{Q}\}$. The proof is given in Appendix A. On this curve, f is rational when g is in the following set: $\{2, 18, 128, 882, 6050, 41472, \dots\} = \{g_n \mid g_{n+2} = 7g_{n+1} - g_n + 4, g_0 = 0, g_1 = 2, n \geq 1\}$. The pair (g_n, f_n) can be described as $(2F_{2n}^2, F_{2n}/F_{2n+1})$, where F_n are *Fibonacci* numbers defined as $\{F_n \mid F_{n+2} = F_{n+1} + F_n, F_0 = 0, F_1 = 1, n \geq 1\} = \{1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, \dots\}$. The $g \in \mathbb{N}$ and their corresponding values of f are summarized as follows.

- (a) When $g = 1$, for the branching diagram without the rule, f is $(\sqrt{13} - 1)/6 \sim 0.434259$; for the linear without the rule, f is $-1/2 + \{(9 + \iota\sqrt{687})^{1/3} + (9 - \iota\sqrt{687})^{1/3}\}/12^{2/3} \sim 0.557454$; and for the linear with the rule, f is $(1 + \sqrt{17})/8 \sim 0.640388$.
- (b) When $g > 1$ and $g \in \mathbb{N}$, for the branching diagram without the rule, f is a rational number if g is in $\{2F_{2n}^2 \mid n \geq 1\}$, otherwise f is a quadratic irrational number. For the linear diagram without and with the rule, f is always a quadratic irrational number (see also Appendix A). A quadratic irrational number is a solution to the quadratic equation $ax^2 + bx + c = 0$, where a, b, c are integers, and $b^2 - 4ac$ is positive and not a perfect square. The cell chain associated with a quadratic irrational number is quasiperiodic [11, 16]. Further, based on [9], we can reconstruct the rewriting rules that produce the pattern associated with a given quadratic irrational or rational number f via its continued fraction. We have constructed the sequences in the case of $g = 1, f = (1 + \sqrt{17})/8 = [0; 1, 1, 1, 3, 1, 1, 3, 1, 1, 3, 1, 1, 3, \dots]$ in the linear diagram with the rule; and $g = 2, f = 1/2$ in the branching diagram, respectively, as shown in Fig. 3. In this figure, we laid the two-dimensional space with 40,000 colored spiral cells, where the red, green, and blue boxes denote $A-$, $B-$, and $C-$ cell types. Figures 3 (B) and (C) show the quasiperiodic and periodic patterns, respectively.

Further, let us consider the region of small $g < 1, 1/g \in \mathbb{N}$. The curve \ddagger in Table 2 is transformed into the minimal model $y^2 + xy = x^3 - 7x + 9$ by birational transformation:

$$\begin{cases} f = (x - 2)/(x + 2), \\ g = (2 - x)(3 + y)/(2x(2 + x)), \\ x = 2(1 + f)/(1 - f), \\ y = (-3f + 3f^2 - 4g - 4fg)/((1 - f)f). \end{cases}$$

Likewise, the curve \dagger is transformed into $y^2 + xy + y = x^3 - x^2 - 9x + 9$ by:

$$\begin{cases} f = (3x + y - 1)/(x + y - 3), \\ g = (3x + y - 1)^2(x^2 - 4x + 3 - 2y)/(2(x + y - 3)(x - y + 5)(2x + y - 2)), \\ x = (f^2(3f + 1) + 2(f + 1)(f - 2)g)/(f^2(f - 1)), \\ y = 2(f + 1)(2f^2 - (f - 2)(f - 3)g)/(f^2(f - 1)^2). \end{cases}$$

These curves are *elliptic curves* [18, Ch. 3]. These two curves have Mordell–Weil rank = 1 and no torsion points over \mathbb{Q} . Concerning points such that $1/g \in \mathbb{N}$ and $f \in \mathbb{Q}$ ($g > 0, 0 < f < 1$), we have proved that such a point does not exist in the branching diagram and that only $(g, f) = (1/7, 1/4)$ and $(1/4, 1/3)$ exist on the curve † (the linear diagram without the rule) and the curve ‡ (the linear diagram with the rule), respectively. The finiteness of these points follows from Siegel’s theorem on the finiteness of integral points [18, Ch. 9]. In the above curves † and ‡, we transformed the curves into quartic elliptic equations in which one can compute all integral points based on [19, 20].

We have summarized in Table 3 the relations between f and g , showing that there are Fibonacci-related periodic and quasiperiodic patterns in the $g > 1, g \in \mathbb{N}$ region and that there are only two elliptic curve-related periodic patterns in $g < 1, 1/g \in \mathbb{N}$.

6 Conclusion

In this paper, using symbolic computation, we have derived relations between the measure of cell-type diversity and the cell-type ratio constraint over the bio-inspired Lindenmayer-system model. We have obtained the five equations between the measure and the constraint. Analysis of points on these equations has revealed frequent quasiperiodicity; and Fibonacci- and elliptic curve-related periodicity.

Acknowledgments

We wish to express our gratitude to Professor Christopher W. Brown and Professor Masato Kuwata for helpful calculations and useful suggestions on QEPCAD-B and on the Elliptic Surface Calculator. Author H.Y. was supported by a Grant-in-Aid for Young Scientists (B) (grant 21740077) from the Ministry of Education, Culture, Sports, Science and Technology of Japan (MEXT). This study was also supported in part by the Program for Improvement of Research Environment for Young Researchers from Special Coordination Funds for Promoting Science and Technology (SCF) commissioned by the Japan Science and Technology Agency (JST).

Appendix A: Proof of the existence or non-existence of rational numbers f

- For the branching cell-lineage diagram without the cell-type order conservation rule, we obtain the equation between f and g :

$$(g + 2)f^2 + gf - g = 0. \quad (12)$$

The discriminant of the above equation is $D = 5g^2 + 8g$. We survey (g_n, f_n) such that g_n and f_n are natural and positive rational numbers, respectively. If and only if $D = m^2, m \in \mathbb{N}$, f is rational. $D = m^2$ yields $(5g + 4)^2 - 5m^2 = 4^2$. Let X be $5g + 4$. Then the equation $X^2 - 5m^2 = 4^2$ is a Pell's equation. It follows that $X + \sqrt{5}m = \pm 4 \left((3 + \sqrt{5})/2 \right)^n, n = 0, 1, 2, \dots$ yields a pair $\{(X, m) | X \in \mathbb{Z}, m \in \mathbb{Z}\}$. Therefore, from $X = 5g + 4, g + \sqrt{5}m' = 4 \left\{ \left((3 + \sqrt{5})/2 \right)^{2n} - 1 \right\} / 5, n = 1, 2, 3, \dots$ yields a natural number sequence $\{g_n | g_n \in \mathbb{N}\}$. Moreover, g_n can more explicitly be described as $2 \left\{ \left((3 + \sqrt{5})/2 \right)^{2n} + \left((3 - \sqrt{5})/2 \right)^{2n} - 2 \right\} / 5$. It follows from the golden ratio $\varphi = (1 + \sqrt{5})/2$ that g_n is $2(\varphi^{4n} + \varphi^{-4n} - 2) / 5$. Because *Fibonacci* number F_n can be described as $(\varphi^n - (-\varphi)^{-n}) / \sqrt{5}, g_n = 2F_{2n}^2 \square$

The recurrence formula of g_n can be derived as follows:

$$\begin{aligned} g_{n+2} &= 2F_{2(n+2)}^2 = 2(F_{2n+3} + F_{2(n+1)})^2 \\ &= 2(2F_{2(n+1)} + F_{2n+1})^2 \\ &= 2(3F_{2(n+1)} - F_{2n})^2 \\ &= 2(9F_{2(n+1)}^2 + F_{2n}^2 - 6F_{2(n+1)}F_{2n}) \\ &\quad \text{From the formula } 3F_{2(n+1)}F_{2n} = F_{2(n+1)}^2 + F_{2n}^2 - 1, \\ &= 2(7F_{2(n+1)}^2 - F_{2n}^2 + 2) \\ &= 7g_{n+1} - g_n + 4. \quad \square \end{aligned}$$

The corresponding f_n with respect to g_n on the curve (12) is calculated as:

$$f_n = \frac{F_{2n} \left(\sqrt{4 + 5F_{2n}^2} - F_{2n} \right)}{2(1 + F_{2n}^2)}.$$

The formulas on *Fibonacci* and *Lucas* numbers F_n and L_n [12, Ch. 5]:

$$F_{n+1}F_{n-1} - F_n^2 = (-1)^n, 5F_n^2 - L_n^2 = 4(-1)^{n+1}, F_n + L_n = 2F_{n+1}$$

reveal that $f_n = F_{2n}/F_{2n+1} \square$

- With respect to the cell-lineage diagram without the rule, we obtain:

$$2(3g - 1)f^2 + (3g - 1)f - 3g = 0. \quad (13)$$

The discriminant of the above equation is $D = 81g^2 - 30g + 1$. We survey (g_n, f_n) such that g_n and f_n are natural and positive rational numbers, respectively. If and only if $D = m^2, m \in \mathbb{N}$, f is rational. $D = m^2$ yields $(27g - 5)^2 - (3m)^2 = 4^2$, which can be factorized as $(27g + 3m - 5)(27g - 3m - 5) = 4^2$. From $g \geq 1$ and $m \geq 1$, $27g + 3m - 5 \geq 5^2$. Therefore, there exists no rational f_n for natural number g_n \square

- For the linear diagram with the rule, we obtain:

$$2gf^2 + (g - 1)f - g = 0. \quad (14)$$

We survey (g_n, f_n) such that g_n and f_n are natural numbers ($g > 0$) and positive rational numbers ($0 < f < 1$), respectively.

We prove the non-existence of such pairs in $0 < g, 0 < f < 1$. Assume that $f = n/m$, $\gcd(m, n) = 1$, $m \in \mathbb{N}, n \in \mathbb{N}$ because it is a positive rational number. Then, the equation (14) can be transformed into:

$$g(2n - m)(m + n) = mn. \quad (15)$$

From $0 < g$ and $0 < f < 1$, it follows that $m/2 < n < m$. Furthermore, from (15), $2n^2g \equiv 0 \pmod{m}$ and $m^2g \equiv 0 \pmod{n}$. It follows from $\gcd(m, n) = 1$ that $2g \equiv 0 \pmod{m}$ and $g \equiv 0 \pmod{n}$, revealing $2g \equiv 0 \pmod{mn}$. This fact, together with $g = mn/((2n - m)(m + n)), m/2 < n < m$, implies that the only possible value of g is $mn/2$. $g = mn/2$ yields an equation $(2n - m)(m + n) = 2$, which has a single solution $m = n = 1$, contradicting the assumption $f = n/m < 1$ \square

References

- [1] B. Alberts, A. Johnson, J. Lewis, M. Raff, K. Roberts, P. Walter, *Molecular Biology of the Cell*, 5th ed., Garland Science, New York, 2008.
- [2] P. J. Bryant, S. V. Bryant, V. French, Biological regeneration and pattern formation, *Sci. Am.* 237 (1977) 67–81.
- [3] B. F. Caviness, J. R. Johnson, *Quantifier Elimination and Cylindrical Algebraic Decomposition*, Springer-Verlag, Wien, 1998.
- [4] P. Eichhorst, F. Ruskey, On unary stochastic Lindenmayer systems, *Inf. Control* 48 (1) (1981) 1–10.
- [5] P. Eichhorst, W. J. Savitch, Growth functions of stochastic Lindenmayer systems, *Inf. Control* 45 (3) (1980) 217–228.
- [6] V. French, Leg regeneration in the cockroach, *Blattella germanica* I. regeneration from a congruent tibial graft/host junction, *Roux Arch. Dev. Biol.* 179 (1976) 57–76.
- [7] V. French, P. J. Bryant, S. V. Bryant, Pattern regulation in epimorphic fields, *Science* 193 (1976) 969–980.
- [8] S. F. Gilbert, *Developmental biology*, 8th ed., Sinauer Associates, Sunderland, Mass., 2006.
- [9] K. Iguchi, Theory of quasiperiodic lattices. I. Scaling transformation for a quasiperiodic lattice, *Phys. Rev. B, Condens. matter* 43 (7) (1991) 5915–5918.
- [10] H. Jürgensen, A. Lindenmayer, Inference algorithms for developmental systems with cell lineages, *Bull. Math. Biol.* 49 (1) (1987) 93–123.
- [11] M. Kaneko, T. Odagaki, Selfsimilarity in a class of quadratic-quasiperiodic chains, *J. Phys. Soc. Jpn.* 62 (4) (1993) 1147–1152.
- [12] T. Koshy, *Fibonacci and Lucas numbers with applications*, John Wiley, New York, 2001.
- [13] A. Lindenmayer, Mathematical models for cellular interactions in development. I. filaments with one-sided inputs, *J. Theor. Biol.* 18 (3) (1968) 280–299.
- [14] A. Lindenmayer, Mathematical models for cellular interactions in development. II. simple and branching filaments with two-sided inputs, *J. Theor. Biol.* 18 (3) (1968) 300–315.
- [15] B. Mishra, *Algorithmic Algebra*, Springer-Verlag, New York, 1993.
- [16] T. Odagaki, M. Kaneko, Self-similarity of binary quasiperiodic sequences, *J. Phys. A: Math. Gen.* 27 (1994) 1683–1690.
- [17] G. Rozenberg, A. Salomaa, *The mathematical theory of L system*, Academic

Press, New York, 1980.

- [18] J. H. Silverman, *The arithmetic of elliptic curves*, vol. 106 of Graduate texts in mathematics, Springer-Verlag, 1986.
- [19] R. J. Stroeker, N. Tzanakis, Solving elliptic diophantine equations by estimating linear forms in elliptic logarithms, *Acta Arith.* 67 (1994) 165–190.
- [20] N. Tzanakis, Solving elliptic diophantine equations by estimating linear forms in elliptic logarithms. The case of quartic equations, *Acta Arith.* 75 (1996) 165–190.
- [21] R. Ványi, G. Kókai, Z. Tóth, T. Pető, Grammatical retina description with enhanced methods, in: R. Poli, W. Banzhaf, W. B. Langdon, J. F. Miller, P. Nordin, T. C. Fogarty (eds.), *Genetic Programming, Proceedings of EuroGP*, vol. 1802 of LNCS, Springer-Verlag, 2000.
- [22] T. Yokomori, *Inductive Inference of OL Languages*, chap. 2, Lindenmayer systems: Impacts on theoretical computer science, computer graphics, and developmental biology, Springer-Verlag, 1992, pp. 115–132.
- [23] H. Yoshida, H. Anai, K. Horimoto, Derivation of rigorous conditions for high cell-type diversity by algebraic approach, *Biosystems* 90 (2007) 486–495.
- [24] H. Yoshida, C. Furusawa, K. Kaneko, Selection of initial conditions for recursive production of multicellular organisms, *J. Theor. Biol.* 233 (2005) 501–514.
- [25] H. Yoshida, K. Kaneko, Unified description of regeneration by coupled dynamical systems theory: intercalary/segmented regeneration in insect legs, *Dev. Dyn.*(in press).
- [26] H. Yoshida, T. Yokomori, A. Suyama, A simple classification of the volvocine algae by formal languages, *Bull. Math. Biol.* 67 (2005) 1339–1354.

Figure captions

Fig. 1. Intercalary regeneration in cockroach legs [1]. When mismatched portions of the growing legs are grafted together, new tissue is intercalated to fill the gap so that the noncontiguous positional values disappear. In this figure, we exemplify the rule: $I_8I_4 \rightarrow I_8I_7I_6I_5I_4$.

Fig. 2. The equations between g and f . The gray line denotes the equation in the branching cell lineage diagram without the cell-type order conservation rule. The black and broken lines denote the equations in the linear diagram with and without the rule, respectively. The derivative discontinuities are nearly 1.14254 (g_0) and 1.29661 (g_1). The intersection of the broken and gray lines is nearly 3.28773 (g_2), and that of the black and gray lines is nearly 3.83118 (g_3). The f -value of the broken and black lines approaches $1/2$ as $g \rightarrow \infty$, whereas that of the gray line approaches the reciprocal of the golden ratio $1/\varphi \sim 0.61803$ ($\varphi = (1 + \sqrt{5})/2$) as $g \rightarrow \infty$.

Fig. 3. Spiral tiling of cells. (A) Schematic illustration of spiral tiling of one-dimensional cell chain. (B) $g = 1$, $f = (1 + \sqrt{17})/8$ in the linear diagram with the rule. (C) $g = 2$, $f = 1/2$ in the branching diagram.

Table 1

The three growth matrices. We have calculated the matrices to estimate the composition of two contiguous cell types at step n . The top and middle matrices are of 9×9 form ($AA, AB, BA, AC, CA, BB, BC, CB, CC$) because of the nonexistence of the cell-type conservation rule, whereas the bottom matrix is of 7×7 form ($AA, AB, BA, BB, BC, CB, CC$) with the rule.

Branching diagram without the cell-type order conservation rule

$$\begin{pmatrix} m_{1,1} & m_{1,2} & m_{1,3} & m_{1,2} & p_{1,2}^2 & p_{1,2}p_{1,3} & m_{1,3} & p_{1,2}p_{1,3} & p_{1,3}^2 \\ p_{1,1} & m_{2,2} & 0 & 0 & p_{1,2} + p_{2,2} & 0 & 0 & p_{1,3} & 0 \\ p_{1,1} & 0 & m_{2,2} & 0 & 0 & p_{1,2} & 0 & 0 & p_{1,3} + p_{3,3} \\ p_{1,1} & 0 & 0 & m_{2,2} & p_{1,2} + p_{2,2} & p_{1,3} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 + 2p_{2,2} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & p_{2,2} & 1 & 0 & 0 & p_{3,3} \\ p_{1,1} & 0 & 0 & 0 & 0 & 0 & m_{2,2} & p_{1,2} & p_{1,3} + p_{3,3} \\ 0 & 0 & 0 & 0 & p_{2,2} & 0 & 0 & 1 & p_{3,3} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 + 2p_{3,3} \end{pmatrix}$$

$$\text{with } m_{1,1} = 2p_{1,1} + (1 - p_{1,2} - p_{1,3})^2, m_{1,2} = p_{1,2}(1 - p_{1,2} - p_{1,3}),$$

$$m_{1,3} = p_{1,3}(1 - p_{1,2} - p_{1,3}), m_{2,2} = 1 - p_{1,2} - p_{1,3}.$$

Linear diagram without the rule

$$\begin{pmatrix} m'_{1,1} & m'_{1,2} & 0 & m'_{1,2} & p_{1,2}^2 & 0 & 0 & 0 & 0 \\ p_{1,1} & m'_{2,2} & p_{2,3} - p_{1,2}p_{2,3} & 0 & m'_{2,5} & p_{1,2}p_{2,3} & 0 & 0 & 0 \\ p_{1,1} & 0 & 1 - p_{1,2} & 0 & 0 & p_{1,2} & 0 & 0 & p_{3,3} \\ p_{1,1} & 0 & 0 & m'_{2,2} & m'_{2,5} & 0 & p_{2,3} - p_{1,2}p_{2,3} & p_{1,2}p_{2,3} & 0 \\ 0 & 0 & 0 & 0 & m'_{5,5} & m'_{5,6} & 0 & m'_{5,6} & p_{2,3}^2 \\ 0 & 0 & 0 & 0 & p_{2,2} & 1 - p_{2,3} & 0 & 0 & p_{2,3} + p_{3,3} \\ p_{1,1} & 0 & 0 & 0 & 0 & 0 & 1 - p_{1,2} & p_{1,2} & p_{3,3} \\ 0 & 0 & 0 & 0 & p_{2,2} & 0 & 0 & 1 - p_{2,3} & p_{2,3} + p_{3,3} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 + 2p_{3,3} \end{pmatrix}$$

$$\text{with } m'_{1,1} = p_{1,1} + (1 - p_{1,2})^2, m'_{1,2} = p_{1,2}(1 - p_{1,2}), m'_{2,2} = (1 - p_{1,2})(1 - p_{2,3}),$$

$$m'_{2,5} = p_{1,2} + p_{2,2} - p_{1,2}p_{2,3}, m'_{5,5} = 2p_{2,2} + (1 - p_{2,3})^2, m'_{5,6} = p_{2,3}(1 - p_{2,3}).$$

Linear diagram with the rule

$$\begin{pmatrix} m''_{1,1} & m''_{1,2} & m''_{1,2} & p_{1,2}^2 & 0 & 0 & 0 \\ p_{1,1} & 1 - p_{1,2} & 0 & m''_{2,4} & p_{2,3} & 0 & 0 \\ p_{1,1} & 0 & 1 - p_{1,2} & m''_{2,4} & 0 & p_{2,3} & 0 \\ 0 & 0 & 0 & m''_{3,4} & m''_{3,5} & m''_{3,5} & p_{2,3}^2 \\ 0 & 0 & 0 & p_{2,2} & 1 - p_{2,3} & 0 & p_{2,3} + p_{3,3} \\ 0 & 0 & 0 & p_{2,2} & 0 & 1 - p_{2,3} & p_{2,3} + p_{3,3} \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 + 2p_{3,3} \end{pmatrix}$$

$$\text{with } m''_{1,1} = 2p_{1,1} + (1 - p_{1,2})^2, m''_{1,2} = (1 - p_{1,2})p_{1,2}, m''_{2,4} = p_{1,2} + p_{2,2} - p_{1,2}p_{2,3},$$

$$m''_{3,4} = 2p_{2,2} + (1 - p_{2,3})^2, m''_{3,5} = (1 - p_{2,3})p_{2,3}.$$

Table 2

The algebraic equations between g and f ; g denotes the parameter in the constraint (5) and f the measure of cell-type diversity. $root_{-1}$ denotes the largest real root of the equation $2(g+1)f^4 + g(g+1)f^3 - g(2g+5)f^2 - g^2f + 2g^2 = 0$ in g . g_1 and is nearly 1.29661, which is exactly the intersection of the two curves within $0 < g, 0 < f < 1$, the largest real root of $1 - 67g + 1224g^2 - 4008g^3 + 4599g^4 - 1701g^5 = 0$. Likewise, g_0 is nearly 1.14254 and is exactly the real root of $1 - 3g + 11g^2 - 8g^3 = 0$.

Branching without the cell-type order conservation rule

$$(g+2)f^2 + gf - g = 0$$

Linear without the rule

$$\begin{cases} g = root_{-1}2(g+1)f^4 + g(g+1)f^3 - g(2g+5)f^2 - g^2f + 2g^2 \dagger, & (0 \leq g < g_1) \\ 2(3g-1)f^2 + (3g-1)f - 3g = 0 & (g \geq g_1) \end{cases}$$

Linear with the rule

$$\begin{cases} 2(g-1)f^3 - g(g+3)f^2 + gf + g^2 = 0 \ddagger, & (0 \leq g < g_0) \\ 2gf^2 + (g-1)f - g = 0 & (g \geq g_0) \end{cases}$$

Table 3

The sequences in $g < 1$ and $g > 1$ regions. $g < 1$ and $g > 1$ correspond to the early developmental and mature stages, respectively. F_n denotes Fibonacci numbers. [‡]We should note that because of the mathematical difficulty, we have not completed the survey of quadratic irrational numbers (quasiperiodic patterns) of f in $g < 1, 1/g \in \mathbb{N}$ in the linear diagram without the rule, but we guess that such a number does not exist in this region.

	$g < 1, 1/g \in \mathbb{N}$	$g > 1, g \in \mathbb{N}$
Branching without rule	Quasiperiodic	Periodic ($g \in \{2F_{2n}^2\}$), Quasiperiodic (others)
Linear without rule	Periodic ($g = 1/7$), Indefinite (others [‡])	Quasiperiodic
Linear with rule	Periodic ($g = 1/4$), Quasiperiodic ($g = 1/3$), Indefinite (others)	Quasiperiodic




Fig. 1.





Fig. 2.



B $g=1$ (Linear with the rule)



C $g=2$ (Branching)



Fig. 3.

List of MI Preprint Series, Kyushu University

The Global COE Program
Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p -adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials

- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials
- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings

- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric q -functions of the q -Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications
- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain

- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations