

# Spectral properties of the semigroup for the linearized compressible Navier–Stokes equation around a parallel flow in a cylindrical domain

Aoyama, Reika  
Graduate School of Mathematics, Kyushu University

Kagei, Yoshiyuki  
Faculty of Mathematics, Kyushu University

<https://hdl.handle.net/2324/1500417>

---

出版情報 : MI Preprint Series. 2015-2, 2015-03-31. 九州大学大学院数理学研究院  
バージョン :  
権利関係 :



**MI Preprint Series**

Mathematics for Industry  
Kyushu University

**Spectral properties of the  
semigroup for the linearized  
compressible Navier-Stokes  
equation around a parallel flow  
in a cylindrical domain**

**Reika Aoyama  
& Yoshiyuki Kagei**

**MI 2015-2**

( Received March 31, 2015 )

Institute of Mathematics for Industry  
Graduate School of Mathematics  
Kyushu University  
Fukuoka, JAPAN

# Spectral properties of the semigroup for the linearized compressible Navier-Stokes equation around a parallel flow in a cylindrical domain

Reika Aoyama<sup>1</sup> and Yoshiyuki Kagei<sup>1</sup>

<sup>1</sup> Graduate School of Mathematics,  
Kyushu University,  
Nishi-ku, Motooka 744,  
Fukuoka, 819-0395, JAPAN

<sup>2</sup> Faculty of Mathematics,  
Kyushu University,  
Nishi-ku, Motooka 744,  
Fukuoka 819-0395, Japan

**Abstract:** This paper is concerned with the stability of a parallel flow of the compressible Navier-Stokes equation in a cylindrical domain. The spectrum of the linearized operator is analyzed for the purpose of the study of the nonlinear stability. It is shown that if the Reynolds and Mach numbers are sufficiently small, then the linearized semigroup is decomposed into two parts; one behaves like a solution of a one dimensional heat equation as time goes to infinity and the other one decays exponentially. Some estimates related to the spectral projections are established, which will also be useful for the study of the nonlinear problem.

**Keywords:** Compressible Navier-Stokes equation, parallel flow, cylindrical domain, linearized semigroup, asymptotic behavior

## 1 Introduction

We consider the initial boundary value problem for the equations for a barotropic motion of viscous compressible fluid

$$\partial_t \rho + \operatorname{div}(\rho v) = 0, \tag{1.1}$$

$$\rho(\partial_t v + v \cdot \nabla v) - \mu \Delta v - (\mu + \mu') \nabla \operatorname{div} v + \nabla p(\rho) = \rho g, \tag{1.2}$$

$$v|_{\partial D} = 0, \tag{1.3}$$

$$(\rho, v)|_{t=0} = (\rho_0, v_0) \tag{1.4}$$

in a cylindrical domain  $\Omega = D \times \mathbf{R}$ :

$$\Omega = \{x = (x', x_3); x' = (x_1, x_2) \in D, x_3 \in \mathbf{R}\}.$$

Here  $D$  is a bounded and connected domain in  $\mathbf{R}^2$  with a smooth boundary  $\partial D$ ;  $\rho = \rho(x, t)$  and  $v = {}^T(v^1(x, t), v^2(x, t), v^3(x, t))$  denote the unknown density and velocity at time  $t \geq 0$  and position  $x \in \Omega$ , respectively;  $P(\rho)$  is the pressure that is a smooth function of  $\rho$  and satisfies

$$p'(\rho_*) > 0$$

for a given positive constant  $\rho_*$ ;  $\mu$  and  $\mu'$  are the viscosity coefficients that satisfy

$$\mu > 0, \quad \frac{2}{3}\mu + \mu' \geq 0;$$

and  $g$  is an external force of the form  $g = {}^T(g^1(x'), g^2(x'), g^3(x'))$  with  $g^1$  and  $g^2$  satisfying

$$(g^1(x'), g^2(x')) = (\partial_{x_1}\Phi(x'), \partial_{x_2}\Phi(x')),$$

where  $\Phi$  and  $g^3$  are given smooth functions of  $x'$ . Here and in what follows  ${}^T$  stands for the transposition.

It is known that problem (1.1)-(1.3) has the stationary solution  $\bar{u}_s = {}^T(\bar{\rho}_s(x'), \bar{v}_s(x'))$ ;  $\bar{\rho}_s$  is determined by

$$\begin{cases} \text{Const.} - \Phi(x') = \int_{\rho_*}^{\bar{\rho}_s} \frac{p'(\eta)}{\eta} d\eta, \\ \int_D \bar{\rho}_s - \rho_* dx' = 0; \end{cases}$$

and  $\bar{v}_s$  takes the form

$$\bar{v}_s = {}^T(0, 0, \bar{v}_s^3(x')),$$

where  $\bar{v}_s^3(x')$  is the solution of

$$\begin{cases} -\mu \Delta' \bar{v}_s^3 = \bar{\rho}_s g^3, \\ \bar{v}_s^3|_{\partial D} = 0. \end{cases}$$

Here  $\Delta' = \partial_{x_1}^2 + \partial_{x_2}^2$ . The stationary solution  $\bar{u}_s$  represents a parallel flow in  $\Omega$ .

We are interested in the large time behavior of solutions to problem (1.1)-(1.4) when the initial value  $(\rho, v)|_{t=0} = (\rho_0, v_0)$  is sufficiently close to the stationary solution  $\bar{u}_s = {}^T(\bar{\rho}_s, \bar{v}_s)$ . In [1] the decay estimates of the linearized semigroup for (1.1) - (1.4) were established. In this paper we study the spectral properties of the linearized semigroup in more detail, which play an important role in the analysis of the nonlinear problem.

As for the asymptotic behavior of multidimensional compressible Navier-Stokes equations on unbounded domains, a lot of results have been obtained. See, e.g., [5, 6, 11, 13, 14, 15, 16, 17, 19] and references therein.

For the stability of parallel flow, detailed descriptions of large time behavior of disturbances have been obtained in the case of an  $n$  dimensional infinite layer  $\mathbf{R}^{n-1} \times (0, 1) = \{x = (x_h, x_n); x_h = (x_1, \dots, x_{n-1}) \in \mathbf{R}^{n-1}, 0 < x_n < 1\}$ . It

was proved in [10] that asymptotic behavior of solutions of the linearized problem is described by an  $n - 1$  dimensional linear heat equation, if Reynolds and Mach numbers are sufficiently small. The nonlinear problem was then studied in [9]; it was shown that if Reynolds and Mach numbers are sufficiently small, then the parallel flow is stable under sufficiently small initial disturbances in some Sobolev space. In the case of  $n \geq 3$ , the disturbance  $u(t)$  behaves like a solution of an  $n - 1$  dimensional linear heat equation as  $t \rightarrow \infty$ , while, in the case of  $n = 2$ ,  $u(t)$  behaves like a solution of a one dimensional viscous Burgers equation. See also [2, 3, 4] for the stability of time periodic parallel flow.

As for the case of the cylindrical domain  $\Omega$ , Iooss and Padula [7] studied the linearized stability of a stationary parallel flow under the periodic boundary condition in  $x_3$  with vanishing average condition on the basic period cell for the density-disturbance and proved that if the Reynolds number is suitably small, then the linearized semigroup decays exponentially as  $t \rightarrow \infty$ .

On the other hand, stability under the non-periodic but local disturbances (i.e., belonging to some  $L^2$  Sobolev space on  $\Omega$ ) was studied in [12] in the case of the motionless state  $\tilde{u}_s = T(\rho_*, 0)$ . It was shown that the disturbance decays in  $L^2(\Omega)$  in the order  $t^{-\frac{1}{4}}$  and its asymptotic leading part is given by a solution of a one dimensional linear heat equation. (See also [8] for the analysis in  $L^p(\Omega)$ .)

The linearized stability of parallel flow under non-periodic local disturbances on  $\Omega$  was then studied in [1]. It was shown that the linearized semigroup  $e^{-tL}$  satisfies the decay estimate

$$\|\partial_{x'}^k \partial_{x_3}^l e^{-tL} u_0\|_{L^2(\Omega)} \leq C \left\{ (1+t)^{-\frac{1}{4}-\frac{l}{2}} \|u_0\|_{L^1(\mathbf{R}; L^2(D))} + e^{-dt} \|u_0\|_{H^1(\Omega)} \right\} \quad (1.5)$$

for  $t \geq 0$  and  $0 \leq k + l \leq 1$ , provided that the Reynolds and Mach numbers are sufficiently small and that  $\bar{\rho}_s$  is sufficiently close to  $\rho_*$ . In view of the argument in [9], estimate (1.5) is not enough to show the global in time solvability of the nonlinear problem. The purpose of this paper is to derive more detailed spectral information of the linearized operator which will be available for the nonlinear analysis.

The main results of this paper are summarized as follows. We consider the linearized problem whose non-dimensional form is written as

$$\partial_t u + Lu = 0, \quad u|_{t=0} = u_0. \quad (1.6)$$

Here  $u = T(\phi, w)$  is the unknown; and  $u_0 = T(\phi_0, w_0)$  is a given initial value; and  $L$  denotes the linearized operator on  $L^2(\tilde{\Omega})$  defined by

$$L = \begin{pmatrix} v_s \cdot \nabla & \gamma^2 \operatorname{div}(\rho_s \cdot) \\ \nabla \left( \frac{\tilde{P}'(\rho_s)}{\gamma^2 \rho_s} \cdot \right) + \frac{\nu \Delta v_s}{\gamma^2 \rho_s^2} & -\frac{\nu}{\rho_s} \Delta I_3 - \frac{\nu + \nu'}{\rho_s} \nabla \operatorname{div} + v_s \cdot \nabla + T(\nabla v_s) \end{pmatrix},$$

with domain

$$D(L) = \{u = T(\phi, w) \in L^2(\tilde{\Omega}); w \in H_0^1(\tilde{\Omega}), Lu \in L^2(\tilde{\Omega})\},$$

where  $\tilde{\Omega}$ ,  $\tilde{D}$ ,  $\rho_s$ ,  $v_s$  and  $\tilde{P}(\rho_s)$  are the non-dimensional forms of  $\Omega$ ,  $D$ ,  $\bar{\rho}_s$ ,  $\bar{v}_s$  and  $p(\bar{\rho}_s)$  respectively;  $I_3$  denotes the  $3 \times 3$  identity matrix;  $\nu$ ,  $\nu'$  and  $\gamma$  are some positive constants.

We will show that there exists a bounded projection  $P_0$  satisfying  $P_0 e^{-tL} = e^{-tL} P_0$  such that if Reynolds and Mach numbers are sufficiently small, then

$$\|e^{-tL} P_0 u_0 - (\sigma u^{(0)})(t)\|_{L^2(\Omega)} \leq C(1+t)^{-\frac{3}{4}} \|u_0\|_{L^1(\Omega)}. \quad (1.7)$$

Here  $u^{(0)}$  is some function of  $x'$ , and  $\sigma$  is a function of  $x_3$  and  $t$ ; and  $\sigma$  is a solution of one dimensional linear heat equation

$$\begin{cases} \partial_t \sigma - \kappa_1 \partial_{x_3}^2 \sigma + \kappa_0 \partial_{x_3} \sigma = 0, \\ \sigma|_{t=0} = \int_0^1 \phi_0(x', x_3) dx_3 \end{cases} \quad (1.8)$$

with some constants  $\kappa_0 \in \mathbf{R}$  and  $\kappa_1 > 0$ . Some estimates for operators related to  $P_0$  are established; and, as in [2, 4], we will give a factorization of  $e^{-tL} P_0$  which is useful in the study of nonlinear problem. As for the  $I - P_0$  part of  $e^{-tL}$ , we will establish the exponential decay estimate

$$\|e^{-tL}(I - P_0)u_0\|_{H^1(\Omega)} \leq C e^{-dt} \{ \|u_0\|_{H^1(\Omega) \times \tilde{H}^1(\Omega)} + t^{-\frac{1}{2}} \|w_0\|_{L^2(\Omega)} \} \quad (1.9)$$

for a positive constant  $d$ . Here  $\tilde{H}^1(\Omega)$  is the set of all locally  $H^1$  functions in  $L^2(\Omega)$  whose tangential derivatives near  $\partial\Omega$  belong to  $L^2(\Omega)$ .

The linear problem (1.6) will be investigated through the Fourier transform  $\mathcal{F}$  in  $x_3 \in \mathbf{R}$  that leads to the problem on  $D$ , written in the form,

$$\partial_t u + L_\xi u = 0, \quad u|_{t=0} = u_0,$$

where  $\xi \in \mathbf{R}$  denotes the dual variable. The operator  $L_\xi$  has different properties  $|\xi| \leq r_0$  and  $|\xi| > r_0$ , where  $r_0$  is a positive number sufficiently small.

The spectrum of  $-L_\xi$  for  $|\xi| \leq r_0$  can be regarded as a perturbation from the one with  $\xi = 0$ , and we will show that the spectrum near the origin is given by a simple eigenvalue  $\lambda_0(\xi) = -i\kappa_0\xi - \kappa_1\xi^2 + \mathcal{O}(|\xi|^3)$  as  $|\xi| \rightarrow 0$ . Furthermore, we will establish the boundedness of the eigenprojection  $\Pi(\xi)$  for the eigenvalue  $\lambda_0(\xi)$  in some Sobolev space by investigating the regularity of the corresponding eigenfunctions. Setting  $P_0 = \mathcal{F}^{-1} \mathbf{1}_{\{|\xi| \leq r_0\}} \Pi(\xi) \mathcal{F}$  with a frequency cut-off function  $\mathbf{1}_{\{|\xi| \leq r_0\}}$  such that  $\mathbf{1}_{\{|\xi| \leq r_0\}} = 1$  for  $|\xi| \leq r_0$  and  $\mathbf{1}_{\{|\xi| \leq r_0\}} = 0$  for  $|\xi| > r_0$ , we find the desired asymptotic behavior of  $e^{-tL} P_0$  as described in (1.7) and (1.8). As for the complimentary part  $e^{-tL}(I - P_0)$ , we have already shown in [1] that

$$\|e^{-tL}(I - P_0)u_0\|_{H^1(\Omega)} \leq e^{-dt} \|u_0\|_{H^1(\Omega)}.$$

In this paper we will improve the class of initial value  $u_0$  and will show that the exponential decay estimate holds for  $u_0 \in H^1(\Omega) \times \tilde{H}^1(\Omega)$  as in (1.9). This improvement will be also useful in the study of the nonlinear problem.

This paper is organized as follows. In Section 2 we first rewrite the problem into the system of equations in a non-dimensional form and then present the existence of a stationary solution of parallel flow. We state the main result of this paper in Section 3. In Section 4 we will investigate the spectrum of  $-L_\xi$  for  $|\xi| \leq r_0$ , and in Section 5 we will establish a factorization of  $e^{-tL} P_0$  and prove (1.7). Section 6 is devoted to the proof of (1.9).

## 2 Preliminaries

In this section we introduce notations and then rewrite the problem into a non-dimensional form. In the end of this section we state the existence of stationary solution which represents parallel flow.

### 2.1 Notation

We first introduce some notations which will be used throughout the paper. For  $1 \leq p \leq \infty$  we denote by  $L^p(X)$  the usual Lebesgue space on a domain  $X$  and its norm is denoted by  $\|\cdot\|_{L^p(X)}$ . Let  $m$  be a nonnegative integer.  $H^m(X)$  denotes the  $m$ th order  $L^2$  Sobolev space on  $X$  with norm  $\|\cdot\|_{H^m(X)}$ . In particular, we write  $L^2(X)$  for  $H^0(X)$ .

We denote by  $C_0^m(X)$  the set of all  $C^m$  functions with compact support in  $X$ .  $H_0^m(X)$  stands for the completion of  $C_0^m(X)$  in  $H^m(X)$ . We denote by  $H^{-1}(X)$  the dual space of  $H_0^1(X)$  with norm  $\|\cdot\|_{H^{-1}(X)}$ .

We simply denote by  $L^p(X)$  (resp.,  $H^m(X)$ ) the set of all vector fields  $w = {}^T(w^1, w^2, w^3)$  on  $X$  and its norm is denoted by  $\|\cdot\|_{L^p(X)}$  (resp.,  $\|\cdot\|_{H^m(X)}$ ). For  $u = {}^T(\phi, w)$  with  $\phi \in H^k(X)$  and  $w = {}^T(w^1, w^2, w^3) \in H^m(X)$ , we define  $\|u\|_{H^k(X) \times H^m(X)}$  by  $\|u\|_{H^k(X) \times H^m(X)} = \|\phi\|_{H^k(X)} + \|w\|_{H^m(X)}$ .

When  $X = \Omega$  we abbreviate  $L^p(\Omega)$  as  $L^p$ , and likewise,  $H^m(\Omega)$  as  $H^m$ . The norm  $\|\cdot\|_{L^p(\Omega)}$  is written as  $\|\cdot\|_{L^p}$ , and likewise,  $\|\cdot\|_{H^m(\Omega)}$  as  $\|\cdot\|_{H^m}$ .

In the case  $X = D$  we denote the norm of  $L^p(D)$  by  $|\cdot|_p$ . The norm of  $H^m(D)$  is denoted by  $|\cdot|_{H^m}$ , respectively. The inner product of  $L^2(D)$  is denoted by

$$(f, g) = \int_D f(x') \overline{g(x')} dx', \quad f, g \in L^2(D).$$

Here  $\bar{g}$  denotes the complex conjugate of  $g$ . For  $u_j = {}^T(\phi_j, w_j)$  ( $j = 1, 2$ ), we also define a weighted inner product  $\langle u_1, u_2 \rangle$  by

$$\langle u_1, u_2 \rangle = \frac{1}{\gamma^2} \int_D \phi_1 \bar{\phi}_2 \frac{P'(\rho_s)}{\gamma^2 \rho_s} dx' + \int_D w_1 \cdot \bar{w}_2 \rho_s dx',$$

where  $\rho_s = \rho_s(x')$  is the density of the parallel flow  $u_s$ . As will be seen in Proposition 2.1 below,  $\rho_s(x')$  and  $\frac{P'(\rho_s(x'))}{\rho_s(x')}$  are strictly positive in  $D$ .

For  $f \in L^1(D)$  we denote the mean value of  $f$  over  $D$  by  $\langle f \rangle$ :

$$\langle f \rangle = (f, 1) = \frac{1}{|D|} \int_D f dx',$$

where  $|D| = \int_D dx'$ . For  $u = {}^T(\phi, w) \in L^1(D)$  with  $w = {}^T(w^1, w^2, w^3)$  we define  $\langle u \rangle$  by

$$\langle u \rangle = \langle \phi \rangle + \langle w_1 \rangle + \langle w_2 \rangle + \langle w_3 \rangle.$$

Partial derivatives of a function  $u$  in  $x, x', x_3$  and  $t$  are denoted by  $\partial_x u, \partial_{x'} u, \partial_{x_3} u$  and  $\partial_t u$ . We also write higher order partial derivatives of  $u$  in  $x$  as  $\partial_x^k u = (\partial_x^\alpha u; |\alpha| = k)$ .

We denote the  $n \times n$  identity matrix by  $I_n$ . We define  $4 \times 4$  diagonal matrices  $Q_0$  and  $\tilde{Q}$  by

$$Q_0 = \text{diag}(1, 0, 0, 0), \quad \tilde{Q} = \text{diag}(0, 1, 1, 1).$$

It then follows that for  $u = {}^T(\phi, w)$  with  $w = {}^T(w^1, w^2, w^3)$ ,

$$Q_0 u = \begin{pmatrix} \phi \\ 0 \end{pmatrix}, \quad \tilde{Q} u = \begin{pmatrix} 0 \\ w \end{pmatrix}.$$

We denote the Fourier transform of  $f = f(x_3)$  ( $x_3 \in \mathbf{R}$ ) by  $f$  or  $\mathcal{F}[f]$ :

$$f(\xi) = \mathcal{F}[f](\xi) = \int_{\mathbf{R}} f(x_3) e^{-i\xi x_3} dx_3, \quad \xi \in \mathbf{R}.$$

The inverse Fourier transform is denoted by  $\mathcal{F}^{-1}$ :

$$\mathcal{F}^{-1}[f](x_3) = (2\pi)^{-1} \int_{\mathbf{R}} f(\xi) e^{i\xi x_3} d\xi, \quad x_3 \in \mathbf{R}.$$

We denote the resolvent set of a closed operator  $A$  by  $\rho(A)$  and the spectrum by  $\sigma(A)$ .

We finally introduce a function space which consists of locally  $H^1$  functions in  $L^2(\Omega)$  whose tangential derivatives near  $\partial D$  belong to  $L^2(\Omega)$ . To do so, we first introduce a local curvilinear coordinate system. For any  $\bar{x}'_0 \in \partial D$ , there exist a neighborhood  $\tilde{\mathcal{O}}_{\bar{x}'_0}$  of  $\bar{x}'_0$  and a smooth diffeomorphism map  $\Psi = (\Psi_1, \Psi_2) : \tilde{\mathcal{O}}_{\bar{x}'_0} \rightarrow B_1(0) = \{z' = (z_1, z_2) : |z'| < 1\}$  such that

$$\begin{cases} \Psi(\tilde{\mathcal{O}}_{\bar{x}'_0} \cap D) = \{z' \in B_1(0) : z_1 > 0\}, \\ \Psi(\tilde{\mathcal{O}}_{\bar{x}'_0} \cap \partial D) = \{z' \in B_1(0) : z_1 = 0\}, \\ \det \nabla_{x'} \Psi \neq 0 \quad \text{on} \quad \overline{\tilde{\mathcal{O}}_{\bar{x}'_0} \cap D}. \end{cases}$$

By the tubular neighborhood theorem, there exist a neighborhood  $\mathcal{O}_{\bar{x}'_0}$  of  $\bar{x}'_0$  and a local curvilinear coordinate system  $y' = (y_1, y_2)$  on  $\mathcal{O}_{\bar{x}'_0}$  defined by

$$x' = y_1 a_1(y_2) + \Psi^{-1}(0, y_2) : \mathcal{R} \rightarrow \mathcal{O}_{\bar{x}'_0}, \quad (2.1)$$

where  $\mathcal{R} = \{y' = (y_1, y_2) : |y_1| \leq \tilde{\delta}_1, |y_2| \leq \tilde{\delta}_2\}$  for some  $\tilde{\delta}_1, \tilde{\delta}_2 > 0$ ;  $a_1(y_2)$  is the unit inward normal to  $\partial D$  that is given by

$$a_1(y_2) = \frac{\nabla_{x'} \Psi_1}{|\nabla_{x'} \Psi_1|}.$$

Setting  $y_3 = x_3$  we obtain

$$\nabla_x = e_1(y_2) \partial_{y_1} + J(y') e_2(y_2) \partial_{y_2} + e_3 \partial_{y_3},$$



$$\nabla_y = \begin{pmatrix} T e_1(y_2) \\ \frac{1}{J(y')} T e_2(y_2) \\ T e_3 \end{pmatrix} \nabla_x,$$

where

$$e_1(y_2) = \begin{pmatrix} a_1(y_2) \\ 0 \end{pmatrix}, \quad e_2(y_2) = \begin{pmatrix} a_2(y_2) \\ 0 \end{pmatrix}, \quad e_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}; \quad (2.2)$$

$$J(y') = |\det \nabla_{x'} \Psi|, \quad a_2(y_2) = \frac{-\nabla_{x'}^\perp \Psi_1}{|\nabla_{x'}^\perp \Psi_1|}$$

with  $\nabla_{x'}^\perp \Psi_1 = {}^T(-\partial_{x_2} \Psi_1, \partial_{x_1} \Psi_1)$ . Note that  $\partial_{y_1}$  and  $\partial_{y_2}$  are the inward normal derivative and tangential derivative at  $x' = \Psi^{-1}(0, y_2) \in \partial D \cap \mathcal{O}_{\bar{x}'_0}$ , respectively. Let us denote the normal and tangential derivatives by  $\partial_n$  and  $\partial$ , i.e.,

$$\partial_n = \partial_{y_1}, \quad \partial = \partial_{y_2}.$$

Since  $\partial D$  is compact, there are bounded open sets  $\mathcal{O}_m$  ( $m = 1, \dots, N$ ) such that  $\partial D \subset \cup_{m=1}^N \mathcal{O}_m$  and for each  $m = 1, \dots, N$ , there exists a local curvilinear coordinate system  $y' = (y_1, y_2)$  as defined in (2.1) with  $\mathcal{O}_{\bar{x}'_0}$ ,  $\Psi$  and  $\mathcal{R}$  replaced by  $\mathcal{O}_m$ ,  $\Psi^m$  and  $\mathcal{R}_m = \{y' = (y_1, y_2) : |y_1| < \tilde{\delta}_1^m, |y_2| < \tilde{\delta}_2^m\}$  for some  $\tilde{\delta}_1^m, \tilde{\delta}_2^m > 0$ . At last, we take an open set  $\mathcal{O}_0 \subset D$  such that

$$\cup_{m=0}^N \mathcal{O}_m \supset D, \quad \overline{\mathcal{O}_0} \cap \partial D = \emptyset.$$

We set a local coordinate  $y' = (y_1, y_2)$  such that  $y_1 = x_1, y_2 = x_2$  on  $\mathcal{O}_0$ . We note that if  $h \in H^2(D)$ , then  $h|_{\partial D} = 0$  implies that  $\partial^k h|_{\partial D \cap \mathcal{O}_m} = 0$  ( $k = 0, 1$ ).

Let us introduce a partition of unity  $\{\chi_m\}_{m=0}^N$  subordinate to  $\{\mathcal{O}_m\}_{m=0}^N$ , satisfying

$$\sum_{m=0}^N \chi_m = 1 \text{ on } D, \quad \chi_m \in C_0^\infty(\mathcal{O}_m) \text{ } (m = 0, 1, 2, \dots, N).$$

We denote by  $\tilde{H}^1(\Omega)$  the set of all locally  $H^1$  functions in  $L^2(\Omega)$  whose tangential derivatives near  $\partial\Omega$  belong to  $L^2(\Omega)$ , and its norm is denoted by  $\|w\|_{\tilde{H}^1(\Omega)}$ :

$$\|w\|_{\tilde{H}^1(\Omega)} = \|w\|_2 + \|\partial_{x_3} w\|_2 + \|\chi_0 \partial_{x'} w\|_2 + \sum_{m=1}^N \|\chi_m \partial w\|_2.$$

Note that  $H_0^1(\Omega)$  is dense in  $\tilde{H}^1(\Omega)$ .

## 2.2 Stationary solution

In this subsection we rewrite the problem into the one in a non-dimensional form and state the existence of stationary solution which represents parallel flow. Let  $k_0$  be an integer satisfying  $k_0 \geq 3$ . We introduce the following non-dimensional variables:

$$x = \ell \tilde{x}, \quad v = V \tilde{v}, \quad \rho = \rho_* \tilde{\rho}, \quad t = \frac{\ell}{V} \tilde{t},$$

$$p = \rho_* V^2 \tilde{P}, \quad \Phi = \frac{V^2}{\ell} \tilde{\Phi}, \quad g^3 = \frac{V^2}{\ell} \tilde{g}^3,$$

$$V = |\tilde{v}_s^3|_{C_*^{k_0}(D)} = \sum_{k=0}^{k_0} \sup_{x' \in D} \ell^k |\partial_{x'}^k \tilde{v}_s^3(x')|, \quad \ell = \left( \int_D dx' \right)^{\frac{1}{2}}.$$

The problem (1.1)-(1.4) is then transformed into the following non-dimensional problem on  $\tilde{\Omega} = \tilde{D} \times \mathbf{R}$ :

$$\partial_{\tilde{t}} \tilde{\rho} + \operatorname{div}_{\tilde{x}}(\tilde{\rho} \tilde{v}) = 0, \quad (2.3)$$

$$\tilde{\rho}(\partial_{\tilde{t}} \tilde{v} + \tilde{v} \cdot \nabla_{\tilde{x}} \tilde{v}) - \nu \Delta_{\tilde{x}} \tilde{v} - (\nu + \nu') \nabla_{\tilde{x}} \operatorname{div}_{\tilde{x}} \tilde{v} + \tilde{P}'(\tilde{\rho}) \nabla_{\tilde{x}} \tilde{\rho} = \tilde{\rho} \tilde{g}, \quad (2.4)$$

$$\tilde{v} \mid_{\partial \tilde{D}} = 0, \quad (2.5)$$

$$(\tilde{\rho}, \tilde{v}) \mid_{\tilde{t}=0} = (\tilde{\rho}_0, \tilde{v}_0). \quad (2.6)$$

Here  $\tilde{D}$  is a bounded and connected domain in  $\mathbf{R}^2$ ;  $\tilde{g} = {}^T(\partial_{\tilde{x}_1} \tilde{\Phi}, \partial_{\tilde{x}_2} \tilde{\Phi}, \tilde{g}^3)$ ; and  $\nu$  and  $\nu'$  are non-dimensional parameters:

$$\nu = \frac{\mu}{\rho_* \ell V}, \quad \nu' = \frac{\mu'}{\rho_* \ell V}.$$

We also introduce a parameter  $\gamma$ :

$$\gamma = \sqrt{\tilde{P}'(1)} = \frac{\sqrt{p'(\rho_*)}}{V}.$$

Note that the Reynolds and Mach numbers are given by  $1/\nu$  and  $1/\gamma$ , respectively.

In what follows, for simplicity, we omit tildes of  $\tilde{x}$ ,  $\tilde{t}$ ,  $\tilde{v}$ ,  $\tilde{\rho}$ ,  $\tilde{g}$ ,  $\tilde{P}$ ,  $\tilde{\Phi}$ ,  $\tilde{D}$  and  $\tilde{\Omega}$  and write them as  $x$ ,  $t$ ,  $v$ ,  $\rho$ ,  $g$ ,  $P$ ,  $\Phi$ ,  $D$  and  $\Omega$ . Observe that, due to the non-dimensionalization, we have

$$|D| = \int_D dx' = 1,$$

and thus,

$$\langle f \rangle = \int_D f(x') dx'.$$

Let us state the existence of a stationary solution which represents parallel flow.

**Proposition 2.1.** *If  $\Phi \in C^{k_0}(\overline{D})$ ,  $g^3 \in H^{k_0}(D)$  and  $|\Phi|_{C^{k_0}}$  is sufficiently small, then (2.3)-(2.5) has a stationary solution  $u_s = {}^T(\rho_s, v_s) \in C^{k_0}(\overline{D})$ . Here  $\rho_s$  satisfies*

$$\begin{cases} \text{Const.} - \Phi(x') = \int_1^{\rho_s(x')} \frac{P'(\eta)}{\eta} d\eta, \\ \int_D \rho_s dx' = 1, \quad \rho_1 < \rho_s(x') < \rho_2 \quad (\rho_1 < 1 < \rho_2) \end{cases}$$

for some constants  $\rho_1, \rho_2 > 0$ ; and  $v_s$  is a function of the form  $v_s = {}^T(0, 0, v_s^3)$  with  $v_s^3 = v_s^3(x')$  being the solution of

$$\begin{cases} -\nu \Delta' v_s^3 = \rho_s g^3, \\ v_s^3 \mid_{\partial D} = 0. \end{cases}$$

Furthermore,  $u_s = T(\rho_s, v_s)$  satisfies the estimates:

$$|\rho_s(x') - 1|_{C^k} \leq C|\Phi|_{C^k}(1 + |\Phi|_{C^k})^k,$$

$$|v_s^3|_{C^k} \leq C|v_s^3|_{H^{k+2}} \leq C|\Phi|_{C^k}(1 + |\Phi|_{C^k})^k |g^3|_{H^k}$$

for  $k = 3, 4, \dots, k_0$ .

Proposition 2.1 can be proved in a similar manner to the proof of [18, Lemma 2.1].

### 3 Main results

We set  $\rho = \rho_s + \gamma^{-2}\phi$  and  $v = v_s + w$  in (2.3)-(2.6) (without tildes) and omit the nonlinear terms of  $\phi$  and  $w$ . We then arrive at the linearized problem

$$\partial_t u + Lu = 0, \quad u = T(\phi, w), \quad w|_{\partial D} = 0, \quad u|_{t=0} = u_0, \quad (3.1)$$

where

$$L = \left( \begin{array}{cc} v_s \cdot \nabla & \gamma^2 \operatorname{div}(\rho_s \cdot) \\ \nabla \left( \frac{P'(\rho_s)}{\gamma^2 \rho_s} \cdot \right) + \frac{\nu \Delta' v_s}{\gamma^2 \rho_s^2} & -\frac{\nu}{\rho_s} \Delta I_3 - \frac{\nu + \nu'}{\rho_s} \nabla \operatorname{div} + v_s \cdot \nabla + e_3 \otimes (\nabla v_s^3) \end{array} \right).$$

Here,  $e_3 = T(0, 0, 1)$ ; and, for  $\mathbf{a} = T(a_1, a_2, a_3)$  and  $\mathbf{b} = T(b_1, b_2, b_3)$ ,  $\mathbf{a} \otimes \mathbf{b}$  is the  $3 \times 3$  matrix  $(a_i b_j)$ .

We consider  $L$  as an operator on  $L^2(\Omega)$  with domain

$$D(L) = \{u = T(\phi, w) \in L^2(\Omega); \quad w \in H_0^1(\Omega), \quad Lu \in L^2(\Omega)\}.$$

As was shown in [1] (see also [7]),  $-L$  generates a  $C_0$ -semigroup  $e^{-tL}$  on  $L^2(\Omega)$ . Furthermore, if  $u_0 \in H^1(\Omega) \times \tilde{H}^1(\Omega)$ , then

$$\begin{aligned} e^{-tL} u_0 &\in C([0, T]; H^1(\Omega) \times \tilde{H}^1(\Omega)) \cap C((0, T]; H^1(\Omega) \times H_0^1(\Omega)), \\ \nabla \tilde{Q} e^{-tL} u_0 &\in L^2(0, T; \tilde{H}^1(\Omega)) \end{aligned} \quad (3.2)$$

for all  $T > 0$ . The regularity property (3.2) of  $e^{-tL}$  can be proved as follows. It is not difficult to see that if  $u_0 \in H^1(\Omega) \times H_0^1(\Omega)$ , then  $e^{-tL} u_0$  satisfies

$$\begin{aligned} e^{-tL} u_0 &\in C([0, T]; H^1(\Omega) \times H_0^1(\Omega)), \\ \nabla \tilde{Q} e^{-tL} u_0 &\in L^2(0, T; H^1(\Omega)). \end{aligned} \quad (3.3)$$

Since  $H_0^1(\Omega)$  is dense in  $\tilde{H}^1(\Omega)$ , one can see from (3.3), Lemma 6.5 and Lemma 6.6 below that  $e^{-tL} u_0$  satisfies (3.2) if  $u_0 \in H^1(\Omega) \times \tilde{H}^1(\Omega)$ .

In what follows we set

$$\omega = \|\rho_s - 1\|_{C^{k_0}}.$$

We have the following estimates for  $e^{-tL} u_0$ .

**Theorem 3.1.** *There exist positive constants  $\nu_0$ ,  $\gamma_0$  and  $\omega_0$  such that if  $\nu \geq \nu_0$ ,  $\frac{\gamma^2}{2\nu+\nu'} \geq \gamma_0^2$  and  $\omega \leq \omega_0$ , then  $e^{-tL}u_0$  is decomposed as*

$$e^{-tL}u_0 = e^{-tL}P_0u_0 + e^{-tL}P_\infty u_0.$$

Here  $P_0$  and  $P_\infty$  are projections satisfying

$$P_0 + P_\infty = I, \quad P^2 = P,$$

$$PL \subset LP, \quad Pe^{-tL} = e^{-tL}P$$

for  $P \in \{P_0, P_\infty\}$ ; and  $e^{-tL}P_0$  and  $e^{-tL}P_\infty$  have the following properties.

(i) If  $u_0 \in L^1(\Omega) \cap L^2(\Omega)$ , then  $e^{-tL}P_0u_0$  satisfies the following estimates

$$\|\partial_{x'}^k \partial_{x_3}^l e^{-tL}P_0u_0\|_2 \leq C_{k,l}(1+t)^{-\frac{1}{4}-\frac{l}{2}}\|u_0\|_1 \quad (3.4)$$

uniformly for  $t \geq 0$  and for  $k = 0, 1, \dots, k_0$  and  $l = 0, 1, \dots$ ;

$$\|e^{-tL}P_0u_0 - [\mathcal{H}(t)\langle\phi_0\rangle]u^{(0)}\|_2 \leq Ct^{-\frac{3}{4}}\|u_0\|_1 \quad (3.5)$$

uniformly for  $t > 0$ . Here

$$\mathcal{H}(t)\langle\phi_0\rangle = \mathcal{F}^{-1}[e^{-(i\kappa_0\xi + \kappa_1\xi^2)t}\langle\phi_0\rangle],$$

where  $u^{(0)} = u^{(0)}(x')$  is the function given in Lemma 4.1 below; and  $\kappa_0 \in \mathbf{R}$  and  $\kappa_1 > 0$  are some constants satisfying

$$\kappa_0 = \mathcal{O}(1),$$

$$\kappa_1 = C\frac{\gamma^2}{\nu}\left\{1 + \mathcal{O}\left(\frac{1}{\gamma^2}\right) + \left(\frac{\nu}{\gamma^2} + \frac{1}{\nu^2}\right) \times \mathcal{O}\left(\frac{2\nu+\nu'}{\gamma^2}\right)\right\},$$

where  $C$  is a positive constant.

(ii) If  $u_0 \in H^1(\Omega) \times \tilde{H}^1(\Omega)$ , then there exists a constant  $d > 0$  such that  $e^{-tL}P_\infty u_0$  satisfies

$$\|e^{-tL}P_\infty u_0\|_{H^1} \leq Ce^{-dt}(\|u_0\|_{H^1 \times \tilde{H}^1} + t^{-\frac{1}{2}}\|w_0\|_2) \quad (3.6)$$

uniformly for  $t > 0$ .

**Remark 3.2.** It is well-known that if  $u_0 = {}^T(\phi_0, w_0) \in L^1(\Omega)$ , then  $\|\mathcal{H}(t)\langle\phi_0\rangle\|_2 = \mathcal{O}(t^{-\frac{1}{4}})$ , and  $\sigma = \sigma(x_3, t) = \mathcal{H}(t)\langle\phi_0\rangle$  satisfies

$$\begin{cases} \partial_t \sigma - \kappa_1 \partial_{x_3}^2 \sigma + \kappa_0 \partial_{x_3} \sigma = 0, \\ \sigma|_{t=0} = \int_D \phi_0(x', x_3) dx'. \end{cases}$$

More detailed properties of  $P_0$  and  $e^{-tL}P_0$  will be given in Section 5 below, where we will establish a factorization of  $e^{-tL}P_0$  which will be useful in the nonlinear analysis.

To prove Theorem 3.1, we consider the Fourier transform of (3.1) in  $x_3$  variable which is written as

$$\partial_t \phi + i\xi v_s^3 \phi + \gamma^2 \nabla' \cdot (\rho_s w') + \gamma^2 i\xi \rho_s w^3 = 0, \quad (3.7)$$

$$\partial_t w' - \frac{\nu}{\rho_s} (\Delta' - \xi^2) w' - \frac{\nu + \nu'}{\rho_s} \nabla' (\nabla' \cdot w' + i\xi w^3) + \nabla' \left( \frac{P'(\rho_s)}{\gamma^2 \rho_s} \phi \right) + i\xi v_s^3 w' = 0, \quad (3.8)$$

$$\begin{aligned} \partial_t w^3 - \frac{\nu}{\rho_s} (\Delta' - \xi^2) w^3 - \frac{\nu + \nu'}{\rho_s} i\xi (\nabla' \cdot w' + i\xi w^3) + i\xi \left( \frac{P'(\rho_s)}{\gamma^2 \rho_s} \phi \right) + i\xi v_s^3 w^3 \\ + \frac{\nu}{\gamma^2 \rho_s^2} \Delta' v_s^3 \phi + w' \cdot \nabla' v_s^3 = 0, \end{aligned} \quad (3.9)$$

$$w|_{\partial D} = 0 \quad (3.10)$$

for  $t > 0$ , and

$$T(\phi, w)|_{t=0} = T(\phi_0, w_0) = u_0. \quad (3.11)$$

Therefore, we arrive at the following problem

$$\frac{du}{dt} + L_\xi u = 0, \quad u|_{t=0} = u_0, \quad (3.12)$$

where  $\xi \in \mathbf{R}$  is a parameter. Here  $u = T(\phi(x', t), w(x', t)) \in D(L_\xi)$  ( $x' \in D$ ,  $t > 0$ ),  $u_0$  is a given initial value, and  $L_\xi$  is the operator on  $L^2(D)$  of the form

$$L_\xi = A_\xi + B_\xi + C_0,$$

where

$$\begin{aligned} A_\xi &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & -\frac{\nu}{\rho_s} (\Delta' - |\xi|^2) I_2 - \frac{\nu + \nu'}{\rho_s} \nabla' \nabla' & -i \frac{\nu + \nu'}{\rho_s} \xi \nabla' \\ 0 & -i \frac{\nu + \nu'}{\rho_s} \xi \nabla' & -\frac{\nu}{\rho_s} (\Delta' - |\xi|^2) + \frac{\nu + \nu'}{\rho_s} |\xi|^2 \end{pmatrix}, \\ B_\xi &= \begin{pmatrix} i\xi v_s^3 & \gamma^2 \nabla'(\rho_s \cdot) & i\gamma^2 \rho_s \xi \\ \nabla' \left( \frac{P(\rho_s)}{\gamma^2 \rho_s} \cdot \right) & i\xi v_s^3 I_2 & 0 \\ i\xi \frac{P(\rho_s)}{\gamma^2 \rho_s} & 0 & i\xi v_s^3 \end{pmatrix}, \quad C_0 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \frac{\nu}{\gamma^2 \rho_s^2} \Delta' v_s^3 & T(\nabla' v_s^3) & 0 \end{pmatrix} \end{aligned}$$

with domain

$$D(L_\xi) = \{u = T(\phi, w) \in L^2(D); w \in H_0^1(D), L_\xi u \in L^2(D)\}.$$

Note that  $D(L_\xi) = D(L_0)$  for all  $\xi \in \mathbf{R}$ . Here and in what follows, we denote

$$\Delta' = \partial_{x_1}^2 + \partial_{x_2}^2, \quad \nabla' = T(\partial_{x_1}, \partial_{x_2}).$$

As in the case of  $L$ , we can see that  $-L_\xi$  generates a  $C_0$ -semigroup on  $L^2(D)$ . Furthermore, if  $u_0 \in H^1(D) \times \tilde{H}^1(D)$ , then

$$\begin{aligned} e^{-tL_\xi} u_0 &\in C([0, T]; H^1(D) \times \tilde{H}^1(D)) \cap C((0, T]; H^1(D) \times H_0^1(D)), \\ \partial_{x'} \tilde{Q} e^{-tL_\xi} u_0 &\in L^2(0, T; \tilde{H}^1(D)) \end{aligned} \quad (3.13)$$

for any  $T > 0$ .

In Section 4 we will investigate the spectrum of  $-L_\xi$  for  $|\xi| \ll 1$ . In Section 5 we will give the proof of Theorem 3.1 (i). In Section 6 we will prove Theorem 3.1 (ii).

## 4 Spectrum of $-L_\xi$ for $|\xi| \ll 1$

In this section, we consider the spectrum of  $-L_\xi$  for  $|\xi| \ll 1$ . For simplicity, in what follows, we denote  $\nu + \nu'$  by  $\tilde{\nu}$ , i.e.,

$$\tilde{\nu} = \nu + \nu'.$$

Let us consider the resolvent problem

$$(\lambda + L_\xi)u = f$$

with  $|\xi| \ll 1$ , where  $u = {}^T(\phi, w) \in D(L_\xi) = D(L_0)$  and  $f = {}^T(f^0, g) \in L^2(D)$ .

We introduce the adjoint operator  $L_\xi^*$  of  $L_\xi$  with respect to the weighted inner product  $\langle \cdot, \cdot \rangle$ . The operator  $L_\xi^*$  is given by

$$L_\xi^* = A_\xi^* + B_\xi^* + C_0^*$$

with domain of definition

$$D(L_\xi^*) = \{u = {}^T(\phi, w) \in L^2(D); w \in H_0^1(D), L_\xi^*u \in L^2(D)\}.$$

Here

$$A_\xi^* = A_\xi, \quad B_\xi^* = -B_\xi$$

and

$$C_0^* = \begin{pmatrix} 0 & 0 & \frac{\gamma^2 \nu \Delta' v_s^3}{P'(\rho_s)} \\ 0 & 0 & \nabla' v_s^3 \\ 0 & 0 & 0 \end{pmatrix}.$$

Note that  $D(L_\xi) = D(L_\xi^*)$  for any  $\xi \in \mathbf{R}$ .

We begin with a lemma on the zero eigenvalue of  $L_0$  and  $L_0^*$  which was proved in [1, Lemma 4.1]. Here  $L_0$  and  $L_0^*$  stand for  $L_\xi$  and  $L_\xi^*$  with  $\xi = 0$ , respectively.

**Lemma 4.1.** ([1, Lemma 4.1]) (i) *There exists a constant  $\omega_0 > 0$  such that if  $\omega \leq \omega_0$ , then  $\lambda = 0$  is a simple eigenvalue of  $L_0$  and  $L_0^*$ .*

(ii) *The eigenspaces for  $\lambda = 0$  of  $L_0$  and  $L_0^*$  are spanned by  $u^{(0)}$  and  $u^{(0)*}$ , respectively, where*

$$u^{(0)} = {}^T(\phi^{(0)}, w^{(0)}), \quad w^{(0)} = {}^T(0, 0, w^{(0),3})$$

and

$$u^{(0)*} = {}^T(\phi^{(0)*}, 0).$$

Here

$$\phi^{(0)}(x') = \alpha_0 \frac{\gamma^2 \rho_s(x')}{P'(\rho_s(x'))}, \quad \alpha_0 = \left( \int_D \frac{\gamma^2 \rho_s(x')}{P'(\rho_s(x'))} dx' \right)^{-1};$$

and  $w^{(0),3}$  is the solution of the following problem

$$\begin{cases} -\Delta' w^{(0),3} = -\frac{1}{\gamma^2 \rho_s} \Delta' v_s^3 \phi^{(0)}, \\ w^{(0),3} |_{\partial D} = 0; \end{cases}$$

and

$$\phi^{(0)*}(x') = \frac{\gamma^2}{\alpha_0} \phi^{(0)}(x').$$

Furthermore,  $\phi^{(0)} = O(1)$ ,  $\alpha_0 = O(1)$  and  $w^{(0),3} = O(\gamma^{-2})$  as  $\gamma \rightarrow \infty$ .

(iii) The eigenprojections  $\Pi^{(0)}$  and  $\Pi^{(0)*}$  for  $\lambda = 0$  of  $L_0$  and  $L_0^*$  are given by

$$\Pi^{(0)}u = \langle u, u^{(0)*} \rangle u^{(0)} = \langle Q_0 u \rangle u^{(0)},$$

$$\Pi^{(0)*}u = \langle u, u^{(0)} \rangle u^{(0)*}$$

for  $u = T(\phi, w)$ , respectively.

(iv) Let  $u^{(0)}$  be written as  $u^{(0)} = u_0^{(0)} + u_1^{(0)}$ , where

$$u_0^{(0)} = T(\phi^{(0)}, 0), \quad u_1^{(0)} = T(0, w^{(0)}).$$

Then

$$u^{(0)*} = \frac{\gamma^2}{\alpha_0} u_0^{(0)}$$

and

$$\langle u, u^{(0)} \rangle = \frac{\alpha_0}{\gamma^2} \langle \phi \rangle + (w^3, w^{(0),3} \rho_s)$$

for  $u = T(\phi, w) = T(\phi, w', w^3)$ .

We next establish the resolvent estimate for  $|\xi| \ll 1$ . To do so, let us consider the resolvent problem for  $\xi = 0$

$$(\lambda + L_0)u = f, \tag{4.1}$$

where  $u = T(\phi, w) \in D(L_0)$  and  $f = T(f^0, g) \in L^2(D)$ . Decomposing  $u$  in (4.1) as

$$u = \langle \phi \rangle u^{(0)} + u_1$$

with

$$u_1 = (I - \Pi^{(0)})u,$$

we obtain

$$\lambda(\langle \phi \rangle u^{(0)} + u_1) + L_0 u_1 = f.$$

Applying  $\Pi^{(0)}$  and  $I - \Pi^{(0)}$  to this equation, we have

$$\begin{cases} \lambda \langle \phi \rangle = \langle f^0 \rangle, \\ \lambda u_1 + L_0 u_1 = f_1, \end{cases} \tag{4.2}$$

where  $f_1 = (I - \Pi^{(0)})f$ . We see from the first equation of (4.2) that if  $\lambda \neq 0$ , then

$$\langle \phi \rangle = \frac{1}{\lambda} \langle f^0 \rangle.$$

This implies that

$$|\langle \phi \rangle| \leq \frac{1}{|\lambda|} |f^0|_2. \tag{4.3}$$

On the other hand, the  $u_1$ -part has the following properties. The second equation of (4.2) is written as

$$\begin{cases} \lambda\phi_1 + \gamma^2\nabla' \cdot (\rho_s w_1') = f_1^0, \\ \lambda w_1' - \frac{\nu}{\rho_s}\Delta' w_1' - \frac{\tilde{\nu}}{\rho_s}\nabla'\nabla' \cdot w_1' + \nabla' \left( \frac{P'(\rho_s)}{\gamma^2\rho_s}\phi_1 \right) = g_1', \\ \lambda w_1^3 - \frac{\nu}{\rho_s}\Delta' w_1^3 + \frac{\nu}{\gamma^2\rho_s^2}\Delta' v_s^3\phi_1 + w_1' \cdot \nabla' v_s^3 = g_1^3, \end{cases} \quad (4.4)$$

where  $u_1 = {}^T(\phi_1, w_1) = {}^T(\phi_1, w_1', w_1^3)$  and  $f_1 = {}^T(f_1^0, g_1) = {}^T(f_1^0, g_1', g_1^3)$ .

To state the estimates for the  $u_1$ -part, we introduce a quantity  $\tilde{D}_0[w_1]$  defined by

$$\tilde{D}_0[w_1] = |\nabla' w_1|_2^2 + |\nabla' \cdot w_1'|_2^2$$

for  $w_1 = {}^T(w_1', w_1^3)$ .

**Proposition 4.2.** *There exist constants  $\nu_0 > 0$ ,  $\gamma_0 > 0$  and  $\omega_0 > 0$  and an energy functional  $E_0[u_1]$  such that if  $\nu \geq \nu_0$ ,  $\frac{\gamma^2}{\nu+\tilde{\nu}} \geq \gamma_0^2$  and  $\omega \leq \omega_0$ , then there holds the estimate*

$$(\operatorname{Re}\lambda)E_0[u_1] + c(|\phi_1|_2^2 + \tilde{D}_0[w_1]) \leq C|f_1|_2|u_1|_2,$$

where  $c$  and  $C$  are positive constants independent of  $u_1$  and  $\lambda$ ; and  $E_0[u_1]$  is equivalent to  $|u_1|_2^2$ .

Proposition 4.2 can be proved in a similar manner to the proof of [1, Proposition 4.7] by replacing  $\frac{d}{dt}$  with  $\operatorname{Re}\lambda$  and taking  $\xi = 0$  there.

The Poincaré inequality yields  $\tilde{D}_0[w_1] \geq C|w_1|_2^2$  with a positive constant  $C$ . Therefore, the resolvent estimates for  $-L_0$  now follow from (4.3) and Proposition 4.2.

**Proposition 4.3.** *There exist constants  $\nu_0 > 0$ ,  $\gamma_0 > 0$  and  $\omega_0 > 0$  such that if  $\nu \geq \nu_0$ ,  $\frac{\gamma^2}{\nu+\tilde{\nu}} \geq \gamma_0^2$  and  $\omega \leq \omega_0$ , then there is a positive constant  $c_0 > 0$  such that*

$$\Sigma_0 \equiv \{\lambda \neq 0 : \operatorname{Re}\lambda > -c_0\} \subset \rho(-L_0).$$

Furthermore, the following estimates

$$|(\lambda + L_0)^{-1}f|_2 \leq C\left\{\frac{1}{|\lambda|}|f^0|_2 + \frac{1}{(\operatorname{Re}\lambda + c_0)}|f_1|_2\right\},$$

$$|\partial_{x'}\{\tilde{Q}(\lambda + L_0)^{-1}f\}|_2 \leq C\left\{\frac{1}{|\lambda|}|f^0|_2 + \frac{1}{(\operatorname{Re}\lambda + c_0)^{1/2}}|f_1|_2\right\}$$

hold uniformly for  $\lambda \in \Sigma_0$ . The same assertions also hold for  $-L_0^*$ .

Based on Proposition 4.3, we have the resolvent estimates for  $-L_\xi$  with  $|\xi| \ll 1$ .

**Theorem 4.4.** *There exist constants  $\nu_0 > 0$ ,  $\gamma_0 > 0$  and  $\omega_0 > 0$  such that if  $\nu \geq \nu_0$ ,  $\frac{\gamma^2}{\nu+\tilde{\nu}} \geq \gamma_0^2$  and  $\omega \leq \omega_0$ , then the following assertions hold. For any  $\eta$  satisfying  $0 < \eta \leq \frac{c_0}{2}$  there is a number  $r_0 = r_0(\eta)$  such that*

$$\Sigma_1 \equiv \{\lambda \neq 0 : |\lambda| \geq \eta, \operatorname{Re}\lambda \geq -\frac{c_0}{2}\} \subset \rho(-L_\xi)$$



for  $|\xi| \leq r_0$ . Furthermore, the following estimates

$$|(\lambda + L_\xi)^{-1}f|_2 \leq C|f|_2,$$

$$|\partial_{x'}\{\tilde{Q}(\lambda + L_\xi)^{-1}f\}|_2 \leq C|f|_2$$

hold uniformly for  $\lambda \in \Sigma_1$  and  $\xi$  with  $|\xi| \leq r_0$ . The same assertions also hold for  $-L_\xi^*$ .

**Proof.** Let us decompose  $L_\xi$  as

$$L_\xi = L_0 + \xi L^{(1)} + \xi^2 L^{(2)},$$

where

$$L^{(1)} = i \begin{pmatrix} v_s^3 & 0 & \gamma^2 \rho_s \\ 0 & v_s^3 I_2 & -\frac{\tilde{\nu}}{\rho_s} \nabla' \\ \frac{P'(\rho_s)}{\gamma^2 \rho_s} & -\frac{\tilde{\nu}}{\rho_s} \nabla' & v_s^3 \end{pmatrix}, \quad L^{(2)} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{\nu}{\rho_s} I_2 & 0 \\ 0 & 0 & \frac{\nu + \tilde{\nu}}{\rho_s} \end{pmatrix}.$$

For  $u = {}^T(\phi, w) \in L^2(D) \times H_0^1(D)$  we have

$$|L^{(1)}u|_2 \leq C|u|_{L^2 \times H^1}, \quad |L^{(2)}u|_2 \leq C|u|_2. \quad (4.5)$$

Therefore, we see from Proposition 4.3 that for any  $0 < \eta \leq \frac{c_0}{2}$  there exists  $r_0 > 0$  such that if  $|\xi| \leq r_0$ , then

$$|(\xi L^{(1)} + \xi^2 L^{(2)})(\lambda + L_0)^{-1}f|_2 \leq \frac{1}{2}|f|_2. \quad (4.6)$$

It then follows that

$$\Sigma_1 \equiv \{\lambda : |\lambda| > \eta, \operatorname{Re} \lambda \geq -\frac{c_0}{2}\} \subset \rho(-L_\xi),$$

and that, if  $\lambda \in \Sigma_1$ , then  $(\lambda + L_\xi)^{-1}$  is given by the Neumann series expansion

$$(\lambda + L_\xi)^{-1} = (\lambda + L_0)^{-1} + \sum_{N=0}^{\infty} (-1)^N [(\xi L^{(1)} + \xi^2 L^{(2)})(\lambda + L_0)^{-1}]$$

for  $|\xi| \leq r_0$ , and it holds that

$$|(\lambda + L_\xi)^{-1}f|_2 \leq C|f|_2 \quad (4.7)$$

for  $\lambda \in \Sigma_1$  and  $|\xi| \leq r_0$ . We thus obtain the desired estimates. This completes the proof.  $\square$

As for the spectrum of  $-L_\xi$  near  $\lambda = 0$ , we have the following result.

**Theorem 4.5.** *There exist positive constants  $\nu_0$ ,  $\gamma_0$ ,  $\omega_0$  and  $r_0$  such that if  $\nu \geq \nu_0$ ,  $\frac{\gamma^2}{\nu+\tilde{\nu}} \geq \gamma_0^2$  and  $\omega \leq \omega_0$ , then it holds that*

$$\sigma(-L_\xi) \cap \{\lambda : |\lambda| \leq \frac{c_0}{2}\} = \{\lambda_0(\xi)\}$$

for  $\xi$  with  $|\xi| \leq r_0$ , where  $\lambda_0(\xi)$  is a simple eigenvalue of  $-L_\xi$  that has the form

$$\lambda_0(\xi) = -i\kappa_0\xi - \kappa_1\xi^2 + \mathcal{O}(|\xi|^3)$$

as  $\xi \rightarrow 0$ . Here  $\kappa_0 \in \mathbf{R}$  and  $\kappa_1 > 0$  are the numbers given by

$$\kappa_0 = \langle v_s^3 \phi^{(0)} + \gamma^2 \rho_s w^{(0),3} \rangle = \mathcal{O}(1),$$

$$\kappa_1 = \frac{\gamma^2}{\nu} \left\{ \alpha_0 |(-\Delta')^{-\frac{1}{2}} \rho_s|_2^2 + \mathcal{O}\left(\frac{1}{\gamma^2}\right) + \left(\frac{\nu}{\gamma^2} + \frac{1}{\nu^2}\right) \times \mathcal{O}\left(\frac{\nu+\tilde{\nu}}{\gamma^2}\right) \right\},$$

where  $-\Delta'$  denotes the Laplace operator on  $L^2(D)$  under the zero Dirichlet boundary condition with domain

$$D(-\Delta') = H^2(D) \cap H_0^1(D).$$

**Proof.** For  $u \in L^2(D) \times H_0^1(D)$  we see from Theorem 4.4 and (4.5) that

$$|L^{(1)}u|_2 \leq C(|L_0u|_2 + |u|_2), \quad |L^{(2)}u|_2 \leq C|u|_2.$$

Therefore, since 0 is a simple eigenvalue of  $-L_0$ , we see from the analytic perturbation theory that there exists a positive constant  $r_0$  such that

$$\sigma(-L_\xi) \cap \{\lambda : |\lambda| \leq \frac{c_0}{2}\} = \{\lambda_0(\xi)\}$$

for all  $\xi$  with  $|\xi| \leq r_0$ . Here  $\lambda_0(\xi)$  is a simple eigenvalue of  $-L_\xi$ . Furthermore,  $\lambda_0(\xi)$  and the eigenprojection  $\Pi(\xi)$  for  $\lambda_0(\xi)$  are expanded as

$$\lambda_0(\xi) = \lambda^{(0)} + \xi\lambda^{(1)} + \xi^2\lambda^{(2)} + \mathcal{O}(|\xi|^3),$$

$$\Pi(\xi) = \Pi^{(0)} + \xi\Pi^{(1)} + \mathcal{O}(|\xi|^2) \tag{4.8}$$

with

$$\begin{aligned} \lambda^{(0)} &= 0, \\ \lambda^{(1)} &= -\langle L^{(1)}u^{(0)}, u^{(0)*} \rangle, \\ \lambda^{(2)} &= -\langle L^{(2)}u^{(0)}, u^{(0)*} \rangle + \langle L^{(1)}SL^{(1)}u^{(0)}, u^{(0)*} \rangle, \\ \Pi^{(1)} &= -\Pi^{(0)}L^{(1)}S - SL^{(1)}\Pi^{(0)}, \end{aligned}$$

where

$$S = \left\{ (I - \Pi^{(0)})L_0(I - \Pi^{(0)}) \right\}^{-1}.$$

We first consider  $\lambda^{(1)}$ . Since

$$L^{(1)}u^{(0)} = i \begin{pmatrix} v_s^3 \phi^{(0)} + \gamma^2 \rho_s w^{(0),3} \\ -\frac{\tilde{\nu}}{\rho_s} \nabla' w^{(0),3} \\ \alpha_0 + v_s^3 w^{(0),3} \end{pmatrix},$$

we obtain

$$\lambda^{(1)} = -\langle L^{(1)}u^{(0)}, u^{(0)*} \rangle = -\langle Q_0 L^{(1)}u^{(0)} \rangle = -i\langle v_s^3 \phi^{(0)} + \gamma^2 \rho_s w^{(0),3} \rangle = i\mathcal{O}(1)$$

as  $\gamma^2 \rightarrow \infty$ .

We next consider  $\lambda^{(2)}$ . Since  $Q_0 L^{(2)}u^{(0)} = 0$ , we have

$$\langle L^{(2)}u^{(0)}, u^{(0)*} \rangle = \langle Q_0 L^{(2)}u^{(0)} \rangle = 0.$$

It then follows that

$$\lambda^{(2)} = \langle L^{(1)}SL^{(1)}u^{(0)}, u^{(0)*} \rangle = \langle Q_0 L^{(1)}SL^{(1)}u^{(0)} \rangle.$$

We define  $\tilde{u}$  by

$$\tilde{u} = SL^{(1)}u^{(0)},$$

which satisfies

$$\begin{cases} L_0 \tilde{u} = (I - \Pi^{(0)})L^{(1)}u^{(0)} = L^{(1)}u^{(0)} + \lambda^{(1)}u^{(0)}, \\ \tilde{w} \mid_{\partial D} = 0, \\ \langle \tilde{\phi} \rangle = 0. \end{cases} \quad (4.9)$$

Note that  $\tilde{u} = {}^T(\tilde{\phi}, \tilde{w}) \in i\mathbf{R}^4$  and  $\lambda^{(1)} \in i\mathbf{R}$ . We rewrite  $\lambda^{(2)}$  as

$$\lambda^{(2)} = \langle Q_0 L^{(1)}\tilde{u} \rangle = \langle iv_s^3 \tilde{\phi} + i\gamma^2 \rho_s \tilde{w}^3 \rangle,$$

where  $\tilde{u} = {}^T(\tilde{\phi}, \tilde{w}) = {}^T(\tilde{\phi}, \tilde{w}', \tilde{w}^3)$ . To show the strict negativity of  $\lambda^{(2)}$ , we estimate  $\tilde{u}$ . The problem (4.9) is written as

$$\begin{cases} \gamma^2 \nabla' \cdot (\rho_s \tilde{w}') = i\xi v_s^3 \phi^{(0)} + i\gamma^2 \rho_s w^{(0),3} + \lambda^{(1)} \phi^{(0)}, \\ -\frac{\nu}{\rho_s} \Delta' \tilde{w}' - \frac{\tilde{\nu}}{\rho_s} \nabla' \nabla' \cdot \tilde{w}' + \nabla' \left( \frac{P'(\rho_s)}{\gamma^2 \rho_s} \tilde{\phi} \right) = -i \frac{\tilde{\nu}}{\rho_s} \nabla' w^{(0),3}, \\ -\frac{\nu}{\rho_s} \Delta' \tilde{w}^3 + \frac{\nu \Delta' v_s^3}{\gamma^2 \rho_s^2} \tilde{\phi} + \tilde{w}' \cdot \nabla' v_s^3 = i \frac{P'(\rho_s)}{\gamma^2 \rho_s} \phi^{(0)} + i v_s^3 w^{(0),3} + \lambda^{(1)} w^{(0),3}, \\ \tilde{w} \mid_{\partial D} = 0, \\ \langle \tilde{\phi} \rangle = 0, \end{cases}$$

i.e.,  $\tilde{u} = {}^T(\tilde{\phi}, \tilde{w}) = {}^T(\tilde{\phi}, \tilde{w}', \tilde{w}^3)$  is a solution of

$$\begin{cases} \nabla' \cdot \tilde{w}' = F^0[\tilde{w}'], \\ -\nu \Delta' \tilde{w}' + \nabla' \tilde{\phi} = G'[\tilde{\phi}, \tilde{w}'], \\ \tilde{w}' \mid_{\partial D} = 0, \\ \langle \tilde{\phi} \rangle = 0 \end{cases} \quad (4.10)$$

and

$$\begin{cases} -\nu \Delta' \tilde{w}^3 = G^3[\tilde{\phi}, \tilde{w}'], \\ \tilde{w}^3 \mid_{\partial D} = 0, \end{cases} \quad (4.11)$$

where  $F^0[\tilde{w}']$ ,  $G'[\tilde{\phi}, \tilde{w}']$  and  $G^3[\tilde{\phi}, \tilde{w}']$  are defined as

$$\begin{aligned} F^0[\tilde{w}'] &= \frac{1}{\gamma^2} \{ i v_s^3 \phi^{(0)} + i \gamma^2 \rho_s w^{(0),3} + \lambda^{(1)} \phi^{(0)} \} - \nabla' \cdot ((1 - \rho_s) \tilde{w}'), \\ G'[\tilde{\phi}, \tilde{w}'] &= -i \tilde{\nu} \nabla' w^{(0),3} + \tilde{\nu} \nabla' F^0[\tilde{w}'] + \nabla' ((1 - \rho_s) \tilde{\phi}) \\ &\quad + (\nabla' \rho_s) \tilde{\phi} + \rho_s \nabla' \left\{ \left( 1 - \frac{P'(\rho_s)}{\gamma^2 \rho_s} \right) \right\} \tilde{\phi}, \\ G^3[\tilde{\phi}, \tilde{w}'] &= \rho_s \left\{ i \frac{P'(\rho_s)}{\gamma^2 \rho_s} \phi^{(0)} + i v_s^3 w^{(0),3} + \lambda^{(1)} w^{(0),3} \right\} - \rho_s \left\{ \frac{\nu \Delta' v_s^3}{\gamma^2 \rho_s^2} \tilde{\phi} + \tilde{w}' \cdot \nabla' v_s^3 \right\}. \end{aligned}$$

As for the problem (4.10), since  $\lambda^{(1)} = -i \langle v_s^3 \phi^{(0)} + \gamma^2 \rho_s w^{(0),3} \rangle$ , it holds that  $\langle F^0[\tilde{w}'] \rangle = 0$ . Furthermore, we have

$$\begin{aligned} |F^0[\tilde{w}']|_2 &\leq C \left\{ \frac{1}{\gamma^2} (|\lambda^{(1)}| |\phi^{(0)}|_2 + |\phi^{(0)}|_2 + \gamma^2 |w^{(0),3}|_2) + \omega |\nabla' \tilde{w}'|_2 \right\} \\ &\leq C \omega |\nabla' \tilde{w}'|_2 + \mathcal{O}\left(\frac{1}{\gamma^2}\right), \end{aligned}$$

$$\begin{aligned} |G'[\tilde{\phi}, \tilde{w}']|_{H^{-1}} &\leq C \left\{ \tilde{\nu} |\nabla' w^{(0),3}|_{H^{-1}} + \tilde{\nu} |\nabla' F^0[\tilde{w}']|_{H^{-1}} + |\nabla' ((1 - \rho_s) \tilde{\phi})|_{H^{-1}} \right. \\ &\quad \left. + |\nabla' \rho_s \tilde{\phi}|_{H^{-1}} + |\rho_s ((1 - \frac{P'(\rho_s)}{\gamma^2 \rho_s}) \tilde{\phi})|_{H^{-1}} \right\} \\ &\leq C \omega \{ |\tilde{\phi}|_2 + \tilde{\nu} |\nabla' \tilde{w}'|_2 \} + \mathcal{O}\left(\frac{\tilde{\nu}}{\gamma^2}\right). \end{aligned}$$

Since  $(\tilde{\phi}, \tilde{w}') \in \dot{X} \equiv \{(p, v') \in L^2(D) \times H_0^1(D) : \langle p \rangle = 0\}$  and it is a solution of the Stokes system (4.10), we see from estimate for the Stokes system (see, e.g., [20]) that there holds the estimate

$$\begin{aligned} |\tilde{\phi}|_2^2 + \nu^2 |\nabla' \tilde{w}'|_2^2 &\leq \nu^2 \{ C \omega^2 |\tilde{w}'|_2^2 + \mathcal{O}\left(\frac{1}{\gamma^4}\right) \} + \{ C \omega^2 (|\tilde{\phi}|_2^2 + \tilde{\nu}^2 |\nabla' \tilde{w}'|_2^2) + \mathcal{O}\left(\frac{\tilde{\nu}^2}{\gamma^4}\right) \} \\ &\leq C_1 \omega^2 \{ |\tilde{\phi}|_2^2 + (\nu + \tilde{\nu})^2 |\nabla' \tilde{w}'|_2^2 \} + \mathcal{O}\left(\frac{(\nu + \tilde{\nu})^2}{\gamma^4}\right). \end{aligned}$$

Therefore, if  $\omega$  is so small that  $\omega^2 < \frac{1}{2C_1} \min\{1, (\frac{\nu}{\nu + \tilde{\nu}})^2\}$ , then

$$|\tilde{\phi}|_2^2 + \nu^2 |\nabla' \tilde{w}'|_2^2 = \mathcal{O}\left(\frac{(\nu + \tilde{\nu})^2}{\gamma^4}\right). \quad (4.12)$$

As for the problem (4.11), since

$$\begin{aligned} |G^3[\tilde{\phi}, \tilde{w}']|_2 &\leq C \left\{ |\lambda^{(1)}| |w^{(0),3}|_2 + \frac{1}{\gamma^2} |\phi^{(0)}|_2 + |w^{(0),3}|_2 + \frac{\nu}{\gamma^2} |\tilde{\phi}|_2 + |\tilde{w}'|_2 \right\} \\ &\leq C \left\{ \frac{\nu}{\gamma^2} |\tilde{\phi}|_2 + |\tilde{w}'|_2 \right\} + \mathcal{O}\left(\frac{1}{\gamma^2}\right), \end{aligned}$$

we have  $G^3[\tilde{\phi}, \tilde{w}'] \in L^2(D)$ . It then follows that

$$\tilde{w}^3 = \frac{1}{\nu} (-\Delta')^{-1} G^3[\tilde{\phi}, \tilde{w}'].$$

Since  $\phi^{(0)} = \alpha_0 \frac{\gamma^2 \rho_s}{P'(\rho_s)}$  (see Lemma 4.1 (ii)), we find that

$$\begin{aligned} \langle \rho_s \tilde{w}^3 \rangle &= \frac{1}{\nu} \langle \rho_s (-\Delta')^{-1} G^3[\tilde{\phi}, \tilde{w}'] \rangle \\ &= \frac{1}{\nu} \langle \rho_s (-\Delta')^{-1} (i \alpha_0 \rho_s) \rangle \\ &\quad + \frac{1}{\nu} \langle \rho_s (-\Delta')^{-1} \{ i \rho_s v_s^3 w^{(0),3} + \rho_s \lambda^{(1)} w^{(0),3} - \frac{\nu \Delta' v_s^3}{\gamma^2 \rho_s} \tilde{\phi} - \rho_s \tilde{w}' \cdot \nabla' v_s^3 \} \rangle \\ &= i \frac{\alpha_0}{\nu} |(-\Delta')^{-\frac{1}{2}} \rho_s|_2^2 \\ &\quad + \frac{1}{\nu} \langle \rho_s (-\Delta')^{-1} \{ i \rho_s v_s^3 w^{(0),3} + \rho_s \lambda^{(1)} w^{(0),3} - \frac{\nu \Delta' v_s^3}{\gamma^2 \rho_s} \tilde{\phi} - \rho_s \tilde{w}' \cdot \nabla' v_s^3 \} \rangle. \end{aligned}$$

Furthermore, since  $\tilde{u} = {}^T(\tilde{\phi}, \tilde{w}') \in i\mathbf{R}^4$  and  $\lambda^{(1)} \in i\mathbf{R}$ , we see from (4.12) that

$$\begin{aligned} & \langle \rho_s (-\Delta')^{-1} \{ i \rho_s v_s^3 w^{(0),3} + \rho_s \lambda^{(1)} w^{(0),3} - \frac{\nu \Delta' v_s^3}{\gamma^2 \rho_s} \tilde{\phi} - \rho_s \tilde{w}' \cdot \nabla' v_s^3 \} \rangle \\ &= i \mathcal{O}\left(\frac{1}{\gamma^2}\right) + i \left(\frac{\nu}{\gamma^2} + \frac{1}{\nu^2}\right) \times \mathcal{O}\left(\frac{\nu+\tilde{\nu}}{\gamma^2}\right). \end{aligned}$$

It then follows that

$$\langle \rho_s \tilde{w}^3 \rangle = i \frac{\alpha_0}{\nu} |(-\Delta')^{-\frac{1}{2}} \rho_s|_2^2 + i \frac{1}{\nu} \left\{ \mathcal{O}\left(\frac{1}{\gamma^2}\right) + \left(\frac{\nu}{\gamma^2} + \frac{1}{\nu^2}\right) \times \mathcal{O}\left(\frac{\nu+\tilde{\nu}}{\gamma^2}\right) \right\}.$$

By (4.12) we also have

$$\langle v_s^3 \tilde{\phi} \rangle = i \mathcal{O}\left(\frac{\nu+\tilde{\nu}}{\gamma^2}\right).$$

We conclude that

$$\begin{aligned} \lambda^{(2)} &= \langle i v_s^3 \tilde{\phi} + i \gamma^2 \rho_s \tilde{w}^3 \rangle \\ &= i \gamma^2 \left[ i \frac{\alpha_0}{\nu} |(-\Delta')^{-\frac{1}{2}} \rho_s|_2^2 + i \frac{1}{\nu} \left\{ \mathcal{O}\left(\frac{1}{\gamma^2}\right) + \left(\frac{\nu}{\gamma^2} + \frac{1}{\nu^2}\right) \times \mathcal{O}\left(\frac{\nu+\tilde{\nu}}{\gamma^2}\right) \right\} \right] + i \cdot i \mathcal{O}\left(\frac{\nu+\tilde{\nu}}{\gamma^2}\right) \\ &= -\frac{\gamma^2}{\nu} \left[ \alpha_0 |(-\Delta')^{-\frac{1}{2}} \rho_s|_2^2 + \left\{ \mathcal{O}\left(\frac{1}{\gamma^2}\right) + \left(\frac{1}{\nu^2} + \frac{\nu}{\gamma^2}\right) \times \mathcal{O}\left(\frac{\nu+\tilde{\nu}}{\gamma^2}\right) \right\} \right] \\ &< 0 \end{aligned}$$

for sufficiently small  $\frac{1}{\nu}$  and  $\frac{\nu+\tilde{\nu}}{\gamma^2}$ . We thus obtain the desired estimates. This completes the proof.  $\square$

We next establish some estimates related to  $\Pi(\xi)$  in  $H^k(D)$ . We first consider estimates for higher order derivatives of  $(\lambda + L_0)^{-1} f$ .

**Proposition 4.6.** *For any  $f = {}^T(f^0, g) \in H^k(D) \times H^{k-1}(D)$ . There exist positive constants  $\nu_0, \gamma_0, \omega_0$  and  $c_1$  such that if  $\nu \geq \nu_0, \frac{\gamma^2}{\nu+\tilde{\nu}} \geq \gamma_0^2, \omega \leq \omega_0$  and  $\lambda \in \Sigma_2 \equiv \{\lambda \neq 0 : |\lambda| \leq c_1\}$ , then  $(\lambda + L_0)^{-1} f \in H^k(D) \times (H^{k+1}(D) \cap H_0^1(D))$  for  $k = 0, 1, \dots, k_0$ . Furthermore, the following estimate holds:*

$$|(\lambda + L_0)^{-1} f|_{H^k \times H^{k+1}} \leq C \left(1 + \frac{1}{|\lambda|}\right) |f|_{H^k \times H^{k-1}},$$

where  $C$  is a positive constant independent of  $\lambda \in \Sigma_2$ . The same assertions also hold for  $-L_0^*$ .

**Proof.** For a given  $f = {}^T(f^0, g) \in H^k(D) \times H^{k-1}(D)$ , we consider the problem

$$\begin{cases} (\lambda + \mathcal{L}_0)U = f, \\ W|_{\partial D} = 0 \end{cases} \quad (4.13)$$

for  $U = {}^T(\Phi, W)$ . Here  $\mathcal{L}_0$  is differential operator given by

$$\mathcal{L}_0 U = \begin{pmatrix} \gamma^2 \nabla' \cdot (\rho_s W') \\ -\frac{\nu}{\rho_s} \Delta' W' - \frac{\tilde{\nu}}{\rho_s} \nabla' \nabla' \cdot W' + \nabla' \left( \frac{P'(\rho_s)}{\gamma^2 \rho_s} \Phi \right) \\ -\frac{\nu}{\rho_s} \Delta' W^3 + \frac{\nu \Delta' v_s^3}{\gamma^2 \rho_s^2} \Phi + W' \cdot \nabla' v_s^3 \end{pmatrix}$$

for  $U = {}^T(\Phi, W)$ . To solve the problem (4.13), we decompose  $\Phi$  and  $f^0$  as

$$\Phi = \Phi_1 + \sigma, \quad f^0 = f_1^0 + \langle f^0 \rangle,$$

where  $\sigma = \langle \Phi \rangle$ ,  $\Phi_1 = \Phi - \sigma$  and  $f_1^0 = f^0 - \langle f^0 \rangle$ . Note that

$$\langle \Phi_1 \rangle = 0, \quad \langle f_1^0 \rangle = 0.$$

Then (4.13) is equivalent to the problem

$$\lambda \sigma = \langle f^0 \rangle, \quad (4.14)$$

$$\lambda \Phi_1 + \gamma^2 \nabla' \cdot (\rho_s W') = f_1^0, \quad (4.15)$$

$$\lambda W' - \frac{\nu}{\rho_s} \Delta' W' - \frac{\tilde{\nu}}{\rho_s} \nabla' \nabla' \cdot W' + \nabla' \left( \frac{P'(\rho_s)}{\gamma^2 \rho_s} (\sigma + \Phi_1) \right) = g', \quad (4.16)$$

$$\lambda W^3 - \frac{\nu}{\rho_s} \Delta' W^3 + \frac{\nu \Delta' v_s^3}{\gamma^2 \rho_s^2} (\sigma + \Phi_1) - W' \cdot \nabla' v_s^3 = g^3 \quad (4.17)$$

with  $W|_{\partial D} = 0$ . If  $\lambda \neq 0$ , then we find from (4.14) that

$$\sigma = \frac{1}{\lambda} \langle f^0 \rangle. \quad (4.18)$$

Substituting  $\sigma = \frac{1}{\lambda} \langle f^0 \rangle$  into (4.16) and (4.17), we obtain

$$\begin{cases} \lambda \Phi_1 + \gamma^2 \nabla' \cdot (\rho_s W') = f_1^0, \\ \lambda W' - \frac{\nu}{\rho_s} \Delta' W' - \frac{\tilde{\nu}}{\rho_s} \nabla' \nabla' \cdot W' + \nabla' \left( \frac{P'(\rho_s)}{\gamma^2 \rho_s} \Phi_1 \right) = g' - \frac{1}{\lambda} \langle f^0 \rangle \nabla' \left( \frac{P'(\rho_s)}{\gamma^2 \rho_s} \right), \\ \lambda W^3 - \frac{\nu}{\rho_s} \Delta' W^3 + \frac{\nu \Delta' v_s^3}{\gamma^2 \rho_s^2} \Phi_1 - W' \cdot \nabla' v_s^3 = g^3 - \frac{1}{\lambda} \langle f^0 \rangle \frac{\nu \Delta' v_s^3}{\gamma^2 \rho_s^2} \end{cases} \quad (4.19)$$

with  $W|_{\partial D} = 0$ . Let us write the problem (4.19) as

$$\begin{cases} \nabla' \cdot W' = F^0[\Phi_1, W' : f_1^0], \\ -\nu \Delta' W' + \nabla' \Phi_1 = G'[\Phi_1, W' : f^0, g'], \\ W'|_{\partial D} = 0 \end{cases} \quad (4.20)$$

and

$$\begin{cases} -\nu \Delta' W^3 = G^3[\Phi_1, W', W^3 : f^0, g^3], \\ W^3|_{\partial D} = 0. \end{cases} \quad (4.21)$$

Here

$$\begin{aligned} F^0[\Phi_1, W' : f_1^0] &= -\frac{1}{\gamma^2} \lambda \Phi_1 + \nabla' \cdot ((1 - \rho_s) W') + \frac{1}{\gamma^2} f_1^0, \\ G'[\Phi_1, W' : f^0, g'] &= -\lambda \rho_s W' + \tilde{\nu} \nabla' F^0[\Phi_1, W' : f_1^0] + \nabla' ((1 - \rho_s) \Phi_1) + \nabla' \rho_s \Phi_1 \\ &\quad - \frac{1}{\lambda} \langle f^0 \rangle \rho_s \nabla' \left( \frac{P'(\rho_s)}{\gamma^2 \rho_s} \right) + \rho_s \nabla' \left( \left( 1 - \frac{P'(\rho_s)}{\gamma^2 \rho_s} \right) \Phi_1 \right) + \rho_s g', \\ G^3[\Phi_1, W', W^3 : f^0, g^3] &= -\lambda \rho_s W^3 - \frac{\nu \Delta' v_s^3}{\gamma^2 \rho_s^2} \frac{1}{\lambda} \langle f^0 \rangle - \frac{\nu \Delta' v_s^3}{\gamma^2 \rho_s^2} \Phi_1 - \rho_s W' \cdot \nabla' v_s^3 + \rho_s g^3. \end{aligned}$$

We now define a set  $\dot{X}_k$  by

$$\dot{X}_k = \{(p, v') \in H^k(D) \times (H^{k+1}(D) \cap H_0^1(D)) : \langle p \rangle = 0\}$$

with norm

$$|(p, v')|_{\dot{X}_k} = |p|_{H^k} + \nu|v'|_{H^{k+1}}.$$

For a given  $(\tilde{\Phi}_1, \tilde{W}') \in \dot{X}_k$ , we consider the problem

$$\begin{cases} \nabla' \cdot W' = F^0[\tilde{\Phi}_1, \tilde{W}' : f_1^0], \\ -\nu \Delta' W' + \nabla' \Phi_1 = G'[\tilde{\Phi}_1, \tilde{W}' : f^0, g'], \\ W' |_{\partial D} = 0. \end{cases} \quad (4.22)$$

It holds that

$$\langle F^0[\tilde{\Phi}_1, \tilde{W}' : f_1^0] \rangle = 0, \quad F^0[\tilde{\Phi}_1, \tilde{W}' : f_1^0] \in H^k(D),$$

$$G'[\tilde{\Phi}_1, \tilde{W}' : f^0, g'] \in H^{k-1}(D).$$

In fact, we see that

$$\langle F^0[\tilde{\Phi}_1, \tilde{W}' : f_1^0] \rangle = -\frac{1}{\gamma^2} \lambda \langle \tilde{\Phi}_1 \rangle + \langle \nabla' \cdot ((1 - \rho_s) \tilde{W}') \rangle + \frac{1}{\gamma^2} \langle f_1^0 \rangle = 0,$$

$$|F^0[\tilde{\Phi}_1, \tilde{W}' : f_1^0]|_{H^k} \leq C \left\{ \frac{1}{\gamma^2} |\lambda| |\tilde{\Phi}_1|_{H^k} + \omega |\tilde{W}'|_{H^{k+1}} + \frac{1}{\gamma^2} |f_1^0|_{H^k} \right\}$$

and

$$\begin{aligned} & |G'[\tilde{\Phi}_1, \tilde{W}' : f^0, g']|_{H^{k-1}} \\ & \leq C \left\{ |\lambda| |\tilde{W}'|_{H^{k-1}} + \tilde{\nu} |F^0[\tilde{\Phi}_1, \tilde{W}' : f_1^0]|_{H^k} + \omega |\tilde{\Phi}_1|_{H^k} + \frac{1}{|\lambda|} |\langle f^0 \rangle| + |g'|_{H^{k-1}} \right\} \\ & \leq C \left\{ \left( \frac{\tilde{\nu}}{\gamma^2} |\lambda| + \omega \right) |\tilde{\Phi}_1|_{H^k} + \nu \left( \frac{1}{\nu} |\lambda| + \frac{\tilde{\nu}}{\nu} \omega \right) |\tilde{W}'|_{H^{k+1}} + \left( \frac{\tilde{\nu}}{\gamma^2} + \frac{1}{|\lambda|} \right) |f^0|_{H^k} + |g'|_{H^{k-1}} \right\} \end{aligned}$$

for a positive constant  $C$  independent of  $\lambda$ . From [20], we see that there is a unique solution  $(\Phi_1, W') \in \dot{X}_k$  of (4.22) and there holds the estimate

$$\begin{aligned} & |\Phi|_{H^k} + \nu |W'|_{H^{k+1}} \\ & \leq C \left\{ \nu |F^0[\tilde{\Phi}_1, \tilde{W}' : f_1^0]|_{H^k} + |G'[\tilde{\Phi}_1, \tilde{W}' : f^0, g']|_{H^{k-1}} \right\} \\ & \leq C \left\{ \left( \frac{\nu + \tilde{\nu}}{\gamma^2} |\lambda| + \omega \right) |\tilde{\Phi}_1|_{H^k} + \nu \left( \frac{1}{\nu} |\lambda| + \frac{\nu + \tilde{\nu}}{\nu} \omega \right) |\tilde{W}'|_{H^{k+1}} \right. \\ & \quad \left. + \left( \frac{\nu + \tilde{\nu}}{\gamma^2} + \frac{1}{|\lambda|} \right) |f^0|_{H^k} + |g'|_{H^{k-1}} \right\} \end{aligned} \quad (4.23)$$

for a positive constant  $C$  independent of  $\lambda$ . Let us define a map  $\Gamma_1 : \dot{X}_k \rightarrow \dot{X}_k$  such that

$$\Gamma_1(\tilde{\Phi}_1, \tilde{W}') = (\Phi_1, W'),$$

where  $(\Phi_1, W') \in \dot{X}_k$  is a solution of (4.22). From (4.23), for  $(\tilde{\Phi}_{1,1}, \tilde{W}'_1), (\tilde{\Phi}_{1,2}, \tilde{W}'_2) \in \dot{X}_k$ , the estimate

$$\begin{aligned} & |\Gamma_1(\tilde{\Phi}_{1,1}, \tilde{W}'_1) - \Gamma_1(\tilde{\Phi}_{1,2}, \tilde{W}'_2)|_{H^k \times H^{k+1}} \\ & \leq C_1 \left\{ \left( \frac{\nu + \tilde{\nu}}{\gamma^2} + \frac{1}{\nu} \right) |\lambda| + \left( \frac{\nu + \tilde{\nu}}{\nu} + 1 \right) \omega \right\} |(\tilde{\Phi}_{1,1} - \tilde{\Phi}_{1,2}, \tilde{W}'_1 - \tilde{W}'_2)|_{\dot{X}_k} \end{aligned}$$

holds for a positive constant  $C_1$  independent of  $\lambda$ . If  $\omega$  and  $|\lambda|$  are so small that  $\omega < \frac{1}{2C_1} \frac{\nu}{\nu+\bar{\nu}}$  and  $|\lambda| < \frac{1}{2C_1}$ , then  $\Gamma_1 : \dot{X}_k \rightarrow \dot{X}_k$  is a contraction map. This implies that there is a unique  $(\Phi_1, W') \in \dot{X}_k$  such that  $\Gamma_1(\Phi_1, W') = (\Phi_1, W')$ , i.e., there is a unique solution  $(\Phi_1, W') \in \dot{X}_k$  of (4.20). Furthermore, from (4.23),  $(\Phi_1, W')$  satisfies the estimate

$$|\Phi_1|_{H^k} + |W'|_{H^{k+1}} \leq C \left\{ \left(1 + \frac{1}{|\lambda|}\right) |f^0|_{H^k} + |g'|_{H^{k-1}} \right\}, \quad (4.24)$$

where  $C$  is a positive constant independent of  $\lambda$ .

As for (4.21), for a given  $\widetilde{W}^3 \in H^{k+1}(D) \cap H_0^1(D)$ , we consider the problem

$$\begin{cases} -\nu \Delta' W^3 = G^3[\Phi_1, W', \widetilde{W}^3 : f^0, g^3], \\ W^3|_{\partial D} = 0, \end{cases} \quad (4.25)$$

where  $(\Phi_1, W') \in \dot{X}_k$  is a solution of (4.20). It holds that

$$G^3[\Phi_1, W', \widetilde{W}^3 : f^0, g^3] \in H^{k-1}(D).$$

In fact, we have

$$\begin{aligned} & |G^3[\Phi_1, W', \widetilde{W}^3 : f^0, g^3]|_{H^{k-1}} \\ & \leq C \left\{ |\lambda| |\widetilde{W}^3|_{H^{k-1}} + |\Phi_1|_{H^{k-1}} + |W'|_{H^{k-1}} + |g^3|_{H^{k-1}} + \frac{1}{|\lambda|} |\langle f^0 \rangle| \right\} \\ & \leq C_2 \left\{ |\lambda| |\widetilde{W}^3|_{H^{k-1}} + \left(1 + \frac{1}{|\lambda|}\right) |f^0|_{H^k} + |g|_{H^{k-1}} \right\} \end{aligned} \quad (4.26)$$

for a positive constant  $C_2$  independent of  $\lambda$ . If  $|\lambda|$  is sufficiently small satisfying  $|\lambda| < \min\{\frac{1}{2C_1}, \frac{1}{C_2}\}$ , then there is a unique solution  $W^3 \in H^{k+1}(D) \cap H_0^1(D)$  of (4.21). Furthermore, from (4.26),  $W^3$  satisfies the estimate

$$|W^3|_{H^{k+1}} \leq C \left\{ \left(1 + \frac{1}{|\lambda|}\right) |f^0|_{H^k} + |g|_{H^{k-1}} \right\}, \quad (4.27)$$

where  $C$  is a positive constant independent of  $\lambda$ .

Now we set

$$\Sigma_2 \equiv \left\{ \lambda \neq 0 : |\lambda| < \min\left\{ \frac{1}{2C_1}, \frac{1}{C_2} \right\} \right\}.$$

Since  $\Phi = \sigma + \Phi_1$ , we see that if  $\omega < \frac{1}{2C_1} \frac{\nu}{\nu+\bar{\nu}}$  and  $\lambda \in \Sigma_2$ , then there is a unique solution  $(\Phi, W) \in H^k(D) \times (H^{k+1}(D) \cap H_0^1(D))$  of (4.13). Moreover, from (4.18), (4.24) and (4.27),  $\Phi$  and  $W$  satisfies the estimate

$$\begin{aligned} |\Phi|_{H^k} + |W|_{H^{k+1}} & \leq |\sigma| + |\Phi_1|_{H^k} + |W'|_{H^{k+1}} + |W^3|_{H^{k+1}} \\ & \leq C \left\{ \left(1 + \frac{1}{|\lambda|}\right) |f^0|_{H^k} + |g|_{H^{k-1}} \right\} \end{aligned}$$

for a positive constant  $C$  independent of  $\lambda \in \Sigma_2$ .

Since  $D(L_0) \supset H^k(D) \times (H^{k+1}(D) \cap H_0^1(D))$ , we can replace  $\mathcal{L}_0$  with  $L_0$ ; and we find that if  $\omega < \frac{1}{2C_1} \frac{\nu}{\nu+\bar{\nu}}$  and  $\lambda \in \Sigma_2$ , then  $(\lambda + L_0)^{-1} f \in H^k(D) \times (H^{k+1}(D) \cap H_0^1(D))$ . Furthermore,  $(\lambda + L_0)^{-1} f$  satisfies the estimate

$$|(\lambda + L_0)^{-1} f|_{H^k \times H^{k+1}} \leq C \left\{ \left(1 + \frac{1}{|\lambda|}\right) |f^0|_{H^k} + |g|_{H^{k-1}} \right\},$$



where  $C$  is a positive constant independent of  $\lambda \in \Sigma_2$ . We thus obtain the desired estimates. The assertions for  $L_0^*$  can be proved in a similar manner. This completes the proof.  $\square$

We finally obtain the following estimates for the eigenfunctions  $u_\xi$  and  $u_\xi^*$  associated with  $\lambda_0(\xi)$  and  $\bar{\lambda}_0(\xi)$ , respectively, which yields the boundedness of  $\Pi(\xi)$  on  $H^k(D)$ .

**Theorem 4.7.** *There exist positive constants  $\nu_0$ ,  $\gamma_0$  and  $\omega_0$  such that if  $\nu \geq \nu_0$ ,  $\frac{\gamma^2}{\nu+\bar{\nu}} \geq \gamma_0^2$  and  $\omega \leq \omega_0$ , then there exists a positive constant  $r_0$  such that for any  $\xi \in \mathbf{R}$  with  $|\xi| \leq r_0$  the following assertions hold. There exist  $u_\xi$  and  $u_\xi^*$  eigenfunctions associated with  $\lambda_0(\xi)$  and  $\bar{\lambda}_0(\xi)$ , respectively, that satisfy*

$$\langle u_\xi, u_\xi^* \rangle = 1,$$

and the eigenprojection  $\Pi(\xi)$  for  $\lambda_0(\xi)$  is given by

$$\Pi(\xi)u = \langle u, u_\xi^* \rangle u_\xi.$$

Furthermore,  $u_\xi$  and  $u_\xi^*$  are written in the form

$$\begin{aligned} u_\xi(x') &= u^{(0)}(x') + i\xi u^{(1)}(x') + |\xi|^2 u^{(2)}(x', \xi), \\ u_\xi^*(x') &= u^{*(0)}(x') + i\xi u^{*(1)}(x') + |\xi|^2 u^{*(2)}(x', \xi), \end{aligned}$$

and the following estimates hold

$$|u|_{H^{k+2}} \leq C_{k,r_0}$$

for  $u \in \{u_\xi, u_\xi^*, u^{(1)}, u^{*(1)}, u^{(2)}, u^{*(2)}\}$  and  $k = 0, 1, \dots, k_0$ : and a positive constant  $C_{k,r_0}$  depending on  $k$  and  $r_0$ .

We can prove Theorem 4.7 by using Proposition 4.6, similarly to the proof of [9, Lemma 4.3]. We thus omit the proof.

## 5 Spectral properties of $e^{-tL}P_0$

In this section we give a factorization of  $e^{-tL}P_0$  and prove Theorem 3.1 (i).

We denote the characteristic function of a set  $\{\xi \in \mathbf{R} : |\xi| \leq r_0\}$  by  $\mathbf{1}_{\{|\xi| \leq r_0\}}$ , i.e.,

$$\mathbf{1}_{\{|\eta| \leq r_0\}}(\xi) = \begin{cases} 1, & |\xi| \leq r_0, \\ 0, & |\xi| > r_0. \end{cases}$$

We define the projection  $P_0$  by

$$P_0 = \mathcal{F}^{-1} \mathbf{1}_{\{|\xi| \leq r_0\}} \Pi(\xi) \mathcal{F}.$$

$P_0$  is a bounded projection on  $L^2(\Omega)$  satisfying

$$P_0 L \subset L P_0, \quad P_0 e^{-tL} = e^{-tL} P_0.$$

As in [2, 4], to investigate  $e^{-tL} P_0$ , we introduce operators related to  $u_\xi$  and  $u_\xi^*$ . We define the operators  $\mathcal{T} : L^2(\mathbf{R}) \rightarrow L^2(\Omega)$ ,  $\mathcal{P} : L^2(\Omega) \rightarrow L^2(\mathbf{R})$  and  $\Lambda : L^2(\mathbf{R}) \rightarrow L^2(\mathbf{R})$  by

$$\begin{aligned} \mathcal{T}\sigma &= \mathcal{F}^{-1}[\mathcal{T}_\xi \sigma], & \mathcal{T}_\xi \sigma &= \mathbf{1}_{\{|\xi| \leq r_0\}} u_\xi \sigma; \\ \mathcal{P}u &= \mathcal{F}^{-1}[\mathcal{P}_\xi u], & \mathcal{P}_\xi u &= \mathbf{1}_{\{|\xi| \leq r_0\}} \langle u, u_\xi^* \rangle; \\ \Lambda\sigma &= \mathcal{F}^{-1}[\mathbf{1}_{\{|\xi| \leq r_0\}} \lambda_0(\xi) \sigma] \end{aligned}$$

for  $u \in L^2(\Omega)$  and  $\sigma \in L^2(\mathbf{R})$ . It then follows that

$$P_0 = \mathcal{T}\mathcal{P}, \quad e^{-tL} P_0 = \mathcal{T} e^{t\Lambda} \mathcal{P}.$$

We investigate boundedness properties of  $\mathcal{T}$ ,  $\mathcal{P}$  and  $e^{t\Lambda}$ .

As for  $\mathcal{T}$ , we have the following

**Proposition 5.1.** *The operator  $\mathcal{T}$  has the following properties:*

- (i)  $\partial_{x_3}^l \mathcal{T} = \mathcal{T} \partial_{x_3}^l$  for  $l = 0, 1, \dots$ .
- (ii)  $\|\partial_{x_3}^k \partial_{x_3}^l \mathcal{T} \sigma\|_2 \leq C \|\sigma\|_{L^2(\mathbf{R})}$  for  $k = 0, 1, \dots, k_0$ ,  $l = 0, 1, \dots$  and  $\sigma \in L^2(\mathbf{R})$ .
- (iii)  $\mathcal{T}$  is decomposed as

$$\mathcal{T} = \mathcal{T}^{(0)} + \partial_{x_3} \mathcal{T}^{(1)} + \partial_{x_3}^2 \mathcal{T}^{(2)}.$$

Here  $\mathcal{T}^{(j)} \sigma = \mathcal{F}^{-1}[\mathcal{T}^{(j)} \sigma]$  ( $j = 0, 1, 2$ ) with

$$\begin{aligned} \mathcal{T}^{(0)} \sigma &= \mathbf{1}_{\{|\xi| \leq r_0\}} \sigma u^{(0)}, \\ \mathcal{T}^{(1)} \sigma &= \mathbf{1}_{\{|\xi| \leq r_0\}} \sigma u^{(1)}(\cdot), \\ \mathcal{T}^{(2)} \sigma &= -\mathbf{1}_{\{|\xi| \leq r_0\}} \sigma u^{(2)}(\cdot, \xi), \end{aligned}$$

where  $u^{(j)}$  ( $j = 0, 1, 2$ ) are the functions given in Theorem 4.7. The assertions (i) and (ii) hold with  $\mathcal{T}$  replaced by  $\mathcal{T}^{(j)}$  ( $j = 0, 1, 2$ ).

**Proof.** It is clear that (i) is true. As for (ii), we can prove the estimates by using the properties of  $u_\xi$  in Theorem 4.7 and the Sobolev inequality. From the expansion of  $u_\xi$  given in Theorem 4.7, we can expand  $\mathcal{T}$  as in (iii).  $\square$

As for  $\mathcal{P}$ , we have the following properties.

**Proposition 5.2.** *The operator  $\mathcal{P}$  has the following properties:*

- (i)  $\partial_{x_3}^l \mathcal{P} = \mathcal{P} \partial_{x_3}^l$  for  $l = 0, 1, \dots$ .
- (ii)  $\|\partial_{x_3}^k \mathcal{P} u\|_{L^2(\mathbf{R})} \leq C \|u\|_2$  for  $k = 0, 1, \dots, k_0$ ,  $l = 0, 1, \dots$  and  $u \in L^2(\Omega)$ .  
Furthermore,  $\|\mathcal{P} u\|_{L^2(\mathbf{R})} \leq C \|u\|_1$  for  $u \in L^1(\Omega)$ .
- (iii)  $\mathcal{P}$  is decomposed as

$$\mathcal{P} = \mathcal{P}^{(0)} + \partial_{x_3} \mathcal{P}^{(1)} + \partial_{x_3}^2 \mathcal{P}^{(2)}.$$

Here  $\mathcal{P}^{(j)}u = \mathcal{F}^{-1}[\mathcal{P}^{(j)}u]$  ( $j = 0, 1, 2$ ) with

$$\begin{aligned}\mathcal{P}^{(0)}u &= \mathbf{1}_{\{|\xi| \leq r_0\}} \langle u, u^{*(0)} \rangle = \mathbf{1}_{\{|\xi| \leq r_0\}} \langle Q_0 u \rangle, \\ \mathcal{P}^{(1)}u &= \mathbf{1}_{\{|\xi| \leq r_0\}} \langle u, u^{*(1)} \rangle, \\ \mathcal{P}^{(2)}u &= -\mathbf{1}_{\{|\xi| \leq r_0\}} \langle u, u^{*(2)}(\xi) \rangle,\end{aligned}$$

where  $u^{(j)*}$  ( $j = 0, 1, 2$ ) are the functions given in Theorem 4.7. The assertions (i) and (ii) hold with  $\mathcal{P}$  replaced by  $\mathcal{P}^{(j)}$  ( $j = 0, 1, 2$ ).

**Proof.** It is clear that (i) holds true. As for (ii), we can prove the estimates by using the properties of  $u_\xi^*$  in Theorem 4.7 and the Sobolev inequality. From the expansion of  $u_\xi^*$  given in Theorem 4.7, we can expand  $\mathcal{P}$  as in (iii).  $\square$

As for  $\Lambda$ , we have the following decay estimates for  $e^{t\Lambda}$ .

**Proposition 5.3.** *The operator  $e^{t\Lambda}$  satisfies the following decay estimates.*

- (i)  $\|\partial_{x_3}^l e^{t\Lambda} \mathcal{P}u\|_{L^2(\mathbf{R})} \leq C(1+t)^{-\frac{1}{4}-\frac{l}{2}} \|u\|_1,$
  - (ii)  $\|\partial_{x_3}^l e^{t\Lambda} \mathcal{P}^{(j)}u\|_{L^2(\mathbf{R})} \leq C(1+t)^{-\frac{1}{4}-\frac{l}{2}} \|u\|_1, \quad j = 0, 1, 2,$
  - (iii)  $\|\partial_{x_3}^l (\mathcal{T} - \mathcal{T}^{(0)}) e^{t\Lambda} \mathcal{P}u\|_2 \leq C(1+t)^{-\frac{3}{4}-\frac{l}{2}} \|u\|_1,$
- for  $u \in L^1(\Omega)$  and  $l = 0, 1, 2, \dots$ .

**Proof.** Since  $\lambda_0(\xi) = -i\kappa_0\xi - \kappa_1\xi^2 + \mathcal{O}(|\xi|^3)$ , we see from Theorem 4.7 that

$$\begin{aligned}\|\partial_{x_3}^l e^{t\Lambda} \mathcal{P}^{(j)}u\|_{L^2(\mathbf{R})} &\leq C \int_{\mathbf{R}} \mathbf{1}_{\{|\xi| \leq r_0\}} |\xi|^{2l} e^{-t(i\kappa_0\xi + \kappa_1\xi^2)} |\langle u(\xi), u^{*(j)} \rangle|^2 d\xi \\ &\leq C \int_{\mathbf{R}} \mathbf{1}_{\{|\xi| \leq r_0\}} |\xi|^{2l} e^{-t(i\kappa_0\xi + \kappa_1\xi^2)} |u(\xi)|_1^2 d\xi \\ &\leq C \begin{cases} \|u\|_1^2, \\ t^{-\frac{1}{2}-l} \|u\|_1^2. \end{cases}\end{aligned}\tag{5.1}$$

This implies (i) and (ii). As for (iii), since  $\mathcal{T} - \mathcal{T}^{(0)} = \partial_{x_3} \mathcal{T}^{(1)} + \partial_{x_3}^2 \mathcal{T}^{(2)}$ , we obtain the desired estimate from (i) and Proposition 5.1.  $\square$

The estimate (3.4) in Theorem 3.1 follows from Propositions 5.1 and 5.3.

We next investigate the asymptotic behavior of  $e^{-tL}$ . Recall that  $\mathcal{H}(t)$  is defined by

$$\mathcal{H}(t)\sigma = \mathcal{F}^{-1}[e^{-(i\kappa_0\xi + \kappa_1\xi^2)t}\sigma]$$

for  $\sigma \in L^2(\mathbf{R})$ , where  $\kappa_0 \in \mathbf{R}$  and  $\kappa_1 > 0$  are given in Theorem 4.5. We first introduce the well-known decay estimate for  $\mathcal{H}(t)$ .

**Proposition 5.4.** *There holds the estimate*

$$\|\partial_{x_3}^l (\mathcal{H}(t)\sigma)\|_{L^2(\mathbf{R})} \leq Ct^{-\frac{1}{4}-\frac{l}{2}} \|\sigma\|_{L^1(\mathbf{R})} \quad (l = 0, 1, \dots)$$

for  $\sigma \in L^1(\mathbf{R})$ .

We next consider the asymptotic behavior of  $e^{t\Lambda}$ . The asymptotic leading part of  $e^{t\Lambda} \mathcal{P}$  is given by  $\mathcal{H}(t)$ . In fact, we have the following

**Proposition 5.5.** For  $u \in L^2(\Omega)$ , we set  $\sigma = \langle Q_0 u \rangle$ . If  $u \in L^1(\Omega)$ , then there holds the estimate

$$\|\partial_{x_3}^l (e^{t\Lambda} \mathcal{P}u - \mathcal{H}(t)\sigma)\|_{L^2(\mathbf{R})} \leq Ct^{-\frac{3}{4}-\frac{l}{2}} \|u\|_1 \quad (l = 0, 1, \dots).$$

**Proof.** By Proposition 5.2 we have

$$e^{t\Lambda} \mathcal{P} = e^{t\Lambda} \mathcal{P}^{(0)} + \partial_{x_3} e^{t\Lambda} \mathcal{P}^{(1)} + \partial_{x_3}^2 e^{t\Lambda} \mathcal{P}^{(2)}.$$

Set  $\sigma = \langle Q_0 u \rangle$ . Since  $e^{t\Lambda} \mathcal{P}^{(0)} u = \mathcal{F}^{-1}[\mathbf{1}_{\{|\xi| \leq r_0\}} e^{\lambda_0(\xi)t} \sigma]$ , we see that

$$\mathcal{F}[e^{t\Lambda} \mathcal{P}^{(0)} u - \mathcal{H}(t)\sigma] = (\mathbf{1}_{\{|\xi| \leq r_0\}} - 1) e^{-(i\kappa_0\xi + \kappa_1\xi^2)t} \sigma + \mathbf{1}_{\{|\xi| \leq r_0\}} (e^{\lambda_0(\xi)t} - e^{-(i\kappa_0\xi + \kappa_1\xi^2)t}) \sigma.$$

By using the relation

$$\lambda_0(\xi) + (i\kappa_0\xi + \kappa_1\xi^2) = \mathcal{O}(|\xi|^3)$$

we obtain

$$\begin{aligned} e^{\lambda_0(\xi)t} - e^{-(i\kappa_0\xi + \kappa_1\xi^2)t} &= e^{-(i\kappa_0\xi + \kappa_1\xi^2)t} (e^{(\lambda_0(\xi) + i\kappa_0\xi + \kappa_1\xi^2)t} - 1) \\ &= e^{-(i\kappa_0\xi + \kappa_1\xi^2)t} \mathcal{O}(|\xi|^3)t. \end{aligned}$$

It then follows that

$$\begin{aligned} \int_{|\xi| \leq r_0} |\xi|^{2l} |e^{\lambda_0(\xi)t} - e^{-(i\kappa_0\xi + \kappa_1\xi^2)t}|^2 d\xi &\leq C \int_{|\xi| \leq r_0} |\xi|^{2(l+3)} t^2 e^{-2\kappa_1\xi^2 t} d\xi \|\sigma\|_{L^1(\mathbf{R})}^2 \\ &\leq C \int_{|\xi| \leq r_0} (|\xi|^2 t)^2 e^{-\kappa_1\xi^2 t} |\xi|^{2(l+1)} e^{-\kappa_1\xi^2 t} d\xi \|\sigma\|_{L^1(\mathbf{R})}^2 \\ &\leq C \int_{|\xi| \leq r_0} |\xi|^{2(l+1)} e^{-\kappa_1\xi^2 t} d\xi \|\sigma\|_{L^1(\mathbf{R})}^2 \\ &\leq Ct^{-\frac{3}{2}-l} \|\sigma\|_{L^1(\mathbf{R})}^2. \end{aligned}$$

On the other hand, we also have

$$\int_{|\xi| \leq r_0} |\xi|^{2l} |e^{\lambda_0(\xi)t} - e^{-(i\kappa_0\xi + \kappa_1\xi^2)t}|^2 d\xi \leq C \|\sigma\|_{L^1(\mathbf{R})}^2.$$

We thus obtain

$$\int_{|\xi| \leq r_0} |\xi|^{2l} |e^{\lambda_0(\xi)t} - e^{-(i\kappa_0\xi + \kappa_1\xi^2)t}|^2 d\xi \leq C(1+t)^{-\frac{3}{2}-l} \|\sigma\|_{L^1(\mathbf{R})}^2.$$

Similarly, we have

$$\|(\mathbf{1}_{\{|\xi| \leq r_0\}} - 1) e^{-(i\kappa_0\xi + \kappa_1\xi^2)t} \sigma\|_2^2 \leq Ct^{-\frac{1}{2}-l} e^{-\kappa_1 r_0^2 t} \|\sigma\|_{L^1(\mathbf{R})}^2.$$

We thus see that

$$\|e^{t\Lambda} \mathcal{P}^{(0)} u - \mathcal{H}(t)\sigma\|_{L^2(\mathbf{R})} \leq Ct^{-\frac{3}{4}-\frac{l}{2}} \|u\|_1.$$

This estimate and Proposition 5.3 (ii) give the desired estimate. This completes the proof.  $\square$

We are now in a position to prove estimate (3.5) in Theorem 3.1 (i). In fact, we have

$$e^{-tL}P_0u - [\mathcal{H}(t)\sigma]u^{(0)} = (\mathcal{T} - \mathcal{T}^{(0)})e^{t\Lambda}\mathcal{P}u + [e^{t\Lambda}\mathcal{P}u - \mathcal{H}(t)\sigma]u^{(0)}.$$

This, together with Proposition 5.3 (iii) and Proposition 5.5, yields the desired estimate (3.5).

We finally state the estimates for the projection  $P_0$ .

**Theorem 5.6.** *The projection  $P_0$  has the following properties:*

- (i)  $\partial_{x_3}^l P_0 = P_0 \partial_{x_3}^l$  for  $l = 0, 1, \dots$ .
- (ii)  $\|\partial_{x'}^k \partial_{x_3}^l P_0 u\|_2 \leq C_k \|u\|_1$  for  $k = 0, 1, \dots, k_0$ ,  $l = 0, 1, \dots$  and  $u \in L^1(\Omega)$ .
- (iii)  $P_0$  is decomposed as

$$P_0 = P_0^{(0)} + \partial_{x_3} P_0^{(1)} + \partial_{x_3}^2 P_0^{(2)},$$

where  $P_0^{(j)}u = \mathcal{F}^{-1}[P_0^{(j)}u]$  ( $j = 0, 1, 2$ ) with

$$P_0^{(0)} = \mathcal{T}^{(0)}\mathcal{P}^{(0)} = \mathbf{1}_{\{|\xi| \leq r_0\}}\Pi^{(0)}, \quad (5.2)$$

$$P_0^{(1)} = \mathcal{T}^{(0)}\mathcal{P}^{(1)} + \mathcal{T}^{(1)}\mathcal{P}^{(0)} = -i\mathbf{1}_{\{|\xi| \leq r_0\}}\Pi^{(1)}, \quad (5.3)$$

$$P_0^{(2)} = \mathcal{T}^{(0)}\mathcal{P}^{(2)} + \mathcal{T}^{(1)}\{\mathcal{P}^{(1)} + \partial_{x_3}\mathcal{P}^{(2)}\} + \mathcal{T}^{(2)}\{\mathcal{P}^{(0)} + \partial_{x_3}\mathcal{P}^{(1)} + \partial_{x_3}^2\mathcal{P}^{(2)}\}. \quad (5.4)$$

Furthermore,  $P_0^{(j)}$  ( $j = 0, 1, 2$ ) satisfy assertions (i) and (ii) by replacing  $P_0$  with  $P_0^{(j)}$ .

**Proof.** It is clear that (i) is true. Estimates in (ii) are given by Propositions 5.1, 5.2. As for (iii), it is easy to see that  $\partial_{x_3}^l P_0^{(j)} = P_0^{(j)} \partial_{x_3}^l$ . The estimates

$$\|\partial_{x'}^k \partial_{x_3}^l P_0^{(j)}u\|_2 \leq C_k \|u\|_1$$

can also be obtained by Propositions 5.1, 5.2. The relations (5.3) and (5.4) can be verified by equating the coefficients of each power of  $\xi$  in the expansions of  $\Pi(\xi)$  in (4.8) and  $\langle u, u_\xi^* \rangle u_\xi$ . This completes the proof.  $\square$

## 6 Decay estimate for $e^{-tL}(I - P_0)$

In this section we prove Theorem 3.1 (ii). We set

$$P_\infty = I - P_0.$$

To prove Theorem 3.1 (ii), we first introduce the decay estimate of  $e^{-tL}P_\infty u_0$  for  $u_0 \in H^1(\Omega) \times H_0^1(\Omega)$ .

**Proposition 6.1.** *There exist constants  $\nu_0$ ,  $\gamma_0$  and  $\omega_0$  such that if  $\nu \geq \nu_0$ ,  $\frac{\gamma^2}{\nu+\bar{\nu}} \geq \gamma_0^2$  and  $\omega \leq \omega_0$ , then  $e^{-tL}P_\infty u_0$  have the following properties. If  $u_0 \in H^1(\Omega) \times H_0^1(\Omega)$ , then there exists a constant  $d > 0$  such that  $e^{-tL}P_\infty u_0$  satisfies*

$$\|e^{-tL}P_\infty u_0\|_{H^1} \leq Ce^{-dt}\|u_0\|_{H^1} \quad (6.1)$$

for  $t \geq 0$ .

**Proof.**  $P_\infty$  is written as

$$P_\infty = P_{\infty,0} + \tilde{P}_\infty,$$

where

$$\begin{aligned} P_{\infty,0}u &= \mathcal{F}^{-1}[P_{\infty,0}u], & P_{\infty,0}u &= \mathbf{1}_{\{|\xi| \leq r_0\}}(I - P_0)u, \\ \tilde{P}_\infty u &= \mathcal{F}^{-1}[\tilde{P}_\infty u], & \tilde{P}_\infty u &= (1 - \mathbf{1}_{\{|\xi| \leq r_0\}})u. \end{aligned}$$

The estimate  $\|e^{-tL}\tilde{P}_\infty u_0\|_{H^1} \leq Ce^{-dt}\|u_0\|_{H^1}$  was proved in [1, Theorem 3.3]. As for  $P_{\infty,0}$  part, since  $\rho(-L_\xi|_{(I-\Pi_0(\xi))L^2}) \subset \{\lambda \in \mathbf{C} : \text{Re}\lambda \geq -\frac{c_0}{2}\}$  by Theorem 4.4, we have

$$|e^{-tL_\xi}P_{\infty,0}u_0|_2 \leq Ce^{-\frac{c_0}{4}t}|u_0|_2. \quad (6.2)$$

We now apply the argument of the proof of [1, Proposition 4.20] to  $u(t) = e^{-tL}P_{\infty,0}u_0$ . Due to (6.2), one can replace  $e^{-\frac{d_2}{2}|\xi|^2 t}|u_0|_2^2$  in the inequality (4.72) of [1] by  $e^{-\frac{c_0}{2}t}|u_0|_2^2$  to obtain  $E_{4,1}^{(0)}[u](t) \leq Ce^{-2\tilde{d}_1 t}|u_0|_{H^1}^2$  for a positive constant  $\tilde{d}_1$ . Integrating this over  $|\xi| \leq r_0$  and using the Plancherel Theorem, we have

$$\|e^{-tL}P_{\infty,0}u_0\|_{H^1} \leq Ce^{-\tilde{d}t}\|u_0\|_{H^1}$$

for a positive constant  $\tilde{d}$ . Combining the estimates for  $e^{-tL}\tilde{P}_\infty u_0$  and  $e^{-tL}P_{\infty,0}u_0$  we obtain the desired estimate. This completes the proof.  $\square$

We next consider the estimate for  $e^{-tL}u$  for  $0 < t \leq 1$ .

**Proposition 6.2.** *Let  $T > 0$ . If  $u_0 \in H^1(\Omega) \times \tilde{H}^1(\Omega)$ , then  $e^{-tL}u_0$  satisfies  $e^{-tL}u_0 \in H^1(\Omega) \times H_0^1(\Omega)$  for  $t > 0$  and*

$$\|e^{-tL}u_0\|_{H^1} \leq C_T\{\|u_0\|_{H^1 \times \tilde{H}^1} + t^{-\frac{1}{2}}\|w_0\|_2\} \quad (6.3)$$

for  $0 < t \leq T$ .

Let  $u_0 \in H^1(\Omega) \times \tilde{H}^1(\Omega)$ . Applying Proposition 6.2 with  $t = 1$ , we have  $u_1 = e^{-tL}u_0|_{t=1} \in H^1(\Omega) \times H_0^1(\Omega)$  and

$$\|u_1\|_{H^1} \leq C\|u_0\|_{H^1 \times \tilde{H}^1}.$$

This, together with Proposition 6.1 and Proposition 6.2, implies Theorem 3.1 (ii).

It remains to prove Proposition 6.2.

**Lemma 6.3.** *Let  $T > 0$ . There hold the following estimates for  $0 \leq t \leq T$ :*

(i) for  $\ell = 0, 1$ ,

$$\|\partial_{x_3}^\ell u(t)\|_2^2 + c \int_0^t \|\nabla \partial_{x_3}^\ell w\|_2^2 + \|\operatorname{div} \partial_{x_3}^\ell w\|_2^2 d\tau \leq C_T \|\partial_{x_3}^\ell u_0\|_2^2,$$

(ii)

$$\begin{aligned} & \|\chi_0 \partial_{x'} u(t)\|_2^2 + c \int_0^t \|\chi_0 \nabla \partial_{x'} w(\tau)\|_2^2 + \|\chi_0 \operatorname{div} \partial_{x'} w\|_2^2 d\tau \\ & \leq C_T \left\{ \|u_0\|_2^2 + \|\partial_{x_3} u_0\|_2^2 + \|\chi_0 \partial_{x'} u_0\|_2^2 + \int_0^t \|\partial_{x'} \phi(\tau)\|_2^2 d\tau \right\}, \end{aligned}$$

(iii) for  $1 \leq m \leq N$ ,

$$\begin{aligned} & \|\chi_m \partial u(t)\|_2^2 + c \int_0^t \|\chi_m \nabla \partial w\|_2^2 + \|\chi_m \operatorname{div} \partial w\|_2^2 d\tau \\ & \leq C_T \left\{ \|u_0\|_2^2 + \|\partial_{x_3} u_0\|_2^2 + \|\chi_m \partial u_0\|_2^2 + \int_0^t \|\partial_{x'} \phi\|_2^2 d\tau \right\}. \end{aligned}$$

Lemma 6.3 can be proved by the energy method as those in the proof of [1, Propositions 4.7, 4.15, 4.17]. Note that here are no restrictions on  $\nu$ ,  $\tilde{\nu}$  and  $\gamma$  but  $C_T$  depends on  $T$ .

We next consider the  $L^2$  estimate of the normal derivative for  $\phi$ .

**Lemma 6.4.** *Let  $T > 0$ . For  $1 \leq m \leq N$ , there holds the estimate for  $0 \leq t \leq T$ :*

$$\begin{aligned} & \|\chi_m \partial_n \phi(t)\|_2^2 \\ & \leq C_T \left\{ \|u_0\|_2^2 + \|\partial_{x_3} u_0\|_2^2 + \|\chi_m \partial u_0\|_2^2 + \|\chi_m \partial_n \phi_0\|_2^2 + \int_0^t \|\partial_{x'} \phi\|_2^2 d\tau \right\}. \end{aligned}$$

**Proof.** Let us transform a scalar field  $p(x')$  on  $D \cap \mathcal{O}_m$  as

$$\tilde{p}(y') = p(x') \quad (y' = \Psi^m(x'), \quad x' \in D \cap \mathcal{O}_m),$$

where  $\Psi^m(x')$  is a function given in Section 2. Similarly we transform a vector field  $h(x') = {}^T(h^1(x'), h^2(x'), h^3(x'))$  into  $\tilde{h}(y') = {}^T(\tilde{h}^1(y'), \tilde{h}^2(y'), \tilde{h}^3(y'))$  as

$$h(x') = E(y') \tilde{h}(y')$$

where  $E(y') = (e_1(y'), e_2(y'), e_3)$  with  $e_1(y')$ ,  $e_2(y')$  and  $e_3$  given in Section 2. From the proof of [1, Proposition 4.16], we have

$$\partial_\tau \partial_{y_1} \tilde{\phi} + (a + b \partial_{y_3}) \partial_{y_1} \tilde{\phi} = \tilde{\rho}_s I - \frac{\gamma^2 \tilde{\rho}_s^2}{\nu + \tilde{\nu}} \partial_\tau \tilde{w}^1, \quad (6.4)$$

where

$$a(y') = \frac{\tilde{\rho}_s \tilde{P}'(\tilde{\rho}_s)}{\nu + \tilde{\nu}}, \quad b(y') = \tilde{v}_s^3(y'),$$

$$I = -\frac{\gamma^2}{\nu+\tilde{\nu}}\left\{\nu(\operatorname{rot}_y \operatorname{rot}_y \tilde{w})^1 + \tilde{\rho}_s \partial_{y_1} \left( \frac{\tilde{P}'(\tilde{\rho}_s)}{\gamma^2 \tilde{\rho}_s} \right) \tilde{\phi} + \frac{\nu}{\gamma^2} \tilde{\rho}_s (\Delta_{y'} \tilde{v}_s)^1 \tilde{\phi} + \tilde{\rho}_s \tilde{v}_s^3 \partial_{y_3} \tilde{w}^1 \right\} \\ - \left\{ \frac{1}{\tilde{\rho}_s} \partial_{y_1} \tilde{v}_s^3 \partial_{y_3} \tilde{\phi} + \gamma^2 \frac{1}{\tilde{\rho}_s} \partial_{y_1} (\operatorname{div}_y (\tilde{\rho}_s \tilde{w})) - \gamma^2 \partial_{y_1} \operatorname{div}_y \tilde{w} \right\}.$$

Here  $(\operatorname{rot}_y \tilde{w})^1$  denotes the  $e_1(y')$  component of  $\operatorname{rot}_y \tilde{w}$ , and so on. We note that  $(\operatorname{rot}_y \operatorname{rot}_y \tilde{w})^1$  does not contain  $\partial_{y_1}^2$ . See the proof of [1, Proposition 4.16]. We also note that there is a positive constant  $a_0$  such that

$$a(y') \geq a_0 > 0$$

for any  $y' \in \Psi^m(D)$ .

We denote by  $e^{-t(a+b\partial_{y_3})}$  the semigroup generated by  $-(a+b\partial_{y_3})$ , i.e.,

$$e^{-t(a+b\partial_{y_3})} \tilde{\phi}_0 = \mathcal{F}^{-1} [e^{-(a(y')+i\xi b(y'))t} \widehat{\tilde{\phi}_0}].$$

Then it is easy to see that

$$\|\tilde{\chi}_m e^{-t(a+b\partial_{y_3})} \tilde{\phi}_0\|_2 \leq e^{-a_0 t} \|\tilde{\chi}_m \tilde{\phi}_0\|_2.$$

In terms of  $e^{-t(a+b\partial_{y_3})}$ ,  $\partial_{y_1} \tilde{\phi}$  is written as

$$\begin{aligned} \partial_{y_1} \tilde{\phi}(t) &= e^{-t(a+b\partial_{y_3})} \partial_{y_1} \tilde{\phi}_0 + \int_0^t e^{-(t-\tau)(a+b\partial_{y_3})} \tilde{\rho}_s \tilde{I}(\tau) d\tau \\ &\quad - \frac{\gamma^2 \tilde{\rho}_s^2}{\nu + \tilde{\nu}} \int_0^t e^{-(t-\tau)(a+b\partial_{y_3})} \partial_\tau \tilde{w}^1 d\tau \\ &\equiv J_1 + J_2 + J_3. \end{aligned}$$

As for  $J_1$  and  $J_2$ , we have

$$\begin{aligned} \|\tilde{\chi}_m J_1\|_2 &\leq e^{-a_0 t} \|\tilde{\chi}_m \partial_{y_1} \tilde{\phi}_0\|_2, \\ \|\tilde{\chi}_m J_2\|_2 &\leq C \int_0^t e^{-a_0(t-\tau)} \|\tilde{\chi}_m \tilde{I}(\tau)\|_2 d\tau. \end{aligned}$$

As for  $J_3$ , integrating by parts, we have

$$J_3 = \frac{\gamma^2 \tilde{\rho}_s^2}{\nu + \tilde{\nu}} \left[ e^{-t(a+b\partial_{y_3})} \tilde{w}_0^1 - \tilde{w}^1(t) + (a+b\partial_{y_3}) \int_0^t e^{-(t-\tau)(a+b\partial_{y_3})} \tilde{w}^1(\tau) d\tau \right].$$

We thus obtain

$$\|\tilde{\chi}_m J_3\|_2 \leq C \left\{ e^{-a_0 t} \|\tilde{\chi}_m \tilde{w}_0^1\|_2 + \|\tilde{\chi}_m \tilde{w}^1(t)\|_2 + \int_0^t e^{-a_0(t-\tau)} \|\tilde{\chi}_m \partial_{y_3} \tilde{w}^1(\tau)\|_2 d\tau \right\}.$$

Furthermore, we have

$$\begin{aligned} \|\tilde{\chi}_m I(\tau)\|_2 &\leq C \left\{ \|\tilde{\chi}_m \tilde{\phi}(\tau)\|_2 + \|\tilde{\chi}_m \partial_{y_3} \tilde{\phi}(\tau)\|_2 + \|\tilde{\chi}_m \tilde{w}(\tau)\|_2 \right. \\ &\quad \left. + \|\tilde{\chi}_m \nabla_y \tilde{w}(\tau)\|_2 + \|\tilde{\chi}_m \nabla_y \partial_{y_2} \tilde{w}(\tau)\|_2 + \|\tilde{\chi}_m \nabla_y \partial_{y_3} \tilde{w}(\tau)\|_2 \right\}. \end{aligned}$$



It then follows that

$$\begin{aligned} \|\tilde{\chi}_m \partial_{y_1} \tilde{\phi}(t)\|_2 &\leq C \left[ e^{-a_0 t} (\|\tilde{\chi}_m \partial_{y_1} \tilde{\phi}_0\|_2 + \|\tilde{\chi}_m \tilde{w}_0^1\|_2) + \|\tilde{\chi}_m \tilde{w}^1(t)\|_2 \right. \\ &\quad + \int_0^t e^{-a_0(t-\tau)} \left\{ \|\tilde{\chi}_m \tilde{\phi}(\tau)\|_2 + \|\tilde{\chi}_m \partial_{y_3} \tilde{\phi}(\tau)\|_2 + \|\tilde{\chi}_m \tilde{w}(\tau)\|_2 \right. \\ &\quad \left. \left. + \|\tilde{\chi}_m \nabla_y \tilde{w}(\tau)\|_2 + \|\tilde{\chi}_m \nabla_y \partial_{y_2} \tilde{w}(\tau)\|_2 + \|\tilde{\chi}_m \nabla_y \partial_{y_3} \tilde{w}(\tau)\|_2 \right\} d\tau \right]. \end{aligned}$$

Inverting to the original coordinates  $x'$  and noting that  $\partial_{y_1} = \partial_n$ ,  $\partial_{y_2} = \partial$ , we see that

$$\begin{aligned} \|\chi_m \partial_n \phi(t)\|_2 &\leq C \left\{ e^{-a_0 t} (\|\chi_m \partial_n \phi_0\|_2 + \|\chi_m w_0^1\|_2) + \|\chi_m w^1(t)\|_2 \right. \\ &\quad + \int_0^t \left( \|\chi_m \phi(\tau)\|_2 + \|\chi_m \partial_{x_3} \phi(\tau)\|_2 + \|\chi_m w(\tau)\|_2 \right. \\ &\quad \left. \left. + \|\chi_m \nabla w(\tau)\|_2 + \|\chi_m \nabla \partial w(\tau)\|_2 + \|\chi_m \nabla \partial_{x_3} w(\tau)\|_2 \right) d\tau \right\}. \end{aligned}$$

This, together with Lemma 6.3, yields the desired estimate. This completes the proof.  $\square$

By Lemma 6.3 and Lemma 6.4, we have the following estimate.

**Lemma 6.5.** *Let  $T > 0$ . There exists a positive constant  $c$  such that the estimate*

$$\begin{aligned} &\|u(t)\|_{H^1 \times \tilde{H}^1}^2 + c \int_0^t \left( \|\nabla w(\tau)\|_2^2 + \|\operatorname{div} w(\tau)\|_2^2 + \|\nabla \partial_{x_3} w(\tau)\|_2^2 + \|\operatorname{div} \partial_{x_3} w(\tau)\|_2^2 \right. \\ &\quad \left. + \|\chi_0 \nabla \partial_{x'} w(\tau)\|_2^2 + \|\chi_0 \operatorname{div} \partial_{x'} w(\tau)\|_2^2 + \sum_{m=1}^N \left\{ \|\chi_m \nabla \partial w(\tau)\|_2^2 + \|\chi_m \operatorname{div} \partial w(\tau)\|_2^2 \right\} \right) d\tau \\ &\leq C_T \|u_0\|_{H^1 \times \tilde{H}^1}^2 \end{aligned}$$

holds for  $0 \leq t \leq T$ .

We finally consider the  $L^2$  estimate for  $\partial_{x'} w$ .

**Lemma 6.6.** *Let  $T > 0$ . There holds the estimate*

$$\|\partial_{x'} w(t)\|_2 \leq C_T \{ \|u_0\|_{H^1 \times \tilde{H}^1} + t^{-\frac{1}{2}} \|w_0\|_2 \}$$

for  $0 < t \leq T$ .

**Proof.** We see that  $w$  satisfies the equation

$$\partial_t w + \bar{A} w + \bar{B} u = 0,$$

where  $\bar{A}$  is the  $3 \times 3$  operator defined by

$$\bar{A} = -\frac{\nu}{\rho_s} \Delta - \frac{\nu + \tilde{\nu}}{\rho_s} \nabla \operatorname{div},$$

$\bar{B}$  is the  $3 \times 4$  operator defined by

$$\bar{B} = \begin{pmatrix} \nabla' \left( \frac{P(\rho_s)}{\gamma^2 \rho_s} \cdot \right) & v_s^3 \partial_{x_3} I_2 & 0 \\ \partial_{x_3} \left( \frac{P'(\rho_s)}{\gamma^2 \rho_s} \cdot \right) + \frac{\nu \Delta' v_s^3}{\gamma^2 \rho_s^2} & T(\nabla' v_s^3) & v_s^3 \partial_{x_3} \cdot \end{pmatrix}$$

We write  $w(t)$  as

$$w(t) = e^{-t\bar{A}} w_0 + \int_0^t e^{-(t-\tau)\bar{A}} \bar{B} u(\tau) d\tau.$$

Then

$$\nabla' w(t) = \nabla' e^{-t\bar{A}} w_0 + \int_0^t \nabla' e^{-(t-\tau)\bar{A}} \bar{B} u(\tau) d\tau. \quad (6.5)$$

Since  $\bar{A}$  is strongly elliptic, we have

$$\|\nabla' e^{-t\bar{A}} w_0\|_2 \leq C t^{-\frac{1}{2}} \|w_0\|_2$$

for  $0 < t \leq T$ . Furthermore, we see from Lemma 6.3 and Lemma 6.5 that

$$\begin{aligned} \left\| \int_0^t \nabla' e^{-(t-\tau)\bar{A}} \bar{B} u(\tau) d\tau \right\|_2 &\leq C \int_0^t (t-\tau)^{-\frac{1}{2}} \|\bar{B} u(\tau)\|_2 d\tau \\ &\leq C \int_0^t (t-\tau)^{-\frac{1}{2}} \|u(\tau)\|_{H^1 \times \tilde{H}^1} d\tau \\ &\leq C \|u_0\|_{H^1 \times \tilde{H}^1} \int_0^t (t-\tau)^{-\frac{1}{2}} d\tau \\ &\leq C T^{\frac{1}{2}} \|u_0\|_{H^1 \times \tilde{H}^1} \end{aligned} \quad (6.6)$$

for  $0 < t \leq T$ . It then follows from (6.5) and (6.6) that

$$\|\partial_{x'} w(t)\|_2 \leq C_T \{ \|u_0\|_{H^1 \times \tilde{H}^1} + t^{-\frac{1}{2}} \|w_0\|_2 \} \quad (6.7)$$

for  $0 < t \leq T$ . This completes the proof.  $\square$

**Proof of Proposition 6.2.** Let  $u(t) = e^{-tL} u_0$ . It is not difficult to see that if  $u_0 \in H^1(\Omega) \times H_0^1(\Omega)$ , then  $u(t)$  satisfies

$$u \in C([0, T]; H^1(\Omega) \times H_0^1(\Omega)), \quad \tilde{Q}u \in L^2(0, T; H^2(\Omega)). \quad (6.8)$$

Using Lemma 6.5 and Lemma 6.6, we obtain the estimate

$$\|u(t)\|_{H^1}^2 + c \int_0^t \bar{D}_1[w](\tau) d\tau \leq C_T \{ \|u_0\|_{H^1 \times \tilde{H}^1}^2 + t^{-1} \|w_0\|_2^2 \}$$

for  $0 < t \leq T$ . Here

$$\begin{aligned} \bar{D}_1[w] &= (\|\nabla w\|_2^2 + \|\operatorname{div} w\|_2^2) + (\|\nabla \partial_{x_3} w\|_2^2 + \|\operatorname{div} \partial_{x_3} w\|_2^2) \\ &\quad + (\|\chi_0 \nabla \partial_{x'} w\|_2^2 + \|\chi_0 \operatorname{div} \partial_{x'} w\|_2^2) + \sum_{m=1}^N (\|\chi_m \nabla \partial w\|_2^2 + \|\chi_m \operatorname{div} \partial w\|_2^2). \end{aligned}$$

We thus obtain estimate (6.3) if  $u_0 \in H^1(\Omega) \times H_0^1(\Omega)$ . Since  $H_0^1(\Omega)$  is dense in  $\tilde{H}^1(\Omega)$ , one can see from Lemma 6.5, (6.3) and (6.8) that if  $u_0 \in H^1(\Omega) \times \tilde{H}^1(\Omega)$ , then  $u(t)$  satisfies

$$u \in C([0, T]; H^1(\Omega) \times \tilde{H}^1(\Omega)) \cap C((0, T]; H^1(\Omega) \times H_0^1(\Omega))$$

and estimate (6.3). This completes the proof.  $\square$

**Acknowledgements.** Y. Kagei was partly supported by JSPS KAKENHI Grant Number 24340028, 22244009, 24224003.

## References

- [1] R. Aoyama, Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a parallel flow in a cylindrical domain, to appear in Kyushu J. Math.
- [2] J. Březina, Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a time-periodic parallel flow, SIAM J. Math. Anal. **45** (2013), pp. 3514-3574.
- [3] J. Březina, Y. Kagei, Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow, Math. Models Meth. Appl. Sci. **22** (2012), 1250007, pp. 1-53.
- [4] J. Březina, Y. Kagei, Spectral properties of the linearized compressible Navier-Stokes equation around time-periodic parallel flow, J. Differential Equations **255** (2013), pp. 1132-1195.
- [5] K. Deckelnick, Decay estimates for the compressible Navier-Stokes equations in unbounded domains, Math. Z. **209** (1992), pp. 115-130.
- [6] D. Hoff, K. Zumbrun, Multi-dimensional diffusion waves for the Navier-Stokes equations of compressible flow, Indiana Univ. Math. J. **44** (1995), pp. 604-676.
- [7] G. Iooss, M. Padula, Structure of the linearized problem for compressible parallel fluid flows. Ann. Univ. Ferrara Sez. VII **43** (1997), pp. 157-171.
- [8] Y. Ishihara, Y. Kagei, Large time behavior of the semigroup on  $L^p$  spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain. J. Differential Equations **248** (2010), pp. 252-286.
- [9] Y. Kagei, Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow, Arch. Rational Mech. Anal. **205** (2012), pp. 585-650.

- [10] Y. Kagei, Y. Nagahuchi, T. Sudou, Decay estimate on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow, J. Math-for-Ind. **2A** (2010), pp. 39-56; Correction to "Decay estimate on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow" in J. Math-for-Ind. **2A** (2010), pp. 39-56 J. Math-for-Ind. **2B** (2010), pp. 235.
- [11] Y. Kagei, T. Kobayashi, Asymptotic behavior of solutions to the compressible Navier-Stokes equations on the half space, Arch. Rational Mech. Anal. **177** (2005), pp. 231-330.
- [12] Y. Kagei, T. Nukumizu, Asymptotic behavior of solutions to the compressible Navier-Stokes equation in a cylindrical domain, Osaka J. Math. **45** (2008), pp. 987-1026.
- [13] S. Kawashima, Systems of a hyperbolic-parabolic composite type, with applications to the equations of magnethydrodynamics, Ph. D. Thesis, Kyoto University (1983).
- [14] T. Kobayashi, Some estimates of solutions for the equations of motion of compressible viscous fluid in an exterior domain in  $\mathbf{R}^3$ , J. Differential Equations **184** (2002), pp. 587-619.
- [15] T. Kobayashi, Y. Shibata, Decay estimates of solutions for the equations of motion of compressible viscous and heat-conductive gases in an exterior domain in  $\mathbf{R}^3$ , Comm. Math. Phys. **200** (1999), pp. 621-659.
- [16] A. Matsumura, An energy method for the equations of motion of compressible viscous and heat-conductive fluids, University of Wisconsin-Madison, MRC Technical Summary Report # 2194 (1981), pp. 1-16.
- [17] A. Matsumura, T. Nishida, The initial value problem for the equations of motion of compressible viscous and heat-conductive fluids, Proc. Japan Acad. Ser. A **55** (1979), pp. 337-342.
- [18] A. Matsumura, T. Nishida, Initial Boundary Value Problem for the Equations of Motion of General Fluids, Comput. Meth. Appl. Sci. Eng. **V** (1982), pp. 389-406.
- [19] A. Matsumura, T. Nishida, Initial Boundary Value Problems for the Equations of Motion of Compressible Viscous and Heat-Conductive Fluids, Comm. Math. Phys. **89** (1983), pp. 445-464.
- [20] H. Sohr, *The Navier-Stokes Equations. An Elementary Functional Analytic approach*, Birkhäuser, Basel (2001).

# List of MI Preprint Series, Kyushu University

## The Global COE Program

### Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI  
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA  
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO  
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU  
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI  
Torsion points of abelian varieties with values in infinite extensions over a p-adic field
- MI2008-6 Yoshiyuki TOMIYAMA  
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI  
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA  
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA  
Alpha-determinant cyclic modules and Jacobi polynomials
- MI2008-10 Sangyeol LEE & Hiroki MASUDA  
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA  
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO  
On the  $L^2$  a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA  
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials

- MI2008-14 Takashi NAKAMURA  
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA  
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO  
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI  
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI  
Variable selection for functional regression model via the  $L_1$  regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI  
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI  
Flat modules and Groebner bases over truncated discrete valuation rings
- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA  
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI  
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI  
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA  
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO  
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA  
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA  
Hypergeometric  $\tau$ -functions of the q-Painlevé system of type  $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA  
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA  
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications

- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI  
Large time behavior of the semigroup on  $L^p$  spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain
- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE  
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO  
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO  
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI  
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI  
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO  
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA  
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO  
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI  
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO  
On very accurate enclosure of the optimal constant in the a priori error estimates for  $H_0^2$ -projection
- MI2009-26 Manabu YOSHIDA  
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI  
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA  
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA  
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance

- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA  
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI  
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA  
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA  
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA  
Projective reduction of the discrete Painlevé system of type  $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI  
Finite element computation for scattering problems of micro-hologram using DtN map
- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA  
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA  
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA  
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA  
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI  
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI  
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI  
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI  
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA  
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO  
Abstract collision systems on groups



- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA  
An algebraic approach to underdetermined experiments
- MI2010-10 Kei HIROSE & Sadanori KONISHI  
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA  
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU  
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA  
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI  
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO  
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO  
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE  
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU  
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA  
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE  
Lagrangian approach to weakly nonlinear stability of an elliptical flow
- MI2010-21 Hiroki MASUDA  
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU  
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiro FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI  
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU  
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra

- MI2010-25 Toshimitsu TAKAESU  
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA  
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA  
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA  
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE  
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI  
Spin-spin correlation functions of the  $q$ -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI  
Nonlinear regression modeling and spike detection via Gaussian basis expansions
- MI2010-32 Nobutaka NAKAZONO  
Hypergeometric  $\tau$  functions of the  $q$ -Painlevé systems of type  $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI  
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI  
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI  
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA  
CAP representations of inner forms of  $Sp(4)$  with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER  
Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide FUKUMOTO & Alexander B. SAMOKHIN  
Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI  
Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI  
Anderson-Darling test and the Malliavin calculus
- MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI  
Nonlinear regression modeling via Compressed Sensing

- MI2011-5 Hiroshi INOUE  
Implications in Compressed Sensing and the Restricted Isometry Property
- MI2011-6 Daeju KIM & Sadanori KONISHI  
Predictive information criterion for nonlinear regression model based on basis expansion methods
- MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI  
Group variable selection via relevance vector machine
- MI2011-8 Jan BREZINA & Yoshiyuki KAGEI  
Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow  
Group variable selection via relevance vector machine
- MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC  
Recursive structures in the multispecies TASEP
- MI2011-10 Kazunori YASUTAKE  
On projective space bundle with nef normalized tautological line bundle
- MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA  
An explicit formula for the discrete power function associated with circle patterns of Schramm type
- MI2011-12 Yoshiyuki KAGEI  
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow
- MI2011-13 Vladimír CHALUPECKÝ & Adrian MUNTEAN  
Semi-discrete finite difference multiscale scheme for a concrete corrosion model: approximation estimates and convergence
- MI2011-14 Jun-ichi INOBUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA  
Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves
- MI2011-15 Hiroshi INOUE  
A generalization of restricted isometry property and applications to compressed sensing
- MI2011-16 Yu KAWAKAMI  
A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space
- MI2011-17 Naoyuki KAMIYAMA  
Matroid intersection with priority constraints
- MI2012-1 Kazufumi KIMOTO & Masato WAKAYAMA  
Spectrum of non-commutative harmonic oscillators and residual modular forms
- MI2012-2 Hiroki MASUDA  
Mighty convergence of the Gaussian quasi-likelihood random fields for ergodic Levy driven SDE observed at high frequency

- MI2012-3 Hiroshi INOUE  
A Weak RIP of theory of compressed sensing and LASSO
- MI2012-4 Yasuhide FUKUMOTO & Youich MIE  
Hamiltonian bifurcation theory for a rotating flow subject to elliptic straining field
- MI2012-5 Yu KAWAKAMI  
On the maximal number of exceptional values of Gauss maps for various classes of surfaces
- MI2012-6 Marcio GAMEIRO, Yasuaki HIRAOKA, Shunsuke IZUMI, Miroslav KRAMAR, Konstantin MISCHAIKOW & Vidit NANDA  
Topological Measurement of Protein Compressibility via Persistence Diagrams
- MI2012-7 Nobutaka NAKAZONO & Seiji NISHIOKA  
Solutions to a  $q$ -analog of Painlevé III equation of type  $D_7^{(1)}$
- MI2012-8 Naoyuki KAMIYAMA  
A new approach to the Pareto stable matching problem
- MI2012-9 Jan BREZINA & Yoshiyuki KAGEI  
Spectral properties of the linearized compressible Navier-Stokes equation around time-periodic parallel flow
- MI2012-10 Jan BREZINA  
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a time-periodic parallel flow
- MI2012-11 Daeju KIM, Shuichi KAWANO & Yoshiyuki NINOMIYA  
Adaptive basis expansion via the extended fused lasso
- MI2012-12 Masato WAKAYAMA  
On simplicity of the lowest eigenvalue of non-commutative harmonic oscillators
- MI2012-13 Masatoshi OKITA  
On the convergence rates for the compressible Navier- Stokes equations with potential force
- MI2013-1 Abuduwaili PAERHATI & Yasuhide FUKUMOTO  
A Counter-example to Thomson-Tait-Chetayev's Theorem
- MI2013-2 Yasuhide FUKUMOTO & Hirofumi SAKUMA  
A unified view of topological invariants of barotropic and baroclinic fluids and their application to formal stability analysis of three-dimensional ideal gas flows
- MI2013-3 Hiroki MASUDA  
Asymptotics for functionals of self-normalized residuals of discretely observed stochastic processes
- MI2013-4 Naoyuki KAMIYAMA  
On Counting Output Patterns of Logic Circuits
- MI2013-5 Hiroshi INOUE  
RIPless Theory for Compressed Sensing

- MI2013-6 Hiroshi INOUE  
Improved bounds on Restricted isometry for compressed sensing
- MI2013-7 Hidetoshi MATSUI  
Variable and boundary selection for functional data via multiclass logistic regression modeling
- MI2013-8 Hidetoshi MATSUI  
Variable selection for varying coefficient models with the sparse regularization
- MI2013-9 Naoyuki KAMIYAMA  
Packing Arborescences in Acyclic Temporal Networks
- MI2013-10 Masato WAKAYAMA  
Equivalence between the eigenvalue problem of non-commutative harmonic oscillators and existence of holomorphic solutions of Heun's differential equations, eigenstates degeneration, and Rabi's model
- MI2013-11 Masatoshi OKITA  
Optimal decay rate for strong solutions in critical spaces to the compressible Navier-Stokes equations
- MI2013-12 Shuichi KAWANO, Ibuki HOSHINA, Kazuki MATSUDA & Sadanori KONISHI  
Predictive model selection criteria for Bayesian lasso
- MI2013-13 Hayato CHIBA  
The First Painleve Equation on the Weighted Projective Space
- MI2013-14 Hidetoshi MATSUI  
Variable selection for functional linear models with functional predictors and a functional response
- MI2013-15 Naoyuki KAMIYAMA  
The Fault-Tolerant Facility Location Problem with Submodular Penalties
- MI2013-16 Hidetoshi MATSUI  
Selection of classification boundaries using the logistic regression
- MI2014-1 Naoyuki KAMIYAMA  
Popular Matchings under Matroid Constraints
- MI2014-2 Yasuhide FUKUMOTO & Youichi MIE  
Lagrangian approach to weakly nonlinear interaction of Kelvin waves and a symmetry-breaking bifurcation of a rotating flow
- MI2014-3 Reika AOYAMA  
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Parallel flow in a cylindrical domain
- MI2014-4 Naoyuki KAMIYAMA  
The Popular Condensation Problem under Matroid Constraints

- MI2014-5 Yoshiyuki KAGEI & Kazuyuki TSUDA  
Existence and stability of time periodic solution to the compressible Navier-Stokes equation for time periodic external force with symmetry
- MI2014-6 This paper was withdrawn by the authors.
- MI2014-7 Masatoshi OKITA  
On decay estimate of strong solutions in critical spaces for the compressible Navier-Stokes equations
- MI2014-8 Rong ZOU & Yasuhide FUKUMOTO  
Local stability analysis of azimuthal magnetorotational instability of ideal MHD flows
- MI2014-9 Yoshiyuki KAGEI & Naoki MAKIO  
Spectral properties of the linearized semigroup of the compressible Navier-Stokes equation on a periodic layer
- MI2014-10 Kazuyuki TSUDA  
On the existence and stability of time periodic solution to the compressible Navier-Stokes equation on the whole space
- MI2014-11 Yoshiyuki KAGEI & Takaaki NISHIDA  
Instability of plane Poiseuille flow in viscous compressible gas
- MI2014-12 Chien-Chung HUANG, Naonori KAKIMURA & Naoyuki KAMIYAMA  
Exact and approximation algorithms for weighted matroid intersection
- MI2014-13 Yusuke SHIMIZU  
Moment convergence of regularized least-squares estimator for linear regression model
- MI2015-1 Hidetoshi MATSUI  
Sparse regularization for multivariate linear models for functional data
- MI2015-2 Reika AOYAMA & Yoshiyuki KAGEI  
Spectral properties of the semigroup for the linearized compressible Navier-Stokes equation around a parallel flow in a cylindrical domain