Stable Matchings with Ties, Master Preference Lists, and Matroid Constraints

Kamiyama, Naoyuki

Institute of Mathematics for Industry, Kyushu University | JST, PRESTO

<http://hdl.handle.net/2324/1500418>

出版情報 : Lecture Notes in Computer Science. 9347, pp.3-14, 2015. Springer Berlin Heidelberg
バージョン :
権利関係 :

MI Preprint Series

**Mathematics for Industry
Kyushu University**

Stable Matchings with Ties, Master Preference Lists, and Matroid Constraints

Naoyuki Kamiyama

MI 2015-3

(Received April 23, 2015)

Institute of Mathematics for Industry
Graduate School of Mathematics
Kyushu University
Fukuoka, JAPAN

Stable Matchings with Ties, Master Preference Lists, and Matroid Constraints

Naoyuki Kamiyama^{*†‡}

Abstract

In this paper, we consider a matroid generalization of the hospitals/residents problem with ties and master lists. In this model, the capacity constraints for hospitals are generalized to matroid constraints. By generalizing the algorithms of O'Malley for the hospitals/residents problem with ties and master lists, we give polynomial-time algorithms for deciding whether there exist a super-stable matching and a strongly stable matching in our model, and finding such matchings, if they exist.

1 Introduction

The stable matching problem introduced by Gale and Shapley [4] is one of the most popular mathematical models of a matching problem in which agents have preferences. It is known [4] that if preference lists do not contain ties, then there always exists a stable matching and we can find one in polynomial time. However, if there exist ties in preference lists, then the situation dramatically changes. In the stable matching problem with ties, three stability concepts were proposed by Irving [5] (see also Chapter 3 of [11] for a survey of the stable matching problem with ties). The first one is called the weak stability. This stability concept guarantees that there exists no unmatched pair each of whom prefers the other to the current partner. It is known [5] that there always exists a weakly stable matching and we can find one in polynomial time by slightly modifying the algorithm of [4]. The second one is called the strong stability. This stability concept guarantees that there exists no unmatched pair such that (i) there exists an agent a in this pair that prefers the other to the current partner, and (ii) the agent in this pair other than a prefers a to the current partner, or is indifferent between a and the current partner. The last one is called the super-stability. This stability concept guarantees that there exists no unmatched pair each of whom prefers the other to the current partner or is indifferent between the other and the current partner.

One of the most notable differences between the last two concepts and the stability concept in the stable matching problem without ties is that there may exist no stable matching [5]. Thus, from the algorithmic viewpoint, it is important to reveal whether we can check the existence of matchings satisfying such stability concepts in polynomial time. For the one-to-one setting (i.e., the stable matching problem with ties), Irving [5] gave polynomial-time algorithms for finding a super-stable matching and a strongly stable matching (see also [10]). For the many-to-one setting (i.e., the hospitals/residents problem with ties), Irving, Manlove, and Scott [6] gave a polynomial-time algorithm for finding a super-stable matching, and Irving, Manlove, and

^{*}Institute of Mathematics for Industry, Kyushu University, Fukuoka, Japan

[†]JST, PRESTO, Saitama, Japan

[‡]kamiyama@imi.kyushu-u.ac.jp

Scott [7] a polynomial-time algorithm for finding a strongly stable matching. It should be noted that Kavitha, Mehlhorn, Michail, and Paluch [9] proposed a faster algorithm for this setting.

In this paper, we consider a matroid generalization of the hospitals/residents problem with ties. In our model, the capacity constraints for hospitals are generalized to matroid constraints. In contrast to the stable matching problem without ties (and the weak stability) [3], to the best of our knowledge, it is open whether we can extend the results about the super-stability and the strong stability to the matroid setting. In this paper, as a step toward the settlement of this question, we focus on the situation in which we are given a master list and the preference list of each hospital over residents is derived from this master list. For the stable matching problem with ties and master lists, Irving, Manlove, and Scott [8] gave simple polynomial-time algorithms for finding a super-stable matching and a strongly stable matching. Furthermore, O'Malley [12] gave polynomial-time algorithms for finding a super-stable matching and a strongly stable matching in the hospitals/residents problem with ties and master lists. In this paper, by generalizing the algorithms of [12], we give polynomial-time algorithms for finding a super-stability matching and a strongly stable matching in a matroid generalization of the hospitals/residents problem with ties and master lists, i.e., a partial positive answer for the above question.

It should be noted that Abraham, Irving, and Manlove [1] posed an open problem related to the super-stability and the strongly stability in the student-project allocation problem that is a variation of the hospitals/residents problem having hierarchical capacity constraints (see also [11, p.268]). As pointed in [1], these hierarchical capacity constraints can be expressed by using a matroid. However, since the definitions of the super-stability and the strong stability in the student-project allocation problem with ties were not explicitly given in [1], we do not discuss about the relationship between their model and our model. Instead, we give a concrete example in the student-project allocation setting that our model can express.

The rest of this paper is organized as follows. In Section 2, we formally define our problem and give basics of matroids. In Sections 3 and 4, we give polynomial-time algorithms for finding a super-stable matching and a strongly stable matching in our model, respectively. In Section 5, we give examples of concrete models that our model can express.

2 Preliminaries

For each set X and each element x , we define $X + x := X \cup \{x\}$ and $X - x := X \setminus \{x\}$. A pair $\mathbf{M} = (U, \mathcal{I})$ is called a *matroid*, if U is a finite set and $\mathcal{I}$ is a family of subsets of U satisfying the following conditions.

(I0) $\emptyset \in \mathcal{I}$.

(I1) If $I \in \mathcal{I}$ and $J \subseteq I$, then $J \in \mathcal{I}$.

(I2) If $I, J \in \mathcal{I}$ and $|I| < |J|$, then $I + u \in \mathcal{I}$ for some element u in $J \setminus I$.

2.1 Problem formulation

In this paper, we are given a simple bipartite graph $G = (R \cup H, E)$ such that its vertex set is partitioned into two disjoint subsets R and H , and every edge in E connects a vertex in R and a vertex in H . We call a vertex in R (resp., H) a *resident* (resp., a *hospital*). For each vertex v in $R \cup H$ and each subset F of E , we denote by $F(v)$ the set of edges in F incident to v . For each resident r in R and each hospital h in H , if there exists an edge in E connecting r and h , then we denote by (r, h) this edge. Furthermore, we are given a matroid $\mathbf{N} = (E, \mathcal{F})$ such that

$\{e\} \in \mathcal{F}$ for every edge e in E . The matroid $\mathbf{N}$ represents constraints on assignment of residents to hospitals. For example, it can express capacity constraints in the hospitals/residents problem (see Section 5). For each resident r in R , we are given a reflexive and transitive binary relation $\lesssim_r$ on $E(r) \cup \{\emptyset\}$ such that at least one of $e \lesssim_r f$ and $f \lesssim_r e$ holds for every edges e, f in $E(r)$, and $e \lesssim_r \emptyset$ and $\emptyset \not\lesssim_r e$ for every edge e in $E(r)$. For each resident r in R , the binary relation $\lesssim_r$ represents the preference list of r . Namely, for each resident r in R and each edges e, f in $E(r)$, $e \lesssim_r f$ means that r prefers e to f , or is indifferent between e and f . For each resident r in R and each edges e, f in $E(r)$, we write $e \succ_r f$ (resp., $e \sim_r f$), if $e \lesssim_r f$ and $e \not\lesssim_r f$ (resp., $e \lesssim_r f$ and $f \lesssim_r e$). Furthermore, we are given a reflexive and transitive binary relation $\lesssim_H$ on R such that at least one of $r \lesssim_H s$ and $s \lesssim_H r$ holds for every residents r, s in R . The binary relation $\lesssim_H$ represents the master preference list of hospitals in H . We define the notations $\succ_H$ and $\sim_H$ in the same way as $\succ_r$ and $\sim_r$ for a resident r in R .

A subset M of E is called a *matching* in G , if

(M1) $|M(r)| \leq 1$ for every resident r in R , and

(M2) $M \in \mathcal{F}$.

For each matching M in G and each resident r in R such that $M(r) \neq \emptyset$, we do not distinguish between $M(r)$ and its unique element. For each matching M in G and each edge $e = (r, h)$ in $E \setminus M$, we say that r *weakly* (resp., *strictly*) *prefers* e on M , if $e \lesssim_r M(r)$ (resp., $e \succ_r M(r)$). In addition, for each matching M in G and each edge $e = (r, h)$ in $E \setminus M$, we say that H *weakly* (resp., *strictly*) *prefers* e on M , if

(P1) $M + e \in \mathcal{F}$, and/or

(P2) there exists an edge $f = (s, p)$ in M such that $M + e - f \in \mathcal{F}$ and $r \lesssim_H s$ (resp., $r \succ_H s$).

A matching M in G is said to be *super-stable*, if there exists no edge (r, h) in $E \setminus M$ such that r and H weakly prefer (r, h) on M . Furthermore, a matching M in G is said to be *strongly stable*, if there exists no edge (r, h) in $E \setminus M$ such that

(S1) r and H weakly prefer (r, h) on M , and

(S2) at least one of r and H strictly prefers (r, h) on M .

Unfortunately, it is known [5] that there may exist no matching satisfying the above conditions. Thus, our goal is to give a polynomial-time algorithm for deciding whether there exists a matching satisfying the above conditions, and find such a matching, if one exists.

2.2 Basics of matroids

Let $\mathbf{M} = (U, \mathcal{I})$ be a matroid. A subset of U belonging to $\mathcal{I}$ is called an *independent set* of $\mathbf{M}$. A subset C of U is called a *circuit* of $\mathbf{M}$, if C is not an independent set of $\mathbf{M}$, but every proper subset of C is an independent set of $\mathbf{M}$. Notice that the condition (I1) implies that for every subset X of U , if there exists a circuit of $\mathbf{M}$ that is a subset of X , then $X \notin \mathcal{I}$. Let I and u be an independent set of $\mathbf{M}$ and an element in $U \setminus I$ such that $I + u \notin \mathcal{I}$, respectively. It is known [13, Proposition 1.1.6] that there exists a unique circuit of $\mathbf{M}$ that is a subset of $I + u$, and u belongs to this unique circuit. This circuit is called the *fundamental circuit* of u , and denoted by $C_{\mathbf{M}}(u, I)$. Define

$$C_{\mathbf{M}}^-(u, I) := C_{\mathbf{M}}(u, I) - u.$$

It is known [13, p.20, Exercise 5] that we have

$$\mathbf{C}_{\mathbf{M}}(u, I) = \{w \in I + u \mid I + u - w \in \mathcal{I}\}.$$

It should be noted that for each subset M of E and each edge e in $E \setminus M$ such that $M + e \notin \mathcal{F}$, the condition (P2) can be restated as follows.

- There exists an edge $f = (s, p)$ in $\mathbf{C}_{\mathbf{N}}^-(e, M)$ such that $r \succsim_H s$ (resp., $r \succ_H s$).

A maximal independent set of $\mathbf{M}$ is called a *base* of $\mathbf{M}$. The condition (I2) implies that every base of $\mathbf{M}$ has the same size. For each subset X of U , we define

$$\mathcal{I}|X := \{J \subseteq X \mid J \in \mathcal{I}\}, \quad \mathbf{M}|X := (X, \mathcal{I}|X).$$

It is known [13, p.20] that for each subset X of U , $\mathbf{M}|X$ is a matroid. For each subset X of U , we define $r_{\mathbf{M}}(X)$ as the size of a base of $\mathbf{M}|X$. It is not difficult to see that for every subset B of U , B is a base of $\mathbf{M}$ if and only if $|B| = r_{\mathbf{M}}(U)$. For each disjoint subsets X, J of U , we define

$$p(J; X) := r_{\mathbf{M}}(J \cup X) - r_{\mathbf{M}}(X).$$

For each subset X of U , we define

$$\mathcal{I}/X := \{J \subseteq U \setminus X \mid p(J; X) = |J|\}, \quad \mathbf{M}/X := (U \setminus X, \mathcal{I}/X).$$

It is known [13, Proposition 3.1.6] that for each subset X of U , $\mathbf{M}/X$ is a matroid.

The following property of circuits is used in the sequel.

Theorem 1 (See, e.g., [13, p.15, Exercise 14]). *Assume that we are given a matroid $\mathbf{M} = (U, \mathcal{I})$ and circuits C_1, C_2 of $\mathbf{M}$ such that $C_1 \cap C_2 \neq \emptyset$ and $C_1 \setminus C_2 \neq \emptyset$. Then, for every element u in $C_1 \cap C_2$ and every element w in $C_1 \setminus C_2$, there exists a circuit C of $\mathbf{M}$ such that $w \in C$ and C is a subset of $(C_1 \cup C_2) - u$.*

Although the following lemmas easily follow, we give their proofs for completeness.

Lemma 2. *Assume that we are given a matroid $\mathbf{M} = (U, \mathcal{I})$ and independent sets I, J of $\mathbf{M}$ such that $I \subseteq J$. Then, for every element u in $U \setminus J$ such that $I + u \notin \mathcal{I}$, we have $J + u \notin \mathcal{I}$ and $\mathbf{C}_{\mathbf{M}}(u, I) = \mathbf{C}_{\mathbf{M}}(u, J)$.*

Proof. Let u be an element in $U \setminus J$ such that $I + u \notin \mathcal{I}$. Since $I \subseteq J$, $\mathbf{C}_{\mathbf{M}}(u, I)$ is a subset of $J + u$, which implies that $J + u \notin \mathcal{I}$. Define $C_1 := \mathbf{C}_{\mathbf{M}}(u, I)$ and $C_2 := \mathbf{C}_{\mathbf{M}}(u, J)$. Assume that $C_1 \neq C_2$. Since u belongs to $C_1 \cap C_2$, Theorem 1 implies that there exists a circuit C of $\mathbf{M}$ such that C is a subset of $(C_1 \cup C_2) - u$. In addition, since $C_1 - u \subseteq I \subseteq J$ and $C_2 - u \subseteq J$, we have $C \subseteq J$. This contradicts the fact that $J \in \mathcal{I}$. $\square$

Lemma 3. *Assume that we are given a matroid $\mathbf{M} = (U, \mathcal{I})$, circuits $C, C_1, C_2, \dots, C_k$ of $\mathbf{M}$, and elements $u_1, u_2, \dots, u_k$ in U such that $u_i \in C \cap C_i$ for every $i = 1, 2, \dots, k$. In addition, we assume that there exists an element w in U such that $w \in C$ and $w \notin C_i$ for every $i = 1, 2, \dots, k$. Then, there exists a circuit C' of $\mathbf{N}$ such that $C' \subseteq (C \cup C_1 \cup C_2 \cup \dots \cup C_k) \setminus \{u_1, u_2, \dots, u_k\}$.*

Proof. We consider the following procedure.

Step 1. Set $t := 1$ and $K_0 := C$.

Step 2. If $t \leq k$, then do the following steps. Otherwise, go to **Step 3**.

- (2-a) If $u_t \notin K_{t-1}$, then go to **Step (2-c)**.
- (2-b) Find a circuit K_t of $\mathbf{N}$ such that $w \in K_t$ and K_t is a subset of $(K_{t-1} \cup C_t) - u_t$.
- (2-c) Update $t := t + 1$ and go back to the beginning of **Step 2**.

Step 3. Output K_k and halt.

Notice that Theorem 1 implies that in **Step (2-b)** there exists a circuit K_t satisfying the above properties. It is not difficult to see that K_k is a subset of $C \cup C_1 \cup C_2 \cup \dots \cup C_k$ and $u_i \notin K_k$ for every $i = 1, 2, \dots, k$. This completes the proof. $\square$

Lemma 4. Assume that we are given a matroid $\mathbf{M} = (U, \mathcal{I})$ and a subset X of U .

1. Let B be an arbitrary base of $\mathbf{M}|X$. For every subset I of $U \setminus X$, I is an independent set of $\mathbf{M}/X$ if and only if $I \cup B$ is an independent set of $\mathbf{M}$.
2. For every base B_1 of $\mathbf{M}|X$ and every base B_2 of $\mathbf{M}/X$, $B_1 \cup B_2$ is a base of $\mathbf{M}$.
3. For every subset I of U such that $I \cap X$ is a base of $\mathbf{M}|X$, if $I \setminus X$ is not an independent set of $\mathbf{M}/X$, then I is not an independent set of $\mathbf{M}$.

Proof. The statement (1) follows from Proposition 3.1.7 of [13]. Next we consider the statement (2). Assume that $B_1 \cup B_2$ is not a base of $\mathbf{M}$. Then, there exists an element u in $U \setminus (B_1 \cup B_2)$ such that $(B_1 \cup B_2) + u$ is an independent set of $\mathbf{M}$. If u belongs to X , then this contradicts the fact that B_1 is a base of $\mathbf{M}|X$. Thus, u does not belong to X . Since B_1 is a base of $\mathbf{M}$, the statement (1) implies that $B_2 + u$ is an independent set of $\mathbf{M}/X$. This contradicts the fact that B_2 is a base of $\mathbf{M}/X$. Lastly we prove the statement (3). Assume that $I \in \mathcal{I}$. Since $I \cap X$ is a base of $\mathbf{M}|X$, the statement (1) implies that $I \setminus X$ is an independent set of $\mathbf{M}/X$. This contradicts the fact that $I \setminus X$ is not an independent set of $\mathbf{M}/X$. $\square$

Let $\mathbf{M}_1 = (U, \mathcal{I}_1)$ and $\mathbf{M}_2 = (U, \mathcal{I}_2)$ be matroids on the same ground set. A subset I of U is called a *common independent set* of $\mathbf{M}_1$ and $\mathbf{M}_2$, if I belongs to $\mathcal{I}_1 \cap \mathcal{I}_2$. It is known that we can find a maximum-size common independent set in time bounded by a polynomial in $|U|$ and γ , where γ is the time required to decide whether X is a common independent set of $\mathbf{M}_1$ and $\mathbf{M}_2$ for one subset X of U . If we use the algorithm proposed by Cunningham [2], then we can find a maximum-size common independent set in $O(|U|^{2.5}\gamma)$ time.

3 Super-Stable Matchings

In this section, we consider the problem of deciding whether there exists a super-stable matching in G , and find such a matching, if one exists. We first partition R into non-empty disjoint subsets $R_1, R_2, \dots, R_n$ such that

1. $r \sim_H s$ for every $i = 1, 2, \dots, n$ and every residents r, s in R_i , and
2. $r \succ_H s$ for every $i, j = 1, 2, \dots, n$ such that $i < j$ and every residents r in R_i and s in R_j .

For each $i, j = 1, 2, \dots, n$, we define $R_{i,j} := R_i \cup R_{i+1} \cup \dots \cup R_j$.

Our algorithm **SuperM** is described as follows.

Algorithm SuperM

Step 1. Define $i := 1$, $M_0 := \emptyset$, and $D_0 := \emptyset$.

Step 2. If $i \leq n$, then do the following steps. Otherwise, go to **Step 3**.

(2-a) For each resident r in R_i , we define T_r by

$$T_r := \left\{ e \in E(r) \setminus D_{i-1} \mid \forall f \in E(r) \setminus D_{i-1}: e \succsim_r f \right\}.$$

(2-b) If there exists a resident r in R_i such that $|T_r| > 1$, then output **null** and halt (i.e., there exists no super-stable matching in G).

(2-c) Define $E_i := \cup_{r \in R_i} T_r$. If $M_{i-1} \cup E_i \notin \mathcal{F}$, then output **null** and halt.

(2-d) Define $M_i := M_{i-1} \cup E_i$.

(2-e) Define

$$L_i := \left\{ (r, h) \in E \setminus D_{i-1} \mid r \in R_{i+1,n}, M_i + (r, h) \notin \mathcal{F} \right\}.$$

Furthermore, define $D_i := D_{i-1} \cup L_i$.

(2-f) Update $i := i + 1$ and go back to the beginning of **Step 2**.

Step 3. Output M_n and halt (i.e., M_n is a super-stable matching in G).

End of Algorithm

In the sequel, we prove the correctness of the algorithm **SuperM**.

Lemma 5. Assume that the algorithm **SuperM** halts when $i = \delta$. Then, for any edge e in $D_{\delta-1}$, there does not exist a super-stable matching N in G such that $e \in N$.

Proof. An edge e in $D_{\delta-1}$ is said to be *bad*, if there exists a super-stable matching N in G such that $e \in N$. We prove this lemma by contradiction. Assume that there exists a bad edge in $D_{\delta-1}$. Define Δ as the set of integers ℓ in $\{1, 2, \dots, \delta - 1\}$ such that there exists a bad edge in L_ℓ . Let j be the minimum integer in Δ . Let $e = (r, h)$ and N be a bad edge in L_j and a super-stable matching such that $e \in N$, respectively. Since e belongs to L_j , $M_j + e \notin \mathcal{F}$. Define $C := C_N(e, M_j)$. If C is a subset of N , then this contradicts the fact that $N \in \mathcal{F}$. Thus, $C \setminus N$ is not empty. For every edge (s, p) in $C \setminus N$, since $r \in R_{j+1,k}$ and $s \in R_{1,j}$, we have $s \succ_H r$.

We first assume that there exists an edge $f = (s, p)$ in $C \setminus N$ such that $N(s) \succ_s f$. Since e belongs to $C \cap N$, we have $e \neq f$. This implies that f belongs to M_j , i.e., $f \in T_s$. Thus, the definition of the algorithm **SuperM** implies that $N(s)$ belongs to D_{j-1} , i.e., $N(s) \in L_k$ for an integer k in $\{1, 2, \dots, j - 1\}$. Furthermore, since $D_{j-1} \subseteq D_{\delta-1}$ and N is super-stable, $N(s)$ is bad. This contradicts the minimality of j .

Next we assume that $f \succsim_s N(s)$ for every edge $f = (s, p)$ in $C \setminus N$. If there exists an edge f in $C \setminus N$ such that $N + f \in \mathcal{F}$, then this contradicts the fact that N is super-stable. Thus, we assume that $N + f \notin \mathcal{F}$ for every edge f in $C \setminus N$. For each edge f in $C \setminus N$, we define C_f as $C_N(f, N)$. Since N is super-stable, $t \succ_H s$ for every edge $f = (s, p)$ in $C \setminus N$ and every edge $g = (t, q)$ in $C_f - f$. For every edge $f = (s, p)$ in $C \setminus N$, since $s \succ_H r$, this implies that $e \notin C_f$. In addition, $f \in C \cap C_f$ for every edge f in $C \setminus N$. Thus, Lemma 3 implies that there exists a circuit C' of $\mathbf{N}$ such that C' is a subset of $(C \cup C^*) \setminus (C \setminus N)$, where $C^* = \cup_{f \in C \setminus N} C_f$. Thus, since $C_f - f$ is a subset of N , C' is a subset of N . This contradicts the fact that $N \in \mathcal{F}$. $\square$

Lemma 6. Assume that the algorithm **SuperM** halts when $i = \delta$, and we are given a super-stable matching N in G . Then, for every resident r in $R_{1,\delta}$ such that $T_r = \emptyset$ (resp., $T_r \neq \emptyset$), we have $N(r) = \emptyset$ (resp., $N(r) \in T_r$).

Proof. For every resident r in $R_{1,\delta}$ such that $T_r = \emptyset$, every edge in $E(r)$ belongs to $D_{\delta-1}$. Thus, Lemma 5 implies that $N(r) = \emptyset$ for every resident r in $R_{1,\delta}$ such that $T_r = \emptyset$.

From here, we consider a resident r in $R_{1,\delta}$ such that $T_r \neq \emptyset$. We assume that r belongs to R_j for an integer j in $\{1, 2, \dots, \delta\}$. If $N(r) \in T_r$, then the proof is done. Thus, we assume that $N(r)$ does not belong to T_r , and we prove this lemma by contradiction. If $N(r) \succsim_r e$ for an edge e in T_r (notice that $f \sim_r g$ for every edges f, g in T_r), then the definition of T_r implies that $N(r)$ belongs to $D_{\delta-1}$, which contradicts Lemma 5. Thus, we assume that $e \succ_r N(r)$ for an edge e in T_r . In addition, we assume that there exists no resident s in $R_{1,j-1}$ such that $T_s \neq \emptyset$ and $e \succ_s N(s)$ for an edge e in T_s . Notice that such a resident r clearly exists.

Let e be an edge in T_r . Since e belongs to T_r , e does not belong to D_{j-1} . Since this implies that $e \notin D_{j-2}$ and $e \notin L_{j-1}$, we have $M_{j-1} + e \in \mathcal{F}$. Since $N(r)$ does not belong to T_r , e does not belong to N . Furthermore, since N is super-stable, $N + e \notin \mathcal{F}$ and $s \succ_H r$ for every edge (s, p) in $\mathbf{C}_{\mathbf{N}}^-(e, N)$. Here we prove that $\mathbf{C}_{\mathbf{N}}^-(e, N)$ is a subset of $M_{j-1} + e$. This contradicts the fact that $M_{j-1} + e \in \mathcal{F}$. Let $f = (s, p)$ be an edge in $\mathbf{C}_{\mathbf{N}}^-(e, N)$. Since $s \succ_H r$, s belongs to R_k for an integer k in $\{1, 2, \dots, j-1\}$. This and the definition of the algorithm **superM** imply that $|T_s| \leq 1$. If $T_s = \emptyset$, then in the same way as above we can prove that $N(s) = \emptyset$, which contradict $f = N(s)$. Thus, $|T_s| = 1$. In this case, the unique edge in T_s belongs to M_{j-1} . Thus, if $f \in T_s$, then the proof is done. Assume that f does not belong to T_s . In this case, since $f = N(s)$, the definition of r implies that $N(s) \succsim_s g$ for an edge g in T_s . This implies that f belongs to $D_{\delta-1}$. Since f belongs to a super-stable matching N , this contradicts Lemma 5. $\square$

Lemma 7. *If the algorithm **SuperM** outputs M_n , then M_n is a super-stable matching in G .*

Proof. For every $\ell = 1, 2, \dots, n$ and every resident r in R_ℓ , $|E_\ell(r)| \leq 1$. Thus, since $M_\ell \in \mathcal{F}$ for every $\ell = 1, 2, \dots, n$, M_n is a matching in G . What remains is to prove that M_n is super-stable. Let $e = (r, h)$ be an edge in $E \setminus M_n$. We prove that at least one of r and H does not weakly prefer e on M_n . If $M_n(r) \succ_r e$, then the proof is done. Thus, we consider the case where $e \succsim_r M_n(r)$. Assume that r belongs to R_j for an integer j in $\{1, 2, \dots, n\}$. Since $M_n(r)$ belongs to T_r and $|T_r| = 1$, the definition of the algorithm **SuperM** implies that e belongs to L_k for an integer k in $\{1, 2, \dots, j-1\}$. Thus, $M_k + e \notin \mathcal{F}$ and $s \succ_H r$ for every edge (s, p) in $\mathbf{C}_{\mathbf{N}}^-(e, M_k)$. Since M_k is a subset of M_n , Lemma 2 implies that $M_n + e \notin \mathcal{F}$ and $\mathbf{C}_{\mathbf{N}}^-(e, M_k) = \mathbf{C}_{\mathbf{N}}^-(e, M_n)$. Thus, $s \succ_H r$ for every edge $f = (s, p)$ in $\mathbf{C}_{\mathbf{N}}^-(e, M_n)$. This completes the proof. $\square$

Lemma 8. *If the algorithm **SuperM** outputs **null**, then there exists no super-stable matching in G .*

Proof. We assume that the algorithm **SuperM** outputs **null** when $i = \delta$. We prove this lemma by contradiction. Assume that there exists a super-stable matching N in G .

We first consider the case where in **Step (2-b)** the algorithm **SuperM** outputs **null**. In this case, there exists a resident r in R_δ such that $|T_r| > 1$. Lemma 6 implies that $N(r)$ belongs to T_r . Let e be an edge in T_r such that $e \neq N(r)$. Since e and $N(r)$ belong to T_r , we have $e \sim_r N(r)$. Thus, what remains is to prove that $N + e \in \mathcal{F}$ and/or there exists an edge $f = (s, p)$ in N such that $N + e - f \in \mathcal{F}$ and $r \succsim_H s$, which contradicts the fact that N is super-stable. If $N + e \in \mathcal{F}$, then proof is done. Thus, we assume that $N + e \notin \mathcal{F}$ and $s \succ_H r$ (i.e., s belongs to $R_{1,\delta-1}$) for every edge (s, p) in $\mathbf{C}_{\mathbf{N}}^-(e, N)$. Since $e \in T_r$, $M_{\delta-1} + e \in \mathcal{F}$. Lemma 6 and the definition of the algorithm **SuperM** imply that $N(s) = M_{\delta-1}(s)$ for every resident s in $R_{1,\delta-1}$. These imply that $\mathbf{C}_{\mathbf{N}}^-(e, N)$ is a subset of $M_{\delta-1} + e$, which contradicts the fact that $M_{\delta-1} + e \in \mathcal{F}$.

Next we consider the case where in **Step (2-c)** the algorithm **SuperM** outputs **null**. That is, $M_{\delta-1} \cup E_\delta \notin \mathcal{F}$. Lemma 6 and the definition of the algorithm of **SuperM** imply that $M_{\delta-1}(r) =$

$N(r)$ for every resident r in $R_{1,\delta-1}$ and $E_\delta(r) = N(r)$ for every resident r in R_δ . These imply that $M_{\delta-1} \cup E_\delta$ is a subset of N , which contradicts the fact that $N \in \mathcal{F}$. $\square$

Lemmas 7 and 8 imply the following theorem.

Theorem 9. *The algorithm **SuperM** can decide whether there exists a super-stable matching in G , and find such a matching, if one exists.*

Here we consider the time complexity of **SuperM**. Define $m := |E|$, and we denote by γ the time required to decide whether $F \in \mathcal{F}$ for one subset F of E . For simplicity, we assume that $\gamma = \Omega(m)$ and $m \geq |R|$. Since the number of iterations of **Step 2** is at most m , we consider the time complexity of one iteration of **Step 2**. Clearly, **Step (2-e)** is the bottleneck and the time complexity of this step is $O(m\gamma)$. Thus, the time complexity of the algorithm **SuperM** is $O(m^2\gamma)$.

4 Strongly Stable Matchings

In this section, we give an algorithm for the problem of deciding whether there exists a strongly stable matching in G , and find such a matching, if one exists. We define subsets $R_1, R_2, \dots, R_n$ of R and the notation $R_{i,j}$ in the same way as in Section 3.

Our algorithm **StrongM** is described as follows.

Algorithm StrongM

Step 1. Define $i := 1$, $M_0 := \emptyset$, $D_0 := \emptyset$, and $P_0 := \emptyset$.

Step 2. If $i \leq n$, then do the following steps. Otherwise, go to **Step 3**.

(2-a) For each resident r in R_i , we define T_r by

$$T_r := \left\{ e \in E(r) \setminus D_{i-1} \mid \forall f \in E(r) \setminus D_{i-1}: e \succ_r f \right\}.$$

(2-b) Define

$$E_i := \bigcup_{r \in R_i} T_r, \quad P_i := \bigcup_{\ell=1,2,\dots,i} E_\ell, \quad \mathbf{N}_i := (\mathbf{N} \setminus P_i) \setminus P_{i-1}, \quad R_i^* := \{a \in R_i \mid T_a \neq \emptyset\}.$$

(2-c) If $r_{\mathbf{N}_i}(E_i) > |R_i^*|$, then output **null** and halt.

(2-d) Define

$$\mathcal{U}_i := \left\{ F \subseteq E_i \mid \forall r \in R_i: |F(r)| \leq 1 \right\}, \quad \mathbf{A}_i := (E_i, \mathcal{U}_i).$$

Notice that $\mathbf{A}_i$ is a matroid. Then, find a maximum-size common independent set F_i of $\mathbf{A}_i$ and $\mathbf{N}_i$.

(2-e) If $|F_i| < |R_i^*|$, then output **null** and halt. Otherwise, define $M_i := M_{i-1} \cup F_i$.

(2-f) Define

$$L_i := \left\{ (r, h) \in E \setminus D_{i-1} \mid r \in R_{i+1,n}, M_i + (r, h) \notin \mathcal{F} \right\}.$$

Furthermore, define $D_i := D_{i-1} \cup L_i$.

(2-g) Update $i := i + 1$ and go to the beginning of **Step 2**.

Step 3. Output M_n and halt.

End of Algorithm

In the sequel, we prove the correctness of the algorithm **StrongM**.

Lemma 10. *Assume that the algorithm **StrongM** halts when $i = \delta$. Then,*

1. F_ℓ is a base of $\mathbf{N}_\ell$ for every $\ell = 1, 2, \dots, \delta - 1$, and
2. M_ℓ is a base of $\mathbf{N}|P_\ell$ for every $\ell = 1, 2, \dots, \delta - 1$.

Proof. We first consider the statement (1). Let us fix an integer ℓ in $\{1, 2, \dots, \delta - 1\}$. Since the algorithm **StrongM** does not output **null** when $i = \ell$, we have $r_{\mathbf{N}_\ell}(E_\ell) \leq |R_\ell^*|$ and $|F_\ell| \geq |R_\ell^*|$. These imply that $r_{\mathbf{N}_\ell}(E_\ell) \leq |F_\ell|$. Furthermore, since F_ℓ is an independent set of $\mathbf{N}_\ell$, we have $|F_\ell| \leq r_{\mathbf{N}_\ell}(E_\ell)$. These imply that $|F_\ell| = r_{\mathbf{N}_\ell}(E_\ell)$, i.e., F_ℓ is a base of $\mathbf{N}_\ell$.

Next we prove the statement (2) by induction on ℓ . The statement (2) for $\ell = 1$ is equivalent to the statement (1) for $\ell = 1$. Let ξ be an integer in $\{1, 2, \dots, \delta - 2\}$. Assume that the statement (2) holds for the case of $\ell = \xi$, and we consider the case of $\ell = \xi + 1$. Since $\mathbf{N}|P_\xi = (\mathbf{N}|P_{\xi+1})|P_\xi$ and $F_{\xi+1}$ is a base of $\mathbf{N}_{\xi+1} = (\mathbf{N}|P_{\xi+1})/P_\xi$, the induction assumption and Lemma 4(2) imply that $M_{\xi+1}$ is a base of $\mathbf{N}|P_{\xi+1}$. $\square$

Lemma 11. *Assume that the algorithm **StrongM** halts when $i = \delta$. Then, for any edge e in $D_{\delta-1}$, there does not exist a strongly stable matching N in G such that $e \in N$.*

Proof. We say that an edge e in $D_{\delta-1}$ is *bad*, if there exists a strongly stable matching N in G such that $e \in N$. We prove this lemma by contradiction. That is, we assume that there exists a bad edge in $D_{\delta-1}$. Define Δ as the set of integers ℓ in $\{1, 2, \dots, \delta - 1\}$ such that there exists a bad edge in L_ℓ . Let j be the minimum integer in Δ . Furthermore, let $e = (r, h)$ and N be a bad edge in L_j and a strongly stable matching such that $e \in N$, respectively. Since e belongs to L_j , $M_j + e \notin \mathcal{F}$. Define $C := \mathbf{C}_{\mathbf{N}}(e, M_j)$ (notice that $M_j \in \mathcal{F}$ follows from Lemma 10(2)). If C is a subset of N , then this contradicts the fact that $N \in \mathcal{F}$. Thus, $C \setminus N$ is not empty. For every edge (s, p) in $C \setminus N$, since $r \in R_{j+1, k}$ and $s \in R_{1, j}$, we have $s \succ_H r$.

We first assume that there exists an edge $f = (s, p)$ in $C \setminus N$ such that $N(s) \succ_s f$. Since e belongs to N , $e \neq f$. This implies that f belongs to M_j , i.e., $f \in T_s$. Thus, the definition of the algorithm **StrongM** implies that $N(s)$ belongs to D_{j-1} , i.e., $N(s) \in L_k$ for an integer k in $\{1, 2, \dots, j - 1\}$. Furthermore, since $D_{j-1} \subseteq D_{\delta-1}$ and N is strongly stable, $N(s)$ is bad, which contradicts the minimality of j .

Next we assume that $f \not\succ_s N(s)$ for every edge $f = (s, p)$ in $C \setminus N$. If there exists an edge f in $C \setminus N$ such that $N + f \in \mathcal{F}$, then this contradicts the fact that N is strongly stable. Thus, we assume that $N + f \notin \mathcal{F}$ for every edge f in $C \setminus N$. For each edge f in $C \setminus N$, we define C_f as $\mathbf{C}_{\mathbf{N}}(f, N)$. Since N is strongly stable, $t \succ_H s$ for every edge $f = (s, p)$ in $C \setminus N$ and every edge $g = (t, q)$ in $C_f - f$. For every edge $f = (s, p)$ in $C \setminus N$, since $s \succ_H r$, this implies that $e \notin C_f$. In addition, $f \in C \cap C_f$ for every edge f in $C \setminus N$. Thus, Lemma 3 implies that there exists a circuit C' of $\mathbf{N}$ such that C' is a subset of N . This contradicts the fact that $N \in \mathcal{F}$. $\square$

Lemma 12. *Assume that the algorithm **StrongM** halts when $i = \delta$, and we are given a strongly stable matching N in G . Then, for every resident r in $R_{1, \delta}$ such that $T_r = \emptyset$ (resp., $T_r \neq \emptyset$), we have $N(r) = \emptyset$ (resp., $N(r) \in T_r$).*

Proof. For every resident r in $R_{1,\delta}$ such that $T_r = \emptyset$, every edge in $E(r)$ belongs to $D_{\delta-1}$. Thus, Lemma 11 implies that $N(r) = \emptyset$ for every resident r in $R_{1,\delta}$ such that $T_r = \emptyset$.

From here, we consider a resident r in $R_{1,\delta}$ such that $T_r \neq \emptyset$. We assume that r belongs to R_j for an integer j in $\{1, 2, \dots, \delta\}$. If $N(r) \in T_r$, then the proof is done. Thus, we assume that $N(r)$ does not belong to T_r , and we prove this lemma by contradiction. If $N(r) \succ_r e$ for an edge e in T_r , then the definition of T_r implies that $N(r)$ belongs to $D_{\delta-1}$, which contradicts Lemma 11. Thus, we assume that $e \succ_r N(r)$ for an edge e in T_r . Furthermore, we assume that there exists no resident s in $R_{1,j-1}$ such that $T_s \neq \emptyset$ and $e \succ_s N(s)$ for an edge e in T_s . Notice that such a resident r clearly exists.

Let e be an edge in T_r . Since e belongs to T_r , $M_{j-1} + e \in \mathcal{F}$. Since $N(r)$ does not belong to T_r , e does not belong to N . In addition, since N is strongly stable, $N + e \notin \mathcal{F}$ and $s \succ_H r$ for every edge (s, p) in $C_N^-(e, N)$. Define $C := C_N(e, N)$. Since $M_{j-1} + e \in \mathcal{F}$, C is not a subset of $M_{j-1} + e$. Let $f = (s, p)$ be an edge in $C \setminus (M_{j-1} + e)$. Since $s \succ_H r$, $s \in R_k$ for an integer k in $\{1, 2, \dots, j-1\}$. In the same way as above, we can prove that $f \in T_s$ or $g \succ_s f$ for an edge g in T_s . The assumption of r implies that f belongs to T_s , i.e., $f \in E_k$. Since $f \notin (M_{j-1} + e)$ implies that f does not belong to F_k and Lemma 10(1) implies that F_k is a base of $\mathbf{N}_k$, $F_k + f$ is not an independent set of $\mathbf{N}_k = (\mathbf{N}|P_k)/P_{k-1}$. Thus, since Lemma 10(2) implies that M_{k-1} is a base of $(\mathbf{N}|P_k)/P_{k-1}$, Lemma 4(3) implies that $M_k + f \notin \mathcal{F}$. Since $M_k \subseteq M_{j-1}$, $M_{j-1} + f \notin \mathcal{F}$. For each edge f in $C \setminus (M_{j-1} + e)$, we define $C_f := C_N(f, M_{j-1})$. For every edge f in $C \setminus (M_{j-1} + e)$, since $e \notin M_{j-1}$ and $f \neq e$, e does not belong to C_f . Furthermore, $f \in C \cap C_f$ for every edge f in $C \setminus (M_{j-1} + e)$. Thus, Lemma 3 implies that there exists a circuit C' of $\mathbf{N}$ that C' is a subset of $M_{j-1} + e$, which contradicts the fact that $M_{j-1} + e \in \mathcal{F}$. $\square$

Lemma 13. *Assume that the algorithm StrongM halts when $i = \delta$, and we are given a strongly stable matching M . Then, for every $\ell = 1, 2, \dots, \delta$, the following statements hold.*

1. *If $\ell > 1$, then $N \cap P_{\ell-1}$ is a base of $\mathbf{N}|P_{\ell-1}$.*
2. *$N \cap E_\ell$ is a base of $\mathbf{N}_\ell$.*

Proof. We prove this lemma by induction on ℓ . We first consider the case of $\ell = 1$, and we prove this case by contradiction. Assume that $N \cap E_1$ is not a base of $\mathbf{N}_1 = \mathbf{N}|E_1$. Then, since $N \in \mathcal{F}$, $N \cap E_1$ is an independent set of $\mathbf{N}_1$. Since $N \cap E_1$ is not a base of $\mathbf{N}_1$, there exists an edge $e = (r, h)$ in $E_1 \setminus N$ such that $(N \cap E_1) + e \in \mathcal{F}$. Since $D_0 = \emptyset$ and $e \in T_r$, we have $e \succ_r N(r)$. Thus, if $N + e \in \mathcal{F}$, then this contradicts the fact that N is strongly stable. Thus, we assume that $N + e \notin \mathcal{F}$. Then, since $(N \cap E_1) + e \in \mathcal{F}$, $C_N(e, N)$ is not a subset of $(N \cap E_1) + e$. Let $f = (s, p)$ be an edge in $C_N(e, N) \setminus ((N \cap E_1) + e)$. If $s \in R_j$ for an integer j in $\{2, 3, \dots, n\}$, then $r \succ_H s$. This contradicts the fact that N is strongly stable. Thus, we have $s \in R_1$. Since $f \in E(s)$ (i.e., $E(s) \neq \emptyset$) and $D_0 = \emptyset$, we have $T_s \neq \emptyset$. Thus, since $f \in N$, Lemma 12 implies that $f \in T_s$, i.e., $f \in E_1$. This contradicts the fact that $f \notin N \cap E_1$.

Let ξ be an integer in $\{1, 2, \dots, \delta - 2\}$, and assume that this lemma holds for the case of $\ell = \xi$, and we consider the case of $\ell = \xi + 1$. We first prove the statement (1). If $\xi = 1$, then the statement (1) for $\ell = \xi + 1$ is equivalent to the statement (2) for the case of $\ell = 1$. Assume that $\xi \geq 2$. The induction assumption implies that $N \cap P_{\xi-1}$ is a base of $(\mathbf{N}|P_\xi)/P_{\xi-1}$ and $N \cap E_\xi$ is a base of $(\mathbf{N}|P_\xi)/P_{\xi-1}$. Thus, Lemma 4(2) implies that $N \cap P_\xi$ is a base of $\mathbf{N}|P_\xi$, which completes the proof. Next we prove the statement (2) by contradiction. Assume that $N \cap E_{\xi+1}$ is not a base of $\mathbf{N}_{\xi+1}$. Then, there exists an edge $e = (r, h)$ in $E_{\xi+1} \setminus N$ such that $(N \cap E_{\xi+1}) + e$ is an independent set of $\mathbf{N}_{\xi+1} = (\mathbf{N}|P_{\xi+1})/P_\xi$. Since the statement (1) for the case of $\ell = \xi + 1$ implies that $N \cap P_\xi$ is a base of $(\mathbf{N}|P_{\xi+1})/P_\xi$, Lemma 4(1) implies that $(N \cap P_{\xi+1}) + e \in \mathcal{F}$. Since e belongs to $E_{\xi+1}$ (i.e., T_r), Lemma 12 implies that $e \sim_r N(r)$. Thus, if $N + e \in \mathcal{F}$, then

this contradicts the fact that N is strongly stable. Thus, we consider the case where $N + e \notin \mathcal{F}$. Since $(N \cap P_{\xi+1}) + e \in \mathcal{F}$, $\mathbf{C}_N(e, N)$ is not a subset of $(N \cap P_{\xi+1}) + e$. Let $f = (s, p)$ be an edge in $\mathbf{C}_N(e, N) \setminus ((N \cap P_{\xi+1}) + e)$. Since $f = N(s)$, Lemma 12 implies that f belongs to T_s . This and $f \notin P_{\xi+1}$ imply that $s \in R_j$ for an integer j in $\{\xi + 2, \xi + 3, \dots, n\}$. Thus, $r \succ_H s$, which contradicts the fact that N is strongly stable. $\square$

Lemma 14. *If the algorithm **StrongM** outputs M_n , then M_n is a strongly stable matching in G .*

Proof. Since Lemma 10(2) implies that $M_n \in \mathcal{F}$ and $|M_n(r)| \leq 1$ clearly holds for every resident r in R , M_n is a matching in G . What remains is to prove that M_n is strongly stable.

Let $e = (r, h)$ be an edge in $E \setminus M_n$. If $M_n(r) \succ_r e$, then r does not weakly prefer e on M . Thus, we do not need to consider this case. Assume that $e \succ_r M_n(r)$ and $r \in R_j$ for an integer j in $\{1, 2, \dots, n\}$. We first consider the case where $e \succ_r M_n(r)$. In this case, since $M_n(r) \in T_r$, the definition of the algorithm **StrongM** implies that $e \in L_k$ for an integer k in $\{1, 2, \dots, j-1\}$. Thus, $M_k + e \notin \mathcal{F}$ and $s \succ_H r$ for every edge (s, p) in $\mathbf{C}_N^-(e, M_k)$. Since $M_k \subseteq M_n$, Lemma 2 implies that $\mathbf{C}_N(e, M_k) = \mathbf{C}_N(e, M_n)$. Thus, $s \succ_H r$ for every edge $f = (s, p)$ in $\mathbf{C}_N^-(e, M_n)$. This implies that H does not weakly prefer e on M . Lastly we consider the case where $e \sim_r M_n(r)$. We first assume that $e \notin T_r$. In this case, the definition of the algorithm **StrongM** implies that $e \in L_k$ for an integer k in $\{1, 2, \dots, j-1\}$. Thus, we can treat this case in the same way as above. Next we assume that $e \in T_r$. Notice that $e \in E_j \setminus F_j$. Since Lemma 10(1) implies that F_j is a base of $\mathbf{N}_j$, $F_j + e$ is not an independent set of $(\mathbf{N}|P_j)/P_{j-1}$. In addition, Lemma 10(2) implies that M_{j-1} is a base of $(\mathbf{N}|P_j)|P_{j-1}$. Thus, Lemma 4(3) implies that $M_j + e \notin \mathcal{F}$. Since $M_j \subseteq P_j$, we have $s \succ_H r$ for each edge (s, p) in $\mathbf{C}_N(e, M_j)$. Since $M_j \subseteq M_n$, Lemma 2 implies that $M_n + e \notin \mathcal{F}$ and $\mathbf{C}_N(e, M_j) = \mathbf{C}_N(e, M_n)$. Thus, we have $s \succ_H r$ for every edge $f = (s, p)$ in $\mathbf{C}_N^-(e, M_n)$. This completes the proof. $\square$

Lemma 15. *If the algorithm **StrongM** outputs **null**, then there exists no strongly stable matching in G .*

Proof. We assume that the algorithm **StrongM** outputs **null** when $i = \delta$. We prove this lemma by contradiction. Assume that there exists a strongly stable matching N in G .

We first consider the case where in **Step (2-c)** the algorithm **StrongM** outputs **null**. In this case, we have $r_{\mathbf{N}_\delta}(E_\delta) > |R_\delta^*|$. Since Lemma 13(2) implies that $N \cap E_\delta$ is a base of $\mathbf{N}_\delta$, we have $|N \cap E_\delta| = r_{\mathbf{N}_\delta}(E_\delta)$, which implies that $|N \cap E_\delta| > |R_\delta^*|$. However, since N is a matching and E_δ is a subset of $\cup_{r \in R_\delta^*} E(r)$, $|N \cap E_\delta| \leq |R_\delta^*|$, which contradicts the fact that $|N \cap E_\delta| > |R_\delta^*|$.

Next we consider the case where in **Step (2-e)** the algorithm **StrongM** outputs **null**. In this case, we have $|F_\delta| < |R_\delta^*|$. Since N is a matching in G , $|N(r)| \leq 1$ for every resident r in R_δ . Furthermore, Lemma 13(2) implies that $N \cap E_\delta$ is an independent set of $\mathbf{N}_\delta$. Thus, $N \cap E_\delta$ is a common independent set of $\mathbf{A}_\delta$ and $\mathbf{N}_\delta$. Since F_δ is a maximum-size common independent set of $\mathbf{A}_\delta$ and $\mathbf{N}_\delta$, we have $|N \cap E_\delta| \leq |F_\delta| < |R_\delta^*|$. This implies that there exists a resident r in R_δ^* such that $N(r) \notin E_\delta$. However, since $T_r \subseteq E_\delta$, this contradicts Lemma 12. $\square$

Lemmas 14 and 15 imply the following theorem.

Theorem 16. *The algorithm **StrongM** can decide whether there exists a strongly stable matching in G , and find such a matching, if one exists.*

Here we consider the time complexity of **StrongM**. Define $m := |E|$, and we denote by γ the time required to decide whether $F \in \mathcal{F}$ for one subset F of E . For simplicity, we assume that $\gamma = \Omega(m)$ and $m \geq |R|$. Since the number of iterations of **Step 2** is at most m , we consider the time complexity of one iteration of **Step 2**. Clearly, **Step (2-d)** is the bottleneck. Lemma 4(1)

implies that by computing bases of $\mathbf{N}|P_\ell$ for all $\ell = 1, 2, \dots, n$ in $O(m^2\gamma)$ time, we can decide in $O(\gamma)$ time whether F is an independent set of $\mathbf{N}_\ell$ for an integer ℓ in $\{1, 2, \dots, n\}$ and one subset F of E_ℓ . Thus, the time complexity of this step is $O(m^{2.5}\gamma)$, and the time complexity of **SuperM** is $O(m^{3.5}\gamma)$.

5 Examples

In this section, we give examples of concrete models that our model can express.

5.1 Hospitals/residents setting

Assume that for each hospital h in H , we are given a positive integer q_h . Then, we define

$$\mathcal{F} := \left\{ F \subseteq E \mid \forall h \in H: |F(h)| \leq q_h \right\}.$$

It is not difficult to see that $\mathbf{N} = (E, \mathcal{F})$ is a matroid. When we define the matroid $\mathbf{N}$ as above, a subset M of E is a matching if and only if

1. $|M(r)| \leq 1$ for every resident r in R , and
2. $|M(h)| \leq q_h$ for every hospital h in H .

Furthermore, for each matching M in G and each edge $e = (r, h)$ in $E \setminus M$, H weakly (strictly) prefers e on M if and only if

1. $|M(h)| < q_h$, and/or
2. there exists an edge $f = (s, h)$ in $M(h)$ such that $r \succsim_H s$ (resp., $r \succ_H s$).

Thus, our problem is equivalent to the hospitals/residents problem with ties and master lists [12].

5.2 Student-project allocation setting

Assume that H is partitioned into non-empty subsets $H_1, H_2, \dots, H_k$, and we are given positive integers c_h and d_i for each hospital h in H and each $i = 1, 2, \dots, k$, respectively. Then, we define

$$\mathcal{F} := \left\{ F \subseteq E \mid \forall h \in H: |F(h)| \leq c_h, \right. \\ \left. \forall i = 1, 2, \dots, k: \sum_{h \in H_i} |F(h)| \leq d_i \right\}.$$

It is not difficult to see that $\mathbf{N} = (E, \mathcal{F})$ is a matroid. When we define the matroid $\mathbf{N}$ as above, a subset M of E is a matching if and only if

1. $|M(r)| \leq 1$ for every resident r in R ,
2. $|M(h)| \leq c_h$ for every hospital h in H , and
3. $\sum_{h \in H_i} |M(h)| \leq d_i$ for every $i = 1, 2, \dots, k$.

These constraints are exactly the same as the capacity constraints of the student-project allocation problem [1]. For each matching M in G and each edge $e = (r, h)$ in $E \setminus M$ such that $h \in H_i$ for an integer i in $\{1, 2, \dots, k\}$, H weakly (strictly) prefers e on M if and only if at least one of the following conditions is satisfied.

1. $|M(h)| < c_h$ and $\sum_{h \in H_i} |M(h)| < d_i$.
2. $|M(h)| < c_h$ and there exists an edge $f = (s, p)$ in M such that $p \in H_i$ and $r \succsim_H s$ (resp., $r \succ_H s$).
3. There exists an edge $f = (s, h)$ in $M(h)$ such that $r \succsim_H s$ (resp., $r \succ_H s$).

It should be noted that in the above definition, if $|M(h)| < c_h$ and $M(r) \in H_i$, then H weakly prefers e on M (i.e., the condition (2) is satisfied), but it does not mean that H strictly prefers e on M since $r \not\succ_H r$.

References

- [1] D. J. Abraham, R. W. Irving, and D. Manlove. Two algorithms for the student-project allocation problem. *Journal of Discrete Algorithms*, 5(1):73–90, 2007.
- [2] W. H. Cunningham. Improved bounds for matroid partition and intersection algorithms. *SIAM Journal on Computing*, 15(4):948–957, 1986.
- [3] T. Fleiner. A fixed-point approach to stable matchings and some applications. *Mathematics of Operations Research*, 28(1):103–126, 2003.
- [4] D. Gale and L. S. Shapley. College admissions and the stability of marriage. *The American Mathematical Monthly*, 69(1):9–15, 1962.
- [5] R. W. Irving. Stable marriage and indifference. *Discrete Applied Mathematics*, 48(3):261–272, 1994.
- [6] R. W. Irving, D. F. Manlove, and S. Scott. The hospitals/residents problem with ties. In *Proceedings of the 7th Scandinavian Workshop on Algorithm Theory*, volume 1851 of *Lecture Notes in Computer Science*, pages 259–271, 2000.
- [7] R. W. Irving, D. F. Manlove, and S. Scott. Strong stability in the hospitals/residents problem. In *Proceedings of the 20th Annual Symposium on Theoretical Aspects of Computer Science*, volume 2607 of *Lecture Notes in Computer Science*, pages 439–450, 2003.
- [8] R. W. Irving, D. F. Manlove, and S. Scott. The stable marriage problem with master preference lists. *Discrete Applied Mathematics*, 156(15):2959–2977, 2008.
- [9] T. Kavitha, K. Mehlhorn, D. Michail, and K. Paluch. Strongly stable matchings in time $O(nm)$ and extension to the hospitals-residents problem. *ACM Transactions on Algorithms*, 3(2):Article 15, 2007.
- [10] D. F. Manlove. Stable marriage with ties and unacceptable partners. Technical Report TR-1999-29, The University of Glasgow, Department of Computing Science, 1999.
- [11] D. F. Manlove. *Algorithmics of matching under preferences*. World Scientific, 2013.
- [12] G. O’Malley. *Algorithmic aspects of stable matching problems*. PhD thesis, The University of Glasgow, 2007.
- [13] J. G. Oxley. *Matroid theory*. Oxford University Press, 2nd edition, 2011.

List of MI Preprint Series, Kyushu University

The Global COE Program

Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p-adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials
- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials

- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings
- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric τ -functions of the q-Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications

- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain
- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection
- MI2009-26 Manabu YOSHIDA
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance

- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA
Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI
Finite element computation for scattering problems of micro-hologram using DtN map
- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO
Abstract collision systems on groups

- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA
An algebraic approach to underdetermined experiments
- MI2010-10 Kei HIROSE & Sadanori KONISHI
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE
Lagrangian approach to weakly nonlinear stability of an elliptical flow
- MI2010-21 Hiroki MASUDA
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiro FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra

- MI2010-25 Toshimitsu TAKAESU
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI
Spin-spin correlation functions of the q -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and spike detection via Gaussian basis expansions
- MI2010-32 Nobutaka NAKAZONO
Hypergeometric τ functions of the q -Painlevé systems of type $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA
CAP representations of inner forms of $Sp(4)$ with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER
Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide FUKUMOTO & Alexander B. SAMOKHIN
Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI
Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI
Anderson-Darling test and the Malliavin calculus
- MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling via Compressed Sensing

- MI2011-5 Hiroshi INOUE
Implications in Compressed Sensing and the Restricted Isometry Property
- MI2011-6 Daeju KIM & Sadanori KONISHI
Predictive information criterion for nonlinear regression model based on basis expansion methods
- MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI
Group variable selection via relevance vector machine
- MI2011-8 Jan BREZINA & Yoshiyuki KAGEI
Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow
Group variable selection via relevance vector machine
- MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC
Recursive structures in the multispecies TASEP
- MI2011-10 Kazunori YASUTAKE
On projective space bundle with nef normalized tautological line bundle
- MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA
An explicit formula for the discrete power function associated with circle patterns of Schramm type
- MI2011-12 Yoshiyuki KAGEI
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow
- MI2011-13 Vladimír CHALUPECKÝ & Adrian MUNTEAN
Semi-discrete finite difference multiscale scheme for a concrete corrosion model: approximation estimates and convergence
- MI2011-14 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves
- MI2011-15 Hiroshi INOUE
A generalization of restricted isometry property and applications to compressed sensing
- MI2011-16 Yu KAWAKAMI
A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space
- MI2011-17 Naoyuki KAMIYAMA
Matroid intersection with priority constraints
- MI2012-1 Kazufumi KIMOTO & Masato WAKAYAMA
Spectrum of non-commutative harmonic oscillators and residual modular forms
- MI2012-2 Hiroki MASUDA
Mighty convergence of the Gaussian quasi-likelihood random fields for ergodic Levy driven SDE observed at high frequency

- MI2012-3 Hiroshi INOUE
A Weak RIP of theory of compressed sensing and LASSO
- MI2012-4 Yasuhide FUKUMOTO & Youich MIE
Hamiltonian bifurcation theory for a rotating flow subject to elliptic straining field
- MI2012-5 Yu KAWAKAMI
On the maximal number of exceptional values of Gauss maps for various classes of surfaces
- MI2012-6 Marcio GAMEIRO, Yasuaki HIRAOKA, Shunsuke IZUMI, Miroslav KRAMAR, Konstantin MISCHAIKOW & Vidit NANDA
Topological Measurement of Protein Compressibility via Persistence Diagrams
- MI2012-7 Nobutaka NAKAZONO & Seiji NISHIOKA
Solutions to a q -analog of Painlevé III equation of type $D_7^{(1)}$
- MI2012-8 Naoyuki KAMIYAMA
A new approach to the Pareto stable matching problem
- MI2012-9 Jan BREZINA & Yoshiyuki KAGEI
Spectral properties of the linearized compressible Navier-Stokes equation around time-periodic parallel flow
- MI2012-10 Jan BREZINA
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a time-periodic parallel flow
- MI2012-11 Daeju KIM, Shuichi KAWANO & Yoshiyuki NINOMIYA
Adaptive basis expansion via the extended fused lasso
- MI2012-12 Masato WAKAYAMA
On simplicity of the lowest eigenvalue of non-commutative harmonic oscillators
- MI2012-13 Masatoshi OKITA
On the convergence rates for the compressible Navier- Stokes equations with potential force
- MI2013-1 Abuduwaili PAERHATI & Yasuhide FUKUMOTO
A Counter-example to Thomson-Tait-Chetayev's Theorem
- MI2013-2 Yasuhide FUKUMOTO & Hirofumi SAKUMA
A unified view of topological invariants of barotropic and baroclinic fluids and their application to formal stability analysis of three-dimensional ideal gas flows
- MI2013-3 Hiroki MASUDA
Asymptotics for functionals of self-normalized residuals of discretely observed stochastic processes
- MI2013-4 Naoyuki KAMIYAMA
On Counting Output Patterns of Logic Circuits
- MI2013-5 Hiroshi INOUE
RIPless Theory for Compressed Sensing

- MI2013-6 Hiroshi INOUE
Improved bounds on Restricted isometry for compressed sensing
- MI2013-7 Hidetoshi MATSUI
Variable and boundary selection for functional data via multiclass logistic regression modeling
- MI2013-8 Hidetoshi MATSUI
Variable selection for varying coefficient models with the sparse regularization
- MI2013-9 Naoyuki KAMIYAMA
Packing Arborescences in Acyclic Temporal Networks
- MI2013-10 Masato WAKAYAMA
Equivalence between the eigenvalue problem of non-commutative harmonic oscillators and existence of holomorphic solutions of Heun's differential equations, eigenstates degeneration, and Rabi's model
- MI2013-11 Masatoshi OKITA
Optimal decay rate for strong solutions in critical spaces to the compressible Navier-Stokes equations
- MI2013-12 Shuichi KAWANO, Ibuki HOSHINA, Kazuki MATSUDA & Sadanori KONISHI
Predictive model selection criteria for Bayesian lasso
- MI2013-13 Hayato CHIBA
The First Painleve Equation on the Weighted Projective Space
- MI2013-14 Hidetoshi MATSUI
Variable selection for functional linear models with functional predictors and a functional response
- MI2013-15 Naoyuki KAMIYAMA
The Fault-Tolerant Facility Location Problem with Submodular Penalties
- MI2013-16 Hidetoshi MATSUI
Selection of classification boundaries using the logistic regression
- MI2014-1 Naoyuki KAMIYAMA
Popular Matchings under Matroid Constraints
- MI2014-2 Yasuhide FUKUMOTO & Youichi MIE
Lagrangian approach to weakly nonlinear interaction of Kelvin waves and a symmetry-breaking bifurcation of a rotating flow
- MI2014-3 Reika AOYAMA
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Parallel flow in a cylindrical domain
- MI2014-4 Naoyuki KAMIYAMA
The Popular Condensation Problem under Matroid Constraints

- MI2014-5 Yoshiyuki KAGEI & Kazuyuki TSUDA
Existence and stability of time periodic solution to the compressible Navier-Stokes equation for time periodic external force with symmetry
- MI2014-6 This paper was withdrawn by the authors.
- MI2014-7 Masatoshi OKITA
On decay estimate of strong solutions in critical spaces for the compressible Navier-Stokes equations
- MI2014-8 Rong ZOU & Yasuhide FUKUMOTO
Local stability analysis of azimuthal magnetorotational instability of ideal MHD flows
- MI2014-9 Yoshiyuki KAGEI & Naoki MAKIO
Spectral properties of the linearized semigroup of the compressible Navier-Stokes equation on a periodic layer
- MI2014-10 Kazuyuki TSUDA
On the existence and stability of time periodic solution to the compressible Navier-Stokes equation on the whole space
- MI2014-11 Yoshiyuki KAGEI & Takaaki NISHIDA
Instability of plane Poiseuille flow in viscous compressible gas
- MI2014-12 Chien-Chung HUANG, Naonori KAKIMURA & Naoyuki KAMIYAMA
Exact and approximation algorithms for weighted matroid intersection
- MI2014-13 Yusuke SHIMIZU
Moment convergence of regularized least-squares estimator for linear regression model
- MI2015-1 Hidetoshi MATSUI
Sparse regularization for multivariate linear models for functional data
- MI2015-2 Reika AOYAMA & Yoshiyuki KAGEI
Spectral properties of the semigroup for the linearized compressible Navier-Stokes equation around a parallel flow in a cylindrical domain
- MI2015-3 Naoyuki KAMIYAMA
Stable Matchings with Ties, Master Preference Lists, and Matroid Constraints