

## Sap-flow velocity reduction by soil water deficit observed in a *Lithocarpus edulis* forest on Kyushu Island, Japan

**Komatsu, Hikaru**

Division of Forest Ecosphere and Management, Department of Forestry and Forest Products Science, Faculty of Agriculture, Kyushu University

**Katayama, Ayumi**

Division of Forest Ecosphere and Management, Department of Forestry and Forest Products Science, Faculty of Agriculture, Kyushu University

**Kume, Tomonori**

Division of Forest Ecosphere and Management, Department of Forestry and Forest Products Science, Faculty of Agriculture, Kyushu University

**Otsuki, Kyoichi**

Division of Forest Ecosphere and Management, Department of Forestry and Forest Products Science, Faculty of Agriculture, Kyushu University

<https://doi.org/10.15017/15050>

---

出版情報：九州大学農学部演習林報告. 88, pp.21-32, 2007-03. 九州大学農学部附属演習林  
バージョン：  
権利関係：



## Original Article

**Sap-flow velocity reduction by soil water deficit observed in a *Lithocarpus edulis* forest on Kyushu Island, Japan \***

Hikaru KOMATSU\*\*, Ayumi KATAYAMA \*\*, Tomonori KUME \*\*, and Kyoichi OTSUKI \*\*

**Abstract**

This paper examines transpiration reduction due to soil water deficits observed in a *Lithocarpus edulis* forest on Kyushu Island, south-western Japan. Continuous sap flow measurements were performed on six trees to monitor changes in tree transpiration rates. A reduction in sap flow velocity was observed in the period between late-September and late-October 2003. Precipitation amounts for this period were limited resulting in a corresponding low soil matric potential. A clear reduction in sap flow velocity occurred when soil matric potential at a 10 cm depth was  $< -70$  (kPa). Sap flow velocity scaled for 1.0 kPa vapor pressure deficit declined by 42 % when soil matric potential at a 10 cm depth was  $< -70$  (kPa) compared to  $\geq -70$  (kPa). In addition, intensive leaf-scale physiological measurements were performed on two days that had contrasting soil matric potentials. The intensive measurements showed lower leaf transpiration rates and stomatal conductance on the day with the lower soil matric potential. This finding suggests soil water deficits caused stomatal closure and therefore reduction in tree transpiration. Sap flow velocity and leaf water potential data showed that stem hydraulic conductance did not decline with low soil matric potential when compared to the day with higher soil matric potential.

**Key words:** forest canopy; *Lithocarpus edulis* ; sap flow; soil water deficit; transpiration

---

\* 九州のマテバシイ林で計測された土壌水分減少による樹液流速の低下

\*\* Division of Forest Ecosphere and Management, Department of Forestry and Forest Products Science, Faculty of Agriculture, Kyushu University, Fukuoka 811-2415 九州大学大学院農学研究院森林資源科学部門森林生態圏管理学講座

## 1. Introduction

As forests are the predominant land-cover type in Japan (Fujimori, 2000), examining the forest water cycle is crucial for water resources management (Komatsu, 2007). Forest canopy transpiration is a critical component affecting water yield from forested watersheds (Vertessy *et al.*, 2001; Komatsu *et al.*, 2007b). Thus, numerous researchers have investigated forest canopy transpiration in Japan (Komatsu *et al.*, 2006a,b; Kumagai *et al.*, 2007).

Soil water availability is a factor affecting forest canopy transpiration. Reduction in forest canopy transpiration by soil water deficit has been reported by several international studies (Granier *et al.*, 2000; Law *et al.*, 2001) and is included in forest water cycle models (Granier *et al.*, 1999; Williams *et al.*, 2001). However, very few studies (Hattori *et al.*, 1993; Takimoto *et al.*, 1994; Tanaka *et al.*, 2002) have examined reduction in forest canopy transpiration by soil water deficit in Japan. Furthermore, water cycle models developed in Japan (Kondo *et al.*, 1992; Komatsu *et al.*, 2005, 2007b) usually do not include this phenomenon. Thus, reduction in forest canopy transpiration by soil water deficit may not be a common event in Japan.

Nevertheless, examining the effect of soil water deficit on forest canopy transpiration is significant for the following two reasons. First, soil water deficit could greatly affect forest canopy transpiration during infrequent severe droughts brought on by inter-annual precipitation variability or a possible climate change. Second, examining the forest water cycle during a severe drought is important because water resources management becomes critical. This study examined reduction in forest canopy transpiration by soil water deficit in Japan.

## 2 Materials and methods

### 2.1 The site

The experiments were executed in the Kasuya Research Forest of Kyushu University, on Kyushu Island, south-western Japan (33° 38'N, 130° 31'E, elevation 50 m). The study site was located in a 28-year old evergreen broad-leaved plantation forest (*Lithocarpus edulis*) with an approximate canopy height of 9 m. Surface soil was layered (Imura *et al.*, 2000); the first layer, containing humus, ranged between 0 and 25 cm depth; the second layer, consisting of clay, ranged between 25 and 80 cm depth; and the soil layer located > 80 cm depth consisted of hard bedrock. An examination by Hosokawa *et al.* (2001) in 1998 revealed the majority of the tree roots were distributed between the 0 to 20 cm range. The saturated hydraulic conductivity was  $1.5 \times 10^{-2} \text{ cm s}^{-1}$  for the first layer and  $3.0 \times 10^{-5} \text{ cm s}^{-1}$  for the second layer as shown in a study by Imura *et al.* (2000) conducted in 1998. The annual mean temperature is about 16 °C and the annual total precipitation is about 1800 mm at

this site (Hirose *et al.*, 2005). Further details regarding the description of the site are available in research papers by Imura *et al.* (2000), Sato *et al.* (2003, 2004) and Hirose *et al.* (2005).

The site included two plots with differing pedestrian trampling intensity to examine the effect, if any, on the forest water cycle. Our previous study (Komatsu *et al.*, 2007a) found a significant difference in sap flow velocity between the plots. However, this difference did not alter the conclusions of this study. Thus, the data from the two plots were not differentiated.

## 2.2 Measurements

Tree transpiration was monitored by measuring sap flow on six trees. The measurements were performed during the period of September 6 to December 1 in 2003. The diameter at breast height (DBH) of the sample trees ranged between 13.6 and 15.3 cm.

Sap flow measurements ( $n = 6$ ) were performed based on the heat-pulse method (Closs, 1958; Marshall, 1958; Swanson, 1994). Instrumentation for the heat-pulse method was comprised by a set of three sensors (HP-3, Hayasi Denko Co.) consisting of a heater probe and two thermistor probes (diameter: 2.0 mm; length: 60 mm). The three sensors were installed at a height of 120 cm in holes drilled to a depth of 10 mm.

A heat-pulse tracer was released for a duration of 1.5 s every 20 min and the temperature difference between the thermistor probes was measured every 0.25 s. The time delay for the same temperature increase to occur at both thermistor probes was recorded with a solid-state memory module (CR10X, Campbell Scientific).

We measured meteorological and soil water conditions concurrently. This data was used later to interpret the sap-flow velocity data. Precipitation was measured using a tipping bucket gauge situated in an open space adjacent to the forest. Above-canopy air temperature and relative humidity were measured using a thermistor (VHE, Vaisala) positioned 14 m from the ground surface. Soil matric potential was measured at 10 cm depth at two different points using tensiometers (DIK-3021, Daiki Rika Kogyo). These two measurement points were about 18 m apart.

In addition, we took intensive measurements of leaf stomatal conductance and leaf transpiration rates using a porometer (Li-6400, Li-Cor) and leaf water potential using a pressure chamber (Model 600, PMS instrument Co.). These measurements were conducted on two leaves of four different trees, on three of which sap flow measurements were performed. These intensive measurements were undertaken on October 7 and November 14, days which displayed contrasting soil water conditions. Data from these measurements were used to examine the processes of transpiration reduction.

### 3 Results and discussion

Monthly precipitation amounts were 64, 7, and 122 mm for September, October, and November, respectively. Monthly precipitation amounts at the site, averaged for the previous 30 years, were 175, 81, and 81 mm for September, October, and November, respectively. Thus, precipitation amounts in 2003 were much lower than usual in September and October. Figure 1a shows daily precipitation during the period that sap flow measurements were performed. The amount of precipitation was quite small for the period between mid-September and late-October.




Fig. 1. Meteorological conditions, daily heat-pulse velocities (HPVs), and soil matric potential at a 10 cm depth. (a) Precipitation and air temperature; (b) HPVs and vapor pressure deficit (VPD); (c) soil matric potential at two observation points in the forest. HPV values were normalized by the value on September 6 (i.e. the first day of sap flow measurements) and averaged for the number of sample trees ( $n = 6$ ). Vertical bars indicate standard deviation.

Figure 1b shows daily heat-pulse velocities (HPVs) and vapor pressure deficit (VPD). HPVs were normalized by the value on September 6 (i.e. the first day of sap flow measurements) and then averaged for the number of sample trees ( $n = 6$ ). Normalized HPVs corresponded well to VPD in early-September and in November. During the period between late-September and late-October, normalized HPVs did not correspond well to VPD. This implies that soil water deficit limited transpiration during this period and which is supported by the soil matric potential data. Figure 1c displays 10 cm depth soil matric potential at two different observation points. Soil matric potential tended to be lower at both observation points in the period between late-September and late-October.




Fig. 2. Relationship between vapor pressure deficit (VPD) and normalized heat-pulse velocities (HPVs) classified according to soil matric potential (average of data from Observation Points 1 and 2).

Figure 2 shows the relationship between VPD and the normalized HPV, classified by soil matric potential (Fig. 1c). Regression lines were determined by the least-squares method. The normalized HPV increased with increasing VPD regardless of soil matric potential class. However, the slope of the regression line was greater when

soil matric potential was  $\geq -70$  (kPa) ( $p < 0.001$ ; t-test). HPV scaled for 1.0 kPa VPD declined by 42% for soil matric potential at 10 cm depth  $< -70$  (kPa). Thus, sap-flow velocity reduction was significant when the soil matric potential was  $< -70$  (kPa). This agrees with Komatsu *et al.*'s (2007a) examination on the relationship between soil matric potential and water content. They showed that reduction in soil water content was drastic when soil matric potential was  $< -70$  (kPa) (Figure 5b of Komatsu *et al.*, 2007a), indicating that available soil water greatly reduced when soil matric potential became  $< -70$  (kPa).




Fig. 3. Diurnal course of (a) solar radiation, (b) air temperature, (c) vapor pressure deficit (VPD), (d) leaf transpiration rates, (e) leaf stomatal conductance, and (f) leaf water potential on October 7 and November 14. Leaf transpiration rates and stomatal conductance were not measured in the early morning of November 14 due to heavy dew. October 7 was in the period with heat-pulse velocity reduction, while November 14 lies outside the period.

Figure 3 shows the diurnal course of meteorological factors (Figs. 3a, 3b, and 3c), leaf transpiration rates (Fig. 3d), leaf stomatal conductance (Fig. 3e), and leaf water potential (Fig. 3f) on October 7 and November 14. October 7 was included in the period with sap-flow velocity reduction while November 14 lies outside this period (Fig. 1b). Transpiration rates were lower for October 7 than for November 14 (Fig. 3d) which is consistent with the HPV data in Fig. 1b. Stomatal conductance and predawn water potential were lower on October 7 than November 14 (Figs. 3e and 3f). This indicates a lower soil water availability caused lower stomatal conductance and therefore lower transpiration rates on October 7.




Fig. 4. Relationship between heat-pulse velocities (HPVs) and leaf water potential on October 7 and November 14. October 7 was in the period with heat-pulse velocity reduction while November 14 lies outside the period. Bars indicate standard deviation.

Figure 4 shows the relationships between HPV and leaf water potential on October 7 and November 14. Regression lines were determined based on the least-squares method. The slope for October 7 was not significantly different from that for November 14 ( $p = 0.39$ ; t-test). This suggests that stem hydraulic conductance did not decline on October 7 compared to November 14.

## **4 Conclusions**

Sap flow was monitored for six trees in a broad-leaved forest on Kyushu Island located in south-western Japan. Sap flow velocity was clearly reduced when the precipitation amount, and therefore the soil matric potential, was low. Reduction in sap flow velocity was significant when the soil matric potential at a 10 cm depth was  $< -70$  (kPa): HPV scaled for 1.0 kPa VPD was reduced by 42% for soil matric potential at a 10 cm depth  $< -70$  (kPa). Leaf-scale physiological measurements suggested tree transpiration reduction was caused by physiological stomatal control due to soil water deficit.

Our study is important as few studies have reported on such reductions in Japan. Further research will clarify the forest water cycle during the infrequent severe droughts that can be brought on by inter-annual precipitation variability or possible future climate change.

## **Acknowledgements**

We would like to thank Dr. Atsushi Kume (University of Toyama, Japan) and Dr. Shigeki Hirose (T & F, Japan) for their instruction and assistance in measurements. Thanks are also due to two anonymous reviewers whose comments greatly improved the manuscript. This research was supported by Grant-in-Aid for Scientific Research (#17380096 and #18810023) from the Ministry of Education, Science, and Culture of Japan.

## References

- Closs R.H. (1958)** The heat pulse method for measuring rate of sap flow in a plant stem. *N. Z. J. Sci.* **1**: 281-288.
- Fujimori, T. (2000)** Living with Forests. Maruzen, Tokyo. (in Japanese).
- Granier A., Breda N., Biron P., and Villetle S. (1999)** A lumped water balance model to evaluate duration and intensity of drought constraints in forest stands *Ecol. Model.* **116**: 269-283.
- Granier A., Biron P., and Lemoine D. (2000)** Water balance, transpiration and canopy conductance in two beech stands. *Agric. For. Meteorol.* **100**: 291-308
- Hattori S., Tamai K., and Abe T. (1993)** Effects of soil moisture and vapor pressure deficit on evapotranspiration in a hinoki plantation. *J. Jpn. For. Soc.* **75**: 216-224 (in Japanese with English summary).
- Hirose S., Kume A., Takeuchi S., Utsumi Y., Otsuki K., and Ogawa S. (2005)** Stem water transport of *Lithocarpus edulis*, an evergreen oak with radial-porous wood. *Tree Physiol.* **25**: 221-228.
- Hosokawa T., Ogawa S., Jinno K., Otsuki K., Takeuchi S., Kumagai T., Fujino K., and Kume A. (2001)** Runoff analysis using a big-leaf model and a tank model considering temporal change in soil matric potential and soil water content in the root zone of *Lithocarpus edulis*. In: Quantification and modelling of water and energy cycling in a forest ecosystem, Ogawa S. (eds.), Kyushu University Forest, 74-85. (in Japanese).
- Imura Y., Hosokawa T., Jinno K., Ogawa S., Otsuki K., Takeuchi S., Kumagai T., and Nishiyama K. (2000)** Characteristics of transpiration and soil water pressure in the root zone of *Pasania Edulis*. *Tech. Rep. Kyushu Univ.* **73**: 679-684 (in Japanese with English summary).
- Komatsu H. (2007)** Comparison of transpiration rates at various forests. In: Forest Hydrology, Kume A. (eds.), Morikita, Tokyo, in press (in Japanese).
- Komatsu H., Kang Y., Kume T., Yoshifuji N., and Hotta N. (2006a)** Transpiration from a *Cryptomeria japonica* plantation forest, part 1: aerodynamic control of transpiration. *Hydrol. Process.* **20**: 1309-1320
- Komatsu H., Kang Y., Kume T., Yoshifuji N., and Hotta N. (2006b)** Transpiration from a *Cryptomeria japonica* plantation forest, part 2: response of canopy conductance to meteorological factors. *Hydrol. Process.* **20**: 1321-1334
- Komatsu H, Katayama A, Hirose S, Kume A, Higashi N, Ogawa S, and Otsuki K. (2007a)** Reduction in soil water availability and of tree transpiration in a forest with pedestrian trampling. *Agric. For. Meteorol.*, accepted.
- Komatsu H., Tanaka N., and Kume T. (2007b)** Do coniferous forests evaporate more water than broad-leaved forest in Japan? *J. Hydrol.*, 336:361-375.
- Komatsu H., Sawano S., Kume T., and Hashimoto S. (2005)** Relationships between forest properties and evapotranspiration rates. *J. Jpn. For. Soc.* **87**: 170-185 (in

Japanese with English summary).

- Kondo J., Watanabe T., and Nakazono M. (1992)** Heat budget evaluation of forest evapotranspiration in Japan. *Tenki* **39**: 113-123 (in Japanese with English summary).
- Kumagai T., Aoki S., Shimizu T. and Otsuki K. (2007)** Sap flow estimates of stand transpiration at two slope positions in a Japanese cedar forest watershed. *Tree Physiol.* **27**: 161-168.
- Law B.E., Goldstein A.H., Anthoni P.M. Unsworth M.H., Panek J.A., Bauer M.R., Fracheboud J.M., and Hultman N. (2001)** Carbon dioxide and water vapor exchange by young and old ponderosa pine ecosystems during a dry summer. *Tree Physiol.* **21**: 299-308.
- Marshall, D.C. (1958)** Measurement of sap flow in conifers by heat transport. *Plant Physiol.* **33**: 385-396.
- Sato, Y., Kume, A., Otsuki, K., and Ogawa, S. (2003)** Effects of difference in canopy structure on the distribution of throughfall — a comparison of throughfall characteristics between the coniferous forest and the broad-leaved forest. *J. Jpn. Soc. Hydrol. & Water Resour.* **16**: 605-616 (in Japanese with English summary).
- Sato, Y., Kumagai, T., Kume, A., Otsuki, K., and Ogawa, S. (2004)** Experimental analysis of moisture dynamics of litter layers — the effects of rainfall conditions and leaf shapes. *Hydrol. Process.* **18**: 3007-3018.
- Swanson, R.H. (1994)** Significant historical developments in thermal methods for measuring sap flow in trees. *Agric. For. Meteorol.* **72**: 113-132.
- Takimoto, H., Horino, H., Watanabe, T., and Maruyama, T. (1994)** Comparison of evapotranspiration properties between a reclaimed farmland and a natural forest — catchment evapotranspiration estimates by the short-term water-budget method —. *J. Jpn Soc. Hydrol. & Water Resour.* **7**: 529-535 (in Japanese with English summary).
- Tanaka K., Kosugi Y., and Nakamura A. (2002)** Impact of leaf physiological characteristics on seasonal variation in CO<sub>2</sub>, latent and sensible heat exchanges over a tree plantation. *Agric. For. Meteorol.* **114**: 103-122.
- Vertessy R.A., Watson F.G.R., and O'Sullivan S.K. (2001)** Factors determining relations between stand age and catchment water balance in mountain ash forests. *For. Ecol. Manage.* **143**: 13-26.
- Williams M., Law B.E., Anthoni P.M., and Unsworth M.H. (2001)** Use of a simulation model and ecosystem flux data to examine carbon-water interactions in ponderosa pine. *Tree Physiol.* **21**: 287-298.

## 九州のマテバシイ林で計測された土壤水分減少による樹液流速の低下

小松 光・片山歩美・久米朋宣・大槻恭一

### 要 旨

本論は、九州のマテバシイ林で計測された土壤水分減少による蒸散低下を調べたものである。蒸散の変化を観測するために森林内の6本の樹木において樹液流計測を行ったところ、2003年9月下旬から10月下旬にかけて樹液流速の低下が認められた。この期間は降水量が少なかったために、土壤マトリックポテンシャルが低下していた。樹液流速の低下は、深度10 cmの土壤マトリックポテンシャルが $-70$  kPa以下のときに顕著であった。大気飽差が $1.0$  kPaのときの樹液流速は、土壤マトリックポテンシャルが $-70$  kPa以下のとき、土壤マトリックポテンシャルが $-70$  kPa以上のときに比べて42%低下していた。樹液流計測に加えて単葉スケールでの樹木生理計測を、土壤マトリックポテンシャルの異なる2日を選んで行った。その結果、単葉の蒸散量と気孔コンダクタンスは、土壤マトリックポテンシャルが低い日において小さかった。このことは、土壤水分減少が気孔の閉鎖を引き起こし、その結果、樹木の蒸散が低下したことを示唆している。樹液流速と水ポテンシャルの関係から、土壤マトリックポテンシャルの低下による幹の通水コンダクタンスの低下は認められなかった。

キーワード: 森林; マテバシイ; 樹液流; 土壤水分減少; 蒸散

