Popular Matchings with Ties and Matroid Constraints

Kamiyama, Naoyuki

Institute of Mathematics for Industry, Kyushu University | JST, PRESTO

<http://hdl.handle.net/2324/1525809>

出版情報 : MI Preprint Series. 2015-6, 2015-05-24. 九州大学大学院数理学研究院
バージョン :
権利関係 :

MI Preprint Series

Mathematics for Industry
Kyushu University

**Popular Matchings with Ties and
Matroid Constraints**

Naoyuki Kamiyama

MI 2015-6

(Received May 24, 2015)

Institute of Mathematics for Industry
Graduate School of Mathematics
Kyushu University
Fukuoka, JAPAN

Popular Matchings with Ties and Matroid Constraints

Naoyuki Kamiyama^{*†‡}

Abstract

In this paper, we consider the popular matching problem with matroid constraints. It is known that if there exists no tie in preference lists of applicants, then this problem can be solved in polynomial time. In this paper, we prove that even if there exist ties in preference lists, this problem can be solved in polynomial time.

1 Introduction

In this paper, we consider some variation of the popular matching problem. The popular matching problem was originally introduced by Abraham, Irving, Kavitha, and Mehlhorn [1]. In this problem, we are given two disjoint sets of applicants and posts, and each applicant has a preference list over posts in which there may exist ties. A matching M between applicants and posts is said to be popular, if there exists no other matching N such that the number of applicants that prefer N to M is larger than the number of applicants that prefer M to N . The concept of popularity was originally proposed by Gärdenfors [10] in the context of matching problems in which agents in both sides have preference lists. The goal of the popular matching problem is to decide whether there exists a popular matching, and find a popular matching if one exists. Abraham, Irving, Kavitha, and Mehlhorn [1] presented polynomial-time algorithms for the popular matching problem with/without ties. Since their seminal paper, several variations of the popular matching problem [12, 19, 21, 24] and related problems [12, 13, 14, 15, 16, 20, 25, 26] have been investigated.

In this paper, we consider a matroid generalization of the popular matching problem. More precisely, in our model, capacity constraints for posts are generalized to matroid constraints. In the study of combinatorial optimization, for example, Fleiner [6] generalized capacity constraints in the stable matching problem to matroid constraints, and Zenklusen [27] considered a matroid generalization of the minimum-cost spanning tree problem with degree constraints. It is known [12] that if there exists no tie in preference lists, then this problem can be solved in polynomial time. In this paper, we prove that even if there exist ties in preference lists, we can solve this problem in polynomial time.

Our algorithm is based on the algorithm of Abraham, Irving, Kavitha, and Mehlhorn [1] for the popular matching problem with ties. However, the algorithm of [1] decomposes

^{*}Institute of Mathematics for Industry, Kyushu University, Fukuoka, Japan

[†]JST, PRESTO, Saitama, Japan

[‡]kamiyama@imi.kyushu-u.ac.jp

the vertex set of an input bipartite graph by using Gallai-Edmonds decomposition [5, 8, 9] (see also [18]) of a bipartite graph based on a maximum-size (one-to-one) matching. Thus, we can not straightforwardly generalize the algorithm of [1] to the matroid setting. Although Manlove and Sng [19] proposed an algorithm for a many-to-one variation of the popular matching with ties, their algorithm copies posts and uses the Gallai-Edmonds decomposition in the one-to-one setting. To overcome this difficulty, we decompose the edge set (instead of the vertex set) of an input bipartite graph bases on a maximum-size common independent set. This decomposition can be regarded as a matroid generalization of the Dulmage-Mendelsohn decomposition [3, 4] of a bipartite graph (see, e.g., [22] for the Dulmage-Mendelsohn type decomposition of the matroid intersection problem).

The rest of this paper is organized as follows. In Section 2, we give the formal definition of our model, and basics of matroids and the matroid intersection problem. In Section 3, we give a characterization of a popular matching in our model. In Section 4, we describe our algorithm, and prove its correctness.

2 Preliminaries

For each set X and each element u , we define $X + u := X \cup \{u\}$ and $X - u := X \setminus \{u\}$. For each sets X and Y , we define $X \Delta Y := (X \setminus Y) \cup (Y \setminus X)$. Assume that we are given a set X and a function $\xi: X \rightarrow \mathbb{Z}$, where $\mathbb{Z}$ is the set of integers. For each subset Y of X , we define

$$\xi(Y) := \sum_{u \in Y} \xi(u).$$

A pair $\mathbf{M} = (U, \mathcal{I})$ is called a *matroid*, if U is a finite set and $\mathcal{I}$ is a family of subsets of U satisfying the following conditions.

(I0) $\emptyset \in \mathcal{I}$.

(I1) If $I \in \mathcal{I}$ and $J \subseteq I$, then $J \in \mathcal{I}$.

(I2) If $I, J \in \mathcal{I}$ and $|I| < |J|$, then $I + u \in \mathcal{I}$ for some element u in $J \setminus I$.

We say that $\mathbf{M}$ is a matroid on the *ground set* U .

In this paper, we are given a finite simple (not necessarily complete) bipartite graph $G = (V, E)$. We assume that V is partitioned into two subsets A and P , and each edge in E connects a vertex in A and a vertex in P . We call a vertex in A an *applicant*, and a vertex in P a *post*. For each applicant a in A and each post p in P , if there exists an edge in E connecting a and p , then we denote by (a, p) this edge. For each vertex v in V and each subset F of E , we define $F(v)$ as the set of edges in F incident to v .

For each applicant a in A , we are given a transitive binary relation $\succsim_a$ on $E(a)$ such that at least one of $e \succsim_a g$ and $g \succsim_a e$ holds for every edges e, g in $E(a)$. Notice that for each applicant a in A and each edges e, g in $E(a)$, there exists a possibility that both $e \succsim_a g$ and $g \succsim_a e$ hold. For each applicant a in A and each edges e, g in $E(a)$, we use the notation $e \succ_a g$, if $e \succsim_a g$ and $g \not\succsim_a e$. For each applicant a in A and each edges e, g in $E(a)$, $e \succ_a g$ means that a prefers e to g . Furthermore, for each applicant a in A and each edges e, g in $E(a)$, if $e \succsim_a g$ and $g \succsim_a e$ hold, then a is indifferent between e and g .

As in [1], we assume that for each applicant a in A , there exists a *last resort post* $\ell(a)$ in P such that $E(\ell(a)) = \{(a, \ell(a))\}$, and $e \succ_a (a, \ell(a))$ for every edge e in $E(a) - (a, \ell(a))$. In addition, we assume that for every applicant a in A , there exists an edge e in $E(a)$ such that $e \neq (a, \ell(a))$. For each post p in P , we are given a matroid $\mathbf{M}_p = (E(p), \mathcal{I}_p)$. We assume that for every edge (a, p) in E , we have $\{(a, p)\} \in \mathcal{I}_p$.

A subset M of E is called a *matching* in G , if

- $|M(a)| = 1$ for every applicant a in A , and
- $M(p) \in \mathcal{I}_p$ for every post p in P .

For each subset F of E and each applicant a in A such that $|F(a)| = 1$, we denote by $\mu_F(a)$ the unique edge in $F(a)$. For each matchings M, N in G , we denote by $\phi(M; N)$ the number of applicants a in A such that $\mu_M(a) \succ_a \mu_N(a)$. A matching M in G is said to be *popular*, if for every matching N in G , $\phi(M; N) \geq \phi(N; M)$. Then, the goal of the *popular matching problem with ties and matroid constraints* (PMTM for short) is to decide whether there exists a popular matching in G , and find a popular matching if one exists.

2.1 Basics of matroids

Let $\mathbf{M} = (U, \mathcal{I})$ be a matroid. A subset of U belonging to $\mathcal{I}$ is called an *independent set* of $\mathbf{M}$. A subset C of U is called a *circuit* of $\mathbf{M}$, if C is not an independent set of $\mathbf{M}$, but every proper subset of C is an independent set of $\mathbf{M}$. Assume that we are given an independent set I of $\mathbf{M}$ and an element u in $U \setminus I$ such that $I + u \notin \mathcal{I}$. It is known [23, Proposition 1.1.6] that there exists a unique circuit of $\mathbf{M}$ that is a subset of $I + u$, and u belongs to this unique circuit. This unique circuit is called the *fundamental circuit* of u , and denoted by $\mathbf{C}_{\mathbf{M}}(u, I)$. It is known [23, p.20, Exercise 5] that we have

$$\mathbf{C}_{\mathbf{M}}(u, I) = \{w \in I + u \mid I + u - w \in \mathcal{I}\}.$$

Notice that if $\{u\} \in \mathcal{I}$, then $\mathbf{C}_{\mathbf{M}}(u, I) - u \neq \emptyset$.

A maximal independent set of $\mathbf{M}$ is called a *base* of $\mathbf{M}$. Notice that (I2) implies that every base of $\mathbf{M}$ has the same size. For each subset X of U , we define $\mathcal{I}|X$ as the family of subsets I of X such that $I \in \mathcal{I}$, and $\mathbf{M}|X := (X, \mathcal{I}|X)$. It is known [23, p.20] that for every subset X of U , $\mathbf{M}|X$ is a matroid. For each subset X of U , we define $\mathbf{r}_{\mathbf{M}}(X)$ as the size of a base of $\mathbf{M}|X$. For each disjoint subsets X, J of U , we define

$$\mathbf{p}(J; X) := \mathbf{r}_{\mathbf{M}}(J \cup X) - \mathbf{r}_{\mathbf{M}}(X).$$

For each subset X of U , we define

$$\begin{aligned} \mathcal{I}/X &:= \{J \subseteq U \setminus X \mid \mathbf{p}(J; X) = |J|\}, \\ \mathbf{M}/X &:= (U \setminus X, \mathcal{I}/X). \end{aligned}$$

It is known [23, Proposition 3.1.6] that for each subset X of U , $\mathbf{M}/X$ is a matroid.

Lemma 1 (see, e.g., [23, Proposition 3.1.7]). *Assume that we are given a matroid $\mathbf{M} = (U, \mathcal{I})$, a subset X of U , and a base B of $\mathbf{M}|X$. Then, for every element u of $U \setminus X$, $\{u\}$ is an independent set of $\mathbf{M}/X$ if and only if $B + u$ is an independent set of $\mathbf{M}$.*

Lemma 2 (See, e.g., [23, p.15, Exercise 14]). Assume that we are given a matroid $\mathbf{M} = (U, \mathcal{I})$ and circuits C_1, C_2 of $\mathbf{M}$ such that $C_1 \cap C_2 \neq \emptyset$ and $C_1 \setminus C_2 \neq \emptyset$. Then, for every element u in $C_1 \cap C_2$ and every element w in $C_1 \setminus C_2$, there exists a circuit C of $\mathbf{M}$ such that $w \in C$ and C is a subset of $(C_1 \cup C_2) - u$.

Lemma 3. Assume that we are given a matroid $\mathbf{M} = (U, \mathcal{I})$, an independent set I, J of $\mathbf{M}$ such that $J \subseteq I$, and an element u in $U \setminus J$ such that $I + u \notin \mathcal{I}$ and $J + u \notin \mathcal{I}$. Then, $\mathbf{C}_{\mathbf{M}}(u, I) = \mathbf{C}_{\mathbf{M}}(u, J)$.

Proof. Define $C_1 := \mathbf{C}_{\mathbf{M}}(u, I)$ and $C_2 := \mathbf{C}_{\mathbf{M}}(u, J)$. Assume that $C_1 \neq C_2$. Then, Lemma 2 implies that there exists a circuit C of $\mathbf{M}$ such that $C \subseteq (C_1 \cup C_2) - u$. Since $C_1 - u \subseteq I$ and $C_2 - u \subseteq J \subseteq I$, we have $C \subseteq I$. This contradicts the fact that $I \in \mathcal{I}$. $\square$

Lemma 4. Assume that we are given a matroid $\mathbf{M} = (U, \mathcal{I})$, an independent set I of $\mathbf{M}$, and a subset J of $U \setminus I$ such that $I \cup J \in \mathcal{I}$. Furthermore, we assume that we are given subsets X, Y of $U \setminus J$ such that $(I \cup X) \setminus Y \in \mathcal{I}$ and $I + x \notin \mathcal{I}$ for every element x in X . Then, $((I \cup X) \setminus Y) \cup J$ is an independent set of $\mathbf{M}$.

Proof. We prove this lemma by contradiction. Assume that $((I \cup X) \setminus Y) \cup J \notin \mathcal{I}$. This implies that there exists a circuit C of $\mathbf{M}$ such that $C \subseteq ((I \cup X) \setminus Y) \cup J$. If $C \cap J = \emptyset$, then this contradicts the fact that $(I \cup X) \setminus Y$ is an independent set of $\mathbf{M}$. Thus, there exists an element u in $J \cap C$. If $C \cap X = \emptyset$, then this contradicts the fact that $I \cup J$ is an independent set of $\mathbf{M}$. Thus, $X \cap C$ is not empty. Assume that $X \cap C = \{x_1, x_2, \dots, x_k\}$. Since $u \in J$, $J \cap X = \emptyset$, and $J \cap I = \emptyset$, we have $u \notin \mathbf{C}_{\mathbf{M}}(x_i, I)$ for every $i = 1, 2, \dots, k$.

Here we consider the following procedure.

Step 1. Set $i := 1$ and $C_0 := C$.

Step 2. If $i \leq k$, then do the following steps.

(2-a) If $x_i \notin C_{i-1}$, then set $C_i := C_{i-1}$ and go to (2-c).

(2-b) Find a circuit C_i of $\mathbf{M}$ such that $u \in C_i$ and $C_i \subseteq (C_{i-1} \cup \mathbf{C}_{\mathbf{M}}(x_i, I)) - x_i$.

(2-c) Update $i := i + 1$ and go back to the beginning of **Step 2**.

Step 3. Output C_k .

In **Step (2-b)**, since $x_i \in C_{i-1} \cap \mathbf{C}_{\mathbf{M}}(x_i, I)$ and $u \in C_{i-1} \setminus \mathbf{C}_{\mathbf{M}}(x_i, I)$, Lemma 2 implies that there exists a circuit C_i of $\mathbf{M}$ such that $u \in C_i$ and $C_i \subseteq (C_{i-1} \cup \mathbf{C}_{\mathbf{M}}(x_i, I)) - x_i$. It is not difficult to see that C_k is a subset of $I \cup J$, which contradicts the fact that $I \cup J \in \mathcal{I}$. This completes the proof. $\square$

Assume that we are given k matroids $\mathbf{M}_1 = (U_1, \mathcal{I}_1), \mathbf{M}_2 = (U_2, \mathcal{I}_2), \dots, \mathbf{M}_k = (U_k, \mathcal{I}_k)$ such that $U_i \cap U_j = \emptyset$ for every distinct $i, j = 1, 2, \dots, k$. Define

$$\bigoplus_{i=1}^k \mathcal{I}_i := \left\{ X \subseteq \bigcup_{i=1}^k U_i \mid X \cap U_i \text{ is an independent set of } \mathbf{M}_i \text{ for every } i = 1, 2, \dots, k \right\}.$$

Furthermore, we define

$$\bigoplus_{i=1}^k \mathbf{M}_i := \left(\bigcup_{i=1}^k U_i, \bigoplus_{i=1}^k \mathcal{I}_i \right).$$

It is not difficult to see that $\bigoplus_{i=1}^k \mathbf{M}_i$ is a matroid.

Let $\mathbf{M} = (U, \mathcal{I})$ and $\mathbf{N} = (U, \mathcal{J})$ be matroids. A subset I of U is called a *common independent set* of $\mathbf{M}$ and $\mathbf{N}$, if $I \in \mathcal{I} \cap \mathcal{J}$. We denote by $\mathcal{B}_{\mathbf{MN}}$ and $\gamma_{\mathbf{MN}}$ the family and the size of maximum-size common independent sets of $\mathbf{M}$ and $\mathbf{N}$, respectively. We can find a maximum-size common independent set of $\mathbf{M}$ and $\mathbf{N}$ in $O(|U|^{2.5}\mathbf{EO})$ by using the algorithm of Cunningham [2], where $\mathbf{EO}$ is the time required to decide whether $X + u$ is an independent set of $\mathbf{L}$ for every matroid $\mathbf{L}$ in $\{\mathbf{M}, \mathbf{N}\}$, every common independent set X of $\mathbf{M}$ and $\mathbf{N}$, and every element u in $U \setminus X$. Furthermore, for every function $\xi: U \rightarrow \mathbb{Z}$, we can find a maximum-size common independent set I of $\mathbf{M}$ and $\mathbf{N}$ such that

$$\xi(I) = \max\{\xi(J) \mid J \in \mathcal{B}_{\mathbf{MN}}\}$$

in $O(|U|^3\mathbf{EO})$ time by using the algorithm of Frank [7].

2.2 Auxiliary graphs and decomposition

Assume that we are given matroids $\mathbf{M} = (U, \mathcal{I})$ and $\mathbf{N} = (U, \mathcal{J})$, and a common independent set I of $\mathbf{M}$ and $\mathbf{N}$. Then, we define a directed graph $\mathbf{D}_{\mathbf{MN}}(I)$ as follows. The vertex set of $\mathbf{D}_{\mathbf{MN}}(I)$ is U . For each elements u of $U \setminus I$ and v of I ,

- there exists an arc from v to u in $\mathbf{D}_{\mathbf{MN}}(I)$ if and only if $I + u \notin \mathcal{I}$ and $v \in \mathbf{C}_{\mathbf{M}}(u, I)$,
- there exists an arc from u to v in $\mathbf{D}_{\mathbf{MN}}(I)$ if and only if $I + u \notin \mathcal{J}$ and $v \in \mathbf{C}_{\mathbf{N}}(u, I)$.

These are all arcs of $\mathbf{D}_{\mathbf{MN}}(I)$. Define

$$\begin{aligned} \mathbf{T}_{\mathbf{M}}(I) &:= \{u \in U \setminus I \mid I + u \in \mathcal{I}\}, \\ \mathbf{T}_{\mathbf{N}}(I) &:= \{u \in U \setminus I \mid I + u \in \mathcal{J}\}. \end{aligned}$$

Lemma 5 (see, e.g., [17, Lemma 13.30]). *Assume that we are given matroids $\mathbf{M}$ and $\mathbf{N}$ on the same ground set U and a common independent set I of $\mathbf{M}$ and $\mathbf{N}$. Then, I is a maximum-size common independent set of $\mathbf{M}$ and $\mathbf{N}$ if and only if there exists no directed path in $\mathbf{D}_{\mathbf{MN}}(I)$ from a vertex in $\mathbf{T}_{\mathbf{M}}(I)$ to a vertex in $\mathbf{T}_{\mathbf{N}}(I)$.*

Assume that I is a maximum-size common independent set of $\mathbf{M}$ and $\mathbf{N}$. We denote by $\Omega_{\mathbf{MN}}^+(I)$ the set of elements u in U such that there exists a directed path in $\mathbf{D}_{\mathbf{MN}}(I)$ from a vertex in $\mathbf{T}_{\mathbf{M}}(I)$ to u . In addition, we define $\Omega_{\mathbf{MN}}^-(I)$ as the set of elements u in U such that there exists a directed path in $\mathbf{D}_{\mathbf{MN}}(I)$ from u to a vertex in $\mathbf{T}_{\mathbf{N}}(I)$. Notice that since I is a maximum-size common independent set of $\mathbf{M}$ and $\mathbf{N}$, Lemma 5 implies that $\Omega_{\mathbf{MN}}^+(I)$ and $\Omega_{\mathbf{MN}}^-(I)$ are disjoint. Define

$$\Omega_{\mathbf{MN}}(I) := U \setminus \left(\Omega_{\mathbf{MN}}^+(I) \cup \Omega_{\mathbf{MN}}^-(I) \right).$$

Lemma 6. Assume that we are given matroids $\mathbf{M}$ and $\mathbf{N}$ on the same ground set U and a common independent set I of $\mathbf{M}$ and $\mathbf{N}$. Furthermore, we assume that we are given a vertex u in $\mathsf{T}_{\mathbf{M}}(I) \cup I$ and a vertex v in $\mathsf{T}_{\mathbf{N}}(I) \cup I$, and a shortest directed path L in $\mathsf{D}_{\mathbf{MN}}(I)$ from u to v . Then, $I \triangle K$ is a common independent set of $\mathbf{M}$ and $\mathbf{N}$, where K is the set of elements in U that L goes through (notice that since L is a shortest directed path, L goes through each vertex at most once).

Proof. This lemma immediately follows from [17, Lemma 13.27] and Lemma 4. $\square$

Lemma 7 (see, e.g., [11, Lemma 4.2]). Assume that we are given matroids $\mathbf{M}$ and $\mathbf{N}$ on the same ground set. Then, for every maximum-size common independent sets I, J of $\mathbf{M}$ and $\mathbf{N}$,

$$J \cap \Omega_{\mathbf{MN}}^-(I) \text{ is a base of } \mathbf{M} | \Omega_{\mathbf{MN}}^-(I), \quad (1)$$

$$J \cap (\Omega_{\mathbf{MN}}(I) \cup \Omega_{\mathbf{MN}}^-(I)) \text{ is a base of } \mathbf{M} | (\Omega_{\mathbf{MN}}(I) \cup \Omega_{\mathbf{MN}}^-(I)), \quad (2)$$

$$J \cap \Omega_{\mathbf{MN}}^+(I) \text{ is a base of } \mathbf{N} | \Omega_{\mathbf{MN}}^+(I), \quad (3)$$

$$J \cap (\Omega_{\mathbf{MN}}^+(I) \cup \Omega_{\mathbf{MN}}(I)) \text{ is a base of } \mathbf{N} | (\Omega_{\mathbf{MN}}^+(I) \cup \Omega_{\mathbf{MN}}(I)). \quad (4)$$

Although the following lemma is well-known, we give its proof for completeness.

Lemma 8. Assume that we are given matroids $\mathbf{M} = (U, \mathcal{I})$ and $\mathbf{N} = (U, \mathcal{J})$. Then, for every maximum-size common independent sets I, J of $\mathbf{M}$ and $\mathbf{N}$, we have

$$\begin{aligned} \Omega_{\mathbf{MN}}^+(I) &= \Omega_{\mathbf{MN}}^+(J), \\ \Omega_{\mathbf{MN}}(I) &= \Omega_{\mathbf{MN}}(J), \\ \Omega_{\mathbf{MN}}^-(I) &= \Omega_{\mathbf{MN}}^-(J). \end{aligned}$$

Proof. For simplicity, we define $\mathsf{D}_{\mathbf{MN}}(\cdot) := \mathsf{D}(\cdot)$, $\Omega_{\mathbf{MN}}^+(\cdot) := \Omega^+(\cdot)$, and $\Omega_{\mathbf{MN}}^-(\cdot) := \Omega^-(\cdot)$. Let I, J be maximum-size common independent sets of $\mathbf{M}$ and $\mathbf{N}$.

We first prove that $\Omega^+(I) = \Omega^+(J)$. For proving this, we first prove that $\Omega^+(J) \subseteq \Omega^+(I)$. Assume that $\mathsf{T}_{\mathbf{M}}(J) \not\subseteq \Omega^+(I)$, and let u be an element in $\mathsf{T}_{\mathbf{M}}(J) \setminus \Omega^+(I)$. Then, $J + u \in \mathcal{I}$ and (I1) imply that

$$(J + u) \cap (\Omega(I) \cup \Omega^-(I)) = (J \cap (\Omega(I) \cup \Omega^-(I))) + u \in \mathcal{I},$$

which contradicts (2). Assume that there exists an arc in $\mathsf{D}(J)$ from a vertex u in $\Omega^+(I)$ to a vertex v in $\Omega(I) \cup \Omega^-(I)$. If u is in I , then $J + v - u \in \mathcal{I}$. This and (I1) imply that

$$(J + v - u) \cap (\Omega(I) \cup \Omega^-(I)) = (J \cap (\Omega(I) \cup \Omega^-(I))) + v \in \mathcal{I}.$$

This contradicts (2). If u is not in I , then $J + u - v \in \mathcal{J}$. This and (I1) imply that

$$(J + u - v) \cap \Omega^+(I) = (J \cap \Omega^+(I)) + u \in \mathcal{J}.$$

This contradicts (3). By exchanging the roles of I and J , we can prove that $\Omega^+(I) \subseteq \Omega^+(J)$, which implies that $\Omega^+(J) = \Omega^+(I)$.

Next we prove that $\Omega^-(J) = \Omega^-(I)$. For proving this, we first prove that $\Omega^-(J) \subseteq \Omega^-(I)$. Assume that $\mathsf{T}_{\mathbf{N}}(J) \not\subseteq \Omega^-(I)$, and let u be an element in $\mathsf{T}_{\mathbf{N}}(J) \setminus \Omega^-(I)$. Then, $J + u \in \mathcal{J}$ and (I1) imply that

$$(J + u) \cap (\Omega^+(I) \cup \Omega(I)) = (J \cap (\Omega^+(I) \cup \Omega(I))) + u \in \mathcal{J}.$$

This contradicts (4). Assume that there exists an arc in $\mathsf{D}(J)$ from u in $\Omega^+(I) \cup \Omega(I)$ to v in $\Omega^-(I)$. If u is in I , then $J + v - u \in \mathcal{I}$. This and (I1) imply that

$$(J + v - u) \cap \Omega^-(I) = (J \cap \Omega^-(I)) + v \in \mathcal{I}.$$

This contradicts (1). If u is not in I , then $J + u - v \in \mathcal{J}$. This and (I1) imply that

$$(J + u - v) \cap (\Omega^+(I) \cup \Omega(I)) = (J \cap (\Omega^+(I) \cup \Omega(I))) + u \in \mathcal{J}.$$

This contradicts (4). By exchanging the roles of I and J , we can prove that $\Omega^-(I) \subseteq \Omega^-(J)$, which implies that $\Omega^-(J) = \Omega^-(I)$. This completes the proof. $\square$

Lemma 8 implies that for every matroids $\mathbf{M}$ and $\mathbf{N}$ on the same ground set, $\Omega_{\mathbf{MN}}^+(I)$, $\Omega_{\mathbf{MN}}(I)$, and $\Omega_{\mathbf{MN}}^-(I)$ do not depend a choice of a maximum-size common independent set I of $\mathbf{M}$ and $\mathbf{N}$. Thus, for each matroids $\mathbf{M}$ and $\mathbf{N}$ on the same ground set and some maximum-size common independent set I , we define

$$\begin{aligned}\Omega_{\mathbf{MN}}^+ &:= \Omega_{\mathbf{MN}}^+(I), \\ \Omega_{\mathbf{MN}} &:= \Omega_{\mathbf{MN}}(I), \\ \Omega_{\mathbf{MN}}^- &:= \Omega_{\mathbf{MN}}^-(I),\end{aligned}$$

respectively.

3 Characterization

In this section, we give a characterization of a popular matching. Define

$$\mathcal{U} := \left\{ F \subseteq E \mid |F(a)| \leq 1 \text{ for every applicant } a \text{ in } A \right\},$$

and $\mathcal{A} := (E, \mathcal{U})$. It is not difficult to see that $\mathcal{A}$ is a matroid. Define

$$\mathbf{P} := \bigoplus_{p \in P} \mathbf{M}_p.$$

Notice that for each subset M of E , M is a matching in G if and only if M is a common independent set of $\mathbf{A}$ and $\mathbf{P}$ such that $|M| = |A|$. For each applicant a in A , we define $f(a)$ by

$$f(a) := \left\{ e \in E(a) \mid e \precsim_a g \text{ for every edge } g \text{ in } E(a) \right\}.$$

Notice that for each applicant a in A , there exists a possibility that $|f(a)| > 1$. Define

$$\begin{aligned}\Gamma &:= \bigcup_{a \in A} f(a), \\ \mathbf{B} &:= \mathbf{A}|\Gamma, \\ \mathbf{Q} &:= \mathbf{P}|\Gamma \quad \left(= \bigoplus_{p \in P} (\mathbf{M}_p|\Gamma(p)) \right).\end{aligned}$$

In addition, we define

$$T := \left\{ e \in E \setminus \Gamma \mid \{e\} \text{ is an independent set of } \mathbf{P}/(\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}}) \right\}.$$

Notice that for each applicant a in A , since there exists the last resort post $\ell(a)$, $T(a) \neq \emptyset$. For each applicant a in A , we define $s(a)$ by

$$s(a) := \left\{ e \in T(a) \mid e \succsim_a g \text{ for every edge } g \text{ in } T(a) \right\}.$$

In addition, we define

$$\begin{aligned}S &:= \bigcup_{a \in A} s(a), \\ \Pi &:= \Gamma \cup S.\end{aligned}$$

The following characterization can be regarded as a generalization of [1, Lemma 3.5].

Theorem 9. *For each matching M in G , M is a popular matching in G if and only if*

(P1) *$M \cap \Gamma$ is a maximum-size common independent set of $\mathbf{B}$ and $\mathbf{Q}$, and*

(P2) *M is a subset of Π .*

We first give lemmas that are necessary for proving the *only if*-part of Theorem 9.

Lemma 10. *Assume that we are given a popular matching M in G . Then, $M \cap \Gamma$ is a maximum-size common independent set of $\mathbf{B}$ and $\mathbf{Q}$.*

Since the proof of Lemma 10 is long, we leave it to Section 3.1.

Lemma 11. *Assume that we are given a popular matching M in G . Then, there exists no applicant a in A such that $e \succ_a \mu_M(a) \succ_a g$ for edges e in $f(a)$ and g in $s(a)$.*

Proof. For proving this lemma by contradiction, we assume that there exists an applicant a in A such that $e \succ_a \mu_M(a) \succ_a g$ for edges e in $f(a)$ and g in $s(a)$. Since M is a popular matching in G , Lemma 10 implies that $M \cap \Gamma$ is a maximum common independent set of $\mathbf{B}$ and $\mathbf{Q}$. This and Lemma 7 imply that $M \cap (\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})$ is a base of $\mathbf{Q}|(\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})$ ($= \mathbf{P}|(\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})$). Define

$$N := (M \cap (\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})) + \mu_M(a). \quad (5)$$

Since $N \subseteq M$ and M is an independent set of $\mathbf{P}$, N is an independent set of $\mathbf{P}$. Thus, (5) and Lemma 1 imply that $\mu_M(a)$ is an independent set of $\mathbf{P}/(\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})$. However, this contradicts that fact that $\mu_M(a) \succ_a g$ for an edge g in $s(a)$. This completes the proof. $\square$

Lemma 12. Assume that we are given a popular matching M in G . Then, there exists no applicant a in A such that $\mu_M(a) \notin \Pi$.

We leave the proof of Lemma 12 to Section 3.2.

Next we give a lemma that is necessary for proving the *if*-part of Theorem 9. For each matching M in G , we denote by $\mathbf{b}(M)$ the set of edges $e = (a, p)$ in E such that $e \succ_a g$ for an edge g in $s(a)$.

Lemma 13. For every matching M in G , we have $\gamma_{\mathbf{BQ}} \geq |\mathbf{b}(M)|$.

Proof. For proving this lemma by contradiction, we assume that there exists a matching M in G such that $\gamma_{\mathbf{BQ}} < |\mathbf{b}(M)|$. Let N be a maximum-size common independent set in $\mathbf{B}$ and $\mathbf{Q}$ such that

$$\forall N' \in \mathcal{B}_{\mathbf{BQ}}: |M \cap N| \geq |M \cap N'|. \quad (6)$$

Since $\gamma_{\mathbf{BQ}} < |\mathbf{b}(M)|$, we have $|N| < |\mathbf{b}(M)|$. In addition, since M is an independent set of $\mathbf{P}$, (I1) implies that $\mathbf{b}(M)$ is an independent set of $\mathbf{P}$. Furthermore, N is an independent set of $\mathbf{P}$. Thus, (I2) implies that there exists an edge $e = (a, p)$ in $\mathbf{b}(M) \setminus N$ such that $N + e$ is an independent set of $\mathbf{P}$. Notice that since e is in $\mathbf{b}(M)$, $e \succ_a g$ for an edge g in $s(a)$.

We first consider the case where $e \notin \Gamma$. Since $N + e$ is an independent set of $\mathbf{P}$, (I1) implies that

$$(N \cap (\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})) + e$$

is an independent set of $\mathbf{P}$. Since it follows from Lemma 7 that $N \cap (\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})$ is a base of $\mathbf{Q} | (\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})$ ($= \mathbf{P} | (\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})$), $\{e\}$ is an independent set of $\mathbf{P} / (\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})$. This contradicts the fact that $e \succ_a g$ for an edge g in $s(a)$.

Next we consider the case where $e \in \Gamma$, i.e., $N + e \subseteq \Gamma$. Since N is a maximum-size common independent set in $\mathbf{B}$ and $\mathbf{Q}$, $N + e$ is not an independent set of $\mathbf{A}$. That is, $|N(a)| = 1$. In addition, since $e \in M$ and M is a matching in G , $\mu_N(a) \notin M$. Thus,

$$|(N + e - \mu_N(a)) \cap M| = |M \cap N| + 1. \quad (7)$$

Since N is an independent set of $\mathbf{B}$, $N + e - \mu_N(a)$ is an independent set of $\mathbf{B}$. Since $N + e$ is an independent set of $\mathbf{Q}$, (I1) implies that $N + e - \mu_N(a)$ is a maximum-size common independent set in $\mathbf{B}$ and $\mathbf{Q}$. These and (7) contradict (6), which completes the proof. $\square$

We are now ready to prove Theorem 9.

Proof of Theorem 9. Since the *only if*-part follows from Lemmas 10 and 12, we prove the *if*-part. Let M be a matching in G satisfying (P1) and (P2), and assume that we are given a matching N in G . We denote by A_M and A_N be the sets of applicants a in A such that $\mu_M(a) \succ_a \mu_N(a)$ and $\mu_N(a) \succ_a \mu_M(a)$, respectively. If we can prove that $|A_M| \geq |A_N|$, then the proof is done. For proving this, it is sufficient to construct an injective function $\tau: A_N \rightarrow A_M$.

Lemma 13 and (P1) imply that there exists an injective function

$$\varphi: \mathbf{b}(N) \setminus (M \cap \Gamma) \rightarrow (M \cap \Gamma) \setminus \mathbf{b}(N).$$

Let a be an applicant in A_N . Since $\mu_N(a) \succ_a \mu_M(a)$, (P2) implies that $\mu_M(a)$ is in $s(a)$. Thus, $\mu_N(a)$ is in $\mathbf{b}(N) \setminus (M \cap \Gamma)$. Here we consider the following procedure.

Step 1. Set $i := 1$, and define $e_1 = (a_1, p_1)$ as $\varphi(\mu_N(a))$.

Step 2. If we have $e_i \succ_{a_i} \mu_N(a_i)$, then we define $\tau(a) := a_i$ and halt. Otherwise, define $e_{i+1} := (a_{i+1}, p_{i+1})$ as $\varphi(\mu_N(a_i))$ and update $i := i + 1$. Then, repeat **Step 2**.

For proving that this procedure is well-defined, it is sufficient to prove in **Step 2**, $\mu_N(a_i)$ is in $\mathbf{b}(N) \setminus (M \cap \Gamma)$. Since $e_i \in (M \cap \Gamma) \setminus \mathbf{b}(N)$, if $e_i \in N$, then $e_i \in f(a_i)$ implies that $e_i \in \mathbf{b}(N)$. This contradicts the fact that $e_i \in (M \cap \Gamma) \setminus \mathbf{b}(N)$. Thus, $e_i \notin N$. In addition, since $e_i \in f(a_i)$, if $e_i \not\succ_{a_i} \mu_N(a_i)$, then $\mu_N(a_i) \in f(a_i)$. Thus, $\mu_N(a_i) \in \mathbf{b}(N) \setminus (M \cap \Gamma)$.

Since G is finite and φ is injective, this procedure halts. Thus, since φ is injective, τ is clearly injective. This completes the proof. $\square$

3.1 Proof of Lemma 10

Here we give a proof of Lemma 10. In the sequel, for each common independent set I of $\mathbf{B}$ and $\mathbf{Q}$, we do not distinguish between each vertex of $\mathbf{D}_{\mathbf{BQ}}(I)$ and the edge in Γ that it corresponds to. That is, we may call a vertex of $\mathbf{D}_{\mathbf{BQ}}(I)$ an edge. Since $\mathbf{D}_{\mathbf{BQ}}(I)$ is a directed graph (i.e., it contains only arcs), it does not make any confusion.

Since M is a matching in G , M is a common independent set of $\mathbf{A}$ and $\mathbf{P}$. This and (I1) imply that $M \cap \Gamma$ is a common independent set of $\mathbf{B}$ and $\mathbf{Q}$. For proving this lemma by contradiction, we assume that $M \cap \Gamma$ is not a maximum-size common independent set of $\mathbf{B}$ and $\mathbf{Q}$. It follows from this and Lemma 5 that there exist an edge $g = (b, q)$ in $\mathbf{T}_{\mathbf{B}}(M \cap \Gamma)$ and an edge $h = (c, r)$ in $\mathbf{T}_{\mathbf{Q}}(M \cap \Gamma)$ such that there exists a directed path in $\mathbf{D}_{\mathbf{BQ}}(M \cap \Gamma)$ from g to h . Let L be a shortest directed path in $\mathbf{D}_{\mathbf{BQ}}(M \cap \Gamma)$ from g to h , and we denote by $K = \{e_1, e_2, \dots, e_k\}$ the set of edges in Γ (i.e., vertices of $\mathbf{D}_{\mathbf{BQ}}(M \cap \Gamma)$) that L goes through. Furthermore, we assume that $e_i := (a_i, p_i)$ for each $i = 1, 2, \dots, k$, and L goes through $e_1, e_2, \dots, e_k$ in this order. Notice that $e_1 = g$, $e_k = h$, k is odd, and $e_i \in M$ for every $i = 2, 4, \dots, k-1$. Define $N := M \triangle K$ and $N_0 := N - \mu_M(b)$. Lemma 6 implies that $N \cap \Gamma (= N_0 \cap \Gamma)$ is an independent set of $\mathbf{Q}$.

For completing a proof of Lemma 10, we first prove necessary lemmas (Section 3.1.1), and then complete a proof of Lemma 3.1 (Section 3.1.2).

3.1.1 Necessary lemmas

Here we prove necessary lemmas for proving Lemma 10.

Lemma 14. *For every post p in $P - r$, we have $N(p) \in \mathcal{I}_p$ and $N_0(p) \in \mathcal{I}_p$.*

Proof. Let p be a post in $P - r$. By using Lemma 4, we prove that $N(p) \in \mathcal{I}_p$. If we can prove this, then (I1) implies that $N_0(p) \in \mathcal{I}_p$.

Define $I := M(p) \cap \Gamma$, $J := M(p) \setminus \Gamma$, $X := K(p) \setminus M$, and $Y := M(p) \cap K$. Then,

$$\begin{aligned} (I \cup X) \setminus Y &= N(p) \cap \Gamma, \\ ((I \cup X) \setminus Y) \cup J &= N(p). \end{aligned}$$

Since M is an independent set of $\mathbf{P}$, the definition of $\mathbf{P}$ implies that $I \cup J \in \mathcal{I}_r$. In addition, the definition of $\mathbf{Q}$ implies that $(I \cup X) \setminus Y$ is an independent set of $\mathbf{M}_p|_{\Gamma(p)}$,

i.e., $\mathbf{M}_p$. Furthermore, for every edge e in X , it follows from the definition of $\mathbf{D}_{\mathbf{BQ}}(M \cap \Gamma)$ that $(M \cap \Gamma) + e$ is not an independent set of $\mathbf{Q}$. Thus, since $M \cap \Gamma$ is an independent set of $\mathbf{Q}$, the definition of $\mathbf{Q}$ implies that $I + e$ is not an independent set of $\mathbf{M}_p|_{\Gamma(p)}$, i.e., $\mathbf{M}_p$. Thus, Lemma 4 implies that $N(p) \in \mathcal{I}_p$. $\square$

Lemma 15. *For every applicant a in $A - b$, (i) $|N(a)| = 1$, and (ii) $\mu_N(a) \preceq_a \mu_M(a)$.*

Proof. For every $i = 2, 4, \dots, k-1$, since there exists an arc in $\mathbf{D}_{\mathbf{BQ}}(M \cap \Gamma)$ from e_i to e_{i+1} , $M + e_{i+1} - e_i$ is an independent set of $\mathbf{B}$. Thus, the definition of $\mathbf{A}$ implies that we have $a_i = a_{i+1}$ for every $i = 2, 4, \dots, k-1$. This implies that for every applicant a in $A - b$, since $|M(a)| = 1$, we have $|N(a)| = 1$. Furthermore, since $K \subseteq \Gamma$, $\mu_N(a) \in f(a)$ for every applicant a in $A - b$, which implies (ii). This completes the proof. $\square$

Lemma 16. (i) $g \succ_b \mu_M(b)$, and (ii) $|N_0(b)| = 1$.

Proof. Since $g \in \mathbf{T}_{\mathbf{B}}(M \cap \Gamma)$, $M(b) \cap \Gamma = \emptyset$. Thus, $\mu_M(b) \notin \Gamma$. This and $g \in \Gamma$ imply that $g \succ_b \mu_M(b)$. As proved in the proof of Lemma 15, $a_i = a_{i+1}$ for every $i = 2, 4, \dots, k-1$. Thus, since $\mu_M(b) \notin \Gamma$, we have $K(b) = \{g\}$. This and $M(b) = \{\mu_M(b)\}$ imply (ii). $\square$

We denote by L' the subpath of L from g to e_{k-1} . Since L is a shortest directed path in $\mathbf{D}_{\mathbf{BQ}}(M \cap \Gamma)$ from g to h , L' is a shortest directed path in $\mathbf{D}_{\mathbf{BQ}}(M \cap \Gamma)$ from g to e_{k-1} . Define $K' := K - h$. Then, $N - h = M \triangle K'$. Lemma 6 implies that $(N - h) \cap \Gamma$ is an independent set of $\mathbf{Q}$.

Lemma 17. $N(r) - h \in \mathcal{I}_r$ and $N_0(r) - h \in \mathcal{I}_r$.

Proof. By using Lemma 4, we prove that $N(r) - h \in \mathcal{I}_r$. If we can prove this, then (I1) implies that $N_0(r) - h \in \mathcal{I}_r$.

Define $I := M(r) \cap \Gamma$, $J := M(r) \setminus \Gamma$, $X := K'(r) \setminus M$, and $Y := M(r) \cap K'$. Then,

$$\begin{aligned} (I \cup X) \setminus Y &= (N(r) - h) \cap \Gamma, \\ ((I \cup X) \setminus Y) \cup J &= N(r) - h. \end{aligned}$$

Since M is an independent set of $\mathbf{P}$, the definition of $\mathbf{P}$ implies that $I \cup J$ is in $\mathcal{I}_r$. In addition, the definition of $\mathbf{Q}$ implies that $(I \cup X) \setminus Y$ is in $\mathcal{I}_r$. For every edge e in X , the definition of $\mathbf{D}_{\mathbf{BQ}}(M \cap \Gamma)$ implies that $(M \cap \Gamma) + e$ is not an independent set of $\mathbf{Q}$. Thus, since $M \cap \Gamma$ is an independent set of $\mathbf{Q}$, this and the definition of $\mathbf{Q}$ imply that $I + e$ is not an independent set of $\mathbf{M}_r$. Thus, Lemma 4 implies that $N(r) - h \in \mathcal{I}_r$. $\square$

3.1.2 Completing a proof

We are now ready to complete a proof of Lemma 10. If $N_0(r) \in \mathcal{I}_r$, then it follows from Lemmas 14, 15(i), and 16(ii) that N_0 is a matching in G . In addition, Lemma 15(ii) and 16(i) imply that $\phi(N_0; M) \geq 1$, which contradicts the fact that M is a popular matching in G . Thus, in the sequel, we can assume that $N_0(r) \notin \mathcal{I}_r$. In this case, since $N_0(r) \subseteq N(r)$, $N(r)$ is not in $\mathcal{I}_r$. Thus, Lemma 17 implies that $\mathbf{C}_{\mathbf{M}_r}(h, N(r) - h)$ and $\mathbf{C}_{\mathbf{M}_r}(h, N_0(r) - h)$ are well-defined. Define

$$\begin{aligned} C &:= \mathbf{C}_{\mathbf{M}_r}(h, N(r) - h), \\ C_0 &:= \mathbf{C}_{\mathbf{M}_r}(h, N_0(r) - h). \end{aligned}$$

Notice that Lemma 3 implies that $C = C_0$.

Since $N_0 \cap \Gamma$ is an independent set of $\mathbf{Q}$, it follows from the definition of $\mathbf{Q}$ that $N_0(r) \cap \Gamma$ is in $\mathcal{I}_r$. Thus, $C_0 \subseteq N_0(r)$ implies that $C_0 \not\subseteq \Gamma$. Let $g_1 = (b_1, r)$ be an edge in $C_0 \setminus \Gamma$. Since $g \in \Gamma$ and $g_1 \notin \Gamma$, we have $g \neq g_1$. Thus, since $g, g_1 \in N_0$, we have $b \neq b_1$. Define $N_1 := N_0 - g_1$. Then, $g_1 \in C_0$ and Lemmas 14, 15(i), and 16(ii) imply that

- $|N_1(a)| = 1$ for every applicant a in $A - b_1$, and $N_1(b_1) = \emptyset$,
- $N_1(p) \in \mathcal{I}_p$ for every post p in P .

Let $h_1 = (b_1, q_1)$ be an edge in $f(b_1)$. We first consider the case where $N_1(q_1) + h_1 \in \mathcal{I}_{q_1}$. Define $N_2 := N_1 + h_1$. In this case, N_2 is a matching in G . Furthermore, since it follows from $g_1 \notin \Gamma$, Lemmas 15(ii) and 16(i) that

$$\begin{aligned}\mu_{N_2}(b) &= g \succ_b \mu_M(b), \\ \mu_{N_2}(b_1) &= h_1 \succ_{b_1} g_1 = \mu_N(b_1) \succ_{b_1} \mu_M(b_1), \\ \forall a \in A \setminus \{b, b_1\} : \mu_{N_2}(a) &= \mu_N(a) \succ_a \mu_M(a),\end{aligned}$$

we have $\phi(N_2; M) \geq 2$. These contradict the fact that M is a popular matching in G .

Next we consider the case where $N_1(q_1) + h_1 \notin \mathcal{I}_{q_1}$. Define

$$C_1 := \mathbf{C}_{\mathbf{M}_{q_1}}(h_1, N_1(q_1)).$$

We first assume that at least one of $q_1 \neq q$ and $C_1 \setminus \{g, h_1\} \neq \emptyset$ holds. Let $g_2 = (b_2, q_1)$ be an edge in $C_1 \setminus \{g, h_1\}$. Since $g_2 \neq g, h_1$ and $g, g_2 \in N_1$, we have $b_2 \neq b, b_1$. Define

$$N_3 := N_2 - g_2 + \ell(b_2).$$

Then, N_3 is a matching in G . Since Lemmas 15(ii) and 16(i) imply that

$$\begin{aligned}\mu_{N_3}(b) &= g \succ_b \mu_M(b), \\ \mu_{N_3}(b_1) &= h_1 \succ_{b_1} g_1 = \mu_N(b_1) \succ_{b_1} \mu_M(b_1), \\ \mu_M(b_2) \succ_{b_2} \ell(b_2) &= \mu_{N_3}(b_2), \\ \forall a \in A \setminus \{b, b_1, b_2\} : \mu_{N_3}(a) &= \mu_N(a) \succ_a \mu_M(a),\end{aligned}$$

we have $\phi(N_3; M) \geq 1$. This contradicts the fact that M is a popular matching in G .

Next we consider the case where $q_1 = q$ and $C_1 = \{g, h_1\}$. Define

$$N_4 := N_2 - g + \mu_M(b).$$

Assume that $\mu_M(b) = (b, q')$. Notice that $q' \neq q$. If $q' = r$, then

$$N_4(r) = N_2(r) + \mu_M(b) = N_1(r) + \mu_M(b) = N_0(r) + \mu_M(b) - g_1 = N(r) - g_1.$$

Since $g_1 \in C_0 = C$, $N(r) - g_1 \in \mathcal{I}_r$ and $N_4(r)$ is an independent set of $\mathbf{M}_r$. If $q' \neq r$, then

$$N_4(q') = N_2(q') + \mu_M(b) = N_1(q') + \mu_M(b) = N_0(q') + \mu_M(b) = N(q').$$

Lemma 14 implies that $N_4(q')$ is an independent set of $\mathbf{M}_r$. In both cases, N_4 is a matching in G . Furthermore, since Lemma 15(ii) implies that

$$\begin{aligned}\mu_{N_4}(b) &= \mu_M(b), \\ \mu_{N_4}(b_1) &= h_1 \succ_{b_1} g_1 = \mu_N(b_1) \succ_{b_1} \mu_M(b_1), \\ \forall a \in A \setminus \{b, b_1\}: \mu_{N_4}(a) &= \mu_N(a) \succ_a \mu_M(a),\end{aligned}$$

we have $\phi(N_4; M) \geq 1$. This contradicts the fact that M is a popular matching in G , and completes the proof.

3.2 Proof of Lemma 12

In this subsection, we give a proof of Lemma 12.

For proving this lemma by contradiction, we assume that there exists an applicant b in A such that $\mu_M(b) \notin \Pi$. Then, Lemma 11 implies that $e \succ_b \mu_M(b)$ for an edge e in $s(b)$. Let $g = (b, q)$ be an edge in $s(b)$. Define $N := M + g - \mu_M(b)$. Since $N(q) = M(q) + g$, if $M(q) + g \in \mathcal{I}_q$, then (I1) implies that N is a matching in G . Furthermore, $\phi(N; M) = 1$. This contradicts the fact that M is a popular matching in G . Thus, in the sequel, we can assume that $M(q) + g \notin \mathcal{I}_q$.

We first consider the case where

$$\mathbf{C}_{\mathbf{M}_q}(g, M(q)) - g \not\subseteq \Gamma.$$

Let $e_1 = (a_1, q)$ be an edge in

$$(\mathbf{C}_{\mathbf{M}_q}(g, M(q)) - g) \setminus \Gamma.$$

Since $g \neq e_1$ holds, we have $a_1 \neq b$. Let $e_2 = (a_1, q_1)$ be an edge in $f(a_1)$. Since $e_1 \notin \Gamma$ and $e_2 \in \Gamma$ (i.e., $e_1 \neq e_2$), we have $q_1 \neq q$. Define $N' := N + e_2 - e_1$. If $N(q_1) + e_2 \in \mathcal{I}_{q_1}$, then N' is a matching in G . Furthermore, we have $\phi(N'; M) = 2$, which contradicts the fact that M is a popular matching in G . Assume that $N(q_1) + e_2 \notin \mathcal{I}_{q_1}$. Let $e_3 = (a_2, q_1)$ be an edge in $\mathbf{C}_{\mathbf{M}_{q_1}}(e_2, N(q_1)) - e_2$. Since $q_1 \neq q$, we have $e_3 \neq g$. Define $N'' := N' - e_3 + \ell(a_2)$. Then, N'' is a matching in G and we have $\phi(N''; M) = 1$. This contradicts the fact that M is a popular matching in G .

Next we consider the case where

$$\mathbf{C}_{\mathbf{M}_q}(g, M(q)) - g \subseteq \Gamma. \tag{8}$$

Since g is in $s(a)$, Lemmas 1, 7, and 10 imply

$$(M \cap (\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})) + g$$

is an independent set of $\mathbf{P}$. This and the definition of $\mathbf{P}$ imply that

$$(M(q) \cap (\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})) + g \in \mathcal{I}_q.$$

Thus,

$$\mathbf{C}_{\mathbf{M}_q}(g, M(q)) - g \not\subseteq (M(q) \cap (\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})).$$

This implies that there exists an edge in $\mathbf{C}_{\mathbf{M}_q}(g, M(q)) \cap \Omega_{\mathbf{BQ}}^-$.

Since (8) implies that

$$\mathbf{C}_{\mathbf{M}_q}(g, M(q)) \subseteq (M(q) \cap \Gamma) + g,$$

we have $(M(q) \cap \Gamma) + g \notin \mathcal{I}_q$. Since (I1) and $M(q) \in \mathcal{I}_q$ imply that $M(q) \cap \Gamma \in \mathcal{I}_q$, this and Lemma 3 imply that

$$\mathbf{C}_{\mathbf{M}_q}(g, M(q) \cap \Gamma) = \mathbf{C}_{\mathbf{M}_q}(g, M(q)).$$

Thus,

$$\mathbf{C}_{\mathbf{M}_q}(g, M(q) \cap \Gamma) \cap \Omega_{\mathbf{BQ}}^- \neq \emptyset. \quad (9)$$

Define $\mathbf{C} := \mathbf{A} | (\Gamma + g)$ and $\mathbf{R} := \mathbf{P} | (\Gamma + g)$. It is not difficult to see that $\mathbf{D}_{\mathbf{CR}}(M \cap \Gamma)$ is obtained from $\mathbf{D}_{\mathbf{BQ}}(M \cap \Gamma)$ as follows.

Step 1. Add g to the vertex set of $\mathbf{D}_{\mathbf{BQ}}(M \cap \Gamma)$.

Step 2. Add an arc from g to every edge in $\mathbf{C}_{\mathbf{M}_q}(g, M(q) \cap \Gamma)$.

Furthermore, since $\mu_M(b) \notin f(b)$, we have

$$\begin{aligned} \mathbf{T}_{\mathbf{C}}(M \cap \Gamma) &= \mathbf{T}_{\mathbf{B}}(M \cap \Gamma) + g, \\ \mathbf{T}_{\mathbf{R}}(M \cap \Gamma) &= \mathbf{T}_{\mathbf{Q}}(M \cap \Gamma). \end{aligned}$$

It follows from (9) that there exists an edge h in $\mathbf{T}_{\mathbf{R}}(M \cap \Gamma)$ such that there a directed path in $\mathbf{D}_{\mathbf{CR}}(M \cap \Gamma)$ from g to h . Let L be a shortest directed path in $\mathbf{D}_{\mathbf{CR}}(M \cap \Gamma)$ from g to h . We denote by $K = \{e_1, e_2, \dots, e_k\}$ the set of edges in $\Gamma + g$ that L goes through. Assume that $e_i := (a_i, p_i)$ for each $i = 1, 2, \dots, k$, and L goes through $e_1, e_2, \dots, e_k$ in this order. Notice that $e_1 = g$, $e_k = h$, k is odd, and $e_i \in M$ for every $i = 2, 4, \dots, k-1$. Define $N := M \triangle K$ and $N_0 := N - \mu_M(b)$. Lemma 6 implies that $N \cap (\Gamma + g) (= N_0 \cap (\Gamma + g))$ is an independent set of $\mathbf{R}$.

For completing a proof of Lemma 12, we first necessary lemmas (Section 3.2.1), and complete a proof of Lemma 12 (Section 3.2.2).

3.2.1 Necessary lemmas

Here we give necessary lemmas for completing the proof of Lemma 12.

Lemma 18. *For every post p in $P - r$, we have $N(p) \in \mathcal{I}_p$ and $N_0(p) \in \mathcal{I}_p$.*

Proof. Let p be a post in $P - r$. By using Lemma 4, we prove that $N(p) \in \mathcal{I}_p$. If we can prove this, then (I1) implies that $N_0(p) \in \mathcal{I}_p$.

Define $I := M(p) \cap \Gamma$, $J := M(p) \setminus \Gamma$, $X := K(p) \setminus M$, and $Y := M(p) \cap K$. Then,

$$\begin{aligned} (I \cup X) \setminus Y &= N(p) \cap (\Gamma + g), \\ ((I \cup X) \setminus Y) \cup J &= N(p). \end{aligned}$$

Since M is an independent set of $\mathbf{P}$, the definition of $\mathbf{P}$ implies that $I \cup J$ is in $\mathcal{I}_p$. In addition, the definition of $\mathbf{R}$ implies that $(I \cup X) \setminus Y$ is in $\mathcal{I}_p$. For every edge e in X , the definition of $\mathbf{D}_{\mathbf{CR}}(M \cap \Gamma)$ implies that $(M \cap \Gamma) + e$ is not an independent set of $\mathbf{R}$. Thus, since $M \cap \Gamma$ is an independent set of $\mathbf{R}$, this and the definition of $\mathbf{R}$ imply that $I + e$ is not an independent set of $\mathbf{M}_p$. Thus, Lemma 4 implies that $N(p) \in \mathcal{I}_p$. $\square$

Lemma 19. For every applicant a in $A - b$, (i) $|N(a)| = 1$, and (ii) $\mu_N(a) \succeq_a \mu_M(a)$.

Proof. For every $i = 2, 4, \dots, k-1$, since there exists an arc in $D_{\mathbf{CR}}(M \cap \Gamma)$ from e_i to e_{i+1} , $M + e_{i+1} - e_i$ is an independent set of $\mathbf{C}$. Thus, the definition of $\mathbf{A}$ implies that $a_i = a_{i+1}$ for every $i = 2, 4, \dots, k-1$. This implies that for every applicant a in $A - b$, since $|M(a)| = 1$, we have $|N(a)| = 1$. Furthermore, since $K - g \subseteq \Gamma$, $\mu_N(a) \in f(a)$ for every applicant a in $A - b$, which implies (ii). This completes the proof. $\square$

Lemma 20. (i) $g \succ_b \mu_M(b)$, and (ii) $|N_0(b)| = 1$.

Proof. Since $g \in s(b)$ and $e \succ_b \mu_M(b)$ for an edge e in $s(b)$, we have $g \succ_b \mu_M(b)$. Furthermore, as proved in Lemma 19, we have $a_i = a_{i+1}$ for every $i = 1, 3, \dots, k-1$. Thus, since $\mu_M(b) \notin \Gamma$, we have $K(b) = \{g\}$. This and $M(b) = \{\mu_M(b)\}$ imply (ii). $\square$

We denote by L' the subpath of L from g to e_{k-1} . Since L is a shortest directed path in $D_{\mathbf{CR}}(M \cap \Gamma)$ from g to h , L' is a shortest directed path in $D_{\mathbf{CR}}(M \cap \Gamma)$ from g to e_{k-1} . Define $K' := K - h$. Then, $N - h = M \triangle K'$. Lemma 6 implies that $(N - h) \cap (\Gamma + g)$ is an independent set of $\mathbf{R}$.

Lemma 21. $N(r) - h \in \mathcal{I}_r$ and $N_0(r) - h \in \mathcal{I}_r$.

Proof. By using Lemma 4, we prove that $N(r) - h \in \mathcal{I}_r$. If we can prove this, then (I1) implies that $N_0(r) - h \in \mathcal{I}_r$.

Define $I := M(r) \cap \Gamma$, $J := M(r) \setminus \Gamma$, $X := K'(r) \setminus M$, and $Y := M(r) \cap K'$. Then,

$$\begin{aligned} (I \cup X) \setminus Y &= (N(r) - h) \cap (\Gamma + g), \\ ((I \cup X) \setminus Y) \cup J &= N(r) - h. \end{aligned}$$

Since M is an independent set of $\mathbf{P}$, the definition of $\mathbf{P}$ implies that $I \cup J$ is in $\mathcal{I}_r$. In addition, the definition of $\mathbf{R}$ implies that $(I \cup X) \setminus Y$ is in $\mathcal{I}_r$. For every edge e in X , the definition of $D_{\mathbf{CR}}(M \cap \Gamma)$ implies that $(M \cap \Gamma) + e$ is not an independent set of $\mathbf{R}$. Thus, since $M \cap \Gamma$ is an independent set of $\mathbf{R}$, this and the definition of $\mathbf{R}$ imply that $I + e$ is not an independent set of $\mathbf{M}_r$. Thus, Lemma 4 implies that $N(r) - h \in \mathcal{I}_r$. $\square$

3.2.2 Completing a proof

We are now ready to complete a proof of Lemma 12. If $N_0(r) \in \mathcal{I}_r$, then it follows from Lemmas 18, 19(i), and 20(ii) that N_0 is a matching in G . Furthermore, Lemma 19(ii) and 20(i) imply that $\phi(N_0; M) \geq 1$, which contradicts the fact that M is a popular matching in G . Thus, in the sequel, we can assume that $N_0(r) \notin \mathcal{I}_r$. In this case, since $N_0(r) \subseteq N(r)$, $N(r)$ is not in $\mathcal{I}_r$. Lemma 21 implies that $\mathbf{C}_{\mathbf{M}_r}(h, N(r) - h)$ and $\mathbf{C}_{\mathbf{M}_r}(h, N_0(r) - h)$ are well-defined. Define

$$\begin{aligned} C &:= \mathbf{C}_{\mathbf{M}_r}(h, N(r) - h), \\ C_0 &:= \mathbf{C}_{\mathbf{M}_r}(h, N_0(r) - h). \end{aligned}$$

Lemma 3 implies that $C = C_0$.

Since $N_0 \cap (\Gamma + g)$ is an independent set of $\mathbf{R}$, the definition of $\mathbf{R}$ implies that $N_0(r) \cap (\Gamma + g)$ is in $\mathcal{I}_r$. Thus, if $C_0 \subseteq \Gamma + g$, then $C_0 \subseteq N_0(r)$ implies that $C_0 \subseteq N_0(r) \cap (\Gamma + g)$.

This contradicts the fact that $N_0(r) \cap (\Gamma + g) \in \mathcal{I}_r$. Thus, $C_0 \not\subseteq \Gamma + g$. Let $g_1 = (b_1, r)$ be an edge in $C_0 \setminus (\Gamma + g)$. Notice that since $g \neq g_1$ and $g, g_1 \in N_0$, we have $b \neq b_1$. Define $N_1 := N_0 - g_1$. Then, $g_1 \in C_0$ and Lemmas 18, 19(i), and 20(ii) imply that

- $|N_1(a)| = 1$ for every applicant a in $A - b_1$, and $N_1(b_1) = \emptyset$,
- $N_1(p) \in \mathcal{I}_p$ for every post p in P .

Let $h_1 = (b_1, q_1)$ be an edge in $f(b_1)$. We first consider the case where $N_1(q_1) + h_1 \in \mathcal{I}_{q_1}$. Define $N_2 := N_1 + h_1$. In this case, N_2 is a matching in G . Furthermore, since it follows from $g_1 \notin \Gamma$, Lemmas 19(ii) and 20(i) that

$$\begin{aligned}\mu_{N_2}(b) &= g \succ_b \mu_M(b), \\ \mu_{N_2}(b_1) &= h_1 \succ_{b_1} g_1 = \mu_N(b_1) \succ_{b_1} \mu_M(b_1), \\ \forall a \in A \setminus \{b, b_1\}: \mu_{N_2}(a) &= \mu_N(a) \succ_a \mu_M(a),\end{aligned}$$

we have $\phi(N_2; M) \geq 2$. These contradict the fact that M is a popular matching in G .

Next we consider the case where $N_1(q_1) + h_1 \notin \mathcal{I}_{q_1}$. Define

$$C_1 := \mathbf{C}_{\mathbf{M}_{q_1}}(h_1, N_1(q_1)).$$

We first assume that at least one of $q_1 \neq q$ and $C_1 \setminus \{g, h_1\} \neq \emptyset$ holds. Let $g_2 = (b_2, q_1)$ be an edge in $C_1 \setminus \{g, h_1\}$. Since $g_2 \neq g, h_1$ and $g, g_2 \in N_1$, we have $b_2 \neq b, b_1$. Define

$$N_3 := N_2 - g_2 + \ell(b_2).$$

Then, N_3 is a matching in G . Since Lemmas 19(ii) and 20(i) imply that

$$\begin{aligned}\mu_{N_3}(b) &= g \succ_b \mu_M(b), \\ \mu_{N_3}(b_1) &= h_1 \succ_{b_1} g_1 = \mu_N(b_1) \succ_{b_1} \mu_M(b_1), \\ \mu_{N_3}(b_2) &\succ_{b_2} \ell(b_2) = \mu_{N_3}(b_2), \\ \forall a \in A \setminus \{b, b_1, b_2\}: \mu_{N_3}(a) &= \mu_N(a) \succ_a \mu_M(a),\end{aligned}$$

we have $\phi(N_3; M) \geq 1$. This contradicts the fact that M is a popular matching in G .

Next we consider the case where $q_1 = q$ and $C_1 = \{g, h_1\}$. Define

$$N_4 := N_2 - g + \mu_M(b).$$

Assume that $\mu_M(b) = (b, q')$. Notice that $q' \neq q$. If $q' = r$, then

$$N_4(r) = N_2(r) + \mu_M(b) = N_1(r) + \mu_M(b) = N_0(r) + \mu_M(b) - g_1 = N(r) - g_1.$$

Since $g_1 \in C_0 = C$, $N(r) - g_1 \in \mathcal{I}_r$ and $N_4(r)$ is an independent set of $\mathbf{M}_r$. If $q' \neq r$, then

$$N_4(q') = N_2(q') + \mu_M(b) = N_1(q') + \mu_M(b) = N_0(q') + \mu_M(b) = N(q').$$

Lemma 18 implies that $N_4(q')$ is an independent set of $\mathbf{M}_r$. In both cases, N_4 is a matching in G . Furthermore, since Lemma 19(ii) implies that

$$\begin{aligned}\mu_{N_4}(b) &= \mu_M(b), \\ \mu_{N_4}(b_1) &= h_1 \succ_{b_1} g_1 = \mu_N(b_1) \succ_{b_1} \mu_M(b_1), \\ \forall a \in A \setminus \{b, b_1\}: \mu_{N_4}(a) &= \mu_N(a) \succ_a \mu_M(a),\end{aligned}$$

we have $\phi(N_4; M) \geq 1$, which contradicts the fact that M is a popular matching in G . This completes the proof.

4 Algorithm

In this section, we propose our algorithm for PMTM. Define a function $\xi: \Pi \rightarrow \{0, 1\}$ by

$$\xi(e) := \begin{cases} 1 & \text{if } e \in \Gamma \\ 0 & \text{if } e \in \Pi \setminus \Gamma. \end{cases}$$

Our algorithm PMTM is described as follows.

Algorithm PMTM

Step 1. Compute Π , and define $\mathbf{A}' := \mathbf{A}|\Pi$ and $\mathbf{P}' := \mathbf{P}|\Pi$.

Step 2. Find a maximum-size common independent set M of $\mathbf{A}'$ and $\mathbf{P}'$ such that

$$\xi(M) = \max\{\xi(N) \mid N \in \mathcal{B}_{\mathbf{A}'\mathbf{P}'}\}.$$

Step 3. If $\xi(M) = \gamma_{\mathbf{BQ}}$ and $|M| = |A|$ hold, then output M and halt (in this case, M is a popular matching in G). Otherwise, output **null** and halt (in this case, there exists no popular matching in G).

End of Algorithm

For proving the correctness of the algorithm PMTM, we need the following lemma.

Lemma 22. *Assume that we are given a popular matching M in G . Then, M is a common independent set of $\mathbf{A}'$ and $\mathbf{P}'$ such that $\xi(M) = \gamma_{\mathbf{BQ}}$ and $|M| = |A|$.*

Proof. Since M is a matching in G , M is a common independent set of $\mathbf{A}$ and $\mathbf{P}$ such that $|M| = |A|$. In addition, (P2) of Theorem 9 implies that M is a subset of Π . Thus, M is a common independent set of $\mathbf{A}'$ and $\mathbf{P}'$. Since (P1) of Theorem 9 implies that $M \cap \Gamma$ is a maximum-size common independent set of $\mathbf{B}$ and $\mathbf{Q}$, we have $|M \cap \Gamma| = \gamma_{\mathbf{BQ}}$, i.e., $\xi(M) = \gamma_{\mathbf{BQ}}$. This completes the proof. $\square$

We are now ready to prove the correctness of the algorithm PMTM.

Theorem 23. *The algorithm PMTM can correctly solve PMTM.*

Proof. Let M be a common independent set of $\mathbf{A}'$ and $\mathbf{P}'$ that is found in **Step 2** of the algorithm PMTM. If the algorithm PMTM outputs M , then M is a matching and Theorem 9 implies that M is a popular matching in G .

Assume that the algorithm PMTM outputs **null**. Since M is a common independent set of $\mathbf{A}$ and $\mathbf{P}$, (I1) implies that $M \cap \Gamma$ is a common independent set of $\mathbf{B}$ and $\mathbf{Q}$. Thus, $|M \cap \Gamma| \leq \gamma_{\mathbf{BQ}}$, which implies that $\xi(M) \leq \gamma_{\mathbf{BQ}}$. Furthermore, since $|M(a)| \leq 1$ for every applicant a in A , we have $|M| \leq |A|$. Thus, $\xi(M) < \gamma_{\mathbf{BQ}}$ and/or $|M| < |A|$. We prove that in this case, there exists no popular matching in G by contradiction. Assume that there exists a popular matching N in G . Then, Lemma 22 implies that N is a common independent set of $\mathbf{A}'$ and $\mathbf{P}'$ such that $\xi(N) = \gamma_{\mathbf{BQ}}$ and $|N| = |A|$. If $|M| < |A|$, then the existence of N contradicts the fact that M is a maximum-size common independent

set of $\mathbf{A}'$ and $\mathbf{P}'$. If $\xi(M) < \gamma_{\mathbf{BQ}}$ and $|M| = |A|$, then the existence of N contradicts the fact that

$$\xi(M) = \max\{\xi(N') \mid N' \in \mathcal{B}_{\mathbf{A}'\mathbf{P}'}\}.$$

Thus, there exists no popular matching in G . This completes the proof. $\square$

Here we consider the time complexity of the algorithm **PMTM**. We denote by **EO** the time required to decide whether $I + e \in \mathcal{I}_p$ for every post p in P , every independent set I of $\mathbf{M}_p$, and every edge e in $E(p) \setminus I$. Define $m := |E|$. For simplicity, we assume that $E(p) \neq \emptyset$ for every post p in P and $\mathbf{EO} = \Omega(m)$. Furthermore, we assume that for every applicant a in A and every edges e, g in $E(a)$, we can decide in $O(1)$ time whether $e \succsim_a g$ holds.

We first consider the time complexity of **Step 1**. It is not difficult to see that we can compute $f(a)$ for all applicants a in $O(m)$ time. For computing $s(a)$ for all applicants a in A , we first compute $\Omega_{\mathbf{BQ}}^+$, $\Omega_{\mathbf{BQ}}$, and $\Omega_{\mathbf{BQ}}^-$ in $O(m^{2.5}\mathbf{EO})$ time by finding a maximum-size common independent set of $\mathbf{B}$ and $\mathbf{Q}$ via the algorithm of Cunningham [2]. Then, we find a base B of $\mathbf{P} | (\Omega_{\mathbf{BQ}}^+ \cup \Omega_{\mathbf{BQ}})$ in $O(m\mathbf{EO})$ time. Lemma 1 implies that by using the base B , we can compute $s(a)$ for all applicants a in A in $O(m\mathbf{EO})$ time. Thus, we can compute Π in $O(m^{2.5}\mathbf{EO})$ time.

Next we consider the time complexity of **Step 2**. It is not difficult to see that we can decide in $O(m)$ time whether $M + e$ is an independent set of $\mathbf{A}'$ for every independent set M of $\mathbf{A}'$ and every edge e in $\Pi \setminus M$. Furthermore, it is not difficult to see that we can decide in $O(\mathbf{EO})$ time whether $M + e$ is an independent set of $\mathbf{P}'$ for every independent set M of $\mathbf{P}'$ and every edge e in $\Pi \setminus M$. Thus, in **Step 2**, we can find a desired maximum-size common independent set of $\mathbf{A}'$ and $\mathbf{P}'$ in $O(m^3\mathbf{EO})$ time by using the algorithm of Frank [7]. Thus, the time complexity of the algorithm **PMTM** is $O(m^3\mathbf{EO})$.

References

- [1] D. J. Abraham, R. W. Irving, T. Kavitha, and K. Mehlhorn. Popular matchings. *SIAM Journal on Computing*, 37(4):1030–1045, 2007.
- [2] W. H. Cunningham. Improved bounds for matroid partition and intersection algorithms. *SIAM Journal on Computing*, 15(4):948–957, 1986.
- [3] A. L. Dulmage and N. S. Mendelsohn. Coverings of bipartite graphs. *Canadian Journal of Mathematics*, 10:517–534, 1958.
- [4] A. L. Dulmage and N. S. Mendelsohn. A structure theory of bipartite graphs of finite exterior dimension. *Transactions of the Royal Society of Canada, Section III*, 53:1–13, 1959.
- [5] J. Edmonds. Paths, trees, and flowers. *Canadian Journal of Mathematics*, 17:449–467, 1965.
- [6] T. Fleiner. A fixed-point approach to stable matchings and some applications. *Mathematics of Operations Research*, 28(1):103–126, 2003.

- [7] A. Frank. A weighted matroid intersection algorithm. *Journal of Algorithms*, 2(4):328–336, 1981.
- [8] T. Gallai. Kritische graphen II. *A Magyar Tudományos Akadémia – Matematikai Kutató Intézetének Közleményei*, 8:373–395, 1963.
- [9] T. Gallai. Maximale systeme unabhängiger kanten. *A Magyar Tudományos Akadémia – Matematikai Kutató Intézetének Közleményei*, 9:401–413, 1964.
- [10] P. Gärdenfors. Match making: Assignments based on bilateral preferences. *Behavioral Science*, 20(3):166–173, 1975.
- [11] N. Kamiyama. Matroid intersection with priority constraints. *Journal of the Operations Research Society of Japan*, 56(1):15–25, 2013.
- [12] N. Kamiyama. The popular matching and condensation problems under matroid constraints. In *Proceedings of the 8th Annual International Conference on Combinatorial Optimization and Applications*, volume 8881 of *Lecture Notes in Computer Science*, pages 713–728, 2014.
- [13] T. Kavitha, J. Mestre, and M. Nasre. Popular mixed matchings. *Theoretical Computer Science*, 412(24):2679–2690, 2011.
- [14] T. Kavitha and M. Nasre. Optimal popular matchings. *Discrete Applied Mathematics*, 157(14):3181–3186, 2009.
- [15] T. Kavitha and M. Nasre. Popular matchings with variable item copies. *Theoretical Computer Science*, 412(12):1263–1274, 2011.
- [16] T. Kavitha, M. Nasre, and P. Nimbhorkar. Popularity at minimum cost. *Journal of Combinatorial Optimization*, 27(3):574–596, 2014.
- [17] B. Korte and J. Vygen. *Combinatorial Optimization: Theory and Algorithms*. Springer, 5th edition, 2012.
- [18] L. Lovász and M. D. Plummer. *Matching Theory*. North-Holland, 1986.
- [19] D. F. Manlove and C. T. S. Sng. Popular matchings in the capacitated house allocation problem. In *Proceedings of the 14th European Symposium on Algorithms*, volume 4168 of *Lecture Notes in Computer Science*, pages 492–503, 2006.
- [20] E. McDermid and R. W. Irving. Popular matchings: structure and algorithms. *Journal of Combinatorial Optimization*, 22(3):339–358, 2011.
- [21] J. Mestre. Weighted popular matchings. *ACM Transactions on Algorithms*, 10(1):2:1–2:16, 2014.
- [22] K. Murota. *Matrices and Matroids for Systems Analysis*. Springer, 2000.
- [23] J. G. Oxley. *Matroid theory*. Oxford University Press, 2nd edition, 2011.

- [24] C. T. S. Sng and D. F. Manlove. Popular matchings in the weighted capacitated house allocation problem. *Journal of Discrete Algorithms*, 8(2):102–116, 2010.
- [25] Y.-W. Wu, W.-Y. Lin, H.-L. Wang, and K.-M. Chao. An optimal algorithm for the popular condensation problem. In *Proceedings of the 24th International Workshop on Combinatorial Algorithms*, volume 8288 of *Lecture Notes in Computer Science*, pages 412–422, 2013.
- [26] Y.-W. Wu, W.-Y. Lin, H.-L. Wang, and K.-M. Chao. The generalized popular condensation problem. In *Proceedings of the 25th International Symposium on Algorithms and Computation*, volume 8889 of *Lecture Notes in Computer Science*, pages 606–617, 2014.
- [27] R. Zenklusen. Matroidal degree-bounded minimum spanning trees. In *Proceedings of the 23rd Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1512–1521, 2012.

List of MI Preprint Series, Kyushu University

The Global COE Program

Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p-adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials
- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials

- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings
- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric τ -functions of the q-Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications

- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain
- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection
- MI2009-26 Manabu YOSHIDA
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance

- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA
Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI
Finite element computation for scattering problems of micro-hologram using DtN map
- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO
Abstract collision systems on groups

- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA
An algebraic approach to underdetermined experiments
- MI2010-10 Kei HIROSE & Sadanori KONISHI
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE
Lagrangian approach to weakly nonlinear stability of an elliptical flow
- MI2010-21 Hiroki MASUDA
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiro FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra

- MI2010-25 Toshimitsu TAKAESU
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI
Spin-spin correlation functions of the q -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and spike detection via Gaussian basis expansions
- MI2010-32 Nobutaka NAKAZONO
Hypergeometric τ functions of the q -Painlevé systems of type $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA
CAP representations of inner forms of $Sp(4)$ with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER
Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide FUKUMOTO & Alexander B. SAMOKHIN
Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI
Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI
Anderson-Darling test and the Malliavin calculus
- MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling via Compressed Sensing

- MI2011-5 Hiroshi INOUE
Implications in Compressed Sensing and the Restricted Isometry Property
- MI2011-6 Daeju KIM & Sadanori KONISHI
Predictive information criterion for nonlinear regression model based on basis expansion methods
- MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI
Group variable selection via relevance vector machine
- MI2011-8 Jan BREZINA & Yoshiyuki KAGEI
Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow
Group variable selection via relevance vector machine
- MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC
Recursive structures in the multispecies TASEP
- MI2011-10 Kazunori YASUTAKE
On projective space bundle with nef normalized tautological line bundle
- MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA
An explicit formula for the discrete power function associated with circle patterns of Schramm type
- MI2011-12 Yoshiyuki KAGEI
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow
- MI2011-13 Vladimír CHALUPECKÝ & Adrian MUNTEAN
Semi-discrete finite difference multiscale scheme for a concrete corrosion model: approximation estimates and convergence
- MI2011-14 Jun-ichi INOBUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves
- MI2011-15 Hiroshi INOUE
A generalization of restricted isometry property and applications to compressed sensing
- MI2011-16 Yu KAWAKAMI
A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space
- MI2011-17 Naoyuki KAMIYAMA
Matroid intersection with priority constraints
- MI2012-1 Kazufumi KIMOTO & Masato WAKAYAMA
Spectrum of non-commutative harmonic oscillators and residual modular forms
- MI2012-2 Hiroki MASUDA
Mighty convergence of the Gaussian quasi-likelihood random fields for ergodic Levy driven SDE observed at high frequency

- MI2012-3 Hiroshi INOUE
A Weak RIP of theory of compressed sensing and LASSO
- MI2012-4 Yasuhide FUKUMOTO & Youich MIE
Hamiltonian bifurcation theory for a rotating flow subject to elliptic straining field
- MI2012-5 Yu KAWAKAMI
On the maximal number of exceptional values of Gauss maps for various classes of surfaces
- MI2012-6 Marcio GAMEIRO, Yasuaki HIRAOKA, Shunsuke IZUMI, Miroslav KRAMAR, Konstantin MISCHAIKOW & Vidit NANDA
Topological Measurement of Protein Compressibility via Persistence Diagrams
- MI2012-7 Nobutaka NAKAZONO & Seiji NISHIOKA
Solutions to a q -analog of Painlevé III equation of type $D_7^{(1)}$
- MI2012-8 Naoyuki KAMIYAMA
A new approach to the Pareto stable matching problem
- MI2012-9 Jan BREZINA & Yoshiyuki KAGEI
Spectral properties of the linearized compressible Navier-Stokes equation around time-periodic parallel flow
- MI2012-10 Jan BREZINA
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a time-periodic parallel flow
- MI2012-11 Daeju KIM, Shuichi KAWANO & Yoshiyuki NINOMIYA
Adaptive basis expansion via the extended fused lasso
- MI2012-12 Masato WAKAYAMA
On simplicity of the lowest eigenvalue of non-commutative harmonic oscillators
- MI2012-13 Masatoshi OKITA
On the convergence rates for the compressible Navier- Stokes equations with potential force
- MI2013-1 Abuduwaili PAERHATI & Yasuhide FUKUMOTO
A Counter-example to Thomson-Tait-Chetayev's Theorem
- MI2013-2 Yasuhide FUKUMOTO & Hirofumi SAKUMA
A unified view of topological invariants of barotropic and baroclinic fluids and their application to formal stability analysis of three-dimensional ideal gas flows
- MI2013-3 Hiroki MASUDA
Asymptotics for functionals of self-normalized residuals of discretely observed stochastic processes
- MI2013-4 Naoyuki KAMIYAMA
On Counting Output Patterns of Logic Circuits
- MI2013-5 Hiroshi INOUE
RIPless Theory for Compressed Sensing

- MI2013-6 Hiroshi INOUE
Improved bounds on Restricted isometry for compressed sensing
- MI2013-7 Hidetoshi MATSUI
Variable and boundary selection for functional data via multiclass logistic regression modeling
- MI2013-8 Hidetoshi MATSUI
Variable selection for varying coefficient models with the sparse regularization
- MI2013-9 Naoyuki KAMIYAMA
Packing Arborescences in Acyclic Temporal Networks
- MI2013-10 Masato WAKAYAMA
Equivalence between the eigenvalue problem of non-commutative harmonic oscillators and existence of holomorphic solutions of Heun's differential equations, eigenstates degeneration, and Rabi's model
- MI2013-11 Masatoshi OKITA
Optimal decay rate for strong solutions in critical spaces to the compressible Navier-Stokes equations
- MI2013-12 Shuichi KAWANO, Ibuki HOSHINA, Kazuki MATSUDA & Sadanori KONISHI
Predictive model selection criteria for Bayesian lasso
- MI2013-13 Hayato CHIBA
The First Painleve Equation on the Weighted Projective Space
- MI2013-14 Hidetoshi MATSUI
Variable selection for functional linear models with functional predictors and a functional response
- MI2013-15 Naoyuki KAMIYAMA
The Fault-Tolerant Facility Location Problem with Submodular Penalties
- MI2013-16 Hidetoshi MATSUI
Selection of classification boundaries using the logistic regression
- MI2014-1 Naoyuki KAMIYAMA
Popular Matchings under Matroid Constraints
- MI2014-2 Yasuhide FUKUMOTO & Youichi MIE
Lagrangian approach to weakly nonlinear interaction of Kelvin waves and a symmetry-breaking bifurcation of a rotating flow
- MI2014-3 Reika AOYAMA
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Parallel flow in a cylindrical domain
- MI2014-4 Naoyuki KAMIYAMA
The Popular Condensation Problem under Matroid Constraints

- MI2014-5 Yoshiyuki KAGEI & Kazuyuki TSUDA
Existence and stability of time periodic solution to the compressible Navier-Stokes equation for time periodic external force with symmetry
- MI2014-6 This paper was withdrawn by the authors.
- MI2014-7 Masatoshi OKITA
On decay estimate of strong solutions in critical spaces for the compressible Navier-Stokes equations
- MI2014-8 Rong ZOU & Yasuhide FUKUMOTO
Local stability analysis of azimuthal magnetorotational instability of ideal MHD flows
- MI2014-9 Yoshiyuki KAGEI & Naoki MAKIO
Spectral properties of the linearized semigroup of the compressible Navier-Stokes equation on a periodic layer
- MI2014-10 Kazuyuki TSUDA
On the existence and stability of time periodic solution to the compressible Navier-Stokes equation on the whole space
- MI2014-11 Yoshiyuki KAGEI & Takaaki NISHIDA
Instability of plane Poiseuille flow in viscous compressible gas
- MI2014-12 Chien-Chung HUANG, Naonori KAKIMURA & Naoyuki KAMIYAMA
Exact and approximation algorithms for weighted matroid intersection
- MI2014-13 Yusuke SHIMIZU
Moment convergence of regularized least-squares estimator for linear regression model
- MI2015-1 Hidetoshi MATSUI
Sparse regularization for multivariate linear models for functional data
- MI2015-2 Reika AOYAMA & Yoshiyuki KAGEI
Spectral properties of the semigroup for the linearized compressible Navier-Stokes equation around a parallel flow in a cylindrical domain
- MI2015-3 Naoyuki KAMIYAMA
Stable Matchings with Ties, Master Preference Lists, and Matroid Constraints
- MI2015-4 Reika AOYAMA & Yoshiyuki KAGEI
Large time behavior of solutions to the compressible Navier-Stokes equations around a parallel flow in a cylindrical domain
- MI2015-5 Kazuyuki TSUDA
Existence and stability of time periodic solution to the compressible Navier-Stokes-Korteweg system on R^3
- MI2015-6 Naoyuki KAMIYAMA
Popular Matchings with Ties and Matroid Constraints