Identification and Characterization of a New Species, *Cladobotryum hypsigum*, that Causes Cobweb Disease in Beech Mushroom (*Hypsizygus marmoreus*) in Korea

Back, Chang-Gi

College of Agriculture and Life Sciences, Kyungpook National University

Lee, Seung-Yeol

College of Agriculture and Life Sciences, Kyungpook National University

Asaf, Sajjad

College of Agriculture and Life Sciences, Kyungpook National University

Kang, In-Kyu

College of Agriculture and Life Sciences, Kyungpook National University

他

<https://doi.org/10.5109/1526290>

出版情報：九州大学大学院農学研究院紀要. 60 (1), pp.7-12, 2015-02-27. Faculty of Agriculture, Kyushu University

バージョン：

権利関係：

Identification and Characterization of a New Species, *Cladobotryum hypsigum*, that Causes Cobweb Disease in Beech Mushroom (*Hypsizygus marmoreus*) in Korea

Chang-Gi BACK¹, Seung-Yeol LEE¹, Sajjad ASAF¹, In-Kyu KANG¹,
Hee-Young JUNG^{1*} and Shoji OHGA*

Laboratory of Forest Resources Management, Division of Forest Environmental Sciences,
Department of Agro-environmental Sciences, Faculty of Agriculture,
Kyushu University, Sasaguri, Fukuoka 811–2415, Japan
(Received October 3, 2014 and accepted November 14, 2014)

Symptoms similar to the cobweb disease were observed on beech mushrooms (*Hypsizygus marmoreus*) growing in Cheongdo-gun, Gyeongbuk Province, Korea, in May 2013. The causal agent was collected and cultured on potato dextrose agar media at different temperatures (5–30°C). The growth of the isolated fungus, characterized by the production of numerous sclerotia and red pigments, was temperature dependent, with minimum growth at 5°C and maximum growth at 25°C. A pathogenicity test revealed that the isolated fungus causes cobweb disease in the host mushroom *H. marmoreus* (white) as well as in three other mushrooms—*H. marmoreus* (brown), *Flammulina velutipes*, and *Pleurotus eryngii*—with massive mycelia and few conidia. Phylogenetic analysis of four genes (ITS, *RPB1*, *RPB2*, and *TEF*) showed that this isolate has a unique lineage and is distantly related to other species of *Cladobotryum* that cause the cobweb disease. The results showed that the isolated fungus is a new species, herein named *Cladobotryum hypsigum*, which causes cobweb disease in *H. marmoreus* in Korea.

Key words: Beech mushroom, *Cladobotryum hypsigum*, Cobweb disease, Mushroom disease

INTRODUCTION

Cobweb disease in mushrooms is characterized by the growth of coarse mycelia (Fletcher, *et al.*, 1989). For decades, *Hypomyces rosellus* was the most commonly reported pathogen of cobweb disease (McKay *et al.*, 1998; Bhatt and Singh, 2002; Potocnik *et al.*, 2008); however, during the last decade, the anamorph of *Hypomyces odoratus* has been reported with increasing frequency as the agent responsible for this disease, which has led to significant economic losses in commercial cultivation of mushrooms, mainly *Agaricus bisporus* (McKay *et al.*, 1999; Grogan and Gaze, 2000; Adie *et al.*, 2006; Khan *et al.*, 2008; Back *et al.*, 2010, 2012; Gea *et al.*, 2011, 2012). Reports have suggested that various *Cladobotryum* species such as *Cladobotryum dendroides*, *C. mycophilum*, *C. varium*, *C. multiseptatum*, and *C. verticillatum* cause diseases in several mushrooms (McKay *et al.*, 1999; Adie *et al.*, 2006).

The common symptoms of infection with *Cladobotryum*, the causal agent of cobweb disease, are the production of conidial masses and brown spots on mushroom caps (Back *et al.*, 2010). Studies of the infection pattern have suggested that the cobweb pathogen grows after infecting the casing layer. The rapid growth is assisted by spore dispersal in growing-rooms, covering the casing soil and the mushrooms with white cotton-like mycelium that can turn pinkish over time. The landing spores on developing mushrooms causes spotting,

whereas fruit bodies overgrown by mycelia develop soft, wet rot. The reported infections have mainly been attributed to the aurofusarin-producing fungal species *H. rosellus*, *H. odoratus*, and *C. multiseptatum* (Tamm and Poldmaa, 2013). Other aurofusarin-producing *Cladobotryum* species such as *C. asterophorum*, *C. protrusum*, and *C. paravirescens* have been reported on agaric, decorticated wood, and aphyllophoralean basidiomycete fungi, respectively. Although these species have been isolated from different hosts and are different from one another in conidia shape, size, and sclerotia-like aggregation, they all form the same pale, reddish pigments on potato dextrose agar (PDA) media due to aurofusarin (a secondary metabolite in *Cladobotryum* species) (De Hoog, 1978; Pöldmaa, 2011).

Plant pathologists are continuously seeking ways to suppress diseases such as cobweb disease to prevent economic loss. Chemical fungicides such as benzimidazole are generally used against cobweb disease-causing fungi on mushroom farms. However, *C. mycophilum* and *C. dendroides* became resistant to benzimidazole fungicides in the United Kingdom and are difficult to control (Grogan, 2006).

Hypsizygus marmoreus (beech mushroom) is an asidiomycete fungus cultivated for food in Korea, Japan, and Taiwan (Lee *et al.*, 2007). This mushroom is not only a delicious and nutritious food but also a source of hypsin, a ribosome-inactivating protein with documented medicinal qualities (Lam and Ng, 2001). The natural compounds in *H. marmoreus* have antitumor (Ikekawa *et al.*, 1992, 1995; Tsuchida *et al.*, 1995), antifungal, and antiproliferative activities (Lam and Ng, 2001).

In 2010, Back *et al.* identified *C. varium* as a causal agent of cobweb disease in *H. marmoreus* on commer-

¹ College of Agriculture and Life Sciences, Kyungpook National University, 80 Daehakro, Buk-gu, Daegu 702–701, Korea

* Corresponding author (E-mail: heeyoung@knu.ac.kr, ohga@forest.kyushu-u.ac.jp)

cial mushroom farms in Cheongdo-gun in the Gyeongbuk Province of Korea. However, in 2013, cobweb disease with different symptoms was observed in *H. marmoreus*. The white mycelium-covered fruit bodies of affected mushrooms had fewer conidia and a more scattered infection pattern compared to those infected by *C. varium* (Back *et al.*, 2012). Therefore, the purpose of this study was to isolate and identify the new causal agent of this disease using morphological and molecular characterization.

MATERIALS AND METHODS

Isolation and growth of fungi

The fungal samples were isolated in Cheongdo-gun, Gyeongbuk Province, Korea, in May 2013, from a diseased fruit body of *H. marmoreus* (white) with brown spots on the cap and few conidia, and it was covered with massive mycelium. The isolated fungi were cultured on PDA media in the dark at 25°C for 3 days. To confirm the morphological characteristics, we examined the shape, size, and color of 100 conidia and conidiophores of the isolated fungus using a microscope, Olympus BX-50 (Olympus, Tokyo, Japan). The fungus was then characterized using the descriptions of Gams and Hoozemans (1970) and Pöldmaa (2011). The mycelial plugs (less than 5 mm in diameter) were placed in the center of PDA media plates (90 mm in diameter). The plates were incubated at temperatures of 5°C, 10°C, 15°C, 20°C, 25°C, and 30°C for 8 d to identify the optimum growth temperature. The growth rate of the isolate was observed after 8 d, and the experiment was performed in triplicate.

Pathogenicity tests

Pathogenicity tests were performed at the first pin-head stage on four bottle-cultured mushrooms: *H. marmoreus* (white), *H. marmoreus* (brown), *F. velutipes*, and *P. eryngii*. The inocula were prepared using 10-day-old cultures on PDA media and adjusted to 1×10^2 conidia/mL. The spore suspension was sprayed on *H. marmoreus* (white). The inoculated mushrooms were covered with plastic bags to maintain humidity higher than 80% for 24 h and then incubated at 20°C. The pathogenicity of the isolated fungus was studied by using it to inoculate three additional mushrooms—*H. marmoreus* (brown), *F. velutipes*, and *P. eryngii*—following the procedure used for the initial inoculation.

DNA extraction and polymerase chain reaction (PCR) amplification

Total genomic DNA was extracted from the isolated fungus using lysis buffer according to a procedure described by Liu *et al.* (2000). The resultant total genomic DNA was used to amplify four protein-coding genes for internal transcribed spacer (ITS) region, RNA polymerase II subunit 1 (*RPB1*), RNA polymerase II subunit 2 (*RPB2*), and translation elongation factor (*TEF*) (exon 6 was amplified). These genes were amplified using the following primer pairs: gene for ITS: *ITS1F* (5'-CTT GGT CAT TTA GAG GAA GTA A-3')/*ITS4* (5'-

TCC TCC GCT TAT TGA TAT GC-3') (White *et al.*, 1990); *RPB1*: cRPB1Af (5'-CAY CCW GGY TTY ATC AAG AA-3')/cRPB1Cr (5'-CCN GCD ATN TCR TTR TCC ATR TA-3') (Chaverri *et al.*, 2008); *RPB2*: RPB2-5f (5'-GAY GAY MGW GAT CAY TTY GG-3')/7cR (5'-CCC ATR GCT TGY TTR CCC AT-3') (Liu *et al.*, 1999); *TEF*: EF1-983F (5'-GCY CCY GGH CAY CGT GAY TTY AT-3') (Carbone and Kohn, 1999), EF1-2218R (5'-ATG ACA CCR ACR GCR ACR GTY TG-3') (Rehner, 2001).

PCR amplification was performed in 20 µL of the reaction mixture containing 20 ng of fungal genomic DNA, 5 U of Taq polymerase (BIOFACT, Daejeon, Korea), 2 µL of 10X reaction buffer (100 mM Tris-HCl, 400 mM KCl, 15 mM MgCl₂, pH 9.0), 10 mM dNTP mixture, and 10 pmol of each primer using an Applied Biosystems 2720 thermal cycler (Applied Biosystems, Foster City, CA). The PCR conditions were 94°C for 3 min; 35 cycles of 30 s at 94°C, 30 s at 55°C for ITS and at 57°C for *RPB1*, *RPB2*, and *TEF*, and 1 min at 72°C; and a final extension at 72°C for 7 min. The PCR products were electrophoresed using 1% agarose gel and visualized in an ultraviolet illuminator. Then, the amplified DNA fragments were purified using a HiGene™ Gel & PCR Purification system (BIOFACT).

Sequencing and phylogenetic analysis

The purified DNA fragments were subjected to direct sequencing (BIOFACT). The sequences of the four genes were edited and aligned using DNASTAR (DNASTAR Inc., Madison, Wis.). The sequences were combined, and phylogenetic trees were constructed using the neighbor-joining method in CLUSTAL W (Thompson *et al.*, 1994). The phylogenetic trees for the four combined sequences were constructed from the data using the TreeView program (Win32, ver. 1.6.1). Bootstrap analysis with 1,000 replications was performed to determine support for various clades.

RESULTS AND DISCUSSION

Morphological characteristics of the isolated fungi

The natural symptoms of cobweb disease, including the growth of mycelium covering the affected mushrooms, production of few conidia, and brown spots on the mushroom caps, were observed on *H. marmoreus* in Cheongdo-gun in the Gyeongbuk Province in Korea in May 2013. In the early stages, white mycelia covered the fruit bodies of young mushrooms, whereas the infected fruit bodies rotted and masses of mycelia with few conidia covered them entirely in later stages (Fig. 1A–C). These symptoms differed from those previously reported for cobweb disease in *H. marmoreus* caused by *C. varium* (Back *et al.*, 2012).

According to our results, the isolate formed white mycelia on PDA media during the early stages at 22°C. This mycelium turned yellow after 8 days and then pinkish-red after 15 days (Fig. 1D and E). However, numerous sclerotia were observed after 20 days (Fig. 1E). The conidiophores were branched with two-celled conidia that are 15.9–21.8 µm long, 6.3–9.2 µm thick, and ovoid,

Fig. 1. Morphological characteristics of the isolated fungus. A: Symptoms of cobweb disease on host, B: Fruit body formation on host. C: Enlarged image of B. D: Colony of isolated fungus on potato dextrose agar (PDA) after 20 days at 22°C (front view). E: Development of sclerotia-like bodies (arrows) from the center of the colony (back view). F: Light microscope micrograph showing conidial morphology. G: Conidiophore-bearing conidia. Scale bars: F = 20 μm , G = 100 μm .

Table 1. Comparison of the morphological characteristics of the isolated fungi and other *Cladobotryum* species on potato dextrose agar media

	This study	<i>C. varium</i>	<i>C. asterophorum</i>	<i>C. paravirescens</i>	<i>C. protrusum</i>
Size (conidia)	15.9–21.8 $\times$ 6.3–9.2	8.6–15.8 $\times$ 6.4–8.6	14–23 $\times$ 5.5–7.5	18.0–27.5 $\times$ 6.5–10.0	16.0–27 $\times$ 5.5–9.0
Shape (conidia)	Two-celled	Two-celled	Two-celled	–	–
Shape (conidiospore)	Simple branch	Simple branch	Simple branch	Verticillate	Verticillate, 2–4 branches
Pigment (on plate)	Yellow to red	Cream	Pale crimson	Yellow to crimson	Yellow to brick brown or purple
Form of sclerotium	Formed sclerotium	Absent	Not reported	Sclerotia-like aggregation	Sclerotia-like aggregation
Additional characteristics	Rarely produces spores	Produces many spores	Produces profuse spores	Not reported	Not reported
Reference		Back <i>et al.</i> , 2012	De Hoog, 1978	Pöldmaa, 2011	Pöldmaa, 2011

with large truncate basal hila (Fig. 1F). These morphological characteristics differ from those of a previously reported causal agent of cobweb disease in *H. marmoreus* (Back *et al.*, 2012). The pigment color of the isolated fungi was similar to that of three *Cladobotryum* species (*C. asterophorum*, *C. paravirescens*, and *C. protrusum*; Table 1). However, these species have been isolated from different hosts, not from edible mushrooms. The conidia size of the fungi isolated in the present study (15.9–21.8 $\times$ 6.3–9.2 μm) also differs from those of previously reported *Cladobotryum* species (Table 1). Given these differences, we presumed that the new isolate is a new and important *Cladobotryum* species causing cobweb disease in edible mushrooms. However, only morphological characteristics were insufficient for species identification, and we therefore needed to study additional parameters to confirm the identity of this new isolate.

Effect of temperature on mycelial growth

The effect of temperature on fungal growth was investigated at temperatures ranging from 5°C to 30°C after 15 days of incubation. The criterion for optimum growth was colony diameter, which was maximum (90-mm diameter) at 25°C and minimum (25-mm diameter) at 5°C (Fig. 2). The color of the mycelia also changed to pink and, later, turned reddish after 15 days (Fig. 2). Similar red pigments have been reported in *C. mycophilum* from *P. eryngii*, *A. bisporus*, *C. asterophorum*, *C. paravirescens*, and *C. protrusum*, grown on PDA media and are due to aurofusarin, a secondary metabolite of the *Cladobotryum* species (Pöldmaa, 2011).

Pathogenicity tests

The results of the pathogenicity tests were verified by inoculating healthy *H. marmoreus* (white) with the isolated fungus. The same cobweb-like disease symptoms

Fig. 2. Effect of temperatures between 5°C and 30°C on mycelial growth of the isolated fungus on PDA 15 days after incubation (A: Front view, B: Back view of plate).

Fig. 3. Pathogenicity tests of the isolated fungus on four mushrooms at 20°C. DAI: days after inoculation.

developed on the inoculated mushrooms within 3 days after inoculation (DAI; Fig. 3). Rotted fruit bodies and few spores were observed 10 days later (Fig. 3). The development of the disease on the inoculated mushrooms resembled that on naturally infected mushrooms. The pathogenicity of the isolated fungus was also tested in *H. marmoreus* (brown), *F. velutipes*, and *P. eryngii* using the procedure followed for the initial inoculation. At 3 DAI, the symptoms of the cobweb-like disease were observed in all mushrooms, whereas rotted fruit bodies and mycelia were observed at 10 DAI (Fig. 3). All inoculated mushrooms were entirely rotted at 20 DAI, and reddish fruit bodies, massive mycelia, and few conidiospores were observed (Fig. 3). These symptoms are similar to those of cobweb disease of mushroom farms. However, the characteristics of this disease differ from the previous description of cobweb disease in *H. marmoreus* caused by *C. varium* in the presence of few conidia and reddish rotted fruit bodies; in the previously reported cobweb disease, numerous conidiospores are produced (Back *et al.*, 2012).

Phylogenetic analysis

Molecular characterization of the isolated fungi was performed based on gene encoding the ITS region, *RPB1*, *RPB2*, and *TEF*. Direct sequencing of the PCR products of the amplified gene for the ITS region, *RPB1*, *RPB2*, and *TEF* resulted in sequences of 622 bp, 728 bp, 1,135

bp, and 970 bp, respectively (data not shown). All of the sequences of the identified isolates for each gene were 100% identical to each other. According to our results, these isolate differs from previously reported isolates of *Cladobotryum* sp. in Korea that cause cobweb disease in mushrooms such as *A. bisporus*, *P. eryngii*, *F. velutipes*, and *H. marmoreus* (Kim *et al.*, 1998, 1999; Back *et al.*, 2010, 2012). The sequences obtained for the fungus were deposited in the DNA Data Bank of Japan/GenBank database under the accession numbers AB969671 for the ITS region, AB969673, AB969674 for the *RPB1* and *RPB2*, respectively, and AB969675 for the *TEF*. The phylogenetic analysis of the isolated fungus was carried out based on the combined sequences (ITS region, *RPB1*, *RPB2*, and *TEF*), and the results showed that the isolated fungus has a new lineage that differs from those of the previously reported *Cladobotryum* species (Fig. 4).

In the present study, a new species of *Cladobotryum* was isolated from *H. marmoreus* in Cheongdo-gun, Gyeongbuk Province, Korea. Its morphological characteristics of sclerotia formation, pigmentation, and conidia formation (shape and size) were distinct from those of previously reported *C. varium* species that cause cobweb disease in *H. marmoreus* (Back *et al.*, 2012). The morphological delimitation of the isolated fungi as a unique species was further supported by the phylogenetic analysis of four genes (gene for ITS, *RPB1*, *RPB2*,

Fig. 4. Phylogenetic tree constructed using the neighbor-joining method based on a comparison of four sequences (ITS region, *RPB1*, *RPB2*, and *TEF*) of *Cladobotryum* spp. with those of other *Cladobotryum* species from GenBank. *Hypomyces lactifluorum* and *Hypomyces subiculosus* formed the out group. The *Cladobotryum* species observed in this study are shown in bold font. Numbers on the branches are the confidence values obtained for 100 replicates (only values above 80% are shown). The bar represents a phylogenetic distance of 1%.

and *TEF*). The results showed that the isolated fungi belong to the same clade as *C. protrusum* and *C. paravirescens*; however, it forms a new lineage that is distant from other *Cladobotryum* species. Furthermore, pathogenicity tests revealed that the isolated pathogen also causes cobweb disease in *F. velutipes* and *P. eryngii*; however, natural infection has not yet been reported in these mushrooms.

Although this newly identified pathogen has been isolated less frequently than *C. varium* in Cheongdo-gun, Gyeongbuk Province in Korea, its potential threat to *H. marmoreus* and other mushrooms in Korea remains. The disease caused by this pathogen in *H. marmoreus* is of particular importance because it has been cultivated occasionally on approximately 10 mushroom farms in Korea. The results of the phylogenetic analysis showed that the isolated pathogen forms a new lineage; however, it belongs to same clade as *C. protrusum* and *C. paravirescens*, which have been isolated in decorticated wood and aphyllophoralean basidiomycetes, respectively. Moreover, previous reports have suggested that *H. marmoreus* on mushroom farms is cultivated on substrates consisting of pine sawdust or wood chips (Lee *et*

al., 2011). Therefore, the similar pigment color and phylogenetic analysis of this isolated pathogen suggest that it may exist in the wood chips or sawdust used in mushroom media.

The presence of our isolated pathogen on *H. marmoreus* suggests that it may be adaptive to the environmental conditions in *H. marmoreus* farms, and that it was likely transmitted through wood chips. Furthermore, the cultivation period of *H. marmoreus*, which is three times longer than that of other mushrooms, may help this pathogen grow and cause infection. However, further research is needed to identify the infection pattern of this fungus. On the basis of previously published literature and the results of the present study, we identified this isolated pathogen as a new *C. hypsigum* species that causes cobweb disease in *H. marmoreus* in Korea.

REFERENCES

- Adie, B., H. Grogan, S. Archer and P. Mills 2006 Temporal and spatial dispersal of *Cladobotryum* conidia in the controlled environment of a mushroom growing room. *Appl. Environ. Microbiol.*, **72**: 7212–7217
- Back, C. G., C. Y. Lee, G. S. Seo and H. Y. Jung 2012

- Characterization of species of *Cladobotryum* which cause cobweb disease in edible mushrooms grown in Korea. *Mycobiology*, **40**: 189–194
- Back, C. G., C. Y. Lee, G. S. Seo, H. Y. Jung and S. Oga 2012 First occurrence of cobweb disease on *Hypsizygus marmoreus* caused by *Cladobotryum varium* in Korea. *J. Fac. Agr. Kyushu Univ.*, **57**: 373–377
- Back, C. G., Y. H. Kim, W. S. Jo, H. Chung and H. Y. Jung 2010 Cobweb disease on *Agaricus bisporus* caused by *Cladobotryum mycophilum* in Korea. *J. Gen. Plant Pathol.*, **76**: 232–235
- Bhatt, N. and R. P. Singh 2002 Chemical control of mycoparasites of button mushroom. *Indian. J. Mycol. Plant Pathol.*, **32**: 38–45
- Cabone, I. and L. M. Kohn 1999 A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia*, **91**: 553–556
- Chaverri, P., M. Liu and K. T. Hodge 2008 Neotropical *Hypocrella* (anamorph *Aschersonia*), *Moelleriella*, and *Samuelsia*. *Studies in Mycology*, **60**: 1–68
- De Hoog, G. S. 1978 Notes on some fungicolous hyphomycetes and their relatives. *Persoonia*, **10**: 33–81
- Fletcher, J. T., P. F. White and R. H. Gaze 1989 Mushrooms: pest and disease control. Andover, Hants, England: Intercept. pp. 58–63
- Gams, W. and A. C. M. Hoozemans 1970 *Cladobotryum*–Konidienformen von Hypomyces–Arten. *Persoonia*, **6**: 95–110
- Gea, F. J., M. J. Navarro and L. M. Suz 2011 First report of *Cladobotryum mycophilum* causing cobweb on cultivated king oyster mushroom in Spain. *Plant Dis.*, **95**: 1030
- Grogan, H. M. 2006 Fungicide control of mushroom cobweb disease caused by *Cladobotryum* strains with different benzimidazole resistance profiles. *Pest. Manag. Sci.*, **62**: 153–161
- Grogan, H. M. and R. H. Gaze 2000 Fungicide resistance among *Cladobotryum* spp. – causal agent of cobweb disease of the edible mushroom *Agaricus bisporus*. *Mycol. Res.*, **104**: 357–364
- Ikekawa, T. 1995 Bunashimeji, *Hypsizygus marmoreus*: antitumor activity of extracts and polysaccharides. *Food Rev. Int.*, **11**: 207–209
- Ikekawa, T., H. Saitoh, W. Feng, H. Zhang, L. Li and T. Matsuzawa 1992 Antitumor activity of *Hypsizygus marmoreus*. I. Antitumor activity of extracts and polysaccharides. *Chem. Pharm. Bull.*, **40**: 1954–1957
- Khan, I., F. Shah, S. Bulman and I. Scott 2008 Molecular diagnostic tools for improved mushroom production. In: Abstracts of the XVII International Congress on the Science and Cultivation of Edible and Medicinal Fungi. Cape Town.
- Kim, T. S., H. U. Lee., K. W. Song and W. K. Shin 1998 King oyster mushroom (*Pleurotus eryngii*) white mold disease caused by *Cladobotryum varium*. *KSM News Letter (The Korean Society of Mycology)*, **11**: 46
- Kim, H. K., S. J. Seok, G. P. Kim, B. J. Moon and T. Terashita 1999 Occurrence of disease caused by *Cladobotryum varium* on *Flammulina velutipes* in Korea. *Kor. J. Mycol.*, **27**: 415–419
- Lam, S. K. and T. B. Ng 2001 Hypsin, a novel thermostable ribosome-inactivating protein with antifungal and antiproliferative activities from fruiting bodies of the edible mushroom *Hypsizygus marmoreus*. *Biochem. Biophys. Res. Commun.*, **285**: 1071–1075
- Lee, C. Y., J. E. Park, J. Lee, J. K. Kim and H. S. Ro 2011 Development of new strains and related SCAR markers for an edible mushroom, *Hypsizygus marmoreus*. *FEMS. Microbiol. Lett.*, **327**: 54–59
- Lee, Y. L., M. T. Yen and J. L. Mau 2007 Antioxidant properties of various extracts from *Hypsizygus marmoreus*. *Food Chem.*, **104**: 1–9
- Liu, D., S. Coloe, S. Baird, J. Pedersen 2000 Rapid mini-preparation of fungal DNA for PCR. *J. Clin. Microbiol.*, **38**: 471
- Liu, Y. J., S. Whelen and B. D. Hall 1999 Phylogenetic relationships among ascomycetes, as inferred from RNA polymerase II phylogeny. *Mol. Biol. Evol.*, **16**: 1799–1808
- McKay, G. J., D. Egan, E. Morris and A. E. Brown 1998 Identification of benzimidazole resistance in *Cladobotryum dendroides* using a PCR-based method. *Mycol. Res.*, **102**: 671–676
- McKay, G. J., D. Egan, E. Morris, C. Scott and A. E. Brown 1999 Genetic and morphological characterization of *Cladobotryum* species causing cobweb disease of mushrooms. *Appl. Environ. Microbiol.*, **65**: 606–610
- Pöldmaa, K. 2011 Tropical species of *Cladobotryum* and *Hypomyces* producing red pigments. *Stud. Mycol.*, **68**: 1–34
- Potocnik, I., J. Vukojevic, M. Stajic, B. Tanovic and B. Todorovic 2008 Fungicide sensitivity of selected *Verticillium fungicola* isolates from *Agaricus bisporus* farms. *Arch. Biol. Sci. (Belgrade)*, **60**: 151–157
- Rehner, S. A. 2001 Primers for elongation factor 1- α (*EF1- α*). <http://ocid.nacse.org/research/deephyphae/EF1primer.pdf>
- Tamm, H. and K. Pöldmaa 2013 Diversity, host associations, and phylogeography of temperate aurofusarin-producing *Hypomyces/Cladobotryum* including causal agents of cobweb disease of cultivated mushrooms. *Fungal Biol.*, **117**: 348–367
- Thompson, J. D., D. G. Higgins and T. J. Gibson 1994 CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucl. Acids. Res.*, **22**: 4673–4680
- Tsuchida, K., Y. Aoyagi, S. Odani, T. Mita and M. Isemura 1995 Isolation of a novel collagen-binding protein from the mushroom, *Hypsizygus marmoreus*, which inhibits the Lewis lung carcinoma cell adhesion to type IV collagen. *J. Biol. Chem.*, **270**: 1481–1484
- White, T. J., T. Bruns, S. Lee and J. Taylor 1990 Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR Protocols: A Guide to methods and applications*, Academic Press, San Diego, CA, USA, 315–322