Viability Loss and Oxidation during Desiccation of Recalcitrant *Pachira macrocarpa* Seeds

Shen, Rong-Show

Department of Horticultural Science, National Chiayi University

Lu, Shao-Wei

Department of Horticultural Science, National Chiayi University

Huang, Kuang-Liang

Department of Horticultural Science, National Chiayi University

Chang, Yu-Sen

Department of Horticulture and Landscape Architecture, National Taiwan University

他

<https://doi.org/10.5109/1526292>

出版情報：九州大学大学院農学研究院紀要. 60 (1), pp.23-32, 2015-02-27. Faculty of Agriculture,
Kyushu University

バージョン：

権利関係：

Viability Loss and Oxidation during Desiccation of Recalcitrant *Pachira macrocarpa* Seeds

Rong-Show SHEN^{1*}, Shao-Wei LU¹, Kuang-Liang HUANG¹,
Yu-Sen CHANG² and Ikuo MIYAJIMA³

Institute of Tropical Agriculture, Faculty of Agriculture, Kyushu University,
Fukuoka 812–8581, Japan

(Received October 20, 2014 and accepted November 14, 2014)

This study investigated the correlation between the moisture content and viability of *Pachira macrocarpa* seeds, and observed the changes in seed oxidation and oxidation resistance during the desiccation process. The average moisture content of a fresh mature *P. macrocarpa* seed was approximately $1.93 \text{ gH}_2\text{O}\cdot\text{g}^{-1}$ DW. When the seeds were placed into desiccating environments at 25°C with 60% relative humidity, seed viability and moisture content exhibited highly curve correlation. The R^2 reached 0.98, and seed viability rapidly declined as the moisture content decreased. This test was observed causing seed death occurred at a critical moisture content of $1.4 \text{ gH}_2\text{O}\cdot\text{g}^{-1}$ DW, where the seed germination percentage was 0%. This indicates that *P. macrocarpa* seeds are not only recalcitrant but also sensitive to desiccation. Furthermore, the seed oxidation resistance showed that after 0–5 days of desiccation, the superoxide anion radical (O_2^-) and hydrogen peroxide (H_2O_2) levels increased as the seed moisture content decreased. The activity of the antioxidant enzyme superoxidase dismutase (SOD) and ascorbate peroxidase (APX) also decreased with the seed moisture content. When the seeds were desiccated to a moisture content of $1.22 \text{ gH}_2\text{O}\cdot\text{g}^{-1}$ DW, the SOD activity was reduced to half that of a fresh seed. Although the ascorbate–glutathione cycle continued to function, the seed's malondialdehyde (MDA) content increased 1.5 times, which expedited seed oxidation. However, the reduction of *P. macrocarpa* seed viability during desiccation occurred before significant oxidation, while seed death may have been caused by physical damage not metabolic injury.

Key words: antioxidant enzyme, ascorbate peroxidase, malondialdehyde, superoxidase dismutase

INTRODUCTION

Seedlings production of braiding types in malabar chestnut [*P. macrocarpa* (Cham. & Schl.) Schl.] is the most important export potted foliage in Taiwan, while the seeds are commonly considered recalcitrant or short-lived. When Li *et al.* (2009) desiccated *P. macrocarpa* seeds with silica gel, seed viability decreased with increased desiccation time. However, no reports or clear definitions exist regarding the correlation between the seed viability and moisture content for *P. macrocarpa* seeds, or for the critical moisture content at seed death. Currently, *P. macrocarpa* seeds can only be stored for approximately one month (Sun *et al.*, 2011). Therefore, to overcome the limited factor of *P. macrocarpa* seed storage life, an in-depth understanding of the physiological reaction to desiccation is required to benefit the development of storage technology.

Roberts (1973) established a seed viability equation based on their sensitivity to desiccation, classifying seeds into orthodox and recalcitrant seeds. Subsequent reports have categorized seeds into various desiccation tolerance types based on their sensitivity to desiccation and tem-

perature (Farrant *et al.*, 1988; Ellis *et al.*, 1991; Hong and Ellis, 1996), and even attributed the characteristics of seed longevity to quantitative trait (Berjak and Pammenter, 2001; Win, 2008). Hong *et al.* (1996, 1998) also categorized seeds into orthodox, recalcitrant, and intermediate seeds based on their reaction to desiccation, which is currently the most widely accepted categorization method. However, no effective methods exist to preserve recalcitrant seeds for long terms. The most obvious characteristic of recalcitrant seeds is their sensitivity to desiccation. In addition, regarding the desiccation of *Camellia sinensis* Luntze (Chen *et al.*, 2010), *Telfairia occidentalis* Hook. f. (Ajayi *et al.*, 2006), *Lasia spinosa* (L.) Thwait. (Tang and Long, 2008), *Quercus robur* L. (Bonner, 1996), and *Artocarpus heterophyllus* Lamk. (Wesley-Smith *et al.*, 2001) recalcitrant seeds, the germination percentages all declined as the desiccation time increased or the moisture content decreased. When desiccated to a seed moisture content of 20% to 30%, the majority of the seeds exhibited no germination vigor. This indicates that most recalcitrant seeds cannot tolerate seed moistures below 20%, and seed moisture content is a key factor that affects the viability of recalcitrant seeds.

Furthermore, some plant species have a specific threshold of seed moisture content on storage longevity. Once the moisture content declines below this threshold, the seed no longer possesses germination vigor. Exploring the reason, some reports have indicated that the death of recalcitrant seeds may be caused by an increase in reactive oxygen species (ROS) after desiccation, and the

¹ Department of Horticultural Science, National Chiayi University, Chiayi 60004, Taiwan, ROC.

² Department of Horticulture and Landscape Architecture, National Taiwan University, Taipei 10617, Taiwan, ROC.

³ Institute of Tropical Agriculture, Faculty of Agriculture, Kyushu University, Fukuoka 812–8581, Japan

* Corresponding author (E-mail address: garden@mail.nyu.edu.tw, telephone: +88652717431, fax: +88652775472.)

lack of an anti-oxidation system. For example, following with a reduced seed moisture content, *Acer saccharinum* L. (Pukacka and Ratajczak, 2006), *C. sinensis* Luntze (Chen *et al.*, 2010), *Antiaris toxicaria* L. (Xin *et al.*, 2010), and *Araucaria bidwillii* Hook (Francini *et al.*, 2006) seeds showed an increase in O_2^- , H_2O_2 , free radicals, and other reactive oxygen species. Plants typically possess an anti-oxidation system to eliminate free radicals; however, anti-oxidation enzymes and other anti-oxidants all decline after recalcitrant seeds are desiccated. Resembling the reaction after desiccation, the SOD activity of *P. macrocarpa* and *Theobroma cacao* seeds decreased with the seed moisture content (Li and Sun, 1999); after SOD reduces O_2^- into H_2O_2 , ascorbate peroxidase (APX) reduces H_2O_2 into H_2O . While, the APX activity in both *A. toxicaria* and *T. cacao* recalcitrant seeds decreased with the seed moisture content (Li and Sun, 1999; Xin *et al.*, 2010).

For antioxidant level change, the glutathione (GSH) in *Q. robur* and sugar maple seeds were reduced as the seed moisture content declined (Kranter *et al.*, 2006; Pukacka and Ratajczak, 2006). After sugar maple seeds were desiccated, the GSH and glutathione disulphide (GSSG) ratio declined, and the seeds exhibited oxidation (Pukacka and Ratajczak, 2006). To seem like this, after *Castanea sativa* seeds were dried, the $E_{GSSG/GSH}$ value gradually became positive, indicating that the seed oxidation condition had deteriorated (Roach *et al.*, 2010). If seeds cannot eliminate the reactive oxygen species produced during desiccation, lipid peroxidation may occur. During the desiccation of *A. toxicaria* and *T. cacao* and *A. bidwillii* seeds, the quantity of the lipid peroxidation indicator, that is, thiobarbituric acid reactive substances (TBARS), increased as the seed moisture content decreased (Cheng and Song, 2008; Francini *et al.*, 2006; Li and Sun, 1999), then caused cell membranes to rupture and increasing the ion leakage rate (Kioko *et al.*, 2006; Xin *et al.*, 2010). Thus, during the desiccation process, the anti-oxidation system of recalcitrant seeds becomes imbalanced and oxidation occurs in the seed interior, leading to seed death. However, although *A. toxicaria* seed exhibited oxidation, the level of oxidation was insufficient to cause seed death. Thus, seed death may result from severe mechanical damage to the cells during desiccation (Xin *et al.*, 2010), not metabolic injury.

Therefore, this study investigated the correlation between the moisture content and viability of *P. macrocarpa* seeds, and exploring the seed desiccation tolerance to provide relevant physiology information for future storage of *P. macrocarpa* seeds. We also examined the moisture content of *P. macrocarpa* seeds and its effect on oxidation resistance. The results were adopted to explore the reactive oxygen species, anti-oxidation system, and oxidation situation in recalcitrant *P. macrocarpa* seeds following desiccation, and to understand the physiological aspects for the lack of desiccation tolerance in recalcitrant seeds.

MATERIALS AND METHODS

Seed materials

Mature *P. macrocarpa* fruits were harvested and transported to the laboratory within 1 day, and then placed in shade until they split naturally. After splitting, seeds were extracted and screened based on a visual inspection of their exterior plumpness. All mature seeds with a fresh weight of over 2.5 g were selected for testing.

Desiccation and germination test

The seeds were placed indoors for desiccation at a temperature of $25 \pm 1^\circ\text{C}$ and a relative humidity of 60% for 0 to 12 days. Each treatment was repeated three times using 15 seeds for each replication. The test period was from March 2, 2012, to March 14, 2012.

After 0 to 12 days of indoor desiccation, we selected five seeds from each treatment and followed Wesley-Smith (2001) analysis method for testing. The seeds were dried at 80°C in a hot air oven for 36 h until a constant weight was achieved. The formula for calculating the seed moisture content following desiccation was $[(\text{post-indoor desiccation weight} - \text{dry weight})/\text{dry weight}] \times 100\%$. We then selected 10 seeds from each treatment, the seed coat removed, and sowed the seeds in a plug tray with 35 plugs and plug hole diameter of 7 cm. The plugs were filled with peat-moss (Klasmann-Deilmann Co., Ltd., Germany) as the sowing medium, and the trays were placed in greenhouse provided a day/night temperature of $30/25^\circ\text{C}$ at Department of Horticultural Science of National Chiayi University to investigate seed viability. Seed viability is represented by the germination percentage and seedling rate after sowing. The calculation formula is $\text{germination percentage} = (\text{number of germinations}/\text{number of seeds sown}) \times 100\%$ and $\text{seedling rate} = (\text{number of seedlings}/\text{number of seeds sown}) \times 100\%$.

Microscopy

During this experiment, we also conducted a biopsy of hypocotyl tissues obtained from seeds desiccated for 9 days, and seeds that had not undergone desiccation for control group. The biopsy technique was referencing the paraffin section processing methods by Tsai (1975), fixation, dehydration, infiltration, and embedding/casting were sequentially employed, and a microtome was used to sectioning. The fixation liquid was a solution of 37% formaldehyde, acetic acid, and 50% alcohol in a volume ratio of 5:5:90. Hypocotyl tissue was placed in a 6 mL sample vial for fixation. Tissue coverage by the fixation solution was set as the standard, which involved at least 3 mL of the solution or more. Vacuum suction (15 m per h) was applied in conjunction, and the fixation process conducted for a total of 72 h. After extracting the fixation solution, alcohol was mixed with tertiary butyl alcohol (TBA) into a dehydration solution to conduct dehydration. Finally, the tissue sample was placed in 100% TBA for paraffin infiltration. For the paraffin infiltration process, a small amount of pure paraffin was employed to replace TBA. This was conducted in a 60°C oven until the TBA had completely evaporated. Paper embedding

cassettes were used to conduct paraffin embedding on the paraffin infiltrated tissue. After cooling and solidifying, the paraffin block was trimmed with a razor. A manual rotary microtome (Nanolytik® 4060, Germany) was employed to cut 10 μm sections. The sectioned continuous paraffin ribbon was placed on a glass slide coated with adhesive and formalin dilution, and the transferred to a 38°C paraffin section mounting bath (Sakura SV-C2, Japan) for mounting. Dyeing was performed after two days of mounting. During the dying process, xylene substitute (Thermo Scientific Inc., U.S.A.) was used to melt the paraffin. After a series of alcohol immersions and dying with safranin, the sample underwent a series of immersions in alcohol as well as xylene substitute, thereby completing the procedure. A dyed glass slide was employed with an optical microscope (Zeiss Axio Imager. A1, Carl Zeiss AG, Germany) and lighting equipment (Canon A620, Canon Inc., Japan) to make observations.

Biochemical assays

This experiment employed the same seed material as used for desiccation. The material was placed in the environment described above to desiccate for up to 5 days. After desiccation, five seeds were selected from each treatment. A freeze dryer (Ilshin Lab. Co., Ltd., Korea) was used to freeze dry and grind the seeds for the oxidation resistance experiment. The experimentation period was from March 2, 2012, to March 7, 2012.

The analysis results of the seeds' peroxide level, anti-oxidation enzyme activity, antioxidant quantity, and oxidation following desiccation are explained below.

Reactive Oxygen species (ROS) measurement

(1). Hydrogen peroxide (H_2O_2)

Referencing the method used by Sagisaka (1976), we obtained 0.4 g of freeze-dried *P. macrocarpa* seeds and added 4 mL of 5% trichloroacetic acid (TCA) as the extraction liquid for ground extraction in a cold bath. After the mixture was centrifuged at 18,000 $\times g$ and 4°C for 30 min, 10 μL of the extracted solution was combined with 260 μL of the reacting liquid [containing 10 mM of ferrous ammonium sulfate, 2.5 mM of potassium thiocyanide, and 50% (w/v) TCA] and mixed thoroughly. The solution's absorbency change per unit time at a 480-nm wavelength was measured, and H_2O_2 was used as the standard to estimate H_2O_2 quantity.

(2). Superoxide anion (O_2^-)

Referencing the method used by Liu *et al.* (2012), we obtained 1 g of freeze dried *P. macrocarpa* seed powder, which we combined with 1 mL of potassium phosphate buffer solution (20 mM, pH 6.0) containing 500 μM XTT of sodium salt (XTT). This mixture was placed in a 25°C darkroom for 3 h of oscillating reaction, and then tested for absorptivity changes in 470 nm.

Anti-oxidation enzyme activity analysis

(1). Ascorbate peroxidase (APX)

Based on the method used by Nakano and Asada (1981), we obtained 0.4 g of freeze dried *P. macrocarpa* seed, added 4 mL of sodium phosphate buffer (50 mM,

pH 6.8) solution, and then ground and extracted it in an ice bath. The mixture was centrifuged at 12,000 g and 4°C for 20 min. Then 0.2 mL of the extracted fluid was combined with 2.9 mL of the reacting liquid (containing 1 mL of 100 mM pH 7.0 potassium phosphate buffer, 1 mL 1.5 mM of ascorbate, 0.4 mL 0.75 of mM EDTA, and 0.5 mL 6 mM of H_2O_2) and mixed evenly. A wavelength of 290 nm was used to test the changes in absorptivity. A photoelectric colorimeter absorption coefficient of 2.8 $\text{mM}^{-1}\cdot\text{cm}^{-1}$ was used to estimate the AsA quantity. The enzyme activity was defined as the change in AsA consumption per unit time.

(2). Superoxide dismutase (SOD)

Based on the method used by Paoletti *et al.* (1986), we obtained 0.4 g of freeze dried *P. macrocarpa* seed, added 2 mL of sodium phosphate buffer (50 mM, pH 7.4) solution, and then ground and extracted it in an ice bath. The mixture was centrifuged at 15,000 g and 4°C for 30 min. Subsequently, 0.1 mL of the extracted liquid was combined with 2.73 mL of the reacting liquid [containing 1.6 mL of 100 mM pH 7.4 triethanol amine diethanolamine (Tea-Dea), 80 μL of 7.5 mM reduced-form nicotinamide adenine dinucleotide (NADH), 50 μL of 100 mM pH 7.0 EDTA/ MnCl_2 , and 1 mL of 10 mM 2-mercaptoethanol] and mixed evenly. A wavelength of 340 nm was employed to test the changes in absorptivity, using a molar absorption coefficient of 6.22 $\text{mM}^{-1}\cdot\text{cm}^{-1}$ for estimation. The enzyme activity was measured according to the enzyme activity per gram of fresh weight. The enzyme activity is indicated by the unit activity per gram of fresh weight, and each unit activity was defined as the quantity required for SOD to inhibit the NADH oxidation rate by 50%.

Antioxidant quantity analysis

(1). Glutathione (GSH)

After obtaining 0.4 g of freeze dried *P. macrocarpa* seed, 4 mL of sulfosalicylic acid (5%, w/v) solution was added, and the mixture ground and extracted in an ice bath. The mixture was centrifuged at 15,000 $\times g$ and 4°C for 10 min, according the methods of Smith (1985). The samples obtained 0.25 mL of the extracted sample liquid and then added 0.15 mL of 0.1 M potassium phosphate buffer (pH 7.5). Following oscillation mixing, 1.4 mL of the reacting liquid [containing 1 mL of 0.1 M sodium phosphate buffer (pH 7.5, with 5 mM of ethylenediamine-tetra-acetic acid (EDTA))] and 0.4 mL of 6 mM dithionitro-benzoic acid (DTNB) were added. After reacting for 10 min at 30°C, 0.2 mL of 2 mM NADPH and 0.2 mL of glutathione reductase (2.5 $\text{unit}\cdot\text{mL}^{-1}$) were added, and the solution was evenly mixed. A wavelength of 412 nm was used to test the changes in absorptivity. GSH was used as the standard to estimate GSH quantity. The 0.1 mL of the extracted sample fluid into which 0.15 mL of 0.1 M potassium phosphate buffer (pH 7.5) was mixed before adding 40 μL of 2-vinylpyridine to prevent the GSH from oxidizing into glutathione disulfide (GSSG). After mixing, 60 μL of triethanolamine (35 mM) was added along the tube wall before oscillation mixing. The solution was left at room temperature to react for at least 1 h to eliminate the GSH in the sample. Then 1.5 mL of the

reacting fluid (containing 0.1 M of sodium phosphate buffer pH 7.5 and 5 mM of EDTA) and 0.4 mL of 6 mM dithionitro-benzoic acid (DTNB) were added. After reacting for 10 min at 30°C, 0.2 mL of 2 mM NADPH and 0.2 mL of glutathione reductase (5 unit·mL⁻¹) were mixed into the solution. A wavelength of 412 nm was used to test the changes in absorptivity, and glutathione disulfide (GSSG) was set as the standard for estimating the quantity of GSSG.

(2). Ascorbate (AsA)

The AsA quantity was based on the method employed by Law *et al.* (1983), using 0.4 g of freeze dried *P. macrocarpa* seed. The samples were added 4 mL of 0.1 M pH 6.8 potassium phosphate buffer solution and then ground and extracted the mixture in an ice bath. After centrifugation at 15,000 ×g and 4°C for 30 min, the supernatant became the enzyme extraction liquid. We mixed 100 μL of the extracted sample liquid with 150 mM of pH 7.4 sodium phosphate buffer, added 50 μL of 10 mM DL-dithiothreitol (DTT), and then left the solution at room temperature for 15 min. Subsequently, 50 μL of 0.5% N-ethylmaleimide was added and mixture shaken until evenly mixed. After adding 200 μL of 10% TCA and 200 μL of 44% (v/v) H₃PO₄, the mixture was again shaken until mixed. Next, 200 μL of 4% (w/v) bipyridyl (dissolved in 90% ethanol) was mixed into the solution. Finally, 100 μL of 3% (w/v) FeCl₃ was added, and the solution placed in an oscillator at 37°C to react for 1 h. A wavelength of 525 nm was used to test the changes in absorptivity. The AsA solution was used as the standard to estimate the total AsA quantity. Besides not adding DTT and N-ethylmaleimide to prevent the dehydroascorbate (DHAsA) from reducing to AsA, the procedures for the AsA quantity test were identical to those followed for the total ascorbate analysis.

Determination of lipid peroxidation

Referencing the method used by Heath and Packer (1968), 0.4 g of freeze dried *P. macrocarpa* seed was ground and extracted with 4 mL of 5% TCA extraction liquid in a cold bath. After centrifugation at 18,000 g and 4°C for 30 min, 1 mL of *P. macrocarpa* extraction liquid was added to 4 mL of the reacting liquid [0.5% (w/v) thiobarbituric acid (TBA) containing 20% (w/v) TCA]. After this solution was left in a 95°C bath for 30 min, it was quickly placed in ice for cooling and to stop the reaction. After the solution cooled, it was centrifuged at 3,000 g for 10 min. Wavelengths of 532 nm and 600 nm were used to test the changes in absorptivity. A molar absorption coefficient of 155 mM⁻¹·cm was used to estimate malondialdehyde (MDA) quantity.

Statistical analysis

The experiments adopted a completely randomized design (CRD). The experiment results were analyzed with CoStat 6.4 statistical software (CoHort Software, Monterey, CA, U.S.A.). Analysis of variance (ANOVA) was performed to assess the significances, and the least significant difference (LSD) test was conducted to identify whether significant variances ($p \leq .05$) in the test

procedures existed. Experiment data in percentages were converted using a Bliss conversion table prior to statistical analysis. SigmaPlot® 10.0 statistical software package (Systat software Inc. USA) was used for plotting and comparing the experiment results.

RESULTS

The experiment results showed that the average moisture content of fresh *P. macrocarpa* seeds was approximately 1.93 gH₂O·g⁻¹ DW (Fig. 1A). After sowing, the germination percentage reached 100%. When the seeds were desiccated at an indoor temperature of 25°C and a relative humidity of 60%, the seed moisture content decreased with increases in desiccation time. This decrease in moisture content and increased desiccation time exhibited a high linear correlation ($R^2=0.95$). After 12 day of slow desiccation, the seed moisture content was 1.07 gH₂O·g⁻¹ DW (Fig. 1A). The germination percentage and seedling rate rapidly declined with decreases in the seed moisture content. Simultaneously, the germination percentage and seedling rate exhibited a curve highly correlated with seed moisture content (Fig. 1B). When the coefficient of determination (R^2) for the seed moisture content and germination/seedling rate were 0.98 and 0.97, the mathematical equations were $y = 131x^2 - 284.0x + 148.7$ and $y = 221.7x^2 - 485.3x + 258.5$, respectively. In addition, when the germination percentage declined to

Fig. 1. Correlation between moisture content of *Pachira macrocarpa* seeds and drying time at 25°C and RH 60% (A) and effect of moisture content on germination rate (●) and seedling rate (○) of *Pachira macrocarpa* seeds (B).

50%, the seed moisture content was approximately $1.7 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$. After desiccating for 7 days, when the moisture content had declined to $1.4 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$, the seeds no longer possessed germination abilities (Fig. 1B). The hypocotyl biopsy of *P. macrocarpa* seeds that were not desiccated showed that the cells were organized into neat lines, the cell membrane structure was complete, and clear divisions between cells was evident. After dying with safranin, clear cell nuclear structures were observed, and the cell nuclear membrane maintained completeness. A clear division between the cell nucleus and cytoplasm was also noted (Fig. 2A). While, a hypocotyl biopsy of seeds desiccated for 9 days was performed to observe the cell structure; the results showed unclear divisions between cells. Thereby, the cell membrane structures appeared ruptured, and the division between the cell nucleus and the cytoplasm was unclear. This suggested that the nuclear membrane of the cells had ruptured (Fig. 2B).

Observing the changes in reactive oxygen species after *P. macrocarpa* seeds were desiccated for up to 5 days, superoxide anion radical (O_2^-) and hydrogen peroxide (H_2O_2) increased as the seed moisture content decreased. The O_2^- and H_2O_2 content of the seeds that

were not desiccated was 0.25 and $107.5 \mu\text{g}\cdot\text{g}^{-1} \text{ DW}$, respectively. For seeds desiccated to a moisture content of $1.47 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$, the seed O_2^- content was significantly higher than that of the seeds that were not desiccated. When the moisture content declined to $1.22 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$, the O_2^- content reached a peak of $0.38 \mu\text{g}\cdot\text{g}^{-1} \text{ DW}$. When the seed moisture content declined to $1.87 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$, the H_2O_2 content of desiccated seeds was significantly higher than the $1.89 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$ of seeds that were not desiccated. In addition, when the seeds were desiccated to a moisture content of $1.08 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$, we can observe that H_2O_2 quantity increased to $162.3 \mu\text{g}\cdot\text{g}^{-1} \text{ DW}$ (Fig. 3).

After desiccation, we also observed changes in the anti-oxidation enzyme activity. The SOD activity of seeds that were not desiccated was $3.62 \text{ units}\cdot\text{g}^{-1} \text{ DW}$. With decreases in seed moisture content, SOD activity showed a decreasing trend. When the seed moisture content declined to $1.72 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$, the SOD activity of desiccated seeds was significantly lower than that of the seeds that were not desiccated. When the seeds were desiccated to a moisture content of $1.22 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$, SOD activity was reduced to approximately $2.43 \text{ units}\cdot\text{g}^{-1} \text{ DW}$, half that of seeds that were not desiccated (Fig. 4A). The APX activity in seeds that were not desiccated was approximately $28.95 \mu\text{mol}\cdot\text{g}^{-1} \text{ DW}\cdot\text{min}^{-1}$ and showed an increasing trend until the seeds were desiccated to $1.72 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$. Further desiccation reduced the activity. When the seed moisture content was $1.22 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$, the APX activity was approximately $18.94 \mu\text{mol}\cdot\text{g}^{-1} \text{ DW}\cdot\text{min}^{-1}$, significantly lower than that of the seeds that were not desiccated (Fig. 4B).

In addition, the total quantity of the antioxidant ascorbic acid showed an increase during the early stages of desiccation, and when the moisture content was $1.87 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$, the total ascorbic acid quantity was significantly higher than that during other processes. However, after this peak, the quantity declined as the moisture content decreased. The AsA quantity showed no significant change during the desiccation process. The

Fig. 2. Transverse sections of embryonal axis tissue after 0 day (A) and 9 days (B) desiccation of *Pachira macrocarpa*. The cells arranged in neat rows, and the cell membrane and nucleus appeared distinct (A); After 9-day desiccation to moisture content at $1.2 \text{ gH}_2\text{O}\cdot\text{g}^{-1} \text{ DW}$, cells lose typified cells rank (B), and visible destroyed cell membrane and abnormally shaped nucleus indicated intercellular confusion (arrow-head).

Fig. 3. Effect of seed moisture content on superoxide anion (O_2^-) and hydrogen peroxide (H_2O_2) content in *Pachira macrocarpa*.

^a Means followed by the same letter within each column are not significantly different at the 5% level by LSD test

Fig. 4. Effect of seed moisture content on superoxide dismutase (SOD) and ascorbate peroxidase (APX) activity in *Pachira macrocarpa*.

^z Means followed by the same letter within each column are not significantly different at the 5% level by LSD test

Fig. 5. Effect of seed moisture content on ascorbate (AsA) and dehydroascorbate (DHAsA) content in *Pachira macrocarpa*.

^z Means followed by the same letter within each column are not significantly different at the 5% level by LSD test

changes in DHAsA were similar to that of the total ascorbic acid quantity, which showed a significant increase during the early stages of desiccation, but subsequently declined with decreases in moisture content (Fig. 5A). By converting the AsA and DHAsA quantity into a ratio, we found that when the moisture content was higher than 1.72 gH₂O·g⁻¹ DW, the AsA and DHAsA ratio was lower than 30. This indicates that ascorbic acid has both an oxidation state and a reduction state. Nevertheless, with a moisture content lower than 1.47 gH₂O·g⁻¹ DW, the ratio significantly increased to above 70, which indicates that the ascorbic acid exhibited was primarily in a reduction state (Fig. 5B). Regarding glutathione content, the total glutathione quantity increased as the moisture content decreased. When the moisture content was 1.72 and 1.08 gH₂O·g⁻¹ DW, the glutathione content measured 0.09 and 0.22 mg·g⁻¹ DW, respectively. However, almost no GSSG existed during the desiccation period (Fig. 6A). By converting GSH and GSSG into a ratio, we found that the GSH and GSSG content increased as the moisture content decreased. Thus, as the moisture content decreases, the ratio of GSH and GSSG increases (Fig. 6B).

Finally, the MDA content in the seeds was tested for oxidation following desiccation. During desiccation, the amount of MDA increased with decreases in the moisture content. When the moisture content exceeded 1.47 gH₂O·g⁻¹ DW, the MDA quantity was lower than 25 nmol·g⁻¹ DW. However, when continued desiccation reduced the moisture content to below 1.22 and 1.08 gH₂O·g⁻¹ DW, the MDA content increased to 30.1 and 36.6 nmol·g⁻¹ DW, respectively (Fig. 7).

Fig. 6. Effect of seed moisture content on glutathione (GSH) and glutathione disulfide (GSSG) content in *Pachira macrocarpa*.

^z Means followed by the same letter within each column are not significantly different at the 5% level by LSD test

Fig. 7. Effect of seed moisture content on malondialdehyde (MDA) content in *Pachira macrocarpa*.

^z Means followed by the same letter within each column are not significantly different at the 5% level by LSD test

DISCUSSION

The *P. macrocarpa* seeds examined in this study were desiccated at an indoor temperature of 25°C and a relative humidity of 60%. The germination percentage declined with decreases in the seed moisture content. After 7 days of desiccation, the seed moisture content had declined to less than 1.4 gH₂O·g⁻¹ DW, and the germination percentage was reduced to 0% (Fig. 1B). According to many previous reports have asserted that *P. macrocarpa* seeds have a maximum storage life of 1 month, but they cannot be compared to orthodox seeds longevity (Hong *et al.*, 1998; Sun *et al.*, 2011). The results of this study suggest that *P. macrocarpa* seeds possess the storage characteristics of physiological intolerance to desiccation. However, Win (2008) contended that if a seed has insufficient desiccation, low temperature, or storage tolerance, it exhibits difficult to store property. Therefore, *P. macrocarpa* seeds can be categorized as recalcitrant seeds.

After *T. occidentalis* seeds, which are recalcitrant, were slowly desiccated for 3 days with barium chloride (85% RH), the germination percentage declined with decreased in the seed moisture content (Ajayi *et al.*, 2006). Similarly, after *A. heterophyllum* seeds were slowly desiccated using a supersaturated solution of potassium di-hydrogen phosphate (96% RH) at 25°C for 3 days, the moisture content declined below 0.8 gH₂O·g⁻¹ DW, and the germination percentage decreased drastically. When the seed moisture content of *A. heterophyllum* decreased below 0.4 gH₂O·g⁻¹ DW, the seeds died (Wesley-Smith *et al.*, 2001). In this study, the correlation between the moisture content of *P. macrocarpa* seeds and the germination percentage was similar to the reaction of recalcitrant seeds to desiccation. As the seed moisture content decreased, the germination percentage declined. Although the germination percentage reduction of *T. occidentalis* (Ajayi *et al.*, 2006) seeds can be

divided into three stages, *P. macrocarpa* seeds exhibited a curve reduction (Fig. 1B). This indicates that the reaction of recalcitrant seeds to desiccation differs also depending on the plant species. In addition, when *P. macrocarpa* seeds were desiccated slowly for 6 days, and the moisture content reduced to 1.4 gH₂O·g⁻¹ DW, the seed germination percentage reached 0%. This was significantly higher than the seed death moisture content of *A. heterophyllum* at 0.4 gH₂O·g⁻¹ DW (Wesley-Smith *et al.*, 2001), which indicates that *P. macrocarpa* seeds are more sensitive to desiccation compared to recalcitrant jackfruit (*A. heterophyllum*) seeds. This phenomenon is similar to the argument proposed by Berjak and Pammenter (2001), that is, seed sensitivity to desiccation is not an all-or-none trait, but rather a quantitative trait, exhibiting a different distribution in desiccation sensitivity exists of recalcitrant seeds.

Furthermore, when seeds encounter desiccation, there is a substantial change in cell volume, which causes mechanical harm to the cell structure. This is especially likely to occur in the species with higher seed moisture content. In contrast, when developing orthodox seeds, reductions in vacuole volume or filling the vacuoles with insoluble substances can reduce this type of damage. While, the second type of damage is membrane structure damage. Water content is an important element of the membrane structure, and the presence of water to maintain the integrity of the membrane molecules. Once desiccated, the seed membrane structure loses water molecules, which reduces the completeness of the membrane and causes cellular dehydration damage. The final type of damage is caused by the metabolic system. Some seeds lose metabolic balance in their cells during desiccation, which generates an accumulation of free radicals inside the cells, leading to cell damage. According to previous studies report, seeds that are sensitive to desiccation have reduced metabolic activity during desiccation, which affects the seed's tolerance to desiccation (Pammenter and Berjak, 1999). After desiccating *P. macrocarpa* seeds in this study, a rapid reduction in seed viability was noted based on the germination percentage. In addition, observation of fresh *P. macrocarpa* seed tissue biopsies dyed with safranin showed that the cell nucleus comprised the majority of the cell volume. The cell membrane structure could be clearly observed, and obvious boarders between cells were apparent. After 9 days of desiccation, the cell membrane showed signs of rupture, the cell boarders had disappeared, and the cell nucleus began to rupture, deteriorate, and disappear. This suggested that the membrane structure had sustained damage. This finding was similar to the results of jackfruit and chinaberry seed desiccation. Jackfruit seeds that were not stored maintained a complete cell nucleus structure. The cells contained a number of large vacuoles, which comprised the majority of the cell volume. Plastids and mitochondria with suitable differentiation could be observed, indicating that cellular metabolic activity was normal (Wesley-Smith *et al.*, 2001). Chinaberry seeds exhibited similar characteristics after desiccation. When chinaberry seeds are desiccated to a moisture content of

0.26 g·g⁻¹, plasmolysis may occur between the plasma membrane and the cell wall, and a normal cellular structure cannot be observed (Kioko *et al.*, 2006). After jackfruit seeds were slowly desiccated, the original large volume vacuoles observed in the cell disappeared. The soluble materials originally located in the vacuoles dissolved into the cytoplasm, increasing the osmotic potential pressure in the cell. In addition, the nuclear membranes deteriorated, generating outward expansion (Wesley-Smith *et al.*, 2001). Although this study did not observe the detailed structures of the organelles in the cell's nucleus, we deduced from the cell membrane ruptures that the cell leakages were severe (Fig. 2). Comparing the seed germination percentages, the seeds were found to have already died after 9 days of desiccation. The reaction of recalcitrant seeds to desiccation was as described above.

This study discussed above verified that *P. macrocarpa* seeds are recalcitrant and sensitive to desiccation. After desiccation, the O₂⁻ and H₂O₂ content in the seed increased with decreases in the moisture content. When the seed moisture content declined below 1.7 gH₂O·g⁻¹ DW, the seed germination percentage was lower than 50%. Although the O₂⁻ content in the seed had risen slightly, the statistical deviation standard remained the same as that of fresh seeds. When the metabolic reaction was more downstream, the H₂O₂ content exhibited a clear increase, and when the moisture content declined below 1.4 gH₂O·g⁻¹ DW, the seeds no longer possessed germination abilities (Fig. 1). The O₂⁻ and H₂O₂ content in the seed at this time was 0.33 and 130.7 µg·g⁻¹ DW, respectively, higher than that when the moisture content was at 1.8 gH₂O·g⁻¹ DW (Fig. 3). When the seeds were desiccated to below 1.08 gH₂O·g⁻¹ DW in moisture content, the H₂O₂ quantity reached a peak of 162.3 µg·g⁻¹ DW (Fig. 3). Francini *et al.* (2006) indicated that after *A. Bidwillii* seeds were desiccated, the free radical content increased with decreases in the moisture content. For seeds desiccated to 21% moisture content, the free radical content increased by a maximum of 1.8 times. Quantity changes in reactive oxygen species O₂⁻ and H₂O₂ were observed in *A. saccharinum* L. (Pukacka and Ratajczak, 2006), *C. sinensis* Luntze (Chen *et al.*, 2010), and *A. toxicaria* (Xin *et al.*, 2010) seeds. The results show that as the seed moisture content decreases, the quantity of the reactive oxygen species increased. This was especially true in *C. sinensis* Luntze seeds, where when the seed moisture content declined below 20%, and the H₂O₂ quantity was eight times that of a fresh seed. The results of this study indicated that the increase in reactive oxygen species after desiccation for *P. macrocarpa* seeds conformed to the findings of the above report. SOD is the first line of defense against reactive oxygen species O₂⁻; it converts O₂⁻ into H₂O₂ and oxygen to reduce the toxicity of O₂⁻ (Polle, 2001). In this study, after the *P. macrocarpa* seeds were desiccated, the SOD activity declined with decreases in the moisture content. When the seed moisture content declined below 1.8 gH₂O·g⁻¹ DW, the SOD activity of desiccated *P. macrocarpa* seeds was significantly lower than that of the control group (Fig. 4).

A decrease in the moisture content of *T. cacao* and *P. macrocarpa* seeds also decreased the seed's SOD activity, which subsequently reduced the seed germination percentage and viability index of *P. macrocarpa* (Li and Sun, 1999; Li *et al.*, 2009). These results indicated that desiccation of *P. macrocarpa* seeds reduces SOD activity and seed desiccation tolerance. This result can be explained; desiccation damaged the *P. macrocarpa* seeds to be the first line of defense against oxidative.

After SOD reduces O₂⁻ into H₂O₂, the anti-oxidation enzyme APX reduces H₂O₂ into non-poisonous H₂O, and eliminates the toxicity caused by reactive oxygen species (Bailey, 2004). In this study, APX activity in *P. macrocarpa* seeds during the early stages of desiccation increased. However, as desiccation continued, APX activity decreased (Fig. 4B). After *A. toxicaria* seeds were desiccated, APX activity initially increased with decreases in the moisture content. However, as the seeds were further desiccated, APX activity began to decline (Xin *et al.*, 2010). The change in APX activity for *A. toxicaria* seeds occurred at a moisture content of 0.2–0.4 gH₂O·g⁻¹ DW, lower than that for *P. macrocarpa* seeds. This change in APX activity occurred at a moisture content of below 1.72 gH₂O·g⁻¹ DW (Fig. 4B), significantly higher than that for *A. toxicaria* seeds. Thus, we infer that these two seeds possess differing sensitivities to desiccation, and that *P. macrocarpa* seeds are more sensitive to desiccation.

AsA and GSH are important metabolites in plants. Their main function is to maintain the ascorbate–glutathione cycle by eliminating the H₂O₂ produced by upstream SOD metabolism and to solve cell oxidation problems (Polle, 2001). Ascorbate–glutathione cycle is a cyclic reaction that requires the completion of one cycle to effectively eliminate H₂O₂. The ASA content in *P. macrocarpa* seeds does not vary significantly during desiccation. By contrast, DHAsA increased during the initial stages of desiccation, and then decreased as desiccation continued (Fig. 5). However, the quantity of GSH in the same cycle drastically increased during desiccation (Fig. 6). Based on the ratio of the reduction status and oxidation status, the ascorbic acid and glutathione content during the early desiccation phase possess a higher level of DHAsA and GSSG in the oxidation state (Figs. 5B and 6B). Overall, ascorbate–glutathione cycle has a specific reduction state and oxidation state ratio, which indicates that this cycle can be successfully conducted during the early stages of desiccation. However, when the moisture content declined, the ratio between the two shifted toward a reduction state, indicating that the cycle had stopped. During the desiccation of *Q. robur*, *C. sativa*, and *A. saccharinum* seeds, the GSH quantity in the seed decreased with the moisture content, whereas the GSSG quantity increased. Examining the oxidation E_{GSSG/2GSH} value, the *C. sativa* seed showed increasingly severe oxidation after desiccation. This result indicates that the reactive oxygen species produced in the seed could no longer be metabolized by the ascorbate–glutathione cycle, leading to cell oxidation problems (Kranter *et al.*, 2006; Roach *et al.*, 2010; Pukacka and Ratajczak, 2006). This

result contrasts with that observed when *P. macrocarpa* seeds were desiccated; the ascorbate–glutathione cycle was in a reduction state. The main reason was that the SOD and APX in the upstream reaction of the *P. macrocarpa* anti-oxidation system had already lost activity during the desiccation process (Fig. 4). Consequently, the downstream GSH level increased and maintained a state of reduction during the desiccation process (Fig. 6), although normal operation of the antioxidation system could not be maintained.

Observing the actual oxidation in *P. macrocarpa* seeds after desiccation, the lipid peroxidation product MDA decreased with declines in the moisture content (Fig. 7). During the desiccation process, the thiobarbituric acid reactive substances content in *A. toxicaria* (Cheng and Song, 2008), *A. bidwillii* (Francini *et al.*, 2006), and *T. Cacao* (Li and Sun, 1999) seeds all increased with decreases in the moisture content. This reaction was identical to that observed for *P. macrocarpa* seeds in this study, indicating that metabolic system-induced damage appeared after *P. macrocarpa* seeds were desiccated.

Summarizing the above results, *P. macrocarpa* seeds are recalcitrant, and the germination percentage after the seeds were desiccated exhibited a U-shaped decline, conforming to the desiccation intolerance of recalcitrant seeds. During the desiccation period, the cell membrane system structure in *P. macrocarpa* seeds ruptured, producing physical structure damage. Simultaneously, the reactive oxygen species increased, as demonstrated by the reduced upstream SOD and APX activity in the anti-oxidation system. This produced peroxides in the seed, leading to metabolism induced damages. These results indicated that desiccation of *P. macrocarpa* seeds produced at least two causes that reduced viability. Li *et al.* (2009) asserted that after *P. macrocarpa* seeds were desiccated, SOD activity rapidly declined. This decline caused an inability to eliminate O_2^- , leading to severe oxidation, reduced seed viability, and seed aging and death. However, investigations of why recalcitrant *P. macrocarpa* seeds are desiccation intolerant and die during desiccation cannot verify that *P. macrocarpa* seeds died because of abnormal metabolic activity. The main reason is that abnormalities in the antioxidation system of *P. macrocarpa* seeds primarily occurred when the moisture content was below $1.22\text{ gH}_2\text{O}\cdot\text{g}^{-1}\text{ DW}$. At this point, the amount of SOD and APX activity was significant less than that of fresh seeds (Fig. 4). However, the moisture content threshold for seed death was $1.4\text{ gH}_2\text{O}\cdot\text{g}^{-1}\text{ DW}$ (Fig. 1). The seed's antioxidation system did not show a significant imbalance at this time, indicating that seed death occurred before the appearance of oxidation. Xin *et al.* (2011) focused on the relationship between seed viability and oxidation in *A. toxicaria* seeds after desiccation. Although Xin *et al.* indicated that desiccation affects seed leakage and viability, and also accompanied by an increases in seed oxidation. However, this level of oxidation is not sufficient to cause seed death. *Avicennia marina* seeds exhibited the greatest decline in viability when the seeds were desiccated to a moisture content of

54% to 57% (Greggains *et al.*, 2001). However, lipid peroxidation at this time actually decreased. Thus, the loss of seed viability at high moisture content may be the result of mechanical or physiological damage caused by a drastic reduction in cell volume, not metabolic damage. The reduction in viability of *P. macrocarpa* seeds following desiccation was the same, and primarily accompanied by an increase in reactive oxygen species and an imbalance in the anti-oxidation system. However, the main factors influencing seed intolerance to desiccation and the reasons seeds cannot be stored for as long as orthodox seeds still require further experiments for clarification.

ACKNOWLEDGEMENTS

This study was supported by a grant from the Agriculture and Food Agency, Council of Agriculture, Executive Yuan, Taiwan, ROC. 99AS-4.2.2-FD-Z3 and 100AS-4.2.2-FD-Z3. We are also grateful to Mr. Wu-Chou Chung and Mr. Chi-Hua Lin for providing the seeds materials and to Miss Ying-Jhen Chen for help investigating.

REFERENCES

- Ajaji, S. A., P. Berjak, J. I. Kioko, M. E. Dulloo, R. S. Vodouhe. 2006 Responses of fluted pumpkin (*Telfairia occidentalis* Hook. f., Cucurbitaceae) seeds to desiccation, chilling and hydrated storage. *S. Afr. J. Bot.*, **72**: 544–550
- Bailly, C. 2004 Active oxygen species and antioxidants in seed biology. *Seed Sci. Res.*, **14**: 93–107
- Berjak, P. and N. W. Pammenter. 2001 Seed recalcitrance : current perspectives. *S. Afr. J. Bot.*, **67**: 79–89
- Bonner, F. T. 1996 Responses to drying of recalcitrant seeds of *Quercus nigra* L. *Ann. Bot.*, **78**: 181–187
- Chen, Q., L. Yang, P. Ahmad, X. Wan, X. Hu. 2010 Proteomic profiling and redox status alteration of recalcitrant tea (*Camellia sinensis*) seed in response to desiccation. *Planta*, **233**: 583–592
- Cheng, H. Y. and S. Q. Song. 2008 Possible involvement of reactive oxygen species scavenging enzymes in desiccation sensitivity of *Antiaris toxicaria* seeds and axes. *J. Integr. Plant Biol.*, **50**: 1549–1556
- Ellis, R. H., T. D. Hong, and E. H. Roberts. 1991 An intermediate category of seed storage behaviour? II. Effects of provenance, immaturity, and imbibition on desiccation-tolerance in coffee. *J. Exp. Bot.* **42**: 653–657
- Farrant, J. M., N. W. Pammenter, P. Berjak. 1988 Recalcitrance— a current assessment. *Seed Sci. Tech.*, **16**: 155–166
- Francini, A., L. Galleschi, F. Saviozzi, C. Pinzino, R. Izzo, C. Sgherric, F. Navari-Izzo. 2006 Enzymatic and non-enzymatic protective mechanisms in recalcitrant seeds of *Araucaria bidwillii* subjected to desiccation. *Plant Physiol. Biochem.*, **44**: 556–563
- Greggains, V, W. E. Finch-Savage, N. M. Atherton, P. Berjak. 2001 Viability loss and free radical processes during desiccation of recalcitrant *Avicennia marina* seeds. *Seed Sci. Res.*, **11**: 235–242
- Heath, R. L. and L. Paker. 1968 Photoperoxidation in isolated chloroplasts. I. Kinetics and stoichiometry to fatty acid peroxidation. *Arch. Biochem. Biophys.*, **125**: 189–198
- Hong, T. D. and R. H. Ellis. 1996 A Protocol to determine seed storage behavior. IPGRI Technical Bulletin No. 1. IPGRI., Rome
- Hong, T. D., S. Linington, R. H. Ellis. 1996 Seed storage behavior : a compendium. Handbooks for Genebanks : No. 4. IPGRI.,

- Rome
- Hong, T. D., S. Linington, R. H. Ellis. 1998 Compendium of information on seed storage behavior, Vol. 1 and 2. Royal Botanic Garden, Kew, UK
- Kioko, J. I., P. Berjak, N. W. Pammenter. 2006 Viability and ultrastructural responses of seeds and embryonic axes of *Trichilia emetica* to different dehydration and storage conditions. *S. Afr. J. Bot.*, **72**: 167–176
- Kranner, I., S. Birtić, K. M. Anderson, H. W. Pritchard. 2006 Glutathione half-cell reduction potential: a universal stress marker and modulator of programmed cell death? *Free Radic. Biol. Med.*, **40**: 2155–2156
- Law, M. Y., S. A. Charles, B. Halliwell. 1983 Glutathione and ascorbic acid in spinach (*Spinacia oleracea*) chloroplasts—the effect of hydrogen peroxide and of paraquat. *Biochem. J.*, **210**: 899–903
- Li, C. and W. Q. Sun. 1999 Desiccation sensitivity and activities of free radical scavenging enzymes in recalcitrant *Theobroma cacao* seeds. *Seed Sci. Res.*, **9**: 209–217
- Li, Y. H., Y. M. Ma, L. Han. 2009 Relationship between tolerance to dehydration and superoxide dismutases (SODs) in *Pachira macrocarpa*. *Scientia Silvae Sinica*, **45**: 74–79
- Liu, J., J. Zhou, D. Xing. 2012 Phosphatidylinositol 3-kinase plays a vital role in regulation of rice seed vigor via altering NADPH oxidase activity. *Plos One*, **7**: 1–11
- Nakano, Y. and K. Asada. 1981 Hydrogen peroxide is scavenged by ascorbate specific peroxidase in spinach chloroplasts. *Plant Cell Physiol.*, **22**: 867–880
- Pammenter, N. W. and P. Berjak. 1999 A review of recalcitrant seed physiology in relation to desiccation-tolerance mechanisms. *Seed Sci. Res.* **9**: 13–37
- Paoletti, F., D. Aldinucci, A. Mocali, and A. Capparini. 1986 A sensitive spectrophotometric method for the determination of superoxide dismutase activity in tissue extracts. *Anal. Biochem.*, **154**: 536–541
- Polle, A. 2001 Dissecting the superoxide dismutase-ascorbate-glutathione-pathway in chloroplasts by metabolic modeling. *Plant Physiol.*, **126**: 445–462
- Pukacka, S. and E. Ratajczak. 2006 Antioxidative response of ascorbate-glutathione pathway enzymes and metabolites to desiccation of recalcitrant *Acer saccharinum* seeds. *J. Plant Physiol.*, **163**: 1259–1266
- Roach, T., R. P. Beckett, F. V. Minibayeva, L. Colville, C. Whitaker, H. Chen, C. Bailly, I. Kranner. 2010 Extracellular superoxide production, viability and redox poise in response to desiccation in recalcitrant *Castanea sativa* seeds. *Plant Cell Environ.* **33**: 59–75
- Roberts, E. H. 1973 Predicting the storage life of seeds. *Seed Sci. Tech.*, **1**: 499–514
- Sagisaka, S. 1976 The occurrence of peroxide in perennial plant, *Populus gelrica*. *Plant Physiol.*, **57**: 308–309
- Smith, I. K. 1985 Stimulation of glutathione synthesis in photo respiring plants by catalase inhibitors. *Plant Physiol.*, **79**: 1044–1047
- Sun, Y. W., S. T. Kuo, M. T. Liu, W. W. Liao, W. C. Chung. 2011 The storage and mutagenic technique in *Pachira macrocarpa*. *Seed Sci. Tech.*, **74**: 14–16
- Tang, A. J. and C. L. Long. 2008 Seed germination of *Lasia spinosa* as a function of temperature, light, desiccation, and storage. *Aquat. Bot.*, **89**: 352–365
- Tsai, S. H. 1975 A scheme of plant tissue section technology. Mau-Chang Co. Taipei, Taiwan.
- Wesley-Smith, J., N. W. Pammenter, P. Berjak, C. Walters. 2001 The effects of two drying rates on the desiccation tolerance of embryonic axes of recalcitrant jackfruit (*Artocarpus heterophyllus* Lamk.) seeds. *Ann. Bot.*, **88**: 653–664
- Win, B. 2008 On the compound quantitative characteristic trait of seed recalcitrant. *Acta Bot. Yunnanica*, **30**: 76–88
- Xin, X., X. M. Jing, Y. Liu, and S. Q. Song. 2010 Viability loss pattern under rapid dehydration of *Antiaris toxicaria* axes and its relation to oxidative damage. *J. Integr. Plant Bio.* **52**: 434–441