Three Dimensional Spheroid Culture of Canine Amniotic Fluid Derived Mesenchymal Stem Cells Enhances Differentiation Efficacy

Ha, Yu Na

Department of Animal Science and Biotechnology, College of Agriculture and Life Science, Chungnam National University

Chun, Ju Lan

Department of Animal Science and Biotechnology, College of Agriculture and Life Science, Chungnam National University | Kuju Agricultural Research Center, Faculty of Agriculture, Kyushu University

Kim, Eun Young

Department of Animal Science and Biotechnology, College of Agriculture and Life Science, Chungnam National University

Lee, Ji Hey

Department of Animal Science and Biotechnology, College of Agriculture and Life Science, Chungnam National University

他

<https://doi.org/10.5109/1526398>

出版情報：九州大学大学院農学研究院紀要. 60 (2), pp.377-384, 2015-09-18. Faculty of Agriculture, Kyushu University

バージョン：

権利関係：

Three Dimensional Spheroid Culture of Canine Amniotic Fluid Derived Mesenchymal Stem Cells Enhances Differentiation Efficacy

Yu Na HA^{1**}, Ju Lan CHUN^{1**}, Eun Young KIM¹, Ji Hey LEE¹, Bo Myoung LEE¹,
Kyung-Bon LEE², Takafumi GOTOH, Mina LEE^{3*} and Min Kyu KIM^{1*}

Kuju Agricultural Research Center, Faculty of Agriculture,
Kyushu University, Kuju, Oita, 878-0201, Japan
(Received May 18, 2015 and accepted May 19, 2015)

Amniotic fluid derived mesenchymal stem cells (AFS cells) have been considered as an applicable treatment of stem cell therapy in regenerative medicine. AFS cells have similar differentiation ability to embryonic stem cells with less ethical controversy. For applying in regeneration of injured and damaged tissues there are still considerable hurdles which need to be overcome. One of the crucial problems is how sustains or even amplifies their multi-potent ability till administration in damaged tissue/organ sites. The 2-D adherent culture method provides different environment compared to those of in vivo. 3-D culture method is an alternative manner to culture cells in in vivo-like cell culture condition. Hanging drop is one of simple and effective 3-D culture systems. Here canine AFS cells were cultured in the hanging drops in which cells form spheroids with micro-networks that allow exchanging cellular signalings similar to in vivo. Canine AFS cells were aggregated in tightly spaced spheres in hanging drops supplemented with 20% FBS. In the study of cellular senescence, AFS cells successfully formed and maintained spheroids regardless of their passages. In addition, the multipotent differentiation capacity of AFS cells in shape of 3-D spheroids was enhanced based on MSC differentiation assay after hanging drop culture. Noticeably, the expression levels of cartilage oligomeric matrix protein (COMP) and aggrecan (AGG) were significantly increased in 3-D spheroid cultured AFS cells compared to 2-D adherent cultured controls. Therefore, induction of spheroid formation by hanging drops is a useful method to provide 3-D in vivo-like culture condition without changes of AFS cells differentiation capacity especially to induce chondrogenic differentiation for regenerating damaged cartilages.

Key words: Amniotic fluid derived mesenchymal stem cells, 3-D spheroid culture, Hanging drop, Multipotency, Chondrogenic differentiation

INTRODUCTION

Mesenchymal stem cells (MSCs) have a great ability for tissue engineering because of their unlimited in vitro expansion potential, self-renewal capacity and multipotentiality (Kolf *et al.*, 2007; Sacchetti *et al.*, 2007). For several years, MSCs have been largely studied and used as a new therapeutic tool for clinical applications. It has been suggested that MSCs could reside in virtually all post-natal organs and tissues (da Silva *et al.*, 2006). MSCs were first identified in the bone marrow (Friedenstein *et al.*, 1976) but are now described to exist in connective tissues and particularly in adipose tissue (Zuk *et al.*, 2002), placenta (Fukuchi *et al.*, 2004), umbilical cord (Romanov *et al.*, 2003), dental pulp (Gronthos *et al.*, 2000), tendon (Bi *et al.*, 2007), trabecular bone (Nöth *et al.*, 2002) and synovium (De Bari *et al.*, 2001).

Multi-potent MSCs differentiate into distinctive end-stage cell types such as bone, cartilage, muscle, bone marrow stroma, tendon/ligament, fat, dermis and other connective tissues. Moreover it has been known that MSCs release various soluble factors which are immunoregulatory and support to regenerate microenvironment of tissues. In the damaged tissues, MSCs are especially believed to secrete a broad spectrum of paracrine factors that participate in the regenerative microenvironment and regulate immune infiltration (Fong *et al.*, 2011).

Amniotic fluid is comprised of embryonic or fetal cells derived from three germ layers (In't anker *et al.*, 2004; De Coppi *et al.*, 2007). It has been considered that MSCs in amniotic fluid would be similar to embryonic stem cells as the inherent precursors of all three germ layers. Since the initial identification of human amniotic fluid derived mesenchymal stem cells (hAFS cells), amniotic fluid has been used as a source of stem cells for cell therapy and regenerative medicine (De Coppi *et al.*, 2007). Amniotic fluid derived mesenchymal stem cells express CD90, CD105, and CD73 including Oct4 which is an embryonic stem cell marker (In't Anker *et al.*, 2004; De Gernmis *et al.*, 2006; De Coppi *et al.*, 2007). Oct4 was expressed over 90% of AFS cells. AFS cells have a self-renewal and differentiation potential toward adipocytes, osteoblasts, chondrocytes (Kolambkar *et al.*, 2007), and neurocytes (Rehni *et al.*, 2007) indicating that amniotic fluid may be a promising source of regenerative medicine without any ethical concerns unlike

¹ Department of Animal Science and Biotechnology, College of Agriculture and Life Science, Chungnam National University, 220 Gung-dong, Yuseong-gu, Daejeon 305-764, Korea

² Department of Biology Education, College of Education, Chonnam National University, Gwangju 500-757, Korea

³ Department of Obstetrics and Gynecology Chungnam National University School of Medicine, 266 Munwha-Ro, Jung-Gu, Daejeon 301-747, Korea

* Corresponding author: Min Kyu Kim, DVM, Ph.D. and Mina Lee, MD, Ph.D. E-mail: kminkyu@cnu.ac.kr, minari73@cnuh.co.kr

** Yuna HA and Ju Lan Chun have contributed equally to this work.

embryonic stem cells (Prusa AR *et al.*, 2003; Chung Y *et al.*, 2008).

In general, cell based assay is performed in two-dimensional (2-D) culture system. Monolayer or suspension culture systems (2-D) present a large artificial cellular environment that limits and alter their cellular potential for therapeutic application. It seems that the 2-D culture system provides a non-physiological environment which affects innate cellular functions of cells (Friedrich *et al.*, 2007; Pampaloni *et al.*, 2010). 3-D systems have been used and applied to culture cells *in vitro* to overcome the limitation. 3-D culture systems mimic *in vivo* tissue environment and provide micro-networks of cells which allow cells to communicate each other like *in vivo* that supports them to sustain their function.

The objective of this study was to investigate the efficacy of 3-D culture system using hanging drops to form spheroid of canine AFS (cAFS) cells. Furthermore it was evaluated and compared the multi-lineage differentiation potential and morphological distinction of the cAFS cells between 3-D spheroids and 2-D monolayer. It demonstrated that the hanging drop method offered a microgravity circumstance and conserved constant condition with reliable oxygen and nutrient transfer that allow maintaining their multi-potent differentiation capacity which is involved to regenerate various tissues.

MATERIALS AND METHODS

Isolation and characterization of cAFS cells

All animal experiments were performed by the protocols approved by the Research Ethics Committee and the Institutional Animal Care and Use Committee of Chungnam National University. Canine amniotic fluids were collected from healthy pregnant beagles' amniocentesis using ultrasonographic guidance. Isolation and culture of cAFS cells were performed by the published protocols in Choi *et al.* (2013). Briefly, the cells were isolated from the amniotic fluid by centrifugation at 3,000 rpm for 10 min, and washed twice with Phosphate Buffered Saline (PBS, Gibco). Isolated cells were cultured in low glucose Dulbecco's modified eagle medium (L-DMEM, Sigma) supplemented with 10% fetal bovine serum (FBS, Gibco), 5 ng of fibroblast growth factor (FGF, Sigma), 10 ng of epidermal growth factor (EGF, Sigma) and 1% penicillin-streptomycin (Sigma) at 39°C, 5% CO₂ incubator for 4–5 days.

Spheroid formation and culture in 3-D micro-environment by hanging drop method

cAFS cell spheroids were prepared using the hanging drop method. AFS cells grown as a monolayer were dissociated with trypsin, collected by centrifugation and resuspended in growth medium with 5%, 10% or 20% of fetal bovine serum (FBS, Gibco) at a density of 1×10^6 cells per 1 ml. Hanging drops were formed on the culture dish lid after inverting it using 20 μ l of 20,000 cells in growth medium, and allowed to form cell aggregates at 39°C in 5% CO₂ for 3 days. The culture dish was filled with 10 ml PBS to warrant high humidity (Fig. 1A and

B). On day 3, the hanging drops were harvested, and the cells were transferred to a new culture dish. The spheroids were cultured for 5 days and culture medium was replenished every 3 days. cAFS cells were also cultured in adherent manner and used as 2-D control group. For differentiation assay, cAFS cells were cultured in hanging drop method for 3 days and then placed into adherent culture. Within 3 days of plating, cells initiated to expand from the edge of the attached spheroids.

Adipogenic differentiation

Adipogenic differentiation was induced as described by Pittenger *et al.* (1999). Briefly, 50,000 cells were seeded per well and cultured until confluence. For 3 days, the cells were treated with the adipogenic induction medium consisting of L-DMEM base medium, 10% FBS, 10 ng EGF, 5 ng bFGF, penicillin/streptomycin solution (10,000 IU/10,000 IU/ 100 ml), 1 μ M dexamethasone (Sigma-Alorich, USA), 0.2 mM indomethacin (Sigma-Alorich, USA), 1 mg/ml insulin (Lily, Korea) and 1 mM 3-isobutyl-1-methylxanthin (IBMX) (Sigma-Alorich, USA). Then the medium was changed every 3 days for 21 days. Cells were fixed with formaldehyde solution (Samchun chemical, Korea) (10% v/v in PBS) and washed with 3% (v/v) isopropanol (Amresco, USA). Samples were stained with 0.5% (w/v in 60% isopropanol) Oil Red O (Sigma-Alorich, USA) to determine the presence of oil droplet.

Osteogenic differentiation

40,000 cells were seeded per well in a six-well-plate and cultured until they reach confluence. The osteogenic induction medium was prepared as described by Pittenger *et al.* (1999). The medium is consisted of L-DMEM base medium supplemented with 10% FBS, 10 ng EGF, 5 ng bFGF, penicillin/ streptomycin solution (10,000 IU/10,000 IU/100 ml), 0.1 μ M dexamethasone, 10 mM β -glycerophosphate (Sigma-Alorich, USA) and 50 μ M ascorbate-2-phosphate (Sigma-Alorich, USA). Alizarin Red S (Sigma-Alorich, USA) staining was used to determine the presence of calcium deposition. For Alizarin Red S staining, cells were washed with distilled water (D.W.) two times and fixed in a solution of ice-cold 70% ethanol for 1 hr. After carefully washed 7 times with D.W., cells were stained for 10 min with 40 mM Alizarin Red S and then washed again with D.W. for 2 times in room temperature.

Chondrogenic differentiation

50,000 cells were seeded per well in a six-well-plate and cultured until they reach confluence. For 3 days, the cells were treated with the chondrogenic induction medium including L-DMEM base medium, 10 ng EGF, 5 ng bFGF, penicillin /streptomycin solution (10,000 IU/10,000 IU/ 100 ml) and 10 ng/ml transforming growth factor- β 1 (TGF- β 1) (Sigma-Alorich, USA). And then the cells were cultured in chondrogenic maintenance medium consisting of L-DMEM base medium, penicillin/ streptomycin solution (10,000 IU/10,000 IU/100 ml), TGF- β 1 (10 ng/ml), dexamethasone (0.1 μ M), ascorbate-2-

phosphate (50 μ M), sodium pyruvate (100 μ g/ml) (Sigma–Aldrich, USA), ITS (50 mg/ml) (Sigma–Aldrich, USA), cis–3–Hydroxy–DL–Proline (20 μ g/ml) (Sigma–Aldrich, USA) and 0.1 mg/ml insulin for 21 days. This cycle was repeated six times. Alizarin Red S staining was used to determine the presence of calcium mineralization. For Alizarin Red S staining, cells were washed with D.W. two times and fixed in a solution of ice–cold 70% ethanol for 1 hr. After carefully washed 7 times with D.W., cells were stained for 10 min with 40 mM Alizarin Red S and then washed again with D.W. for 2 times in room temperature.

Reverse–transcription polymerase chain reaction (RT–PCR)

Total RNA was extracted by RNA extract kit following the protocol of manufacturer (MACHEREY–NAGEL). Expression of pluripotent genes (Oct4, NANOG and SOX2) and adipogenic (LPL and Leptin), osteogenic (Runx2 and Osteocalcin), and chondrogenic (Comp and AGG) genes were detected from each samples. β –II microglobulin was used as an internal control. RNA samples (1 μ l of total RNA) were primed with oligo dT primer to synthesize cDNA using iScript reverse transcriptase (BIO–RAD). PCR reactions were consisted of 3 min denaturation at 94°C, followed by 35 cycles at 94°C for 30 sec, 60°C or 58°C for 30 sec and 72°C for 30 sec. A final extension for 5 min was performed at 72°C. β –II microglobulin RT–PCR reactions were modified to 25 cycles with annealing at 60°C. Each amplicon of targeted specific genes was confirmed by DNA sequencing. RT–PCR results were evaluated by Image J software. Primer sequences used were listed in Table 1.

RESULTS

Spheroid formation in 3–D micro–environment by hanging drop method

Using hanging drop method, cAFS cells reproducibly

were aggregated into 3–D spheroids. To characterize and estimate the self–assembling potential, spheroids were grown in different culture conditions. cAFS spheroids were generated in growth medium supplemented with 5%, 10% or 20% of FBS concentration (Fig 1). There were more compacted spheroids found in the medium with 20% FBS compared to (Fig 1. E and H) the medium with 5% (Fig 1. C and F) and 10% (Fig 1. D and G) of FBS, and compacted structure were maintained tightly with 20% FBS than 5% and 10% FBS.

Under the phase contrast microscope, cAFS cells were aggregated in the time dependent manner in hanging drops. Initially, cAFS cells formed a loose connection and then several small aggregates that progressively combined into a single spheroid (Fig 2. A). Once aggregated, the size of spheroids was not changed, and gradually compacted tensely between 48h and 96h after culturing (Fig 2. B).

In addition, it has investigated how the impact of senescence onto forming spheroids of cAFS cells is. Within 3 days of cultivation with 5, 10, and 15 passages of cAFS cells, there was no change of morphology and size enlargement observed. Regardless of the passages, cAFS cells were aggregated compactly, and most of spheroids were generated fairly (Fig 3. C–H).

cAFS cells differentiation

When cultured in adipogenic induction medium for 21 days, spheroid–derived cAFS cells (3–D spheroids in adherent culture) showed morphological changes and appeared as round shaped cells with lipid droplets in their cytoplasm more than those of in 2–D culture (monolayer culture) (Fig 3. A and B). cAFS cells from 3–D spheroids had large number of Oil Red O positive lipid droplets in their cytoplasm and had higher detachment rate than those from 2–D culture during the adipogenic induction (not shown). After induction of osteogenic differentiation cAFS cells were changed their morphology from spindle shape to cuboidal shape (Fig 3. C and D). Deposition of extracellular calcified matrix was increased in cAFS cells from 3–D spheroids compared to that of 2–D adherent culture. Chondrogenic differentiation of 3–D spheroid derived cAFS cells induced rapid proliferation and cAFS cells reached to 90% confluence before cAFS cells from 2–D culture (Fig 3. E and F). Moreover 3–D spheroids showed morphologically more chondrogenic differentiated than 2–D culture.

RT–PCR analysis of cAFS cell differentiation

After induction of MSC differentiation, the specific gene expressions of adipogenic–, osteogenic– and chondrogenic lineages were examined by RT–PCR. The expression of LPL, Leptin (specific markers of adipocytes), Runx2, and Osteocalcin (specific markers of osteoblasts) were detected in both 2–D and 3–D cultured cAFS cells (Fig. 4). Interestingly the expression of COMP and AGG (specific markers of chondrocytes) were barely detectable in 2–D culture. However, under 3–D culture condition COMP and AGG expression were significantly increased (Fig 4. A and B). The 3–D spheroid derived

Table 1. Reverse–transcription polymerase chain reaction primers for adipogenic–, osteogenic–, and chondrogenic–specific genes

Genes	Primer sequences (5'–3')	Product size
LPL	F : AAAACCATCGTGGGCAATTA	204 bp
	R : ACAATTGTGATTCCCAGCAA	
Leptin	F : TTCCACCATCCTGCCACTAT	200 bp
	R : ACCATCTGGAATGCAAGGTC	
Runx2	F : CCCAACTTCCTGTGCTCTGT	197 bp
	R : TCGTTGAACCTTGCTACTTGG	
Osteocalcin	F : AGGGAAGTATGCGAGCTCAA	198 bp
	R : GATGACAAGGACCCCACT	
COMP	F : AGACATACTGGCAGGCGAAT	197 bp
	R : CCAGCCGTAGGATGTCTTGT	
Aggrecan	F : ATCAACAGTGCTTACCAAGACA	130 bp
	R : ATAACCTCACAGCGATAGATCC

Fig. 1. Illustration (A) and macroscopic view (B) of hanging drop culture method. Cell suspension drops were deposited onto the underside of the lid of the culture dish. When the lid is inverted, drops are held in place by surface tension and gradually precipitated. Formation of cAFS spheroids with 20,000 cells/20 μ l with FBS concentration of 5% (C and F), 10% (D and G) and 20% (E and H) following 48 hr and 72 hr of hanging drop culture. (Scale bar, 300 μ m), Magnification : X100

Fig. 2. Time courses of the aggregation of cAFS into a spheroid in a hanging drop (A). Spheroids progressively compacted between 48 and 96 h. (B) Formation of spheroids with cAFS at passage 5 (C and F), 10 (D and G), and 15 (E and H) after 48hr and 72 hr.

Fig. 3. Cytochemical differentiation assay. cAFS differentiation into adipocytes detected by Oil Red O staining (A and B), osteoblasts (C and D), and chondrocytes (E and F) detected by Alizarin Red staining. Adherent cultured (A, C and E) and hanging drop cultured (B, D and F).

cAFS cells expressed 200% and 600% higher levels of COMP and AGG than those of the 2-D cultured cells (Fig 4. B).

DISCUSSION

Mesenchymal stem cells (MSCs; also called mesenchymal stromal cells) are defined as adherent cells which are relatively easy to isolate from donors and patients, and highly proliferate and differentiate into several different cell types in vitro and in vivo (Pittenger *et al.*, 1999; Kolf *et al.*, 2007). A study in 1996 was the first to suggest the possibility of multilineage potential of non-hematopoietic cells present in the amniotic fluid, by showing myogenic conversion of amniocytes (Streubel *et al.*, 1996). Human amniotic cells have been shown their multipotent potential that is able to differentiate at least into neural glial cells and hepatocyte precursors (Sakuragawa *et al.*, 1996; Elwan *et al.*, 1997; Saulnier *et al.*, 2009). The primary advantage of AFS cells is the easiness of harvest, isolation, and expansion in culture with highly advanced differentiation capacity into multi cell lineages that allow them to be a useful source of the stem cell therapy in regenerative medicine.

Most adult stem cell base assays were applied in monolayer or suspension cultures which provide a non-physiological environment to the cells that alters cell-cell and cell-ECM interaction unlike in vivo. For that

reason, it has been accepted that mesenchymal stem cells lose their innate potentials including morphology, proliferation, gene expression, and cellular function after passages in vitro. On the other hand 3-D culture system has been used in several different cell types including tumor cells and human primary cells (Bjerkvig *et al.*, 1997; Kunz-Schughart *et al.*, 1999; Kelm *et al.*, 2004). These studies demonstrated that 3-D cell culture system is reproducible, and would be a definitive method for the formation of 3-D spheroids like the hanging drop method that provides in vivo-like environment with many potential advantages. It seems that 3-D culture provides in vivo-like microenvironment with higher similarity to in vivo conditions in many aspects for tissue engineering and regenerative medicine. For examples, previous studies reported that the spheroid formation method resulted in a beneficial culture environment for reconstruction of liver (Landry *et al.*, 1985), pancreas (Matta *et al.*, 1994), blood vessel (Korff *et al.*, 1998) and bone tissue (Akiyama *et al.*, 2006). Spheroids have also offered as biological models for avascular tumor as drug delivery assays and toxicology. Recently they have gained attention for their potential to serve as a building block for organ reconstruction.

In this study, cAFS cells were induced to form spheroids from aggregates by hanging drop method. After forming spheres, spheroids become gradually compact and were sustained their spheroid structure without dis-

Fig. 4. RT-PCR analysis of lineage specific gene expression for adipocytes, osteoblasts and chondrocytes. Total RNA was extracted from the cells under each differentiation condition. (A) RT-PCR amplicon images and (B) statistical analysis of the relative gene expression. 2-D culture; monolayer culture, 3-D culture; Hanging drop culture, β -II Microglobulin; the internal control.

assembly. To identify the multi-lineage differentiation potential, spheroid derived cAFS cells were induced to differentiate into adipocytes, osteoblasts, and chondrocytes. In MSC differentiation assay, the spheroid-derived cAFS cells acquired adipogenic, osteogenic and chondrogenic morphological changes with typical staining patterns. The spheroid-derived cAFS cells had a higher ratio of positive patterns with Oil-Red-O or Alizarin Red S staining after each differentiation assay compared with 2-D culture system. The cells cultured in 3-D spheroids had remarkable detachment during differentiation compared to cells cultured in 2-D system.

In addition, lineage specific gene expressions were confirmed by RT-PCR analysis. Adipocyte (LPL and Leptin), Osteoblast (Runx2 and Osteocalcin), and chondrocytes (COMP and AGG) specific genes were abundantly expressed in both groups. However, in case of chondrogenic differentiation, AGG was not detected and COMP was barely expressed in the 2-D cultured cells. Under 3-D culture condition COMP and AGG expression were significantly increased.

These results indicated that hanging drop culture method supports to maintain multi-lineage differentiation ability of cAFS cells, and even amplified the differentiation potential of cAFS cells into chondrogenic lineage. The beneficial effect of 3-D spheroids culture system for chondrogenic differentiation has been reported (Kii *et al.*, 2004; Arufe *et al.*, 2009). In the hanging drop culture, cadherin causes cells to adhere intensively each other through homophilic calcium-dependent interaction through their extracellular domains. With that, it is possible to form tight cadherin-mediated cell-cell interactions and spherical cell aggregates. In addition 3-D spheroid formation has a positive effect on cartilaginous matrix production of cells (Sekiya *et al.*, 2002; Shirasawa *et al.*, 2006).

To regenerate osteochondral defects, tissue transplantations were previously performed (Amiel *et al.*, 1985; Yamashita *et al.*, 1985; O'Driscoll *et al.*, 1986; Matsusue *et al.*, 1993). Chondrocytes and mesenchymal stem cells have been applied for transplantations that were contributed in regeneration of hyaline-like cartilage with reduc-

ing clinical symptoms. However, the optimal resource of stem cells for the osteochondral repair has not yet been established. Besides, transplanted single stem cells hardly reside and disappear in the transplanted sites due to joint motion and weight supporting besides the immune cell attack. Studies suggested that the small amounts of survived transplanted 3-D spheroid derived cells could contribute to chondrogenic and osteogenic lineage cells, and they encourage the reconstruction of the defects especially during the beginning of repair process (Kim *et al.*, 1998; No Da *et al.*, 2012). Therefore, 3-D spheroid would be a suitable cell format to transplant highly concentrated stem cells into damaged tissue sites with less possibility of losing cells unlike transplantation of single cells.

In conclusion, this study demonstrated that the simple and economical hanging drop method provides 3-D in vivo-like culture environment that can effectively maintain multi-lineage differentiation potency and even improved chondrogenic differentiation ability of cAFS cells. The results support that 3-D spheroids derived AFS cells would be a potential and applicable source of stem cell therapy in regeneration medicine especially osteochondral defects.

ACKNOWLEDGEMENT

This study was carried out with the support of “Cooperative Research Program for Agriculture Science & Technology Development (Project No. PJ009333)” Rural Development Administration. Republic of Korea.

AUTHOR DISCLOSURE STATEMENT

None of the authors have any competing financial interests to declare.

REFERENCES

- Akiyama, M., H. Nonomura, S. H. Kamil and R. A. Ignatz 2006 Periosteal cell pellet culture system: A new technique for bone engineering. *Cell Transplant*, **15**: 521–532
- Amiel, D., R. D. Coutts, M. Abel, W. Stewart, F. Harwood and W. H. Akeson 1985 Rib perichondrial grafts for the repair of full-thickness articular–cartilage defects. A morphological and biochemical study in rabbits. *J. Bone Joint Surg. Am.*, **67**: 911–920
- Arufe, M. C., A. De la Fuente, I. Fuentes–Boquete, F. J. De Toro and F. J. Blanco 2009 Differentiation of synovial CD–105(+) human mesenchymal stem cells into chondrocyte-like cells through spheroid formation. *J. Cell Biochem.*, **108**: 145–155
- Bi, Y., D. Ehrichtiou, T. M. Kilts, C. A. Inkson, M. C. Embree, W. Sonoyama, L. Li, A. I. Leet, B. M. Seo, L. Zhang, S. Shi and M. F. Young 2007 Identification of tendon stem/progenitor cells and the role of the extracellular matrix in their niche. *Nat. Med.*, **13**: 1219–1227
- Bjerkvig, R., M. Lund–Johansen and K. Edvardson 1997 Tumor cell invasion and angiogenesis in the central nervous system. *Curr. Opin. Oncol.*, **9**: 223–229
- Choi, S. A., H. S. Choi, K. J. Kim, D. S. Lee, J. H. Lee, J. Y. Park, E. Y. Kim, X. Li, H. Y. Oh, D. S. Lee and M. K. Kim 2013 Isolation of canine mesenchymal stem cells from amniotic fluid and differentiation into hepatocyte-like cells. *In Vitro Cell Dev. Biol. Anim.*, **19**: 42–51
- Chung, Y., I. Klimanskaya, S. Becker, T. Li, M. Maserati, S. J. Lu, T. Zdravkovic, D. Ilic, O. Genbacev, S. Fisher, A. Krtolica and R. Lanza 2008 Human embryonic stem cell lines generated without embryo destruction. *Cell Stem Cell*, **2**: 113–117
- da Silva Meirelles, L., P. C. Chagastelles and N. B. Nardi 2006 Mesenchymal stem cells reside in virtually all post-natal organs and tissues. *J. Cell. Sci.*, **119**: 2204–2213
- De Bari, C., F. Dell’Accio, P. Tylzanowski and F. P. Luyten 2001 Multipotent mesenchymal stem cells from adult human synovial membrane. *Arthritis Rheum.*, **44**: 1928–1942
- De Coppi, P., G. Bartsch Jr., M. M. Siddiqui, T. Xu, C. C. Santos, L. Perin, G. Mostoslavsky, A. C. Serre, E. Y. Snyder, J. J. Yoo, M. E. Furth, S. Soker and A. Atala 2007 Isolation of amniotic stem cell lines with potential for therapy. *Nat. Biotechnol.*, **25**: 100–106
- De Coppi, P., A. Callegari, A. Chiavegato, L. Gasparotto, M. Piccoli, J. Taiani, M. Pozzobon, L. Boldrin, M. Okabe, E. Cozzi, A. Atala, P. Gamba and S. Sartore 2007 Amniotic fluid and bone marrow derived mesenchymal stem cells can be converted to smooth muscle cells in the cryo-injured rat bladder and prevent compensatory hypertrophy of surviving smooth muscle cells. *J. Urol.*, **177**: 369–376
- Elwan, M. A. and N. Sakuragawa 1997 Evidence for synthesis and release of catecholamines by human amniotic epithelial cells. *Neuroreport*, **8**: 3435–3438
- Fong, E. L., C. K. Chan and S. B. Goodman 2011 Stem cell homing in musculoskeletal injury. *Biomaterials*, **32**: 395–409
- Friedenstein, A., J. Gorskaja and N. N. Kulagina 1976 Fibroblast precursors in normal and irradiated mouse hematopoietic organs. *Exp. Hematol.*, **4**: 267–274
- Friedrich, J., R. Ebner and L. Kunz–Schughart 2007 Experimental anti-tumor therapy in 3-D: spheroids old hat or new challenge? *Int. J. Radiat. Biol.*, **83**: 849–871
- Fukuchi, Y., H. Nakajima, D. Sugiyama, I. Hirose, T. Kitamura and K. Tsuji 2004 Human placenta-derived cells have mesenchymal stem/progenitor cell potential. *Stem Cells*, **22**: 649–658
- Gronthos, S., M. Mankani, J. Brahimi, P. G. Robey and S. Shi 2000 Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *Proc. Natl. Acad. Sci. USA*, **97**: 13625–13630
- In’t Anker, P. S., S. A. Scherjon, C. Kleijburg–van der Keur, G. M. de Groot–Swings, F. H. Claas, W. E. Fibbe and H. H. Kanhai 2004 Isolation of mesenchymal stem cells of fetal or maternal origin from human placenta. *Stem cells*, **22**: 1338–1345
- Kii, I., N. Amizuka, J. Shimomura, Y. Saga and A. Kudo 2004 Cell–Cell Interaction Mediated by Cadherin–11 Directly Regulates the Differentiation of Mesenchymal Cells Into the Cells of the Osteo–Lineage and the Chondro–Lineage. *J. Bone Miner. Res.*, **19**: 1840–1849
- Kelm, J. M. and M. Fussenegger 2004 Microscale tissue engineering using gravity-enforced cell assembly. *Trends Biotechnol.*, **22**: 195–202
- Kim, S. S., H. Utsunomiya, J. A. Koski, B. M. Wu, M. J. Cima, J. Sohn, K. Mukai, L. G. Griffith and J. P. Vacanti 1998 Survival and function of hepatocytes on a novel three-dimensional synthetic biodegradable polymer scaffold with an intrinsic network of channels. *Ann. Surg.*, **228**: 8–13
- Kolambkar, Y. M., A. Peister, S. Soker, A. Atala and R. S. Guldberg 2007 Chondrogenic differentiation of amniotic fluid-derived stem cells. *J. Mol. Histol.*, **38**: 405–413
- Kolf, C. M., E. Cho and R. S. Tuan 2007 Mesenchymal stromal cells. Biology of adult mesenchymal stem cells: regulation of niche, self-renewal and differentiation. *Arthritis Res. Ther.*, **9**: 204–213
- Korff, T. and H. G. Augustin 1998 Integration of endothelial cells in multicellular spheroids prevents apoptosis and induces differentiation. *J. Cell Biol.*, **143**: 1341–1352
- Kunz–Schughart, L. A. 1999 Multicellular tumor spheroids: intermediates between monolayer culture and in vivo tumor. *Cell Biol. Int.*, **23**: 157–161
- Landry, J., D. Bernier, C. Ouellet, R. Goyette and N. Marceau 1985 Spheroidal aggregate culture of rat liver cells: Histotypic reorganization, biomatrix deposition, and maintenance of functional activities. *J. Cell Biol.*, **101**: 914–923

- Levenberg, S., N. F. Huang, E. Lavik, A. B. Rogers, J. Itskovitz-Eldor and R. Langer 2003 Differentiation of human embryonic stem cells on three-dimensional polymer scaffolds. *Proc. Natl. Acad. Sci. U S A*, **100**: 12741–12746
- Matta, S. G., J. D. Wobken, F. G. Williams and G. E. Bauer 1994 Pancreatic islet cell reaggregation systems: efficiency of cell reassociation and endocrine cell topography of rat islet-like aggregates. *Pancreas*, **9**: 439–449
- Matsusue, Y., T. Yamamuro and H. Hama 1993 Arthroscopic multiple osteochondral transplantation to the chondral defect in the knee associated with anterior cruciate ligament disruption. *Arthroscopy*, **9**: 318–321
- No, da Y., S. A. Lee, Y. Y. Choi, D. Park, J. Y. Jang, D. S. Kim and S. H. Lee 2012 Functional 3D human primary hepatocyte spheroids made by co-culturing hepatocytes from partial hepatectomy specimens and human adipose-derived stem cells. *PLoS One*, **7**: e50723
- Nöth, U., A. M. Osyczka, R. Tuli, N. J. Hickok, K. G. Danielson and R. S. Tuan 2002 Multilineage mesenchymal differentiation potential of human trabecular bone-derived cells. *J. Orthop. Res.*, **20**: 1060–1069
- O'Driscoll, S. W., F. W. Keeley and R. B. Salter 1986 The chondrogenic potential of free autogenous periosteal grafts for biological resurfacing of major full-thickness defects in joint surfaces under the influence of continuous passive motion. An experimental investigation in the rabbit. *J. Bone Joint Surg. Am.*, **68**: 1017–1035
- Pampaloni, F. and E. Stelzer 2010 Three-dimensional cell cultures in toxicology. *Biotechnol. Genet. Eng. Rev.*, **26**: 117–138
- Pittenger, M. F., A. M. Mackay, S. C. Beck, R. K. Jaiswal, R. Douglas, J. D. Mosca, M. A. Moorman, D. W. Simonetti, S. Craig and D. R. Marshak 1999 Multilineage potential of adult human mesenchymal stem cells. *Science*, **284**: 143–147
- Prusa, A. R., E. Marton, M. Rosner, G. Bernaschek and M. Hengstschlager 2003 Oct-4-expressing cells in human amniotic fluid: a new source for stem cell research? *Hum. Reprod.*, **18**: 1489–1493
- Rehni, A. K., N. Singh, A. S. Jaggi and M. Singh 2007 Amniotic fluid derived stem cells ameliorate focal cerebral ischaemia-reperfusion injury induced behavioural deficits in mice. *Behav. Brain Res.*, **183**: 95–100
- Romanov, Y. A., V. A. Svintsitskaya and V. N. Smirnov 2003 Searching for alternative sources of postnatal human mesenchymal stem cells: candidate MSC-like cells from umbilical cord. *Stem Cells*, **21**: 105–110
- Sacchetti, B., A. Funari, S. Michienzi, S. Di Cesare, S. Piersanti, I. Saggio, E. Tagliafico, S. Ferrari, P. G. Robey, M. Riminucci and P. Bianco 2007 Self-renewing osteoprogenitors in bone marrow sinusoids can organize a hematopoietic microenvironment. *Cell*, **131**: 324–336
- Sakuragawa, N., R. Thangavel, M. Mizuguchi, M. Hirasawa and I. Kamo 1996 Expression of markers for both neuronal and glial cells in human amniotic epithelial cells. *Neurosci. Lett.*, **209**: 9–12
- Saulnier, N., W. Lattanzi, M. A. Puglisi, G. Pani, M. Barba, A. C. Piscaglia, M. Giachelia, S. Alfieri, G. Neri, G. Gasbarrini and A. Gasbarrini 2009 Mesenchymal stromal cells multipotency and plasticity: induction toward the hepatic lineage. *Eur. Rev. Med. Pharmacol. Sci.*, **1**: 71–78
- Sekiya, I., J. T. Vuoristo, B. L. Larson and D. J. Prockop 2002 In vitro cartilage formation by human adult stem cells from bone marrow stroma defines the sequence of cellular and molecular events during chondrogenesis. *Proc. Natl. Acad. Sci. U S A*, **99**: 4397–4402
- Shirasawa, S., I. Sekiya, Y. Sakaguchi, K. Yagishita, S. Ichinose and T. Muneta 2006 In vitro chondrogenesis of human synovium-derived mesenchymal stem cells: optimal condition and comparison with bone marrow-derived cells. *J. Cell Biochem.*, **97**: 84–97
- Streubel, B., G. Martucci-Ivessa, T. Fleck and R. E. Bittner 1996 In vitro transformation of amniotic cells to muscle cells—background and outlook. *Wien. Med. Wochenschr.*, **146**: 9–10
- Yamashita, F., K. Sakakida, F. Suzu and S. Takai 1985 The transplantation of an autogeneic osteochondral fragment for osteochondritis dissecans of the knee. *Clin. Orthop. Relat. Res.*, **201**: 43–50
- Zuk, P. A., M. Zhu, P. Ashjian, D. A. De Ugarte, J. I. Huang, H. Mizuno, Z. C. Alfonso, J. K. Fraser, P. Benhaim and M. H. Hedrick 2002 Human adipose tissue is a source of multipotent stem cells. *Mol. Biol. Cell*, **13**: 4279–4295