

Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space

Kawakami, Yu
Faculty of Mathematics, Kyushu University

<https://hdl.handle.net/2324/15425>

出版情報 : MI Preprint Series. 2009-27, 2012. University of Houston
バージョン :
権利関係 : (C) 2012 University of Houston



MI Preprint Series

Kyushu University
The Global COE Program
Math-for-Industry Education & Research Hub

Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space

Y. Kawakami

MI 2009-27

(Received August 11, 2009)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

VALUE DISTRIBUTION OF THE HYPERBOLIC GAUSS MAPS FOR FLAT FRONTS IN HYPERBOLIC THREE-SPACE

YU KAWAKAMI

ABSTRACT. We give an effective estimate for the totally ramified value number of the hyperbolic Gauss maps of complete flat fronts in the hyperbolic three-space. As a corollary, we give the upper bound of the number of exceptional values of them for some topological cases. Moreover, we obtain some new examples for this class.

INTRODUCTION

The study of flat surfaces in the hyperbolic 3-space \mathcal{H}^3 has made a great advance in the last decade. Indeed, Gálvez, Martínez and Milán [GMM] established a Weierstrass-type representation formula for such surfaces. Moreover, Kokubu, Umehara and Yamada ([KUY1], [KUY2]) investigated global properties of flat surfaces in \mathcal{H}^3 with certain kind of singularities, called flat fronts (For precise definition, see Section 1 of this paper). In particular, they gave a representation formula constructing a flat front from a given pair of hyperbolic Gauss maps and an Osserman-type inequality for complete (in the sense of [KUY2], see also Section 1 of this paper) flat fronts. More recently, Kokubu, Rossman, Saji, Umehara and Yamada [KRSUY] gave criteria for a singular point on a flat front in \mathcal{H}^3 be a cuspidal edge or swallowtail and proved the generically flat fronts in \mathcal{H}^3 admit only cuspidal edges and swallowtails. Moreover, Roitman [Ro] and Kokubu, Rossman, Umehara and Yamada [KRUY1] obtained interesting results on flat surfaces or (p-)fronts in \mathcal{H}^3 and their caustics. Furthermore, Kokubu, Rossman, Umehara and Yamada [KRUY2] also investigate the asymptotic behavior of ends of flat fronts in \mathcal{H}^3 . However, we have not seen the study of value distribution of the hyperbolic Gauss maps for complete flat fronts in \mathcal{H}^3 before.

On the other hand, we have recently obtained some results on value distribution of the Gauss map of complete minimal surfaces in Euclidean 3-space \mathbb{R}^3 and the hyperbolic Gauss map of complete constant mean curvature one (CMC-1, for short) surfaces in \mathcal{H}^3 . For instance, we [Ka1] found algebraic minimal surfaces in \mathbb{R}^3 with totally ramified value number of the Gauss map equals 2.5 (By an algebraic minimal surface, we mean a

2000 *Mathematics Subject Classification.* Primary 53A10; Secondary 30D35, 53A35, 53C42.

Key words and phrases. hyperbolic Gauss map, flat fronts, totally ramified value number.

Partly supported by Global COE program (Kyushu university) “Education and Research Hub for Mathematics-for-Industry” and the Grants-in-Aid for Young Scientists (B) No. 21740053, Japan Society for the Promotion of Science.

complete minimal surface with finite total curvature). Moreover, the author, Kobayashi and Miyaoka [KKM] gave an effective estimate for the number of exceptional values and the totally ramified value number of the Gauss map for a wider class of complete minimal surfaces that includes algebraic minimal surfaces (this class is called “pseudo-algebraic”). It also provided new proofs of the Fujimoto [Fu] and the Osserman theorems ([Os1], [Os2]) for this class and revealed the geometric meaning behind it. Furthermore, we [Ka3] gave the definition of “pseudo-algebraic” and “algebraic” CMC-1 surfaces in \mathcal{H}^3 and such an estimate for the hyperbolic Gauss map of these surfaces. These estimates correspond to the defect relation in the Nevanlinna theory ([Ko], [NO]).

The purpose of this paper is to study value distribution of the hyperbolic Gauss maps of flat fronts in \mathcal{H}^3 . In Section 1, we recall the definition and some fundamental properties of flat fronts in \mathcal{H}^3 . In particular, we review a construction of complete flat fronts via a given pair of hyperbolic Gauss maps and an Osserman-type inequality for this class. In Section 2, we give an estimate for the totally ramified value number of the hyperbolic Gauss maps of complete flat fronts in \mathcal{H}^3 (Theorem 2.2). This estimate is effective in the sense that the lower bound which we obtain is described in terms of geometric invariants. We remark that this estimate is similar to the ramification estimate for the Gauss maps of complete minimal surfaces in Euclidean 4-space \mathbb{R}^4 ([Fu], [HO], and [Ka2]). Moreover, as a corollary of this estimate, we give the upper bounds of the number of exceptional values of them for some topological cases. Furthermore, we consider the Fujimoto-Hoffman-Osserman problem for this class, that is, the problem of finding the “common” maximal number of the exceptional values of the hyperbolic Gauss maps for complete flat fronts in \mathcal{H}^3 . In Section 3, we investigate examples of complete flat fronts in \mathcal{H}^3 from the view point of value distribution of the hyperbolic Gauss maps and give some new examples of complete flat fronts in \mathcal{H}^3 .

The author thanks Professors Ryoichi Kobayashi, Masatoshi Kokubu, Pablo Mira, Reiko Miyaoka, Wayne Rossman, Masaaki Umehara and Kotaro Yamada for their useful advice.

1. PRELIMINARIES

In this section, we briefly recall the definition and some basic facts on flat fronts in \mathcal{H}^3 . For details, we refer the reader to [GMM], [KRUY1], [KRUY2], [KUY1] and [KUY2].

Let \mathbb{R}_1^4 be the Lorentz-Minkowski 4-space with the Lorentz metric

$$(1.1) \quad \langle (x_0, x_1, x_2, x_3), (y_0, y_1, y_2, y_3) \rangle = -x_0y_0 + x_1y_1 + x_2y_2 + x_3y_3 .$$

Then the hyperbolic 3-space is

$$(1.2) \quad \mathcal{H}^3 = \{(x_0, x_1, x_2, x_3) \in \mathbb{R}_1^4 \mid -(x_0)^2 + (x_1)^2 + (x_2)^2 + (x_3)^2 = -1, x_0 > 0\}$$

with the induced metric from \mathbb{R}_1^4 , which is a simply connected Riemannian 3-manifold with constant sectional curvature -1 . We identify \mathbb{R}_1^4 with the set of 2×2 Hermitian matrices $\text{Herm}(2) = \{X^* = X\}$ ($X^* := {}^t\bar{X}$) by

$$(1.3) \quad (x_0, x_1, x_2, x_3) \longleftrightarrow \begin{pmatrix} x_0 + x_3 & x_1 + ix_2 \\ x_1 - ix_2 & x_0 - x_3 \end{pmatrix},$$

where $i = \sqrt{-1}$. In this identification, \mathcal{H}^3 is represented as

$$(1.4) \quad \mathcal{H}^3 = \{aa^* \mid a \in SL(2, \mathbb{C})\}$$

with the metric

$$\langle X, Y \rangle = -\frac{1}{2} \text{trace}(X\tilde{Y}), \quad \langle X, X \rangle = -\det(X),$$

where \tilde{Y} is the cofactor matrix of Y . The complex Lie group $PSL(2, \mathbb{C}) := SL(2, \mathbb{C})/\{\pm \text{id}\}$ acts isometrically on \mathcal{H}^3 by

$$(1.5) \quad \mathcal{H}^3 \ni X \longmapsto aXa^*,$$

where $a \in PSL(2, \mathbb{C})$.

Let M be an oriented 2-manifold. A smooth map $f: M \rightarrow \mathcal{H}^3$ is called a *front* if there exists a Legendrian immersion

$$L_f: M \rightarrow T_1^*\mathcal{H}^3$$

into the unit cotangent bundle of \mathcal{H}^3 whose projection is f . Identifying $T_1^*\mathcal{H}^3$ with the unit tangent bundle $T_1\mathcal{H}^3$, we can write $L_f = (f, \nu)$, where $\nu(p)$ is a unit vector in $T_{f(p)}\mathcal{H}^3$ such that $\langle df(p), \nu(p) \rangle = 0$ for each $p \in M$. We call ν a *unit normal vector field* of the front f . By the definition, a front may have singular points (i.e., points of rank $(df) < 2$). A point which is not singular is said to be *regular*, where the first fundamental form is positive definite.

The *parallel front* f_t of a front f at distance t is given by $f_t(p) = \text{Exp}_{f(p)}(t\nu(p))$, where “Exp” denotes the exponential map of \mathcal{H}^3 . In the model for \mathcal{H}^3 as in (1.2), we can write

$$(1.6) \quad f_t = (\cosh t)f + (\sinh t)\nu, \quad \nu_t = (\cosh t)\nu + (\sinh t)f,$$

where ν_t is the unit normal vector field of f_t .

Based on the fact that any parallel surface of a flat surface is also flat at regular points, we define flat fronts as follows; A front $f: M \rightarrow \mathcal{H}^3$ is called a *flat front* if, for each $p \in M$, there exists a real number $t \in \mathbb{R}$ such that the parallel front f_t is a flat immersion at p . By definition, $\{f_t\}$ forms a family of flat fronts. We remark that an equivalent definition of a flat front is that the Gaussian curvature of f vanishes at all regular points. However, there exists the case where this definition is not suitable. For detail, see [KUY2, Remark 2.2].

We assume that f is flat. Then there exists a (unique) complex structure on M and a holomorphic Legendrian immersion

$$(1.7) \quad \mathcal{E}_f: \widetilde{M} \rightarrow SL(2, \mathbb{C})$$

such that f and L_f are projections of \mathcal{E}_f , where \widetilde{M} is the universal covering of M . Here, holomorphic Legendrian map means that $\mathcal{E}_f^{-1}d\mathcal{E}_f$ is off-diagonal (see [GMM], [KUY1], [KUY2]). We call \mathcal{E}_f the *holomorphic Legendrian lift* of f . The map f and its unit normal vector field ν are

$$(1.8) \quad f = \mathcal{E}_f \mathcal{E}_f^*, \quad \nu = \mathcal{E}_f e_3 \mathcal{E}_f^*, \quad e_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

If we set

$$(1.9) \quad \mathcal{E}_f^{-1}d\mathcal{E}_f = \begin{pmatrix} 0 & \theta \\ \omega & 0 \end{pmatrix},$$

the first and second fundamental forms $ds^2 = \langle df, df \rangle$ and $dh^2 = -\langle df, d\nu \rangle$ are given by

$$(1.10) \quad \begin{aligned} ds^2 &= |\omega + \bar{\theta}|^2 = Q + \bar{Q} + (|\omega|^2 + |\theta|^2), & Q &= \omega\theta \\ dh^2 &= |\theta|^2 - |\omega|^2 \end{aligned}$$

for holomorphic 1-forms ω and θ on \widetilde{M} , with $|\omega|^2$ and $|\theta|^2$ on M itself. We call ω and θ the *canonical forms* of f . The holomorphic 2-differential Q appearing in the $(2, 0)$ -part of ds^2 is defined on M , and is called the *Hopf differential* of f . By definition, the umbilic points of f equal the zeros of Q . Defining a meromorphic function on \widetilde{M} by

$$(1.11) \quad \rho = \frac{\theta}{\omega},$$

then $|\rho|: M \rightarrow [0, +\infty]$ is well-defined on M , and $p \in M$ is a singular point if and only if $|\rho(p)| = 1$.

Note that the $(1, 1)$ -part of the first fundamental form

$$(1.12) \quad ds_{1,1}^2 = |\omega|^2 + |\theta|^2$$

is positive definite on M because it is the pull-back of the canonical Hermitian metric of $SL(2, \mathbb{C})$. Moreover, $2ds_{1,1}^2$ coincides with the pull-back of the Sasakian metric on $T_1^*\mathcal{H}^3$ by the Legendrian lift L_f of f (which is the sum of the first and third fundamental forms in this case, see [KUY2, Section 2] for detail). The complex structure on M is compatible with the conformal metric $ds_{1,1}^2$. Note that any flat front is orientable ([KRUY1, Theorem B]). In this paper, for each flat front $f: M \rightarrow \mathcal{H}^3$, we always regard M as a Riemann surface with this complex structure.

The two *hyperbolic Gauss maps* are defined by

$$(1.13) \quad G = \frac{E_{11}}{E_{21}}, \quad G_* = \frac{E_{12}}{E_{22}}, \quad \text{where } \mathcal{E}_f = (E_{ij}).$$

By identifying the ideal boundary \mathbb{S}_∞^2 of \mathcal{H}^3 with the Riemann sphere $\mathbb{C} \cup \{\infty\}$, the geometric meaning of G and G_* is given as follows ([KRUY2, Appendix A], [Ro]): The hyperbolic Gauss maps G and G_* send each point $p \in M$ to the points $G(p)$ and $G_*(p)$ in \mathbb{S}_∞^2 reached by the two oppositely-oriented normal geodesics of \mathcal{H}^3 that start at $f(p)$. In particular, G and G_* are meromorphic functions on M and parallel fronts have the same hyperbolic Gauss maps. The transformation $\mathcal{E}_f \mapsto a\mathcal{E}_f$ by $a = (a_{ij})_{i,j=1,2} \in SL(2, \mathbb{C})$ induces the rigid motion $f \mapsto af a^*$ as in (1.5) and the hyperbolic Gauss maps G and G_* change by the Möbius transformation:

$$(1.14) \quad G \mapsto a \star G = \frac{a_{11}G + a_{12}}{a_{21}G + a_{22}}, \quad G_* \mapsto a \star G_* = \frac{a_{11}G_* + a_{12}}{a_{21}G_* + a_{22}}.$$

Here, we remark the interchangeability of the canonical forms and the hyperbolic Gauss maps. The canonical forms (ω, θ) have the $U(1)$ -ambiguity $(\omega, \theta) \mapsto (e^{is}\omega, e^{-is}\theta)$ ($s \in \mathbb{R}$) which corresponds to

$$(1.15) \quad \mathcal{E}_f \longmapsto \mathcal{E}_f \begin{pmatrix} e^{is/2} & 0 \\ 0 & e^{-is/2} \end{pmatrix}$$

For a second ambiguity, defining the *dual* \mathcal{E}_f^{\natural} of \mathcal{E}_f by

$$\mathcal{E}_f^{\natural} = \mathcal{E}_f \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix},$$

then \mathcal{E}_f^{\natural} is also Legendrian with $f = \mathcal{E}_f^{\natural} \mathcal{E}_f^{\natural*}$. The hyperbolic Gauss maps G^{\natural} , G_*^{\natural} and canonical forms ω^{\natural} , θ^{\natural} of \mathcal{E}_f^{\natural} satisfy

$$G^{\natural} = G_*, \quad G_*^{\natural} = G, \quad \omega^{\natural} = \theta, \quad \theta^{\natural} = \omega.$$

Namely, the operation \natural interchanges the roles of ω and θ and also G and G_* .

Kokubu, Umehara and Yamada gave a representation formula of flat fronts in \mathcal{H}^3 for a given pair of hyperbolic Gauss maps (G, G_*) .

Theorem 1.1 ([KUY1], [KUY2]). *Let G and G_* be nonconstant meromorphic functions on a Riemann surface M such that $G(p) \neq G_*(p)$ for all $p \in M$. Assume that*

$$(1.16) \quad \int_{\gamma} \frac{dG}{G - G_*} \in i\mathbb{R}$$

for every cycle $\gamma \in H_1(M, \mathbb{Z})$. Set

$$(1.17) \quad \xi(z) = c \cdot \exp \int_{z_0}^z \frac{dG}{G - G_*}$$

where $z_0 \in M$ is a reference point and $c \in \mathbb{C} \setminus \{0\}$ is an arbitrary constant. Then

$$(1.18) \quad \mathcal{E} = \begin{pmatrix} G/\xi & \xi G_*/(G - G_*) \\ 1/\xi & \xi/(G - G_*) \end{pmatrix}$$

is a nonconstant meromorphic Legendrian curve defined on \widetilde{M} in $PSL(2, \mathbb{C})$ whose hyperbolic Gauss maps are G and G_* , and the projection $f = \mathcal{E}\mathcal{E}^*$ is single-valued on M . Moreover, f is a front if and only if G and G_* have no common branch points. Conversely, any non-totally-umbilic flat front can be constructed this way.

Throughout this paper, we call the condition (1.16) the *period condition*. The canonical forms ω , θ and the Hopf differential Q of f in Theorem 1.1 are given by

$$(1.19) \quad \omega = -\frac{1}{\xi^2}dG, \quad \theta = \frac{\xi^2}{(G - G_*)^2}dG_*, \quad Q = -\frac{dGdG_*}{(G - G_*)^2}.$$

Note that we can obviously show that there does not exist a flat front in \mathcal{H}^3 whose both hyperbolic Gauss maps are constant.

Remark 1.2. Kokubu, Umehara and Yamada obtained another construction of meromorphic Legendrian curves in $PSL(2, \mathbb{C})$. For detail, see [KUY1].

A front $f: M \rightarrow \mathcal{H}^3$ is said to be *complete* if there exists a symmetric 2-tensor T such that $T = 0$ outside a compact set $C \subset M$ and $ds^2 + T$ is a complete metric of M . In other words, the set of singular points of f is compact and each divergent path has infinite length.

Theorem 1.3 ([GMM], [KUY2]). *Let M be an oriented 2-manifold and $f: M \rightarrow \mathcal{H}^3$ a complete flat front. Then M is biholomorphic to $\overline{M}_\gamma \setminus \{p_1, \dots, p_k\}$, where \overline{M}_γ is a closed Riemann surface of genus γ and $p_j \in \overline{M}_\gamma$ ($j = 1, \dots, k$). Moreover, the Hopf differential Q of f can be extended meromorphically to \overline{M}_γ .*

Each puncture point p_j ($j = 1, \dots, k$) is called an *end* of f . Gálvez, Martínez and Milán study complete ends of flat surfaces in \mathcal{H}^3 . The following fact is essentially proved in [GMM].

Lemma 1.4 ([GMM], [KUY2]). *Let p be an end of complete flat front. The following three conditions are equivalent;*

- (1) *The Hopf differential Q has at most a pole of order 2 at p .*
- (2) *One hyperbolic Gauss map G has at most a pole at p .*
- (3) *The other hyperbolic Gauss map G_* has at most a pole at p .*

If an end of a flat front satisfies one of the three conditions above, it is called a *regular end*. An end that is not regular is called an *irregular end*. An end p is said to be *embedded* if there exists a neighborhood U of $p \in \overline{M}_\gamma$ such that the restriction of the front to $U \setminus \{p\}$ is an embedding.

Lemma 1.5 ([KUY2]). *The two hyperbolic Gauss maps take the same value at a regular end of a complete flat front, that is, $G(p) = G_*(p)$ if p is a regular end.*

By the lemma above and investigation of embedded regular ends of complete flat fronts, Kokubu, Umehara and Yamada showed the following global properties of complete flat fronts.

Theorem 1.6 ([KUY2], Theorem 3.13). *Let $f: \overline{M}_\gamma \setminus \{p_1, \dots, p_k\} \rightarrow \mathcal{H}^3$ be a complete flat front whose ends are all regular. Then*

$$d + d_* \geq k$$

where d is the degree of G considered as a map on \overline{M}_γ (if G has essential singularities, then we define $d = \infty$) and d_* is the degree of G_* considered as the same way. Furthermore, equality holds if and only if all ends are embedded.

We remark that this inequality is analogue of the Osserman inequality for algebraic minimal surfaces in \mathbb{R}^3 ([Os1], [Os2]).

2. AN EFFECTIVE ESTIMATE FOR THE TOTALLY RAMIFIED VALUE NUMBER OF HYPERBOLIC GAUSS MAPS

We first recall the definition of the totally ramified value number of a meromorphic function on a Riemann surface.

Definition 2.1 (Nevanlinna [Ne]). Let M be a Riemann surface and h a meromorphic function on M . We call $b \in \mathbb{C} \cup \{\infty\}$ a *totally ramified value* of h when at all the inverse image points of b , h branches. We regard exceptional values also as totally ramified values. Let $\{a_1, \dots, a_{r_0}, b_1, \dots, b_{l_0}\} \in \mathbb{C} \cup \{\infty\}$ be the set of all totally ramified values of h , where a_j ($j = 1, \dots, r_0$) are exceptional values. For each a_j , put $\nu_j = \infty$, and for each b_j , define ν_j to be the minimum of the multiplicities of h at points $h^{-1}(b_j)$. Then we have $\nu_j \geq 2$. We call

$$\nu_h = \sum_{a_j, b_j} \left(1 - \frac{1}{\nu_j}\right) = r_0 + \sum_{j=1}^{l_0} \left(1 - \frac{1}{\nu_j}\right)$$

the *totally ramified value number* of h .

We next give an effective estimate for the totally ramified value number of the hyperbolic Gauss maps of complete flat fronts in \mathcal{H}^3 .

Theorem 2.2. *Let $f: \overline{M}_\gamma \setminus \{p_1, \dots, p_k\} \rightarrow \mathcal{H}^3$ be a complete flat front. If two hyperbolic Gauss maps G and G_* are nonconstant and $\nu_G > 2$ and $\nu_{G_*} > 2$, then we have*

$$(2.1) \quad \frac{1}{\nu_G - 2} + \frac{1}{\nu_{G_*} - 2} \geq \frac{k}{2\gamma - 2 + k}.$$

Note that the right side of the inequality (2.1) describes in terms of only topological data of $M = \overline{M}_\gamma \setminus \{p_1, \dots, p_k\}$, that is, no data of the degrees of the hyperbolic Gauss maps.

Proof. If f has an irregular end, then G or G_* has an essential singularity there. By the big Picard theorem, we get $\nu_G \leq 2$ or $\nu_{G_*} \leq 2$. Thus we only consider the case where all ends are regular. Assume that G is nonconstant and omits r_0 values. Let d be the degree of G considered as a map on \overline{M}_γ and n_0 be the sum of branching orders at the inverse image of these exceptional values of G . Then we have

$$(2.2) \quad k \geq dr_0 - n_0.$$

Let b_1, \dots, b_{l_0} be the totally ramified values which are not exceptional values. Let n_r be the sum of branching order at the inverse image of b_i ($i = 1, \dots, l_0$) of G . For each b_i , we denote

$$\nu_i = \min_{G^{-1}(b_i)} \{\text{multiplicity of } G(z) = b_i\},$$

then the number of points in the inverse image $G^{-1}(b_i)$ is less than or equal to d/ν_i . Thus we have

$$(2.3) \quad dl_0 - n_r \leq \sum_{i=1}^{l_0} \frac{d}{\nu_i}.$$

This implies

$$(2.4) \quad l_0 - \sum_{i=1}^{l_0} \frac{1}{\nu_i} \leq \frac{n_r}{d}.$$

Let n_G be the total branching order of G on \overline{M}_γ . Then applying the Riemann-Hurwitz theorem to the meromorphic function G on \overline{M}_γ , we obtain

$$(2.5) \quad n_G = 2(d + \gamma - 1).$$

Thus we get

$$(2.6) \quad \nu_G = r_0 + \sum_{i=1}^{l_0} \left(1 - \frac{1}{\nu_i}\right) \leq \frac{n_0 + k}{d} + \frac{n_r}{d} \leq \frac{n_G + k}{d} \leq 2 + \frac{2\gamma - 2 + k}{d}.$$

Similarly, we get

$$(2.7) \quad \nu_{G_*} \leq 2 + \frac{2\gamma - 2 + k}{d_*}.$$

Here we assume that $\nu_G > 2$ and $\nu_{G_*} > 2$. Then we have

$$(2.8) \quad \frac{1}{\nu_G - 2} \geq \frac{d}{2\gamma - 2 + k}, \quad \frac{1}{\nu_{G_*} - 2} \geq \frac{d_*}{2\gamma - 2 + k}.$$

Combining these inequalities and Theorem 1.6, we deduce

$$(2.9) \quad \frac{1}{\nu_G - 2} + \frac{1}{\nu_{G_*} - 2} \geq \frac{d + d_*}{2\gamma - 2 + k} \geq \frac{k}{2\gamma - 2 + k}.$$

□

As a corollary, we can get the upper bounds of the number of exceptional values of the hyperbolic Gauss maps of complete flat fronts in \mathcal{H}^3 for some topological cases. Here, we denote by D_G and D_{G_*} the number of exceptional values of G and G_* , respectively.

Corollary 2.3. *For complete flat fronts in \mathcal{H}^3 , we have the following:*

- (i) *If $\gamma = 0$, $D_G \geq 4$ and $D_{G_*} \geq 4$, then there does not exist such a front.*
- (ii) *If $\gamma = 1$, $D_G \geq 5$ and $D_{G_*} \geq 5$, then there does not exist such a front.*

Proof. When $\gamma = 0$, $D_G > 2$ and $D_{G_*} > 2$, from the inequality (2.1), we have

$$\frac{1}{D_G - 2} + \frac{1}{D_{G_*} - 2} \geq \frac{k}{k - 2} > 1.$$

On the other hand, if $\gamma = 0$, $D_G \geq 4$ and $D_{G_*} \geq 4$, then it holds that

$$\frac{1}{D_G - 2} + \frac{1}{D_{G_*} - 2} \leq 1.$$

Therefore, if $\gamma = 0$, $D_G \geq 4$ and $D_{G_*} \geq 4$, then both G and G_* are constant. However there does not exist such a front. Hence we obtain (i). In the same way, when $\gamma = 1$, $D_G > 2$ and $D_{G_*} > 2$, we have

$$\frac{1}{D_G - 2} + \frac{1}{D_{G_*} - 2} \geq 1.$$

On the other hand, if $\gamma = 1$, $D_G \geq 5$ and $D_{G_*} \geq 5$, then we get

$$\frac{1}{D_G - 2} + \frac{1}{D_{G_*} - 2} < 1.$$

Therefore we obtain (ii). □

Finally, we consider the Fujimoro-Hoffman-Osserman problem, that is, the problem of finding the common maximal number of the exceptional values of two hyperbolic Gauss maps of complete flat fronts in \mathcal{H}^3 . We remark that the common maximal number of the exceptional values of the Gauss maps g_1 and g_2 of nonflat complete minimal surfaces in \mathbb{R}^4 is “4”, that is, $D_{g_1} = D_{g_2} = 4$ ([Fu], [HO] and [Ka2]). By Corollary 2.3, if $\gamma = 0$, then the common maximal number of exceptional values of two hyperbolic Gauss maps is “3”, that is, $D_G = D_{G_*} = 3$. Moreover, if $\gamma = 1$, then the common maximal number of exceptional values of two hyperbolic Gauss maps is “4”, that is, $D_G = D_{G_*} = 4$. Then we get the necessary conditions for the existence of complete flat fronts whose hyperbolic Gauss maps have the common maximal number of exceptional values.

Corollary 2.4. *Let $f: \overline{M}_\gamma \setminus \{p_1, \dots, p_k\} \rightarrow \mathcal{H}^3$ be a complete flat front.*

- (i) *If $\gamma = 0$ and $D_G = D_{G_*} = 3$, then $k \geq 4$.*
- (ii) *If $\gamma = 1$ and $D_G = D_{G_*} = 4$, then all ends are regular and embedded.*

Proof. When $\gamma = 0$, by the inequality (2.1), we have

$$\frac{1}{D_G - 2} + \frac{1}{D_{G_*} - 2} \geq \frac{k}{k - 2}.$$

Moreover, if $D_G = 3$ and $D_{G_*} = 3$, then we have

$$\frac{1}{D_G - 2} + \frac{1}{D_{G_*} - 2} = 2.$$

Therefore, for this case, we get the following inequality.

$$\frac{k}{k - 2} \leq 2.$$

Thus we obtain (i). Next we prove (ii). When $\gamma = 1$, by (2.9), then we get

$$\frac{1}{D_G - 2} + \frac{1}{D_{G_*} - 2} \geq \frac{d + d_*}{k} \geq 1.$$

Moreover, if $D_G = 4$ and $D_{G_*} = 4$, then we have

$$\frac{1}{D_G - 2} + \frac{1}{D_{G_*} - 2} = 1.$$

Therefore, we can get the following equality.

$$d + d_* = k$$

By the virtue of Theorem 1.6, all ends are regular and embedded for this case. \square

3. EXAMPLES OF COMPLETE FLAT FRONTS FROM THE VIEW POINT OF VALUE DISTRIBUTION OF THE HYPERBOLIC GAUSS MAPS

In the first half of this section, we investigate examples of complete flat fronts in \mathcal{H}^3 from the view point of value distribution of the hyperbolic Gauss maps.

Example 3.1 (Example 4.1 of [KUY2]). We set $\overline{M}_0 = \mathbb{C} \cup \{\infty\}$ and consider a pair (G, G_*) of meromorphic functions on \overline{M}_0 given by $G(z) = z$ and $G_*(z) = \alpha z$, for some constant $\alpha \in \mathbb{R} \setminus \{1\}$. We define M by $M = \overline{M}_0 \setminus \{0\}$ for the case where $\alpha = 0$ and $M = \overline{M}_0 \setminus \{0, \infty\}$ for the case where $\alpha \neq 0$, respectively. By Theorem 1.1, we can construct a flat front $f: M \rightarrow \mathcal{H}^3$ whose hyperbolic Gauss maps are G and G_* . Indeed we can easily see that M and (G, G_*) satisfy the period condition and these data give a Legendrian immersion \mathcal{E}_f of f

$$(3.1) \quad \mathcal{E}_f = \left(\begin{array}{cc} \frac{z^{-\alpha/(1-\alpha)}}{c} & \frac{c\alpha z^{1/(1-\alpha)}}{1-\alpha} \\ \frac{z^{-1/(1-\alpha)}}{c} & \frac{c z^{\alpha/(1-\alpha)}}{1-\alpha} \end{array} \right) \text{ for some constant } c.$$

Moreover, the canonical forms ω and θ and the Hopf differential Q of f is given by

$$\omega = -\frac{1}{c^2} z^{-2/(1-\alpha)} dz, \quad \theta = \frac{c^2 \alpha}{(1-\alpha)^2} z^{2\alpha/(1-\alpha)} dz, \quad Q = -\frac{\alpha}{(1-\alpha)^2} z^{-2} dz^2.$$

Thus f is complete. For the case where $\alpha \neq 0$, the hyperbolic Gauss maps G and G_* of f have the same exceptional values 0 and ∞ , that is, $D_G = D_{G_*} = 2$. For the case where $\alpha = 0$, G has one exceptional value 0 and G_* is constant. Note that f is a horosphere if $\alpha = 0$.

We remark that horospheres can be characterized by the hyperbolic Gauss maps as follows:

Theorem 3.2 (Proposition 4.2 of [KUY2]). *If one of two hyperbolic Gauss maps of a complete flat front in \mathcal{H}^3 is constant, then it is a horosphere.*

To our regret, we do not find a complete flat front whose two hyperbolic Gauss maps have the common maximal number of exceptional values for both $\gamma = 0$ and $\gamma = 1$. However, there exists a complete flat front of genus 0 with $(D_G, D_{G_*}) = (3, 2)$.

Example 3.3 (Theorem 4.4 (iii) of [KUY2]). There exists a complete flat front $f: M = \mathbb{C} \setminus \{0, 1\} \rightarrow \mathcal{H}^3$ whose hyperbolic Gauss maps are

$$(3.2) \quad (G, G_*) = (z, z^2).$$

In particular, $D_G = 3$ and $D_{G_*} = 2$ and all ends are regular and embedded.

In the latter half of this section, we give some new examples of complete flat fronts in \mathcal{H}^3 . We first give an example of genus 0 with 4 regular embedded ends and $(\nu_G, \nu_{G_*}) = (3, 2)$.

Proposition 3.4. *There exists a complete flat front $f: M = \mathbb{C} \setminus \{0, \pm 1\} \rightarrow \mathcal{H}^3$ whose hyperbolic Gauss maps are*

$$(3.3) \quad (G, G_*) = \left(z^2, \frac{z(z+a)}{az+1} \right) \quad (a \in \mathbb{R} \setminus \{0, \pm 1\}).$$

In particular, $\nu_G = 3$ and $\nu_{G_*} = 2$ and all ends are regular and embedded.

Proof. By straightforward computation, we see that

$$\frac{dG}{G - G_*} = \frac{2(az+1)}{a(z+1)(z-1)} dz,$$

and it is holomorphic at $z = 0$ and has poles only at $z = \pm 1, \infty$. All of them are simple poles, with residues $(1+a)/a$, $(a-1)/a$, -2 , respectively. By the condition of a , these residues are real. Thus these data satisfy the period condition. Moreover, we can clearly see that G and G_* take the same values at $z = 0, \pm 1, \infty$ and have no common branch points. By Theorem 1.1, we can construct a flat front $f: M \rightarrow \mathcal{H}^3$ whose hyperbolic Gauss maps are (3.3).

On the other hand, the canonical forms ω and θ of f are given by

$$\omega = -\frac{2}{c^2} z(z+1)^{-2(a-1)/a} (z-1)^{-2(a+1)/a} dz, \quad \theta = \frac{c^2}{a^2} z^{-2} (z+1)^{-2/a} (z-1)^{2/a} (az^2 + 2z + a) dz.$$

Furthermore, the Hopf differential of f is given by

$$Q = -\frac{2(az^2 + 2z + a)}{a^2z(z+1)^2(z-1)^2}dz^2.$$

Thus Q has poles only at $z = 0, \pm 1, \infty$ with

$$(\text{ord}_0Q, \text{ord}_1Q, \text{ord}_{-1}Q, \text{ord}_\infty Q) = (-1, -2, -2, -1).$$

Hence f is complete.

All ends of f are regular and embedded because f satisfies the equality of Theorem 1.6. One hyperbolic Gauss map G has three exceptional values $0, 1, \infty$. The other hyperbolic Gauss map G_* has one exceptional value 0 and two totally ramified values. Therefore, we show that $\nu_G = 1 + 1 + 1 = 3$ and $\nu_{G_*} = 1 + (1/2) + (1/2) = 2$. \square

Remark 3.5. By virtue of Theorem 1.6, if a complete flat front has 4 embedded regular ends, then $(d, d_*) = (1, 3), (2, 2)$ or $(3, 1)$. By Example 4.5 of [KUY2] and Proposition 3.4, we see that there exists one example at the least for all cases of $d + d_* = 4$.

We next give an example of complete flat front of genus 0 with $(d, d_*) = (3, 2)$ and 5 regular embedded ends.

Proposition 3.6. *There exists a complete flat front $f: M = \mathbb{C} \setminus \{0, 1, -2, -3/2\} \rightarrow \mathcal{H}^3$ whose hyperbolic Gauss maps are*

$$(3.4) \quad (G, G_*) = \left(z^3, \frac{z(z+6)}{2z+5} \right).$$

In particular, $\nu_G = 3$ and $\nu_{G_} = 1$ and all ends are regular and embedded.*

Proof. By straightforward computation, we see that

$$\frac{dG}{G - G_*} = \frac{3z(2z+5)}{(z-1)(z+2)(2z+3)}dz,$$

and it is holomorphic at $z = 0$ and has poles only at $z = 1, -2, -3/2, \infty$. All of them are simple poles, with residues $7/5, -2, 18/5, -3$, respectively. Thus these data satisfy the period condition. Moreover, we can easily see that G and G_* take the same values at $z = 0, 1, -2, -3/2, \infty$ and have no common branch points. By Theorem 1.1, we can construct a flat front $f: M \rightarrow \mathcal{H}^3$ whose hyperbolic Gauss maps are (3.4).

On the other hand, the canonical forms ω and θ of f are given by

$$\begin{aligned} \omega &= -\frac{3}{c^2}z^2(z-1)^{-14/5}(z+2)^4(2z+3)^{-36/5}dz, \\ \theta &= 2c^2z^{-2}(z-1)^{4/5}(z+2)^{-6}(2z+3)^{26/5}(z^2+6z+15)dz. \end{aligned}$$

Furthermore, the Hopf differential of f is given by

$$Q = -\frac{6(z^2+6z+15)}{(z-1)^2(z+2)^2(2z+3)^2}dz^2.$$

Thus Q has poles only at $z = 1, -2, -3/2$ with

$$(\text{ord}_1 Q, \text{ord}_{-2} Q, \text{ord}_{-3/2} Q) = (-2, -2, -2).$$

Hence f is complete.

All ends of f are regular and embedded because f satisfies the equality of Theorem 1.6. One hyperbolic Gauss map G has two exceptional values $0, \infty$. The other hyperbolic Gauss map G_* has two totally ramified value. Therefore, we show that $\nu_G = 2$ and $\nu_{G_*} = (1/2) + (1/2) = 1$. \square

Finally, we give an example of complete flat front in \mathcal{H}^3 of genus 1 with 5 regular ends. Let \overline{M}_1 be the square torus on which the Weierstrass \wp function satisfies

$$(\wp')^2 = 4\wp(\wp^2 - a^2), \quad a = \wp(1/2).$$

Proposition 3.7. *There exists a complete flat front $f: \overline{M}_1 \setminus \{z; \wp(z)(\wp(z)^2 + a^2) = 0\} \rightarrow \mathcal{H}^3$ whose hyperbolic Gauss maps are*

$$(3.5) \quad (G, G_*) = \left(\frac{\wp'}{\wp}, \frac{2(\wp^2 - 3a^2)}{\wp'} \right)$$

and 5 regular ends.

Proof. For this data, a computation gives

$$\frac{dG}{G - G_*} = d \log \wp.$$

This implies that these data satisfy the period condition. Moreover, G and G_* take the same values on $\{z; \wp(z)(\wp(z)^2 + a^2) = 0\}$ and have no common branch points. By Theorem 1.1, we can construct a flat front $f: M \rightarrow \mathcal{H}^3$ whose hyperbolic Gauss maps are (3.5).

The canonical forms ω, θ and the Hopf differential Q of f are given by

$$\omega = -\frac{2(\wp^2 + a^2)}{c^2 \wp^3} dz, \quad \theta = \frac{c^2 \wp^2 (\wp^4 + 6a^2 \wp^2 - 3a^4)}{(\wp^2 + a^2)^2} dz, \quad Q = \frac{2(\wp^4 + 6a^2 \wp^2 - 3a^4)}{\wp(\wp^2 + a^2)} dz^2$$

from which the completeness of the ends $\{z; \wp(z)(\wp(z)^2 + a^2) = 0\}$ follows. Obviously all ends are regular but not embedded because f does not satisfy the equality of Theorem 1.6. Indeed, we clearly see that $d = 2$ and $d_* = 4$ and $6 = d + d_* > k = 5$. \square

Remark 3.8. There exists a complete flat front of genus 1 with $(d, d_*) = (3, 2)$ and 5 regular embedded ends [KUY2, Example 4.6].

REFERENCES

- [Br] R. Bryant, *Surfaces of mean curvature one in hyperbolic space*, Astérisque **154–155** (1987), 321–347.
[Fu] H. Fujimoto, *On the number of exceptional values of the Gauss map of minimal surfaces*, J. Math. Soc. Japan, **40** (1988), 235 – 247.

- [GMM] J. A. Gálvez, A. Martínez, and F. Milán, *Flat surfaces in hyperbolic 3-space*, Math. Ann., **316** (2000), 419–435.
- [HO] D. A. Hoffman and R. Osserman, The geometry of the generalized Gauss map, Mem. Amer. Math. Soc., **28** (1980), no. 236.
- [Ka1] Y. Kawakami, *On the totally ramified value number of the Gauss map of minimal surfaces*, Proc. Japan Acad., Ser. A Math. Sci., **82** (2006), 1–3.
- [Ka2] Y. Kawakami, *The Gauss map of pseudo-algebraic minimal surfaces in \mathbf{R}^4* , Math. Nachr., **282** (2009), no. 2, 211–218.
- [Ka3] Y. Kawakami, *Ramification estimates for the hyperbolic Gauss map*, to appear in Osaka Journal of Mathematics, arXiv:0804.0470.
- [KKM] Y. Kawakami, R. Kobayashi and R. Miyaoka, *The Gauss map of pseudo-algebraic minimal surfaces*, Forum Math., **20** (2008), no. 6, 1055–1069.
- [Ko] R. Kobayashi, *Toward Nevanlinna theory as a geometric model of Diophantine approximation*, Sugaku Expositions **16** (2003), 39–79.
- [KRSUY] M. Kokubu, W. Rossman, K. Saji, M. Umehara and K. Yamada, *Singularities of flat fronts in hyperbolic space*, Pacific J. Math., **221** (2005), 303–351.
- [KRUY1] M. Kokubu, W. Rossman, M. Umehara and K. Yamada, *Flat fronts in hyperbolic 3-space and their caustics*, J. Math. Soc. Japan, **59**, no. 1 (2007), 265–299.
- [KRUY2] M. Kokubu, W. Rossman, M. Umehara and K. Yamada, *Asymptotic behavior of flat surfaces in hyperbolic 3-space*, J. Math. Soc. Japan, **61**, no. 3 (2009), 799–852.
- [KUY1] M. Kokubu, M. Umehara and K. Yamada, *An elementary proof of Small’s formula for null curves in $PSL(2, \mathbb{C})$ and an analogue for Legendrian curves in $PSL(2, \mathbb{C})$* , Osaka J. Math., **40** (3) (2003), 697–715.
- [KUY2] M. Kokubu, M. Umehara and K. Yamada, *Flat fronts in hyperbolic 3-space*, Pacific J. Math., **216** (2004), 149–175.
- [Ne] R. Nevanlinna, *Analytic Function*, Translated from the second German edition by Phillip Emig. Die Grundlehren der mathematischen Wissenschaften, Springer, New York, 1970.
- [NO] J. Noguchi and T. Ochiai, *Geometric Function Theory in Several Complex Variables*, Transl. Math. Monog. **80**, Amer. Math. Soc., 1990.
- [Os1] R. Osserman, *Global properties of minimal surfaces in E^3 and E^n* , Ann. of Math. **80** (1964), 340–364.
- [Os2] R. Osserman, *A survey of minimal surfaces*, second edition, Dover Publications Inc., 1986.
- [Ro] P. Roitman, *Flat surfaces in hyperbolic 3-space as normal surfaces to a congruence of geodesics*, Tôhoku Math. J., **59** (2007), 21–37.
- [SUY] K. Saji, M. Umehara and K. Yamada, *The geometry of fronts*, Ann. of Math., **169** (2009), 491–529.
- [UY] M. Umehara and K. Yamada, *Complete surfaces of constant mean curvature-1 in the hyperbolic 3-space*, Ann. of Math. **137** (1993), 611–638.

FACULTY OF MATHEMATICS,
 KYUSHU UNIVERSITY,
 6-10-1, HAKOZAKI, HIGASHIKU, FUKUOKA-CITY, 812-8581, JAPAN
E-mail address: kawakami@math.kyushu-u.ac.jp

List of MI Preprint Series, Kyushu University

The Global COE Program
Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p -adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials

- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials
- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings

- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric q -functions of the q -Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications
- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain

- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection

MI2009-26 Manabu YOSHIDA
Ramification of local fields and Fontaine's property (Pm)

MI2009-27 Yu KAWAKAMI
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space