The Number of Proofs for a BCK-Formula

Komori, Yuichi

Hirokawa, Sachio
Computing and Communications Center, Kyushu University

<http://hdl.handle.net/2324/1546528>

出版情報 : The Journal of symbolic logic. 58 (2), pp.626-628, 1993-06. Association for Symbolic Logic

バージョン :

権利関係 :

THE NUMBER OF PROOFS FOR A BCK-FORMULA

YUICHI KOMORI AND SACHIO HIROKAWA¹

In this note, we give a necessary and sufficient condition for a BCK-formula to have the unique normal form proof.

We call implicational propositional formulas formulas for short. BCK-formulas are the formulas which are derivable from axioms $B = (a \rightarrow b) \rightarrow (c \rightarrow a) \rightarrow c \rightarrow b$, $C = (a \rightarrow b \rightarrow c) \rightarrow b \rightarrow a \rightarrow c$, and $K = a \rightarrow b \rightarrow a$ by substitution and modus ponens. It is known that the property of being a BCK-formula is decidable (Jaskowski [11, Theorem 6.5], Ben-Yelles [3, Chapter 3, Theorem 3.22], Komori [12, Corollary 6]). The set of BCK-formulas is identical to the set of provable formulas in the natural deduction system with the following two inference rules.

$$\begin{array}{c} [\gamma] \\ \vdots \\ \delta \\ \hline \gamma \rightarrow \delta \end{array} (\rightarrow I) \qquad \frac{\gamma \rightarrow \delta \quad \gamma}{\delta} (\rightarrow E).$$

Here γ occurs at most once in $(\rightarrow I)$. By the formulae-as-types correspondence [10], this set is identical to the set of type-schemes of closed BCK- λ -terms. (See [5].) A BCK- λ -term is a λ -term in which no variable occurs twice. Basic notions concerning the type assignment system can be found [4]. Uniqueness of normal form proofs has been known for balanced formulas. (See [2, 14].) It is related to the coherence theorem in cartesian closed categories. A formula is balanced when no variable occurs more than twice in it. It was shown in [8] that the proofs of balanced formulas are BCK-proofs. Relevantly balanced formulas were defined in [9], and it was proved that such formulas have unique normal form proofs. Balanced formulas are included in the set of relevantly balanced formulas. We show a necessary and sufficient condition for a BCK-formula to have a unique

Received December 11, 1991; revised April 12, 1992.

¹This work was partially supported by a Grant-in-Aid for Encouragement of Young Scientists No. 02740115 of the Ministry of Education.

normal form proof using the following notion of minimality. The notion of BCK-minimality was introduced by Komori [13]. A formula α is called a *trivial substitution instance* of β iff α is a substitution instance of β and β is a substitution instance of α .

DEFINITION 1. A formula is *BCK-minimal* iff it is a BCK-formula and it is not a nontrivial substitution instance of another BCK-formula. A BCK-formula β is a *minimal formula* of α iff β is BCK-minimal and α is a substitution instance of β .

It is clear that a BCK-minimal formula is a principal type-scheme of a closed BCK- λ -term.

We identify two λ -terms when they are α -convertible. Similarly, two types are identified when one is a trivial substitution instance of the other.

LEMMA 1 ([7]). *If two closed BCK- λ -terms in β -normal form have the same principal type, then they are identical.*

LEMMA 2 ([8]). *A BCK-formula is BCK-minimal iff it is a principal type-scheme of a closed BCK- λ -term in $\beta\eta$ -normal form.*

THEOREM 1. *Given a BCK-formula α , the number of closed BCK- λ -terms in $\beta\eta$ -normal form which has type α is identical to the number of minimal formulas of α .*

PROOF. Let α be a BCK-formula. We denote by $\text{proof}(\alpha)$ the set of closed BCK- λ -terms in $\beta\eta$ -normal form which have type α and we denote by $\text{min}(\alpha)$ the set of minimal formulas of α . We define a function from $\text{proof}(\alpha)$ to $\text{min}(\alpha)$ and show that it is surjective and injective. Let $M \in \text{proof}(\alpha)$. Then M has type α . By the principal type-scheme theorem (Theorem 15.26 of [4]), M has a principal type-scheme. We denote it by $\text{pts}(M)$. Since M is in $\beta\eta$ -normal form, $\text{pts}(M)$ is minimal by Lemma 2. So we have $\text{pts}(M) \in \text{min}(\alpha)$. Thus pts is a function from $\text{proof}(\alpha)$ to $\text{min}(\alpha)$. Injectivity of pts is immediate from Lemma 1. To prove the surjectivity, let $\beta \in \text{min}(\alpha)$ and apply Lemma 2 to β . Then there is a closed BCK- λ -term N in $\beta\eta$ -normal form whose principal type-scheme is β . Therefore pts is surjective. $\square$

One consequence of the theorem is that a BCK-formula α has only a finite number of normal form proofs. In fact, we can enumerate all the minimal formulas instead of λ -terms. Given a formula γ , we denote by $s_0(\gamma)$ the set of formulas β such that γ is a substitution instance of β . Since we identify trivial substitution instances, the set $s_0(\gamma)$ is finite. Next we denote by $s(\gamma)$ the set of BCK-formulas in $s_0(\gamma)$. Since BCK-provability is decidable, we can enumerate the elements of $s(\gamma)$ from $s_0(\gamma)$. Finally note that β is BCK-minimal iff $s(\beta) = \{\beta\}$. Therefore we have

$$\text{min}(\alpha) = \{\beta \in s_0(\alpha) \mid s(\beta) = \{\beta\}\}.$$

Thus we can enumerate all the elements of $\text{min}(\alpha)$.

Akama [1] showed that the number of cut-free proof (in sequent calculus) for a BCK-formula is finite.

COROLLARY 1. *A BCK-formula has a unique proof in $\beta\eta$ -normal form iff it has a unique minimal formula.*

REFERENCES

- [1] Y. AKAMA, *The number of proofs and tree automaton*. II, manuscript, Oct. 1991.
- [2] A. A. BABAEV and S. V. SOLOVIEV, A coherence theorem for canonical morphism in cartesian closed categories, *Zapiski nauchnykh Seminarov Leningradskogo Otdeleniya matematicheskogo Instituta imeni V. A. Steklova Akademii Nauk SSSR*, vol. 88 (1979), pp. 3–29.

- [3] C.⁴B. BEN-YELLES, *Type-assignment in the lambda-calculus*, Ph.D. thesis, University College, Swansea, 1979.
- [4] J. R. HINDLEY and J. P. SELDIN, *Introduction to Combinators and Lambda-Calculus*, Cambridge University Press, London, 1986.
- [5] J. R. HINDLEY, *BCK-combinators and linear lambda-terms have types*, *Theoretical Computer Science*, vol. 64 (1989), pp. 97–105.
- [6] ———, *BCK and BCI logics, condensed detachment and the 2-property*, *Notre Dame Journal of Formal Logic*, to appear.
- [7] S. HIROKAWA, *Principal types of BCK-lambda terms*, *Theoretical Computer Science*, (to appear).
- [8] ———, *Balanced formulas, BCK-minimal formulas and their proofs*, *Proceedings of Symposium on Logical Foundations of Computer Science*, (Nerode and Taitslin, editors), Lecture Notes in Computer Science, vol. 620, Springer-Verlag, Berlin, and New York, 1992, pp. 198–208.
- [9] ———, *BCK-formulas having unique proofs*, *Proceedings of category theory and computer science* (Pitt et al., editors), *Lecture Notes in Computer Science*, vol. 530, Springer-Verlag, Berlin and New York, 1991, pp. 106–120.
- [10] W. A. HOWARD, *The formulae-as-types notion of construction*, *To H. B. Curry, Essays on Combinatory Logic, Lambda Calculus and Formalism* (Hindley and Seldin, editors), Academic Press, Cambridge, Massachusetts, 1980, pp. 479–490.
- [11] S. JASKOWSKI, *Über Tautologien, in welchen keine Variable mehr Als zweimal vorkommt*, *Zeitschrift für Mathematische Logik und Grundlagen der Mathematik*, vol. 9 (1963), pp. 219–228.
- [12] Y. KOMORI, *The class of BCC-algebras is not a variety*, *Mathematica Japonica*, vol. 29 (1984), pp. 391–394.
- [13] ———, *BCK algebras and lambda calculus*, *Proceedings of the 10th Symposium on Semigroups, Sakado 1986*, Josai University, Sakado, 1987, pp. 5–11.
- [14] G. E. MINTS, *A simple proof of the coherence theorem for cartesian closed categories*, manuscript, 1982.

DEPARTMENT OF MATHEMATICS,
SHIZUOKA UNIVERSITY,
SHIZUOKA 422, JAPAN

DEPARTMENT OF COMPUTER SCIENCE,
COLLEGE OF GENERAL EDUCATION,
KYUSHU UNIVERSITY,
FUKUOKA 810, JAPAN

E-mail: hirokawa@ec.kyushu-u.ac.jp