Traveling waves bifurcating from plane Poiseuille flow of the compressible Navier- Stokes equation

Kagei, Yoshiyuki
Faculty of Mathematics, Kyushu University

Nishida, Takaaki
Department of Applied Complex System, Kyoto University

<http://hdl.handle.net/2324/1547351>

出版情報 : MI Preprint Series. 2015-9, 2015-10-16. 九州大学大学院数理学研究院
バージョン :
権利関係 :

MI Preprint Series

Mathematics for Industry
Kyushu University

**Traveling waves bifurcating from
plane Poiseuille flow of the
compressible Navier-Stokes
equation**

**Yoshiyuki Kagei
& Takaaki Nishida**

MI 2015-9

(Received October 16, 2015)

Institute of Mathematics for Industry
Graduate School of Mathematics
Kyushu University
Fukuoka, JAPAN

Traveling waves bifurcating from plane Poiseuille flow of the compressible Navier-Stokes equation

Yoshiyuki Kagei¹ and Takaaki Nishida²

¹ Faculty of Mathematics,
Kyushu University,
Nishi-ku, Motooka 744,
Fukuoka 819-0395, Japan

² Department of Applied Complex System,
Kyoto University,
Yoshida Honmachi, Sakyo-ku,
Kyoto, 606-8317, Japan

Abstract

Plane Poiseuille flow in viscous compressible fluid is known to be asymptotically stable if Reynolds number R and Mach number M are sufficiently small. On the other hand, for R and M being not necessarily small, an instability criterion for plane Poiseuille flow is known; and the criterion says that, when R increases, a pair of complex conjugate eigenvalues of the linearized operator cross the imaginary axis. In this paper it is proved that a spatially periodic traveling wave bifurcates from plane Poiseuille flow when the critical eigenvalues cross the imaginary axis.

Mathematics Subject Classification (2000). 35Q30, 76N15.

Keywords. Compressible Navier-Stokes equation, Poiseuille flow, bifurcation, spatially periodic traveling wave.

1 Introduction

This paper is concerned with the bifurcation of traveling waves from plane Poiseuille flow of the compressible Navier-Stokes equation. We consider the following system of equations

$$\partial_t \rho + \operatorname{div}(\rho v) = 0, \tag{1.1}$$

$$\rho(\partial_t v + v \cdot \nabla v) - \mu \Delta v - (\mu + \mu') \nabla \operatorname{div} v + \nabla P(\rho) = \rho g \tag{1.2}$$

in a 2-dimensional infinite layer $\Omega_\ell = \mathbb{R} \times (0, \ell)$:

$$\Omega_\ell = \{x = (x_1, x_2) : x_1 \in \mathbb{R}, 0 < x_2 < \ell\}.$$

Here $\rho = \rho(x, t)$ and $v = {}^\top(v^1(x, t), v^2(x, t))$ denote the density and velocity, respectively, at time $t \geq 0$ and position $x \in \Omega_\ell$; $P = P(\rho)$ is the pressure that is assumed to be a smooth function of ρ satisfying

$$P'(\rho_*) > 0$$

for a given constant $\rho_* > 0$; μ and μ' are the viscosity coefficients that are assumed to be constants and satisfy

$$\mu > 0, \quad \mu + \mu' \geq 0;$$

div , ∇ and Δ denote the usual divergence, gradient and Laplacian with respect to x ; and $\mathbf{g}$ is a given external force. Here and in what follows ${}^\top \cdot$ stands for the transposition.

We assume that the external force $\mathbf{g}$ takes the form

$$\mathbf{g} = g\mathbf{e}_1,$$

where g is a positive constant and $\mathbf{e}_1 = {}^\top(1, 0) \in \mathbb{R}^2$.

The system (1.1)–(1.2) is considered under the boundary condition

$$v|_{x_2=0,\ell} = 0. \tag{1.3}$$

We also require periodicity of ρ and v in x_1 :

$$\rho(x_1 + \frac{2\pi}{\alpha}, x_2) = \rho(x_1, x_2), \quad v(x_1 + \frac{2\pi}{\alpha}, x_2) = v(x_1, x_2), \tag{1.4}$$

where $\alpha > 0$ is a given wave number.

It is easily seen that (1.1)–(1.4) has a stationary solution $\bar{u}_s = {}^\top(\bar{\rho}_s, \bar{v}_s)$ satisfying

$$\bar{\rho}_s = \rho_*, \quad \bar{v}_s = \frac{\rho_* g}{2\mu} x_2 (\ell - x_2) \mathbf{e}_1,$$

that is the so-called plane Poiseuille flow.

The aim of this paper is to show the bifurcation of traveling wave solutions from plane Poiseuille flow.

The function $\bar{v}_s$ also gives a stationary solution representing parallel flow of the incompressible Navier-Stokes equation. It is known that stationary parallel flow of the incompressible Navier-Stokes equation is stable under any initial perturbations in L^2 if the Reynolds number R is sufficiently small. Furthermore, plane Poiseuille flow is stable under sufficiently small initial perturbations if $R < R_c$ for a critical number $R_c \sim 5772$, and unstable if $R > R_c$ ([9]).

As for the compressible case, the stability of parallel flow in the infinite layer Ω_ℓ was studied in [7]; and it was proved that parallel flow is asymptotically stable under perturbations sufficiently small in some Sobolev space over Ω_ℓ if the Reynolds and Mach numbers are sufficiently small. In [8] an instability criterion was established; plane Poiseuille flow of the compressible Navier-Stokes equation (1.1)–(1.4) is linearly unstable if $\alpha \ll 1$ and

$$\frac{1}{280} > \gamma^2, \quad \frac{1}{280} - \gamma^2 > \frac{\nu}{30\gamma^2}(3\nu + \nu'), \tag{1.5}$$

where $\nu = \frac{\mu}{8\rho_*\ell V_0}$, $\nu' = \frac{\mu'}{8\rho_*\ell V_0}$ and $\gamma = \frac{\sqrt{P'(\rho_*)}}{8V_0}$ with $V_0 = \frac{\rho_* g \ell}{8\mu}$ being the maximum velocity of plane Poiseuille flow $\bar{v}_s$. More precisely, the spectrum of the linearized operator $-L$ consists of simple eigenvalues $\lambda_{\alpha k}$ ($|k| = 1, \dots, n_0$) for some $n_0 \in \mathbb{R}$ such that

$$\lambda_{\alpha k} = -\frac{i}{6}(\alpha k) + \kappa_0(\alpha k)^2 + O(|\alpha k|^3) \quad (\alpha k \rightarrow 0).$$

Here κ_0 is the number given by

$$\kappa_0 = \frac{1}{12\nu} \left[\left(\frac{1}{280} - \gamma^2 \right) - \frac{\nu}{30\gamma^2}(3\nu + \nu') \right].$$

As a consequence, if $\alpha \ll 1$ and (1.5) is satisfied, then $\kappa_0 > 0$ and plane Poiseuille flow $\bar{u}_s = {}^\top(\bar{\rho}_s, \bar{v}_s)$ is linearly unstable. Note that the Reynolds number R and Mach number M are given by $R = \frac{1}{16\nu}$ and $M = \frac{1}{8\gamma}$, respectively. Instability condition (1.5) is thus restated as

$$M > \sqrt{\frac{35}{8}} \sim 2.09, \quad \frac{1}{35} - \frac{1}{8M^2} > \frac{M^2}{15R} \left(\frac{3}{R} + \frac{1}{R'} \right), \quad (1.6)$$

where $R' = \frac{1}{16\nu'}$. Therefore, Reynolds and Mach numbers are not small when (1.5) (i.e., (1.6)) is satisfied. For example, if $M = 2.5$, $R = \frac{173}{16} \sim 10.81$ and $\frac{1}{R'} = -\frac{2}{3R}$ (i.e., $\nu' = -\frac{2\nu}{3}$), then instability condition (1.6) (i.e., (1.5)) is satisfied.¹

When the instability described above occurs, there seems to appear the Hopf bifurcation. In fact, if γ^2 is fixed so that $\frac{1}{280} - \gamma^2 > 0$, one can find the value $\nu_1 > 0$ such that $\kappa_0 < 0$ for $\nu = \nu_1$. When ν is decreased from ν_1 , complex conjugate eigenvalues $\lambda_{\pm\alpha}$ cross the imaginary axis at some $\nu = \nu_0$. We will show that there are traveling wave solutions, which are periodic in x_1 and t , bifurcating from plane Poiseuille flow for $\nu \sim \nu_0$, provided that

$$\sigma(-L) \cap \{\lambda; \operatorname{Re} \lambda = 0\} = \{\lambda_\alpha, \lambda_{-\alpha}\} \text{ at } \nu = \nu_0. \quad (1.7)$$

Since Iooss and Padula ([6]) proved that $\sigma(-L) \cap \{\lambda; \operatorname{Re} \lambda > -c\}$ consists of finite number of eigenvalues with finite multiplicities for some constant $c > 0$, it seems very unlikely that the assumption (1.7) is not satisfied for all $\alpha \ll 1$. We also note that we construct bifurcating solutions from Poiseuille flow when ν and γ are small, which implies that Poiseuille flow is large, in other words, we show the bifurcation from large stationary solution.

The bifurcation problem for compressible fluid was firstly treated by Nishida-Padula-Teramoto [11] (cf., [10]); and the existence of the bifurcating convection solutions was proved for thermal convection problem. The main difficulty in the proof of the bifurcation arises from the convection term $v \cdot \nabla \rho$ in (1.1) which may cause the derivative-loss, in other words, it is not Frechét differentiable in a standard setting. In [11], the effective viscous flux was used to overcome this difficulty

¹The definition of M in [8] should be corrected as the one defined in this paper; in [8], M is defined as $M = \sqrt{P'(\rho_*)}/V_0$; and, in [8, Remark 3.2], the value $M = 8/\gamma = 160$ should be corrected as $M = 1/(8\gamma) = 2.5$ as in the example given here.

and establish the necessary estimates for the proof of the bifurcation of stationary convective patterns. (Cf., [1, 5].) In this paper we will not use the effective viscous flux but employ the iterative method in which the convection term $v \cdot \nabla \rho$ in (1.1) is regarded as a part of the principal part as in the proof of the local solvability of the time evolution problem. The method of this paper will be widely applicable to the bifurcation problem for certain classes of quasilinear hyperbolic-parabolic systems.

To prove the existence of bifurcating traveling waves, we rewrite the time evolution problem to a stationary problem in a moving coordinates. We then decompose the stationary problem into the null space of the linearized operator and its complementary subspace. One of the points of the proof is to establish the solvability in the complementary subspace, for which we apply the Matsumura-Nishida energy method [12] and the results on the resolvent problem for transport equation by Heywood and Padula [4] for a linear system which includes the convective term $v \cdot \nabla \rho$ as in (1.1) with a given velocity v .

This paper is organized as follows. In section 2 we derive a non-dimensional form of system (1.1)–(1.2) and rewrite it into the system of equations for the perturbation. We also introduce notation used in this paper. In section 3 we state the instability result of Poiseuille flow obtained in [8], and in section 4, we state the main result of this paper on the existence of bifurcating traveling waves. Sections 5–8 are devoted to the proof of the main result. In section 5 we first formulate the problem. We then rewrite the time evolution problem to a stationary problem in a moving coordinates, and we give a proof of the main result. In section 6 we prove the solvability in the complementary subspace. Section 7 is devoted to a proof of a periodic version of Bogovskii’s lemma. In section 8 we present a proof of the solvability in the null space of the linearized operator.

2 Preliminaries

In this section we first derive a non-dimensional form of system (1.1)–(1.2) and then give the system of equations for the perturbation. In the end of this section we introduce notations used in this paper.

2.1 Non-dimensionalization

We introduce the following non-dimensional variables:

$$x = \ell \tilde{x}, \quad t = \frac{\ell}{V} \tilde{t}, \quad v = V \tilde{v}, \quad \rho = \rho_* \tilde{\rho}, \quad P = \rho_* V^2 \tilde{p}$$

with

$$V = \frac{\rho_* g \ell^2}{\mu}.$$

Under this transformation, (1.1) and (1.2) on Ω_ℓ are written, by omitting tildes, as

$$\partial_t \rho + \operatorname{div}(\rho v) = 0, \tag{2.1}$$

$$\rho(\partial_t v + v \cdot \nabla v) - \nu \Delta v - (\nu + \nu') \nabla \operatorname{div} v + \nabla p(\rho) = \nu \rho \mathbf{e}_1 \quad (2.2)$$

on the infinite layer $\Omega = \Omega_1$:

$$\Omega = \{x = (x_1, x_2) : x_1 \in \mathbb{R}, 0 < x_2 < 1\}.$$

Here and in what follows we denote $\mathbf{e}_1 = {}^\top(1, 0) \in \mathbb{R}^2$; ν and ν' are the non-dimensional parameters given by

$$\nu = \frac{\mu}{\rho_* \ell V}, \quad \nu' = \frac{\mu'}{\rho_* \ell V}.$$

The assumption $P'(\rho_*) > 0$ is restated as

$$p'(1) > 0.$$

To derive (2.2) we have used the relation $\frac{\ell g}{V^2} = \nu$.

We will show the existence of traveling wave solutions of (2.1)–(2.2) bifurcating from Poiseuille flow. Due to the above non-dimensionalization, the Poiseuille flow is transformed to

$$u_s = {}^\top(\rho_s, v_s),$$

where

$$\rho_s = 1, \quad v_s = {}^\top(v_s^1(x_2), 0), \quad v_s^1(x_2) = \frac{1}{2}(-x_2^2 + x_2).$$

We next derive the system of equations for the perturbation. We substitute $u(t) = {}^\top(\phi(t), w(t)) \equiv {}^\top(\gamma^2(\rho(t) - \rho_s), v(t) - v_s)$ into (2.1) and (2.2), where γ is the non-dimensional number given by

$$\gamma = \sqrt{p'(1)} = \frac{\sqrt{P'(\rho_*)}}{V}.$$

Noting that $\rho_s = 1$, $v_s = {}^\top(v_s^1(x_2), 0)$ and $-\Delta v_s = \mathbf{e}_1$, we obtain the following system of equations

$$\partial_t \phi + v_s^1 \partial_{x_1} \phi + \gamma^2 \operatorname{div} w = f^0, \quad (2.3)$$

$$\partial_t w - \nu \Delta w - \tilde{\nu} \nabla \operatorname{div} w + \nabla \phi - \frac{\nu}{\gamma^2} \phi \mathbf{e}_1 + v_s^1 \partial_{x_1} w + (\partial_{x_2} v_s^1) w^3 \mathbf{e}_1 = f. \quad (2.4)$$

Here $\tilde{\nu} = \nu + \nu'$; and f^0 and $f = {}^\top(f^1, f^2)$ denote the nonlinearities:

$$f^0 = -\operatorname{div}(\phi w),$$

$$f = -w \cdot \nabla w - \frac{\phi}{\gamma^2 + \phi} \left(\nu \Delta w + \frac{\nu}{\gamma^2} \phi \mathbf{e}_1 + \tilde{\nu} \nabla \operatorname{div} w \right) + P^{(1)}(\phi) \phi \nabla \phi$$

where

$$P^{(1)}(\phi) = \frac{1}{\gamma^2 + \phi} \left(1 - \frac{1}{\gamma^2} \int_0^1 P''(1 + \theta \gamma^{-2} \phi) d\theta \right).$$

We consider (2.3)–(2.4) under the boundary conditions

$$w|_{x_2=0,1} = 0, \quad \phi, w: \frac{2\pi}{\alpha}\text{-periodic in } x_1, \quad (2.5)$$

and the initial condition

$$u|_{t=0} = u_0 = {}^\top(\phi_0, w_0). \quad (2.6)$$

Here α is a given positive number.

2.2 Notation

We introduce some notations used in this paper. For given $\alpha > 0$, we denote the basic period cell by

$$\mathcal{P}_\alpha = \left[-\frac{\pi}{\alpha}, \frac{\pi}{\alpha}\right).$$

We set

$$\Omega_\alpha = \mathcal{P}_\alpha \times (0, 1).$$

We denote by $C_{per}^\infty(\overline{\Omega}_\alpha)$ the space of restrictions to $\overline{\Omega}_\alpha$ of functions in $C^\infty(\overline{\Omega})$ which are $\mathcal{P}_\alpha$ -periodic in x_1 . We also denote by $C_{per,0}^\infty(\Omega_\alpha)$ the space of restrictions to $\overline{\Omega}_\alpha$ of functions in $C^\infty(\Omega)$ which are $\mathcal{P}_\alpha$ -periodic in x_1 and vanish near $x_2 = 0, 1$.

We set

$$L_{per}^2(\Omega_\alpha) = \text{the } L^2(\Omega_\alpha)\text{-closure of } C_{per,0}^\infty(\Omega_\alpha),$$

$$H_{per}^k(\Omega_\alpha) = \text{the } H^k(\Omega_\alpha)\text{-closure of } C_{per}^\infty(\overline{\Omega}_\alpha),$$

$$H_{per,0}^1(\Omega_\alpha) = \text{the } H^1(\Omega_\alpha)\text{-closure of } C_{per,0}^\infty(\Omega_\alpha).$$

We note that if $f \in H_{per,0}^1(\Omega_\alpha)$, then $f|_{x_1=-\pi/\alpha} = f|_{x_1=\pi/\alpha}$ and $f|_{x_2=0,1} = 0$. $H_{per}^{-1}(\Omega_\alpha)$ stands for the dual space of $H_{per,0}^1(\Omega_\alpha)$. The inner product of $f_j \in L_{per}^2(\Omega_\alpha)$ ($j = 1, 2$) is denoted by

$$(f_1, f_2) = \int_{\Omega_\alpha} f_1(x) \overline{f_2(x)} dx,$$

where $\bar{z}$ denotes the complex conjugate of z .

The mean value of a function $\phi(x)$ over Ω_α is denoted by $\langle \phi \rangle$:

$$\langle \phi \rangle = \frac{1}{|\Omega_\alpha|} \int_{\Omega_\alpha} \phi(x) dx.$$

The set of all $\phi \in L_{per}^2(\Omega_\alpha)$ with $\langle \phi \rangle = 0$ is denoted by $L_{per,*}^2(\Omega_\alpha)$, i.e.,

$$L_{per,*}^2(\Omega_\alpha) = \{\phi \in L_{per}^2(\Omega_\alpha) : \langle \phi \rangle = 0\}.$$

Furthermore, we set

$$H_{per,*}^k(\Omega_\alpha) = H_{per}^k(\Omega_\alpha) \cap L_{per,*}^2(\Omega_\alpha).$$

For simplicity the set of all vector fields whose components are in $L_{per}^2(\Omega_\alpha)$ (resp. $H_{per,0}^1(\Omega_\alpha)$, $H_{per}^k(\Omega_\alpha)$) is also denoted by $L_{per}^2(\Omega_\alpha)$ (resp. $H_{per,0}^1(\Omega_\alpha)$, $H_{per}^k(\Omega_\alpha)$) if no confusion will occur.

We also use notation $L_{per}^2(\Omega_\alpha)$ for the set of all $u = {}^\top(\phi, w)$ with $\phi \in L_{per}^2(\Omega_\alpha)$ and $w = {}^\top(w^1, w^2) \in L_{per}^2(\Omega_\alpha)$ if no confusion will occur. The inner product of $u_j = {}^\top(\phi_j, w_j) \in L_{per}^2(\Omega_\alpha)$ ($j = 1, 2$) is defined by

$$\langle u_1, u_2 \rangle = \frac{1}{\gamma^2} \int_{\Omega_\alpha} \phi_1(x) \overline{\phi_2(x)} dx + \int_{\Omega_\alpha} w_1(x) \cdot \overline{w_2(x)} dx.$$

In what follows we abbreviate Ω_α in $L_{per}^2(\Omega_\alpha)$, $H_{per}^k(\Omega_\alpha)$, $\dots$, and etc., and write them as L_{per}^2 , H_{per}^k , $\dots$, and etc.

We denote by $L^2(0, 1)$ the usual L^2 space on $(0, 1)$ with norm $|\cdot|_{L^2}$, and, likewise, by $H^k(0, 1)$ the k th order L^2 -Sobolev space on $(0, 1)$ with norm $|\cdot|_{H^k}$. The H^1 -closure of $C_0^\infty(0, 1)$ is denoted by $H_0^1(0, 1)$. As in the case of functions on Ω_α , function spaces of vector fields $w = {}^\top(w^1, w^2)$ and, also, those of $u = {}^\top(\phi, w)$, are simply denoted by $L^2(0, 1)$, $H_0^1(0, 1)$, and so on. We define an inner product $\langle\langle u_1, u_2 \rangle\rangle$ of $u_j = {}^\top(\phi_j, w_j) \in L^2(0, 1)$ ($j = 1, 2$), by

$$\langle\langle u_1, u_2 \rangle\rangle = \frac{1}{\gamma^2} \int_0^1 \phi_1(x_2) \overline{\phi_2(x_2)} dx_2 + \int_0^1 w_1(x_2) \cdot \overline{w_2(x_2)} dx_2.$$

We denote the resolvent set of a closed operator A by $\rho(A)$ and the spectrum of A by $\sigma(A)$. The null space and the range of A are denoted by $N(A)$ and $R(A)$, respectively.

3 Instability of Poiseuille flow

In this section we consider the instability of Poiseuille flow.

Let us consider the linearized problem

$$\partial_t \phi + v_s^1 \partial_{x_1} \phi + \gamma^2 \operatorname{div} w = 0, \quad (3.1)$$

$$\partial_t w - \nu \Delta w - \tilde{\nu} \nabla \operatorname{div} w + \nabla \phi - \frac{\nu}{\gamma^2} \phi \mathbf{e}_1 + v_s^1 \partial_{x_1} w + (\partial_{x_2} v_s^1) w^2 \mathbf{e}_1 = 0, \quad (3.2)$$

$$w|_{x_2=0,1} = 0, \quad \phi, w : \frac{2\pi}{\alpha}\text{-periodic in } x_1, \quad (3.3)$$

$$u|_{t=0} = u_0 = {}^\top(\phi_0, w_0). \quad (3.4)$$

We set

$$X = L_{per,*}^2 \times (L_{per}^2)^2.$$

We define the operator L on X by

$$D(L) = \{u = {}^\top(\phi, w) \in X; w \in (H_{per,0}^1)^2, Lu \in X\},$$

$$L = \begin{pmatrix} v_s^1 \partial_{x_1} & \gamma^2 \operatorname{div} \\ \nabla & -\nu \Delta - \tilde{\nu} \nabla \operatorname{div} \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ -\frac{\nu}{\gamma^2} \mathbf{e}_1 & v_s^1 \partial_{x_1} + (\partial_{x_2} v_s^1) \mathbf{e}_1 {}^\top \mathbf{e}_2 \end{pmatrix}.$$

Recall that $\tilde{\nu} = \nu + \nu' \geq 0$. As in [6] one can show that $-L$ generates a C_0 -semigroup in X .

We state an instability criterion for Poiseuille flow.

Theorem 3.1. ([8]) *There exist constants $r_0 > 0$ and $\eta_0 > 0$ such that if $\alpha \leq r_0$, then*

$$\sigma(-L) \cap \{\lambda \in \mathbb{C} : |\lambda| \leq \eta_0\} = \{\lambda_{\alpha k} : |k| = 1, \dots, n_0\}$$

for some $n_0 \in \mathbb{N}$, where $\lambda_{\alpha k}$ are simple eigenvalues of $-L$ that satisfies

$$\lambda_{\alpha k} = -\frac{i}{6}(\alpha k) + \kappa_0(\alpha k)^2 + O(|\alpha k|^3)$$

as $\alpha k \rightarrow 0$. Here κ_0 is the number given by

$$\kappa_0 = \frac{1}{12\nu} \left[\left(\frac{1}{280} - \gamma^2 \right) - \frac{\nu}{30\gamma^2} (3\nu + \nu') \right].$$

As a consequence, if $\gamma^2 < \frac{1}{280}$ and $\nu(3\nu + \nu') < 30\gamma^2 \left(\frac{1}{280} - \gamma^2 \right)$, then $\kappa_0 > 0$ and plane Poiseuille flow $u_s = {}^T(\phi_s, v_s)$ is linearly unstable.

Remark 3.2. The eigenspace for $\lambda_{\alpha k}$ is spanned by a function of the form $u(x_2)e^{i\alpha k x_1}$ where $u(x_2)$ is an eigenfunction for $\lambda_{\alpha k}$ of $-L_{\eta,k}$. Here $L_{\eta,k}$ is an operator appearing in (5.2) below. See [8, Sections 4–6].

4 Traveling wave solutions

In this section we state the result on the existence of traveling wave solutions bifurcating from the Poiseuille flow when it becomes unstable as in Theorem 3.1.

We fix γ such that $\frac{1}{280} - \gamma^2 > 0$. We will take ν as a bifurcation parameter, and therefore, denote the eigenvalue $\lambda_{\alpha k}$ by $\lambda_{\alpha k}(\nu)$:

$$\lambda_{\alpha k} = \lambda_{\alpha k}(\nu),$$

and the linearized operator L by L_ν :

$$L = L_\nu.$$

Let $\tilde{\nu}_0 > 0$ be the number satisfying $\kappa_0 = 0$, where κ_0 is the coefficient of $(\alpha k)^2$ in $\lambda_{\alpha k}(\nu)$ given in Theorem 3.1. Then, by a perturbation argument, one can see that for each $0 < \alpha \ll 1$, there exists $\nu_0 > 0$ such that

$$\begin{aligned} \operatorname{Re} \lambda_{\pm\alpha}(\nu_0) &= 0; \\ \operatorname{Re} \lambda_{\pm\alpha}(\nu) < 0 &\Leftrightarrow \nu > \nu_0; \\ \operatorname{Re} \lambda_{\pm\alpha}(\nu) > 0 &\Leftrightarrow \nu < \nu_0. \end{aligned}$$

From [8, Section 6], one can see that $\operatorname{Re} \lambda_\alpha(\nu)$ is analytic in α^2 . Setting $\zeta(\alpha^2, \nu) = \operatorname{Re} \lambda_\alpha(\nu)/\alpha^2$, we see that $\partial_\nu \zeta(\alpha^2, \nu) = -\frac{1}{12\nu^2} \left[\left(\frac{1}{280} - \gamma^2 \right) + \frac{\nu^2}{10\gamma^2} \right] + O(\alpha^2) < 0$ for $\alpha \ll 1$, and so $\operatorname{Re} \lambda_\alpha(\nu)$ crosses the imaginary axis from left to right at $\nu = \nu_0$ when ν is decreased.

We make the following assumption:

$$\sigma(-L_{\nu_0}) \cap \{\lambda; \operatorname{Re} \lambda = 0\} = \{\lambda_\alpha(\nu_0), \lambda_{-\alpha}(\nu_0)\}. \quad (4.1)$$

Theorem 4.1. *Assume that (4.1) holds true. Then there is a solution branch $\{\nu, u\} = \{\nu_\varepsilon, u_\varepsilon\}$ ($|\varepsilon| \ll 1$) such that*

$$\begin{aligned}\nu_\varepsilon &= \nu_0 + O(\varepsilon), \\ u_\varepsilon &= u_\varepsilon(x_1 - c_\varepsilon t, x_2), \\ u_\varepsilon(x_1 + \frac{2\pi}{\alpha}, x_2) &= u_\varepsilon(x_1, x_2), \\ u_\varepsilon(x_1, x_2) &= \varepsilon \begin{pmatrix} 1 \\ \frac{1}{2\gamma^2}(-x_2^2 + x_2) \\ 0 \end{pmatrix} \frac{\sqrt{2}}{2} \cos \alpha x_1 (1 + O(\alpha)) + O(\varepsilon^2), \\ c_\varepsilon &= \frac{1}{6} + O(\varepsilon).\end{aligned}$$

Remark 4.2. Iooss and Padula ([6]) showed that, for each ν , there exists a positive number c such that the set

$$\sigma(-L_\nu) \cap \{\lambda; \operatorname{Re} \lambda \geq -c\}$$

consists of a finite number of eigenvalues with finite multiplicities. (See Lemma 6.10 below.) Therefore, it seems very unlikely that assumption (4.1) does not hold true for all $\alpha \ll 1$.

5 Proof of Theorem 4.1

In this section we give a proof of Theorem 4.1.

We set $\eta = \nu - \nu_0$ that will be taken as a new bifurcation parameter. For simplicity, we write L_η for $L_{\eta+\nu_0}$ omitting ν_0 .

5.1 Spectrum of $-L_0$

We first make an observation of the spectrum of $-L_\eta$. Let us consider the resolvent problem

$$\lambda u + Lu = F. \quad (5.1)$$

We expand u and F into the Fourier series in x_1 :

$$\begin{aligned}u &= \sqrt{\frac{\alpha}{2\pi}} \sum_{k \in \mathbb{Z}} u_k(x_2) e^{i\alpha k x_1}, \quad u_k = {}^\top(\phi_k, w_k), \\ F &= \sqrt{\frac{\alpha}{2\pi}} \sum_{k \in \mathbb{Z}} F_k(x_2) e^{i\alpha k x_1}, \quad F_k = {}^\top(f_k^0, f_k)\end{aligned}$$

with $\int_0^1 \phi_0(x_2) dx_2 = \int_0^1 f_0^0(x_2) dx_2 = 0$. Then the problem is reduce to the following problems for $k \in \mathbb{Z}$:

$$(\lambda + L_{\eta,k})u_k = F_k. \quad (5.2)$$

Here $L_{\eta,k}$ is the operator on $L_k^2(0,1) \times L^2(0,1)^2$ obtained by replacing ∂_{x_1} in L by $i\alpha k$ with domain $D(L_{\eta,k}) = \{u_k = {}^\top(\phi_k, w_k) \in L_k^2(0,1) \times L^2(0,1)^2; w_k \in$

$H_0^1(0, 1), L_{\eta, k} u_k \in L_k^2(0, 1) \times L^2(0, 1)^2\}$, where $L_k^2(0, 1) = L^2(0, 1)$ when $k \neq 0$ and $L_0^2(0, 1) = L^2(0, 1) \cap \{\phi; \int_0^1 \phi(x_2) dx_2 = 0\}$.

Let $\tilde{X} = L_{per}^2 \times (L_{per}^2)^2$. We denote by $\tilde{L}$ the extension of L to $\tilde{X}$, more precisely, $\tilde{L}$ is an operator on $\tilde{X}$ with domain $D(\tilde{L}) = \{u = {}^\top(\phi, w) \in \tilde{X}; w \in (H_{per, 0}^1)^2, \tilde{L}u \in \tilde{X}\}$ and $\tilde{L}$ has the same form as L . Similarly, we define an operator $\tilde{L}_{\eta, k}$ on $L^2(0, 1) \times L^2(0, 1)^2$ by the extension of $L_{\eta, k}$ to $L^2(0, 1) \times L^2(0, 1)^2$. Note that $\tilde{L}_{\eta, k} = L_{\eta, k}$ when $k \neq 0$ and $L_{\eta, 0}$ is the restriction of $\tilde{L}_{0, \eta}$ to $L_0^2(0, 1) \times L^2(0, 1)^2$. We also introduce the adjoint operator $\tilde{L}^*$ (with respect to the inner product $\langle \cdot, \cdot \rangle$) which is given by

$$\tilde{L}^* = \begin{pmatrix} -v_s^1 \partial_{x_1} & -\nu^\top \mathbf{e}_1 - \gamma^2 \text{div} \\ -\nabla & -\nu \Delta - \tilde{\nu} \nabla \text{div} - v_s^1 \partial_{x_1} + (\partial_{x_2} v_s^1) \mathbf{e}_2^\top \mathbf{e}_1 \end{pmatrix}.$$

Similarly, the adjoint operators $\tilde{L}_{\eta, k}^*$ of $\tilde{L}_{\eta, k}$ are defined.

Since X is an invariant set of $\tilde{L}$, we see that if λ is an eigenvalue of $-L$, then the eigenprojection for λ of $-L$ is the restriction of the eigenprojection for λ of $-\tilde{L}$. The same also holds for eigenprojections of $L_{\eta, 0}$ and $\tilde{L}_{\eta, 0}$.

Under the assumption (4.1), the following claims are concluded. In what follows we denote the critical eigenvalues $\lambda_{\pm\alpha}(\nu_0)$ by $\pm ia$ with $a = -\frac{\alpha}{6}(1 + O(\alpha^2)) \in \mathbb{R} \setminus \{0\}$:

$$\lambda_{\pm\alpha}(\nu_0) = \pm ia.$$

As for $\sigma(-L_{0, k})$, we have

- $k = \pm 1$:

$$\begin{aligned} \sigma(-L_{0, \pm 1}) \cap \{\lambda; \text{Re } \lambda = 0\} &= \{\pm ia\}, \\ \pm ia &\text{ are isolated simple eigenvalues of } -L_{0, \pm 1}, \\ N(\pm ia + L_{0, \pm 1}) &= \text{span}\{v_{\pm 1}\}, \quad v_{-1} = \overline{v_{+1}}. \end{aligned}$$

- $k \neq \pm 1$: there exists a constant $\beta > 0$ such that $\sigma(-L_{0, k}) \subset \{\lambda; |\text{Re } \lambda| \geq \beta\}$ for all $k \in \mathbb{Z}$ with $k \neq \pm 1$.

The eigenprojections for $\pm ia$ are given in terms of eigenfunctions of the adjoint operator $\tilde{L}_{0, k}^*$. Namely, we have

$$\begin{aligned} \text{the eigenprojections } \Pi_\pm \text{ for } \pm ia &\text{ are given by } \Pi_\pm u = \langle \langle u, v_{\pm 1}^* \rangle \rangle v_{\pm 1}, \\ \text{where } N(\mp ia + \tilde{L}_{0, \pm 1}^*) &= \text{span}\{v_{\pm 1}^*\}, \quad \langle \langle v_{\pm 1}, v_{\pm 1}^* \rangle \rangle = 1. \end{aligned}$$

It then follows that $\sigma(-L_0)$ satisfies

$$\begin{aligned} \sigma(-L_0) \cap \{\lambda; \text{Re } \lambda = 0\} &= \{\pm ia\}, \\ \pm ia &\text{ are isolated simple eigenvalues of } -L_0, \\ N(\pm ia + L_0) &= \text{span}\{V_\pm\}, \\ \text{where } V_\pm &= v_{\pm 1}(x_2) e^{\pm i\alpha x_1}. \end{aligned}$$

Furthermore, $V_{\pm}^* = \frac{\alpha}{2\pi} v_{\pm 1}^*(x_2) e^{\pm i\alpha x_1}$ satisfy

$$-\tilde{L}_0^* V_{\pm}^* = \mp i a V_{\pm}^*, \quad \langle V_{\pm}, V_{\pm}^* \rangle = 1, \quad \langle V_{\pm}, V_{\mp}^* \rangle = 0,$$

and the eigenprojections $P_{\pm}$ for $\pm i a$ of $-L$ are given by

$$P_{\pm} V = \langle V, V_{\pm}^* \rangle V_{\pm}.$$

It was proved in [7] that eigenfunctions $V_{\pm}$ and $V_{\pm}^*$ are smooth and, for each nonnegative integer k , eigenprojections $P_{\pm}$ are bounded from $L_{per,*}^2 \times L_{per}^2$ to $H_{per,*}^k \times H_{per}^k$:

$$\|P_{\pm} u\|_{H^k \times H^k} \leq C_k \|u\|_2.$$

See [7, Lemma 4.3]. These boundedness properties of $P_{\pm}$ will be employed later.

5.2 Traveling wave solution

Let us consider the nonlinear problem

$$\partial_t \tilde{u} + L_{\eta} \tilde{u} = F(\eta, \tilde{u}), \tag{5.3}$$

where $F(\eta, \tilde{u})$ denotes the nonlinear term.

We look for a solution in the form

$$\tilde{u}(x_1, x_2, t) = u(x_1 - ct, x_2).$$

We substitute this into (5.3). Then the problem is rewritten as

$$\mathcal{L}_{c,\eta} u = F(\eta, u), \tag{5.4}$$

where

$$\mathcal{L}_{c,\eta} = L_{\eta} - c \partial_{x_1}.$$

We first investigate the spectrum of $-\mathcal{L}_{c_0,0}$.

5.3 Spectrum of $-\mathcal{L}_{c_0,0}$

The following proposition on the spectrum of $\mathcal{L}_{c_0,0}$ follows from the observation in section 5.1.

Proposition 5.1. *Set $c_0 = -\frac{a}{\alpha}$. Then*

$$\begin{aligned} \sigma(-\mathcal{L}_{c_0,0}) \cap \{\lambda; \operatorname{Re} \lambda = 0\} &= \{0\}, \\ 0 &\text{ is an isolated semisimple eigenvalue of } -\mathcal{L}_{c_0,0}, \\ N(-\mathcal{L}_{c_0,0}) &= \operatorname{span}\{V_+, V_-\}, \quad V_- = \overline{V_+}. \end{aligned}$$

Let us next introduce the eigenprojection for the eigenvalue 0 of $-\mathcal{L}_{c_0,0}$. We set

$$\begin{aligned} V_1 &= \sqrt{2}\operatorname{Re} V_+, \quad V_2 = \sqrt{2}\operatorname{Im} V_+, \\ V_1^* &= \sqrt{2}\operatorname{Re} V_+^*, \quad V_2^* = \sqrt{2}\operatorname{Im} V_+^*. \end{aligned}$$

Then

$$\begin{aligned} N(-\mathcal{L}_{c_0,0}) &= \operatorname{span}\{V_1, V_2\}, \\ \langle V_j, V_k^* \rangle &= \delta_{jk}, \quad j, k = 1, 2. \end{aligned}$$

We introduce the following notation $\llbracket u \rrbracket_j$ ($j = 1, 2$):

$$\llbracket u \rrbracket_j = \langle u, V_j^* \rangle.$$

Proposition 5.2. *Define P , P_1 and P_2 by*

$$Pu = P_1u + P_2u, \quad P_ju = \llbracket u \rrbracket_j V_j \quad (j = 1, 2).$$

Then P is the eigenprojection for eigenvalue 0 of $-\mathcal{L}_{c_0,0}$; and

$$R(P_j) = \operatorname{span}\{V_j\}, \quad P_j^2 = P_j, \quad P_j P_k = O \quad (j \neq k).$$

For each nonnegative integer k , P_j are bounded from $L_{per,}^2 \times L_{per}^2$ to $H_{per,*}^k \times H_{per}^k$:*

$$\|P_j u\|_{H^k \times H^k} \leq C \|u\|_2.$$

Furthermore, $u \in R(I - P_j)$ if and only if $\llbracket u \rrbracket_j = 0$.

5.4 Formulation of the problem

We look for solutions of (5.4) in a neighborhood of $\{c, \eta, u\} = \{c_0, 0, 0\}$ in the form:

$$u = \varepsilon(V_1 + \varepsilon V), \quad V \in R(Q), \quad Q = I - P,$$

$$c = c_0 + \varepsilon \sigma.$$

Here ε is a small parameter. Note that $P_2u = 0$.

We set

$$K_0 = \frac{1}{\eta}(L_\eta - L_0) = \begin{pmatrix} 0 & 0 \\ -\frac{1}{\gamma^2}\mathbf{e}_1 & -\Delta - \nabla \operatorname{div} \end{pmatrix}.$$

Then

$$L_\eta = L_0 + \eta K_0,$$

and

$$\mathcal{L}_{c,\eta} = \mathcal{L}_{c_0,0} - \varepsilon \sigma \partial_{x_1} + \eta K_0.$$

We scale η as

$$\eta = \varepsilon \omega.$$

Problem (5.4) is then written as

$$\mathcal{L}_{c_0,0}V - \sigma \partial_{x_1}(V_1 + \varepsilon V) + \omega K_0(V_1 + \varepsilon V) = \frac{1}{\varepsilon^2} F(\varepsilon\omega, \varepsilon(V_1 + \varepsilon V)). \quad (5.5)$$

We denote the right-hand side by

$$\frac{1}{\varepsilon^2} F(\varepsilon\omega, \varepsilon(V_1 + \varepsilon V)) = -NV_1 + \varepsilon V + G(\varepsilon, \varepsilon\omega, V_1 + \varepsilon V),$$

where

$$N[\tilde{u}]u = {}^\top(\operatorname{div}(\phi\tilde{w}), 0)$$

for $\tilde{u} = {}^\top(\tilde{\phi}, \tilde{w})$ and $u = {}^\top(\phi, w)$, and

$$G(\varepsilon, \omega, u) = {}^\top(0, g(\varepsilon, \omega, u))$$

with

$$\begin{aligned} g(\varepsilon, \omega, u) = & -w \cdot \nabla w - \frac{\phi}{\gamma^2 + \varepsilon\phi} \left((\nu_0 + \omega)\Delta w + \frac{(\nu_0 + \omega)}{\gamma^2} \phi \mathbf{e}_1 + (\tilde{\nu}_0 + \omega)\nabla \operatorname{div} w \right) \\ & + P^{(1)}(\varepsilon\phi)\phi \nabla \phi \end{aligned}$$

for $u = {}^\top(\phi, w)$, where $\tilde{\nu}_0 = \nu_0 + \nu'$.

We decompose (5.5) into the P_j -parts ($j = 1, 2$) and Q -part. Here and in what follows we set

$$Q = I - P = I - P_1 - P_2.$$

We take the inner product of (5.5) with V_j^* ($j = 1, 2$) and apply Q to (5.5). Since

$$\llbracket \partial_{x_1}(V_1 + \varepsilon V) \rrbracket_1 = 0, \quad \llbracket \partial_{x_1}(V_1 + \varepsilon V) \rrbracket_2 = -\alpha,$$

we find that

$$\begin{aligned} \omega \llbracket K_0 V_1 \rrbracket_1 &= -\varepsilon\omega \llbracket K_0 V \rrbracket_1 - \llbracket NV_1 + \varepsilon V \rrbracket_1 \\ &\quad + \llbracket G(\varepsilon, \varepsilon\omega, V_1 + \varepsilon V) \rrbracket_1, \\ \omega \llbracket K_0 V_1 \rrbracket_2 + \alpha\sigma &= -\varepsilon\omega \llbracket K_0 V \rrbracket_2 - \llbracket NV_1 + \varepsilon V \rrbracket_2 \\ &\quad + \llbracket G(\varepsilon, \varepsilon\omega, V_1 + \varepsilon V) \rrbracket_2, \\ \omega Q K_0 V_1 + (\mathcal{L}_{c_0,0} - \varepsilon\sigma Q \partial_{x_1} + \varepsilon Q N[V_1 + \varepsilon V]) V \\ &= -\varepsilon\omega Q K_0 V - Q N[V_1 + \varepsilon V] V_1 + Q G(\varepsilon, \varepsilon\omega, V_1 + \varepsilon V). \end{aligned}$$

We thus arrive at the following problem:

$$T(\varepsilon, \sigma, V)U = \mathcal{F}(\varepsilon, U), \quad (5.6)$$

where

$$U = {}^\top(\omega, \sigma, V) \in \mathbb{R} \times \mathbb{R} \times X^2.$$

Here X^ℓ denotes the function space

$$X^\ell = H_{per,*}^\ell \times [H_{per}^{\ell+1} \cap H_{per,0}^1], \quad \ell = 1, 2,$$

and, for a given $(\tilde{\sigma}, \tilde{V}) \in \mathbb{R} \times X^2$, $T(\varepsilon, \tilde{\sigma}, \tilde{V})$ is the linear map defined by

$$T(\varepsilon, \tilde{\sigma}, \tilde{V}) : \mathbb{R} \times \mathbb{R} \times QX^\ell \rightarrow \mathbb{R} \times \mathbb{R} \times Q(H^\ell \times H^{\ell-1}), \quad \ell = 1, 2,$$

$$T(\varepsilon, \tilde{\sigma}, \tilde{V}) = \begin{pmatrix} \llbracket K_0 V_1 \rrbracket_1 & 0 & 0 \\ \llbracket K_0 V_2 \rrbracket_1 & \alpha & 0 \\ QK_0 V_1 & 0 & \mathcal{L}_{c_0,0} - \varepsilon \tilde{\sigma} Q \partial_{x_1} + \varepsilon QN[V_1 + \varepsilon \tilde{V}] \end{pmatrix}.$$

$\mathcal{F}(\varepsilon, U)$ is the nonlinear map given by

$$\mathcal{F}(\varepsilon, U) = {}^\top(\mathcal{F}_1(\varepsilon, U), \mathcal{F}_2(\varepsilon, U), \mathcal{F}_3(\varepsilon, U)) \quad (U = {}^\top(\omega, \sigma, V)),$$

where

$$\begin{aligned} \mathcal{F}_j(\varepsilon, U) &= -\varepsilon \omega \llbracket K_0 V \rrbracket_j - \llbracket NV_1 + \varepsilon V \rrbracket_j + \llbracket G(\varepsilon, \varepsilon \omega, V_1 + \varepsilon V) \rrbracket_j, \\ &\quad (j = 1, 2), \end{aligned}$$

$$\mathcal{F}_3(\varepsilon, U) = -\varepsilon \omega QK_0 V - QN[V_1 + \varepsilon V]V_1 + QG(\varepsilon, \varepsilon \omega, V_1 + \varepsilon V).$$

Concerning $T(\varepsilon, \tilde{\sigma}, \tilde{V})$ we have the following

Proposition 5.3. (i) $\llbracket K_0 V_1 \rrbracket_1 > 0$.

(ii) For given $M > 0$, there exists $\varepsilon_1 > 0$ such that if $|\varepsilon| \leq \varepsilon_1$ and $|\tilde{\sigma}| + \|\tilde{V}\|_{X^2} \leq M$, then $\mathcal{L}_{c_0,0} - \varepsilon \tilde{\sigma} Q \partial_{x_1} + \varepsilon QN[V_1 + \varepsilon \tilde{V}]$ has a bounded inverse from $Q(H_{per,*}^\ell \times H_{per}^{\ell-1})$ to QX^ℓ ($\ell = 1, 2$).

(iii) Under the assumption of (ii), $T(\varepsilon, \tilde{\sigma}, \tilde{V})$ has a bounded inverse from $\mathbb{R} \times \mathbb{R} \times Q(H_{per,*}^\ell \times H_{per}^{\ell-1})$ to $\mathbb{R} \times \mathbb{R} \times QX^\ell$ ($\ell = 1, 2$), and it holds that for $U = {}^\top(\tilde{\eta}, \sigma, V)$,

$$\|T(\varepsilon, \tilde{\sigma}, \tilde{V})^{-1}U\|_{\mathbb{R} \times \mathbb{R} \times X^\ell} \leq C_1 \|U\|_{\mathbb{R} \times \mathbb{R} \times H^\ell \times H^{\ell-1}}, \quad \ell = 1, 2.$$

We will give a proof of Proposition 5.3 (ii) and (iii) in section 6, and a proof of (i) will be given in section 8.

As for $\mathcal{F}(\varepsilon, U)$, using Sobolev inequalities, we have the following estimates by a straightforward computation.

Proposition 5.4. For given $M \in (0, \frac{\gamma^2}{2C_S}]$, there exists $\varepsilon_2 > 0$ such that if $|\varepsilon| \leq \varepsilon_2$, $\|U\|_{\mathbb{R} \times \mathbb{R} \times X^2} \leq M$ and $\|U^{(j)}\|_{\mathbb{R} \times \mathbb{R} \times X^2} \leq M$ ($j = 1, 2$), then the following estimates hold:

$$\|\mathcal{F}(\varepsilon, U) - \mathcal{F}(0, 0)\|_{\mathbb{R} \times \mathbb{R} \times H^2 \times H^1} \leq C(M)M|\varepsilon|,$$

$$\|\mathcal{F}(\varepsilon, U^{(1)}) - \mathcal{F}(\varepsilon, U^{(2)})\|_{\mathbb{R} \times \mathbb{R} \times H^1 \times H^0} \leq C(M)|\varepsilon| \|U^{(1)} - U^{(2)}\|_{\mathbb{R} \times \mathbb{R} \times X^1},$$

where $C(M) > 0$ is a nondecreasing continuous function of M .

5.5 Iteration

The desired solution branch in Theorem 4.1 can now be obtained by an iteration argument.

We define $U^{(n)} = {}^\top(\omega^{(n)}, \sigma^{(n)}, V^{(n)})$ ($n \geq 1$) in the following way. $U^{(1)}$ is the solution of

$$\begin{aligned} T(0, 0, 0)U^{(1)} &= \mathcal{F}(0, 0) \\ &= {}^\top(\llbracket F(0, V_1) \rrbracket_1, \llbracket F(0, V_1) \rrbracket_2, QF(0, V_1)). \end{aligned}$$

Note that $F(0, V_1) = -N[V_1]V_1 + G(0, 0, V_1)$. By Propositions 5.3 we have

$$\|U^{(1)}\|_{\mathbb{R} \times \mathbb{R} \times X^2} \leq C_1 \|\mathcal{F}(0, 0)\|_{\mathbb{R} \times \mathbb{R} \times H^2 \times H^1} < \infty. \quad (5.7)$$

We set

$$M = 2C_1 \|\mathcal{F}(0, 0)\|_{\mathbb{R} \times \mathbb{R} \times H^2 \times H^1}. \quad (5.8)$$

Let $\varepsilon > 0$ satisfy $|\varepsilon| \leq \min\{\varepsilon_1, \varepsilon_2, \frac{1}{2C_1C(M)}\}$. Then for $n \geq 2$ we can define $U^{(n)}$ by the solution of

$$T(\varepsilon, \sigma^{(n-1)}, V^{(n-1)})U^{(n)} = \mathcal{F}(\varepsilon, U^{(n-1)}), \quad (5.9)$$

and $U^{(n)}$ satisfies

$$\|U^{(n)}\|_{\mathbb{R} \times \mathbb{R} \times X^2} \leq M$$

for all $n \geq 1$. In fact, assume that $\|U^{(n-1)}\|_{\mathbb{R} \times \mathbb{R} \times X^2} \leq M$. Then, $\mathcal{F}(\varepsilon, U^{(n-1)}) \in \mathbb{R} \times \mathbb{R} \times Q(H_{per,*}^2 \times H_{per}^1)$, and thus, Proposition 5.3 implies that (5.9) has a solution $U^{(n)} \in \mathbb{R} \times \mathbb{R} \times X^2$. Furthermore, since

$$T(\varepsilon, \sigma^{(n-1)}, V^{(n-1)})U^{(n)} = \mathcal{F}(0, 0) + (\mathcal{F}(\varepsilon, U^{(n-1)}) - \mathcal{F}(0, 0))$$

and $|\varepsilon| \leq \min\{\varepsilon_1, \varepsilon_2, \frac{1}{2C_1C(M)}\}$, we see from Propositions 5.3 and 5.4 that

$$\|U^{(n)}\|_{\mathbb{R} \times \mathbb{R} \times X^2} \leq \frac{M}{2} + C_1C(M)M|\varepsilon| \leq M.$$

Therefore, with this observation and (5.7), we conclude by induction that $\|U^{(n)}\|_{\mathbb{R} \times \mathbb{R} \times X^2} \leq M$ for all $n \geq 1$.

We next prove that $\{U^{(n)}\}$ is a Cauchy sequence in $\mathbb{R} \times \mathbb{R} \times X^1$. We set

$$\mathcal{D}V = {}^\top(0, 0, \partial_{x_1}V), \quad \mathcal{N}_Q[\tilde{V}]V = {}^\top(0, 0, QN[\tilde{V}]V).$$

Since

$$\begin{aligned} &T(\varepsilon, \sigma^{(n)}, V^{(n)})U^{(n+1)} - T(\varepsilon, \sigma^{(n-1)}, V^{(n-1)})U^{(n)} \\ &= T(\varepsilon, \sigma^{(n)}, V^{(n)})(U^{(n+1)} - U^{(n)}) - \varepsilon(\sigma^{(n)} - \sigma^{(n-1)})\mathcal{D}V^{(n)} \\ &\quad + \varepsilon^2\mathcal{N}_Q[V^{(n)} - V^{(n-1)}]V^{(n)}, \end{aligned}$$

we have

$$\begin{aligned} &T(\varepsilon, \sigma^{(n)}, V^{(n)})(U^{(n+1)} - U^{(n)}) \\ &= \varepsilon(\sigma^{(n)} - \sigma^{(n-1)})\mathcal{D}V^{(n)} - \varepsilon^2\mathcal{N}_Q[V^{(n)} - V^{(n-1)}]V^{(n)} \\ &\quad + (\mathcal{F}(\varepsilon, U^{(n)}) - \mathcal{F}(\varepsilon, U^{(n-1)})), \end{aligned}$$

and by Propositions 5.3 and 5.4,

$$\begin{aligned}
& \|U^{(n+1)} - U^{(n)}\|_{\mathbb{R} \times \mathbb{R} \times X^1} \\
& \leq C_1 \left\{ |\varepsilon| |\sigma^{(n)} - \sigma^{(n-1)}| \|\partial_{x_1} V^{(n)}\|_{H^1 \times H^0} + |\varepsilon|^2 \|QN[V^{(n)} - V^{(n-1)}]V^{(n)}\|_{H^1 \times H^0} \right. \\
& \quad \left. + \|\mathcal{F}(\varepsilon, U^{(n)}) - \mathcal{F}(\varepsilon, U^{(n-1)})\|_{\mathbb{R} \times \mathbb{R} \times H^1 \times H^0} \right\} \\
& \leq C_1 \left\{ CM|\varepsilon| |\sigma^{(n)} - \sigma^{(n-1)}| + CM|\varepsilon|^2 \|V^{(n)} - V^{(n-1)}\|_{X^1} \right. \\
& \quad \left. + C(M)|\varepsilon| \|U^{(n)} - U^{(n-1)}\|_{\mathbb{R} \times \mathbb{R} \times X^1} \right\} \\
& \leq \frac{1}{2} \|U^{(n)} - U^{(n-1)}\|_{\mathbb{R} \times \mathbb{R} \times X^1}
\end{aligned}$$

if $|\varepsilon| \leq \frac{1}{2C_1(2CM+C(M))}$. It then follows that there exists $\varepsilon_0 > 0$ such that if $|\varepsilon| \leq \varepsilon_0$, then $\{U^{(n)}\}$ is a Cauchy sequence in $\mathbb{R} \times \mathbb{R} \times X^1$. We thus conclude if $|\varepsilon| \leq \varepsilon_0$, there exists $U = {}^\top(\omega, \sigma, V) \in \mathbb{R} \times \mathbb{R} \times X^2$ satisfying

$$T(\varepsilon, \sigma, V)U = \mathcal{F}(\varepsilon, U).$$

With this $U = {}^\top(\omega, \sigma, V)$, setting

$$\nu = \nu_0 + \varepsilon\omega, \quad u = \varepsilon V_1(x_1 - ct, x_2) + \varepsilon^2 V(x_1 - ct, x_2), \quad c = c_0 + \varepsilon\sigma,$$

we have the desired traveling wave solutions.

To complete the proof of Theorem 4.1, it remains to prove Proposition 5.3.

6 Proof of Proposition 5.3 (ii), (iii)

In this section we give a proof of Proposition 5.3 (ii), (iii).

By a perturbation argument for $\alpha \ll 1$, one can compute $u_{\pm 1}$ and $u_{\pm 1}^*$ to see assertion (i) $\|K_0 V_1\|_1 > 0$ for $\alpha \ll 1$. See section 8 for the proof of (i). If assertion (ii) holds, then $T(\varepsilon, \tilde{\sigma}, \tilde{V})$ has a bounded inverse $T(\varepsilon, \tilde{\sigma}, \tilde{V})^{-1}$ which is given by

$$T(\varepsilon, \tilde{\sigma}, \tilde{V})^{-1} = \begin{pmatrix} \mathcal{A}^{-1} & 0 \\ -\mathcal{L}(\varepsilon, \tilde{\sigma}, \tilde{V})^{-1} \mathcal{B} \mathcal{A}^{-1} & \mathcal{L}(\varepsilon, \tilde{\sigma}, \tilde{V})^{-1} \end{pmatrix},$$

where

$$\mathcal{A} = \begin{pmatrix} \|K_0 V_1\|_1 & 0 \\ \|K_0 V_2\|_2 & \alpha \end{pmatrix},$$

$$\mathcal{B} = (QK_0 V_1 \quad 0),$$

$$\mathcal{L}(\varepsilon, \tilde{\sigma}, \tilde{V}) = \mathcal{L}_{c_0, 0} - \varepsilon \tilde{\sigma} Q \partial_{x_1} + \varepsilon QN[V_1 + \varepsilon \tilde{V}].$$

Therefore, in the rest of this section we will prove assertion (ii), i.e, $\mathcal{L}(\varepsilon, \tilde{\sigma}, \tilde{V})$ has a bounded inverse.

6.1 Basic estimates

From now on, we simply write $N[\tilde{w}]u$ for $N[\tilde{u}]u$ with $\tilde{u} = {}^\top(\tilde{\phi}, \tilde{w})$:

$$N[\tilde{w}]u = {}^\top(\operatorname{div}(\phi\tilde{w}), 0), \quad u = {}^\top(\phi, w).$$

In this subsection we establish basic a priori estimates of solution u to

$$\lambda u + Lu + N[\tilde{w}]u = F, \quad u \in X^\ell, \quad (6.1)$$

where $\tilde{w}$ is a given function in $H_{per}^3 \cap H_{per,0}^1$ with $\tilde{w}(x) \in \mathbb{R}^2$ and $\lambda \in \mathbb{C}$ is a parameter.

We introduce some notations. We define the new norm $||| \cdot |||_2$ of L_{per}^2 by

$$|||u|||_2 = \left(\frac{1}{\gamma^2} \|\phi\|_2^2 + \|w\|_2^2 \right)^{\frac{1}{2}}$$

for $u = {}^\top(\phi, w)$. We also define $D[w]$ and $\dot{\phi}_\lambda$ by

$$D[w] = \nu \|\nabla w\|_2^2 + \tilde{\nu} \|\operatorname{div} w\|_2^2$$

and

$$\dot{\phi}_\lambda = \lambda \phi + v_s^1 \partial_{x_1} \phi + \operatorname{div}(\phi \tilde{w}),$$

respectively. For operators A and B , we denote by $[A, B]$ the commutator of A and B :

$$[A, B]f = A(Bf) - B(Af).$$

We will prove the following

Proposition 6.1. *There exists a number Λ satisfying $0 < \Lambda \leq \frac{1}{2} \frac{\gamma^2}{\nu + \tilde{\nu}}$ such that if $\operatorname{Re} \lambda \geq -\Lambda$, then*

$$\begin{aligned} & \left(\operatorname{Re} \lambda + \Lambda \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda \right) \left(\|\partial_x u\|_2^2 + \|\partial_x^2 w\|_2^2 + |\lambda|^2 \|u\|_2^2 + \|\dot{\phi}_\lambda\|_{H^1}^2 \right) \\ & \leq C \left\{ \|F\|_{H^1 \times L^2}^2 + \|\tilde{w}\|_{H^3} (1 + \|\tilde{w}\|_{H^3}) \|\phi\|_{H^1}^2 + \|w\|_2^2 + \|\phi\|_{H^{-1}}^2 \right\}, \end{aligned} \quad (6.2)$$

$$\begin{aligned} & \left(\operatorname{Re} \lambda + \Lambda \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda \right) \left(\|\partial_x u\|_2^2 + \|\partial_x^2 \phi\|_2^2 + \|\partial_{x_1}^2 w\|_2^2 + |\lambda|^2 \|u\|_2^2 \right) \\ & \quad + \|\partial_x^2 w\|_2^2 + \|\partial_x^3 w\|_2^2 + |\lambda|^2 \|\nabla u\|_2^2 + \|\dot{\phi}_\lambda\|_{H^2}^2 \\ & \leq C \left\{ \|F\|_{H^2 \times H^1}^2 + |\lambda|^2 \|F\|_2^2 + \|\tilde{w}\|_{H^3} (1 + \|\tilde{w}\|_{H^3}) (\|\phi\|_{H^2}^2 + |\lambda|^2 \|\phi\|_2^2) \right. \\ & \quad \left. + \|w\|_2^2 + \|\phi\|_{H^{-1}}^2 \right\}. \end{aligned} \quad (6.3)$$

To prove Proposition 6.1, we will employ the following Bogovskii lemma.

Lemma 6.2. ([2]) *There exists a bounded operator $\mathcal{B} : L_{per,*}^2 \rightarrow H_{per,0}^1$ such that*

$$\operatorname{div} \mathcal{B}g = g, \quad g \in L_{per,*}^2$$

$$\|\nabla \mathcal{B}g\|_2 \leq C_B \|g\|_2,$$

where C_B is a positive constant depending only on α . Furthermore, if $g = \operatorname{div} \mathbf{g}$ with $\mathbf{g} = {}^\top(g^1, g^2)$ satisfying $g^1|_{x_1=-\frac{\pi}{\alpha}} = g^1|_{x_1=\frac{\pi}{\alpha}}$, $g^2|_{x_2=0,1} = 0$, then

$$\operatorname{div} \mathcal{B}(\operatorname{div} \mathbf{g}) = \operatorname{div} \mathbf{g},$$

$$\|\mathcal{B}(\operatorname{div} \mathbf{g})\|_2 \leq C_B \|\mathbf{g}\|_2.$$

An outline of the proof of Lemma 6.2 will be given in Section 7. We will also employ the Poincaré inequalities

$$\|\phi\|_2 \leq C \|\nabla \phi\|_2, \quad \|w\|_2 \leq \|\nabla w\|_2$$

for $\phi \in H_{per,*}^1$ and $w \in H_{per,0}^1$, and the Sobolev inequality

$$\|f\|_\infty \leq C \|f\|_{H^2}$$

for $f \in H_{per}^2$. Here C is a positive constant depending only on α .

We begin with the following L^2 energy estimates.

Proposition 6.3. *There exists a positive number Λ_0 such that the following inequalities hold uniformly for $\operatorname{Re} \lambda \geq \Lambda_0$.*

$$\begin{aligned} & (\operatorname{Re} \lambda + \Lambda_0) |\lambda|^{2k} \|u\|_2^2 + \frac{1}{4} |\lambda|^{2k} D[w] \\ & \leq C |\lambda|^{2k} \left\{ \|F\|_2 \|u\|_2 + \|\partial_{x_2} v_s^1\|_\infty \|w\|_2^2 + \frac{\nu}{\gamma^4} \|\phi\|_{H^{-1}}^2 + \frac{\|\tilde{w}\|_{H^3}}{\gamma} \|\phi\|_2 \|u\|_2 \right\} \end{aligned} \quad (6.4)$$

for $k = 0, 1$,

$$\begin{aligned} & \left(\operatorname{Re} \lambda + \frac{1}{2} \Lambda_0 \right) \|u\|_2^2 + \frac{1}{8} D[w] + \frac{\nu + \tilde{\nu}}{32\gamma^4} \|\dot{\phi}_\lambda\|_2^2 \\ & \leq C \left\{ \left(\frac{1}{\gamma^2 \Lambda_0} + \frac{\nu + \tilde{\nu}}{\gamma^4} \right) \|f^0\|_2^2 + \frac{1}{\nu} \|f\|_{H^{-1}}^2 + \frac{\|\tilde{w}\|_{H^3}}{\gamma^2} \left(1 + \frac{\|\tilde{w}\|_{H^3}}{\nu} \right) \|\phi\|_2^2 \right. \\ & \quad \left. + \|\partial_{x_2} v_s^1\|_\infty \|w\|_2^2 + \frac{\nu}{\gamma^4} \|\phi\|_{H^{-1}}^2 \right\}, \end{aligned} \quad (6.5)$$

$$\begin{aligned} & \left(\operatorname{Re} \lambda + \frac{1}{2} \Lambda_0 \right) \|\partial_{x_1}^j u\|_2^2 + \frac{1}{16} D[\partial_{x_1}^j w] + \frac{\nu + \tilde{\nu}}{32\gamma^4} \|\partial_{x_1}^j \dot{\phi}_\lambda\|_2^2 \\ & \leq C \left\{ \left(\frac{1}{\gamma^2 \Lambda_0} + \frac{\nu + \tilde{\nu}}{\gamma^4} \right) \|\partial_{x_1}^j f^0\|_2^2 + \frac{1}{\nu} \|\partial_{x_1}^j f\|_{H^{-1}}^2 + \frac{\|\tilde{w}\|_{H^3}}{\gamma^2} \left(1 + \frac{\|\tilde{w}\|_{H^3}}{\nu} \right) \|\phi\|_{H^j}^2 \right. \\ & \quad \left. + \|\partial_{x_2} v_s^1\|_\infty \|\partial_{x_1}^j w\|_2^2 + \frac{\nu}{\gamma^4} \|\partial_{x_1}^j \phi\|_{H^{-1}}^2 \right\} \end{aligned} \quad (6.6)$$

for $j = 1, 2$.

Proof. We follow the argument in [6]. We take the weighted inner product of (6.1) with u . Since

$$\operatorname{Re} \langle Lu, u \rangle = D[w] + \operatorname{Re} \left\{ -\frac{\nu}{\gamma^2} (\phi, w^1) + (\partial_{x_2} v_s^1 w^2, w^1) \right\}$$

and

$$\frac{1}{\gamma^2} \operatorname{Re} (\operatorname{div} (\phi \tilde{w}), \phi) = \frac{1}{2\gamma^2} (\operatorname{div} \tilde{w}, |\phi|^2),$$

we have

$$\begin{aligned} \operatorname{Re} \lambda \|u\|_2^2 + D[w] &= \operatorname{Re} \langle F, u \rangle - \operatorname{Re} \left\{ \frac{1}{2\gamma^2} (\operatorname{div} \tilde{w}, |\phi|^2) - \frac{\nu}{\gamma^2} (\phi, w^1) + (\partial_{x_2} v_s^1 w^2, w^1) \right\} \\ &\leq |\langle F, u \rangle| + \frac{\nu}{\gamma^2} \|\phi\|_{H^{-1}} \|\nabla w\|_2 + \|\partial_{x_2} v_s^1\|_\infty \|w\|_2^2 \\ &\quad + \frac{1}{2\gamma^2} \|\operatorname{div} \tilde{w}\|_\infty \|\phi\|_2^2. \end{aligned} \tag{6.7}$$

We next introduce a new inner product $((u_1, u_2))$ defined by

$$((u_1, u_2)) = \langle u_1, u_2 \rangle - \delta [(w_1, \mathcal{B}\phi_2) + (\mathcal{B}\phi_1, w_2)]$$

for $u_j = {}^\top(\phi_j, w_j)$ ($j = 1, 2$). Here δ is a positive number to be determined later. Note that $((u, u))^{\frac{1}{2}}$ is equivalent to $\|u\|_2$ if $\delta \leq \frac{1}{2C_B\gamma}$. We also write the density and velocity components of Lu as $(Lu)_d$ and $(Lu)_v$, respectively, i.e., $Lu = {}^\top((Lu)_d, (Lu)_v)$. Then, by Lemma 6.2,

$$\begin{aligned} ((Lu)_v, \mathcal{B}\phi) &= \nu(\nabla w, \nabla \mathcal{B}\phi) + \tilde{\nu}(\operatorname{div} w, \operatorname{div} \mathcal{B}\phi) - (\phi, \operatorname{div} \mathcal{B}\phi) \\ &\quad - \frac{\nu}{\gamma^2} (\phi e_1, \mathcal{B}\phi) - (v_s^1 w, \partial_{x_1} \mathcal{B}\phi) + (\partial_{x_2} v_s^1 w^2 e_1, \mathcal{B}\phi) \\ &= \nu(\nabla w, \nabla \mathcal{B}\phi) + \tilde{\nu}(\operatorname{div} w, \phi) - \|\phi\|_2^2 \\ &\quad - \frac{\nu}{\gamma^2} (\phi e^1, \mathcal{B}\phi) - (v_s^1 w, \partial_{x_1} \mathcal{B}\phi) + (\partial_{x_2} v_s^1 w^2 e_1, \mathcal{B}\phi). \end{aligned}$$

Applying Lemma 6.2 again, we have

$$\begin{aligned} -\operatorname{Re} ((Lu)_v, \mathcal{B}\phi) &\geq \|\phi\|_2^2 - \nu C_B \|\nabla w\|_2 \|\phi\|_2 - \tilde{\nu} \|\operatorname{div} w\|_2 \|\phi\|_2 \\ &\quad - \frac{\nu}{\gamma^2} \|\phi\|_{H^{-1}} \|\nabla \mathcal{B}\phi\|_2 - \|v_s^1\|_{C^1} (\|w\|_2 + \|w\|_{H^{-1}}) \|\nabla \mathcal{B}\phi\|_2 \\ &\geq \frac{3}{4} \|\phi\|_2^2 - C \left\{ \nu^2 C_B^2 \|\nabla w\|_2^2 + \tilde{\nu}^2 \|\operatorname{div} w\|_2^2 \right. \\ &\quad \left. - \frac{\nu^2 C_B^2}{\gamma^4} \|\phi\|_{H^{-1}}^2 - C_B^2 \|v_s^1\|_{C^1}^2 \|\nabla w\|_2^2 \right\}. \end{aligned} \tag{6.8}$$

Since $(Lu)_d = \operatorname{div} (\phi v_s + \gamma^2 w + \phi \tilde{w})$, we see from Lemma 6.2 that

$$\begin{aligned} |(\mathcal{B}(Lu)_d, w)| &\leq C_B \|\phi v_s + \gamma^2 w + \phi \tilde{w}\|_2 \|w\|_2 \\ &\leq C_B (\|v_s^1\|_\infty + \|\tilde{w}\|_\infty) \|\phi\|_2 \|w\|_2 + C_B \gamma^2 \|\nabla w\|_2^2 \\ &\leq \frac{1}{4} \|\phi\|^2 + C \left\{ (C_B^2 \|v_s^1\|_\infty^2 + C_B \gamma^2) \|\nabla w\|_2^2 + C_B \|\tilde{w}\|_{H^3} \|\phi\|_2 \|w\|_2 \right\}. \end{aligned} \tag{6.9}$$

Taking $\delta > 0$ such that $\delta \leq \delta_1$ with $\delta_1 = \min\{\frac{1}{2C_B\gamma}, \frac{1}{16CC_B^2\nu}, \frac{\nu}{16CC_B^2\|v_s\|_{C^1}^2}, \frac{\nu}{16CC_B\gamma^2}, \frac{1}{2C\tilde{\nu}}\}$, we deduce from (6.7)–(6.9) that

$$\begin{aligned} & \operatorname{Re} \lambda((u, u)) + \frac{1}{2}D[w] + \frac{\delta}{2}\|\phi\|_2^2 \\ & \leq C\left\{|\langle F, u \rangle| + \delta(|(f, \mathcal{B}\phi)| + |(\mathcal{B}f^0, w)|) \right. \\ & \quad \left. + \frac{\nu}{\gamma^4}\|\phi\|_{H^{-1}}^2 + \|\partial_{x_2}v_s^1\|_\infty\|w\|_2^2 + \frac{\|\tilde{w}\|_{H^3}}{\gamma}\|\phi\|_2\|u\|_2\right\}. \end{aligned} \quad (6.10)$$

By using the Poincaré inequalities, (6.4) follows from (6.10). As for (6.5), we have

$$\begin{aligned} & |\langle F, u \rangle| + \delta(|(f, \mathcal{B}\phi)| + |(\mathcal{B}f^0, w)|) \\ & \leq \frac{1}{\gamma^2}\|f^0\|_2\|\phi\|_2 + \|f\|_{H^{-1}}\|\nabla w\|_2 + \delta\left\{\|f\|_{H^{-1}}\|\nabla \mathcal{B}\phi\|_2 + \|\mathcal{B}f^0\|_2\|w\|_2\right\} \\ & \leq \frac{\delta}{4}\|\phi\|_2^2 + \frac{\nu}{16}\|\nabla w\|_2^2 + C\left\{\left(\frac{1}{\delta\gamma^2} + \frac{\nu}{\gamma^4}\right)\|f^0\|_2^2 + \frac{1}{\nu}\|f\|_{H^{-1}}^2\right\}, \\ & \frac{\|\tilde{w}\|_{H^3}}{\gamma}\|\phi\|_2\|u\|_2 \leq \frac{16C\|\tilde{w}\|_{H^3}}{\gamma^2}\left(1 + \frac{\|\tilde{w}\|_{H^3}}{\nu}\right)\|\phi\|_2^2 + \frac{\nu}{32C}\|\nabla w\|_2^2 \end{aligned}$$

and

$$\|\dot{\phi}_\lambda\|_2^2 = \|- \gamma^2 \operatorname{div} w + f^0\|_2^2 \leq 2\{\gamma^4\|\operatorname{div} w\|_2^2 + \|f^0\|_2^2\}.$$

Combining these inequalities with (6.10), we obtain (6.5). As for (6.6), we observe that

$$\partial_{x_1}^j(Lu)_d = \operatorname{div}(\partial_{x_1}^j\phi v_s + \gamma^2\partial_{x_1}^j w + \partial_{x_1}^j\phi\tilde{w} + [\partial_{x_1}^j, \tilde{w}]\phi)$$

and

$$\|\operatorname{div}([\partial_{x_1}^j, \tilde{w}]\phi)\|_2 + \|[\partial_{x_1}^j, \tilde{w}]\phi\|_2 \leq C\|\tilde{w}\|_{H^3}\|\phi\|_{H^j} \quad (j = 1, 2).$$

Therefore, as in the case of (6.4) and (6.5), we can obtain (6.6). This completes the proof. $\square$

Proposition 6.4. *There holds the inequality*

$$\begin{aligned} & \operatorname{Re} \lambda D[w] + \frac{1}{2}|\lambda|^2\|u\|_2^2 \\ & \leq C\left\{\|f\|_2^2 + \frac{\|v_s^1\|_\infty^2}{\gamma^2}\|\partial_{x_1}\phi\|_2^2 + \frac{\|\tilde{w}\|_{H^3}^2}{\gamma^2}\|\nabla\phi\|_2^2 + \frac{\nu^2}{\gamma^4}\|\phi\|_2^2 \right. \\ & \quad \left. + (\|v_s^1\|_{C^1}^2 + \gamma^2)\|\nabla w\|_2^2\right\}. \end{aligned} \quad (6.11)$$

Proof. We take the inner product of (6.1) with λu . Then the real part of the resulting equation yields

$$\begin{aligned} |\lambda|^2\|u\|_2^2 + \operatorname{Re} \lambda D[w] &= \operatorname{Re} \left\{ \frac{\bar{\lambda}}{\gamma^2}(f^0, \phi) + \bar{\lambda}(f, w) - \frac{\bar{\lambda}}{\gamma^2}(v_s^1\partial_{x_1}\phi, \phi) \right. \\ & \quad \left. - \frac{\bar{\lambda}}{\gamma^2}(\operatorname{div}(\phi\tilde{w}), \phi) - \bar{\lambda}(\operatorname{div} w, \phi) + \bar{\lambda}(\phi, \operatorname{div} w) \right. \\ & \quad \left. + \frac{\nu}{\gamma^2}\bar{\lambda}(\phi, w^1) - \bar{\lambda}(v_s^1\partial_{x_1}w, w) - \bar{\lambda}(\partial_{x_2}v_s^1w^3, w^1) \right\}. \end{aligned}$$

By a direct computation, the right-hand side is bounded by

$$\begin{aligned} & \frac{|\lambda|^2}{2} \|u\|_2^2 + C \left\{ \|f\|_2^2 + \frac{\|v_s^1\|_\infty^2}{\gamma^2} \|\partial_{x_1} \phi\|_2^2 + \frac{\|\tilde{w}\|_{H^3}^2}{\gamma^2} \|\nabla \phi\|_2^2 + \frac{\nu^2}{\gamma^4} \|\phi\|_2^2 \right. \\ & \left. + (\|v_s^1\|_{C^1}^2 + \gamma^2) \|\nabla w\|_2^2 \right\}. \end{aligned}$$

We thus obtain the desired estimate. This completes the proof. $\square$

Proposition 6.5. *Let j and k be integers satisfying $0 \leq j + k \leq 1$. Then there holds the inequality*

$$\begin{aligned} & \left| \operatorname{Re} \lambda + \frac{\gamma^2}{\nu + \tilde{\nu}} \right| \|\partial_{x_1}^j \partial_{x_2}^{k+1} \phi\|_2 \\ & \leq C \left\{ \|\partial_{x_1}^j \partial_{x_2}^{k+1} f^0\|_2 + \frac{\gamma^2}{\nu + \tilde{\nu}} \|\partial_{x_1}^j \partial_{x_2}^k f^2\|_2 \right. \\ & \quad + \|\partial_{x_2} v_s^1\|_{C^k} \|\partial_{x_1}^{j+1} \phi\|_{H^k} + \|\tilde{w}\|_{H^3} \|\phi\|_{H^{j+k+1}} \\ & \quad \left. + \frac{\gamma^2}{\nu + \tilde{\nu}} \left(|\lambda| \|\partial_{x_1}^j \partial_{x_2}^k w\|_2 + \nu \|\partial_{x_1}^{j+1} \partial_{x_2}^k \nabla w\|_2 + \|v_s^1\|_{C^k} \|\partial_{x_1}^{j+1} w\|_{H^k} \right) \right\}. \end{aligned} \quad (6.12)$$

Furthermore, if $\operatorname{Re} \lambda \geq -\frac{1}{2} \frac{\gamma^2}{\nu + \tilde{\nu}}$, then

$$\begin{aligned} & \|\partial_{x_1}^j \partial_{x_2}^{k+1} \dot{\phi}_\lambda\|_2 \\ & \leq C \left\{ \|\partial_{x_1}^j \partial_{x_2}^{k+1} f^0\|_2 + \frac{\gamma^2}{\nu + \tilde{\nu}} \|\partial_{x_1}^j \partial_{x_2}^k f^2\|_2 \right. \\ & \quad + \|\partial_{x_2} v_s^1\|_{C^k} \|\partial_{x_1}^{j+1} \phi\|_{H^k} + \|\tilde{w}\|_{H^3} \|\phi\|_{H^{j+k+1}} \\ & \quad \left. + \frac{\gamma^2}{\nu + \tilde{\nu}} \left(|\lambda| \|\partial_{x_1}^j \partial_{x_2}^k w\|_2 + \nu \|\partial_{x_1}^{j+1} \partial_{x_2}^k \nabla w\|_2 + \|v_s^1\|_{C^k} \|\partial_{x_1}^{j+1} w\|_{H^k} \right) \right\}. \end{aligned} \quad (6.13)$$

Proof. Applying $\partial_{x_1}^j \partial_{x_2}^{k+1}$ to the first component of equation (6.1), we have

$$\begin{aligned} & \lambda \partial_{x_1}^j \partial_{x_2}^{k+1} \phi + v_s^1 \partial_{x_1} \partial_{x_1}^j \partial_{x_2}^{k+1} \phi + \operatorname{div} ((\partial_{x_1}^j \partial_{x_2}^{k+1} \phi) \tilde{w}) + \gamma^2 \partial_{x_1}^j \partial_{x_2}^{k+2} w^2 \\ & = \partial_{x_1}^j \partial_{x_2}^{k+1} f^0 - \left\{ [\partial_{x_2}^{k+1}, v_s^1] \partial_{x_1}^{j+1} \phi + \operatorname{div} ([\partial_{x_1}^j \partial_{x_2}^{k+1}, \tilde{w}] \phi) + \gamma^2 \partial_{x_1}^{j+1} \partial_{x_2}^{k+1} w^1 \right\}. \end{aligned} \quad (6.14)$$

We also apply $\partial_{x_1}^j \partial_{x_2}^k$ to the third component of equation (6.1) to obtain

$$\begin{aligned} & -(\nu + \tilde{\nu}) \partial_{x_1}^j \partial_{x_2}^{k+2} w^2 + \partial_{x_1}^j \partial_{x_2}^{k+1} \phi \\ & = \partial_{x_1}^j \partial_{x_2}^k f^2 - \left\{ \lambda \partial_{x_1}^j \partial_{x_2}^k w^2 - \nu \partial_{x_1}^{j+2} \partial_{x_2}^k w^2 - \tilde{\nu} \partial_{x_1}^{j+1} \partial_{x_2}^{k+1} w^1 + \partial_{x_2}^k (v_s^1 \partial_{x_1}^{j+1} w^2) \right\}. \end{aligned} \quad (6.15)$$

By adding (6.14) and $\frac{\gamma^2}{\nu + \tilde{\nu}} \times (6.15)$ we obtain

$$\lambda \partial_{x_1}^j \partial_{x_2}^{k+1} \phi + \frac{\gamma^2}{\nu + \tilde{\nu}} \partial_{x_1}^j \partial_{x_2}^{k+1} \phi + v_s^1 \partial_{x_1} \partial_{x_1}^j \partial_{x_2}^{k+1} \phi + \operatorname{div} ((\partial_{x_1}^j \partial_{x_2}^{k+1} \phi) \tilde{w}) = H, \quad (6.16)$$

where

$$\begin{aligned} H = & \partial_{x_1}^j \partial_{x_2}^{k+1} f^0 + \frac{\gamma^2}{\nu + \tilde{\nu}} \partial_{x_1}^j \partial_{x_2}^k f^2 - \left\{ [\partial_{x_2}^{k+1}, v_s^1] \partial_{x_1}^{j+1} \phi + \operatorname{div} ([\partial_{x_1}^j \partial_{x_2}^{k+1}, \tilde{w}] \phi) \right\} \\ & - \frac{\gamma^2}{\nu + \tilde{\nu}} \left\{ \lambda \partial_{x_1}^j \partial_{x_2}^k w^2 - \nu \partial_{x_1}^{j+2} \partial_{x_2}^k w^2 + \nu \partial_{x_1}^{j+1} \partial_{x_2}^{k+1} w^1 + \partial_{x_2}^k (v_s^1 \partial_{x_1}^{j+1} w^2) \right\}. \end{aligned}$$

Taking the inner product of (6.16) with $\partial_{x_1}^j \partial_{x_2}^{k+1} \phi$, we have

$$\operatorname{Re} \lambda \|\partial_{x_1}^j \partial_{x_2}^{k+1} \phi\|_2^2 + \frac{\gamma^2}{\nu + \tilde{\nu}} \|\partial_{x_1}^j \partial_{x_2}^{k+1} \phi\|_2^2 = -\frac{1}{2} (\operatorname{div} \tilde{w}, |\partial_{x_1}^j \partial_{x_2}^{k+1} \phi|^2) + \operatorname{Re} (H, \partial_{x_1}^j \partial_{x_2}^{k+1} \phi),$$

from which estimate (6.12) is obtained. As for (6.13), we rewrite (6.16) as

$$\partial_{x_1}^j \partial_{x_2}^{k+1} \dot{\phi}_\lambda = -\frac{\gamma^2}{\nu + \tilde{\nu}} \partial_{x_1}^j \partial_{x_2}^{k+1} \phi + \tilde{H},$$

where

$$\tilde{H} = H + [\partial_{x_2}^{k+1}, v_s^1] \partial_{x_1}^{j+1} \phi + \operatorname{div} ([\partial_{x_1}^j \partial_{x_2}^{k+1}, \tilde{w}] \phi).$$

This, together with (6.12), yields (6.13). This completes the proof. $\square$

We next prepare the following estimate for the Stokes system to estimate the higher order derivatives.

Lemma 6.6. *Let $(\phi, w) \in H_{per,*}^{k+1} \times [H_{per}^{k+2} \cap H_{per,0}^1]$ be a solution of*

$$\operatorname{div} w = h^0,$$

$$-\Delta w + \nabla \phi = h$$

for $(h^0, h) \in H_{per,*}^{k+1} \times H_{per}^k$. Then

$$\|\partial_x^{k+2} w\|_2 + \|\partial_x^{k+1} \phi\|_2 \leq C \left\{ \|h^0\|_{H^{k+1}} + \|h\|_{H^k} \right\}.$$

See, e.g., [3, 13] for the proof. Applying Lemma 6.6 we have the following

Proposition 6.7. *Let j and k be integers satisfying $0 \leq j + k \leq 1$. Then*

$$\begin{aligned} & \|\partial_x^{k+2} \partial_{x_1}^j w\|_2 + \frac{1}{\nu} \|\partial_x^{k+1} \partial_{x_1}^j \phi\|_2 \\ & \leq C \left\{ \frac{\nu + \tilde{\nu}}{\nu \gamma^2} \|\partial_{x_1}^j f^0\|_{H^{k+1}} + \frac{1}{\nu} \|\partial_{x_1}^j f\|_{H^k} + \frac{\nu + \tilde{\nu}}{\nu \gamma^2} \|\partial_{x_1}^j \dot{\phi}_\lambda\|_{H^{k+1}} \right. \\ & \quad + \frac{|\lambda|}{\nu} \|\partial_{x_1}^j w\|_{H^k} + \frac{1}{\gamma^2} \|\partial_{x_1}^j \phi\|_{H^k} \\ & \quad \left. + \frac{1}{\nu} \|v_s^1\|_{C^k} \|\nabla \partial_{x_1}^j w\|_{H^k} + \frac{1}{\nu} \|\partial_{x_2} v_s^1\|_{C^k} \|\partial_{x_1}^j w\|_{H^k} \right\}. \end{aligned} \tag{6.17}$$

Proof. We apply $\partial_{x_1}^j$ to (6.1) and write the resulting equation as

$$\begin{aligned} \operatorname{div} \partial_{x_1}^j w &= \frac{1}{\gamma^2} \partial_{x_1}^j h^0, \\ -\Delta \partial_{x_1}^j w + \nabla \left(\frac{1}{\nu} \partial_{x_1}^j \phi \right) &= \frac{1}{\nu} \partial_{x_1}^j h, \end{aligned}$$

where

$$\begin{aligned} h^0 &= f^0 - \dot{\phi}_\lambda, \\ h &= f - \left\{ \lambda w - \frac{\tilde{\nu}}{\gamma^2} \nabla h^0 - \frac{\nu}{\gamma^2} \phi \mathbf{e}_1 + v_s^1 \partial_{x_1} w + \partial_{x_2} v_s^1 w^2 \mathbf{e}_1 \right\}. \end{aligned}$$

Applying Lemma 6.6 we have the desired estimate. This completes the proof. $\square$

The following proposition follows from the first equation of (6.1).

Proposition 6.8. *There holds the inequality*

$$|\lambda| \|\partial_x^k \phi\|_2 \leq C \left\{ \|\partial_x^k f^0\|_2 + \|v_s^1\|_{C^k} \|\partial_{x_1} \phi\|_{H^k} + \|\tilde{w}\|_{H^3} \|\partial_x \phi\|_{H^k} + \gamma^2 \|\partial_x^k \operatorname{div} w\|_2 \right\} \quad (6.18)$$

for $k = 0, 1$.

We are now in a position to prove Proposition 6.1.

Proof of Proposition 6.1. Observe first that $\|\partial_{x_1} g\|_{H^{-1}} \leq \|g\|_2$. We see from (6.4) with $k = 0$ that

$$\begin{aligned} &(\operatorname{Re} \lambda + \Lambda_0)^2 \|u\|_2^2 \\ &\leq C \left\{ \|F\|_2^2 + \|\partial_{x_2} v_s^1\|_\infty^2 \|w\|_2^2 + \frac{\nu^2}{\gamma^6} \|\phi\|_{H^{-1}}^2 + \frac{\|\tilde{w}\|_{H^3}^2}{\gamma^2} \|\phi\|_2^2 \right\}. \end{aligned} \quad (6.19)$$

We compute (6.19) + (6.5) + $b_1 \times (6.6)|_{j=1}$. Taking b_1 suitably small, we see that if $\operatorname{Re} \lambda > -\Lambda_0/2$, then

$$\begin{aligned} &\left(\operatorname{Re} \lambda + \frac{1}{2} \Lambda_0 \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \frac{1}{2} \Lambda_0 \right) \|\partial_{x_1} u\|_2^2 + \sum_{j=0}^1 (D[\partial_{x_1}^j w] + \|\partial_{x_1}^j \dot{\phi}_\lambda\|_2^2) \\ &\leq C \left\{ \|F\|_{H^1 \times L^2}^2 + \|\tilde{w}\|_{H^3} (1 + \|\tilde{w}\|_{H^3}) \|\phi\|_{H^1}^2 + \|w\|_2^2 + \|\phi\|_{H^{-1}}^2 \right\}. \end{aligned} \quad (6.20)$$

We next consider (6.20) + $b_2 \times (6.11)$. Taking b_2 suitably small, we see that there exists a positive number Λ_1 such that if $\operatorname{Re} \lambda > -\Lambda_1$, then

$$\begin{aligned} &\left(\operatorname{Re} \lambda + \Lambda_1 \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_1 \right) (\|\partial_{x_1} \phi\|_2^2 + \|\partial_x w\|_2^2) \\ &\quad + D[\partial_{x_1} w] + |\lambda|^2 \|u\|_2^2 + \sum_{j=0}^1 \|\partial_{x_1}^j \dot{\phi}_\lambda\|_2^2 \\ &\leq C \left\{ \|F\|_{H^1 \times L^2}^2 + \|\tilde{w}\|_{H^3} (1 + \|\tilde{w}\|_{H^3}) \|\phi\|_{H^1}^2 + \|w\|_2^2 + \|\phi\|_{H^{-1}}^2 \right\}. \end{aligned} \quad (6.21)$$

We then compute $(6.21) + b_3 \times \{(6.12)|_{j=k=0} + (6.13)|_{j=k=0}\}^2$. Taking b_3 suitably small, we see that there exists a positive number Λ_2 such that if $\operatorname{Re} \lambda > -\Lambda_2$, then

$$\begin{aligned} & \left(\operatorname{Re} \lambda + \Lambda_2 \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_2 \right) \|\partial_x u\|_2^2 + D[\partial_{x_1} w] + |\lambda|^2 \|u\|_2^2 + \|\dot{\phi}_\lambda\|_{H^1}^2 \\ & \leq C \left\{ \|F\|_{H^1 \times L^2}^2 + \|\tilde{w}\|_{H^3} (1 + \|\tilde{w}\|_{H^3}) \|\phi\|_{H^1}^2 + \|w\|_2^2 + \|\phi\|_{H^{-1}}^2 \right\}. \end{aligned} \quad (6.22)$$

We next compute $(6.22) + b_4 \times \{(6.17)|_{j=k=0}\}^2$. We take b_4 suitably small to see that there exists a positive number Λ_3 such that if $\operatorname{Re} \lambda > -\Lambda_3$, then

$$\begin{aligned} & \left(\operatorname{Re} \lambda + \Lambda_3 \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_3 \right) \|\partial_x u\|_2^2 + \|\partial_x^2 w\|_2^2 + |\lambda|^2 \|u\|_2^2 + \|\dot{\phi}_\lambda\|_{H^1}^2 \\ & \leq C \left\{ \|F\|_{H^1 \times L^2}^2 + \|\tilde{w}\|_{H^3} (1 + \|\tilde{w}\|_{H^3}) \|\phi\|_{H^1}^2 + \|w\|_2^2 + \|\phi\|_{H^{-1}}^2 \right\}. \end{aligned} \quad (6.23)$$

This shows (6.2).

Let us prove (6.3). We compute $(6.23) + b_5 \times (6.6)|_{j=2}$. Taking b_5 suitably small, we see that there exists a positive number Λ_4 such that if $\operatorname{Re} \lambda > -\Lambda_4$, then

$$\begin{aligned} & \left(\operatorname{Re} \lambda + \Lambda_4 \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_4 \right) (\|\partial_x u\|_2^2 + \|\partial_{x_1}^2 u\|_2^2) \\ & \quad + \|\partial_x^2 w\|_2^2 + \|\nabla \partial_{x_1}^2 w\|_2^2 + |\lambda|^2 \|u\|_2^2 + \|\dot{\phi}_\lambda\|_{H^1}^2 + \|\partial_{x_1}^2 \dot{\phi}_\lambda\|_2^2 \\ & \leq C \left\{ \|F\|_{H^2 \times H^1}^2 + \|\tilde{w}\|_{H^3} (1 + \|\tilde{w}\|_{H^3}) \|\phi\|_{H^2}^2 + \|w\|_2^2 + \|\phi\|_{H^{-1}}^2 \right\}. \end{aligned} \quad (6.24)$$

We next compute $(6.24) + b_6 \times (6.4)|_{k=1}$. Taking b_6 suitably small, we see that there exists a positive number Λ_5 such that if $\operatorname{Re} \lambda > -\Lambda_5$, then

$$\begin{aligned} & \left(\operatorname{Re} \lambda + \Lambda_5 \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_5 \right) (\|\partial_x u\|_2^2 + \|\partial_{x_1}^2 u\|_2^2 + |\lambda|^2 \|u\|_2^2) \\ & \quad + \|\partial_x^2 w\|_2^2 + \|\nabla \partial_{x_1}^2 w\|_2^2 + |\lambda|^2 \|\nabla w\|_2^2 + \|\dot{\phi}_\lambda\|_{H^1}^2 + \|\partial_{x_1}^2 \dot{\phi}_\lambda\|_2^2 \\ & \leq C \left\{ \|F\|_{H^2 \times H^1}^2 + |\lambda|^2 \|F\|_2^2 + \|\tilde{w}\|_{H^3} (1 + \|\tilde{w}\|_{H^3}) (\|\phi\|_{H^2}^2 + |\lambda|^2 \|\phi\|_2^2) \right. \\ & \quad \left. + \|w\|_2^2 + \|\phi\|_{H^{-1}}^2 \right\}. \end{aligned} \quad (6.25)$$

We next consider $(6.25) + b_7 \times \{(6.12)|_{j=1,k=0} + (6.13)|_{j=1,k=0}\}^2$. Taking b_7 suitably small, we see that there exists a positive number Λ_6 such that if $\operatorname{Re} \lambda > -\Lambda_6$, then

$$\begin{aligned} & \left(\operatorname{Re} \lambda + \Lambda_6 \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_6 \right) (\|\partial_x u\|_2^2 + \|\partial_{x_1}^2 u\|_2^2 + \|\partial_{x_1} \partial_{x_2} \phi\|_2^2 + |\lambda|^2 \|u\|_2^2) \\ & \quad + \|\partial_x^2 w\|_2^2 + \|\nabla \partial_{x_1}^2 w\|_2^2 + |\lambda|^2 \|\nabla w\|_2^2 + \|\dot{\phi}_\lambda\|_{H^1}^2 + \|\nabla \partial_{x_1} \dot{\phi}_\lambda\|_2^2 \\ & \leq C \left\{ \|F\|_{H^2 \times H^1}^2 + |\lambda|^2 \|F\|_2^2 + \|\tilde{w}\|_{H^3} (1 + \|\tilde{w}\|_{H^3}) (\|\phi\|_{H^2}^2 + |\lambda|^2 \|\phi\|_2^2) \right. \\ & \quad \left. + \|w\|_2^2 + \|\phi\|_{H^{-1}}^2 \right\}. \end{aligned} \quad (6.26)$$

It then follows from (6.26) + $b_8 \times \{(6.17)|_{j=1,k=0}\}^2$ with suitably small b_8 that there exists a positive number Λ_7 such that if $\operatorname{Re} \lambda > -\Lambda_7$, then

$$\begin{aligned}
& \left(\operatorname{Re} \lambda + \Lambda_7 \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_7 \right) (\|\partial_x u\|_2^2 + \|\partial_{x_1}^2 u\|_2^2 + \|\partial_{x_1} \partial_{x_2} \phi\|_2^2 + |\lambda|^2 \|u\|_2^2) \\
& + \|\partial_x^2 w\|_2^2 + \|\partial_x^2 \partial_{x_1} w\|_2^2 + |\lambda|^2 \|\nabla w\|_2^2 + \|\dot{\phi}_\lambda\|_{H^1}^2 + \|\nabla \partial_{x_1} \dot{\phi}_\lambda\|_2^2 \\
& \leq C \left\{ \|F\|_{H^2 \times H^1}^2 + |\lambda|^2 \|F\|_2^2 + \|\tilde{w}\|_{H^3} (1 + \|\tilde{w}\|_{H^3}) (\|\phi\|_{H^2}^2 + |\lambda|^2 \|\phi\|_2^2) \right. \\
& \quad \left. + \|w\|_2^2 + \|\phi\|_{H^{-1}}^2 \right\}.
\end{aligned} \tag{6.27}$$

We then compute (6.27) + $b_9 \times \{(6.12)|_{j=0,k=1} + (6.13)|_{j=0,k=1}\}^2$ and take b_9 suitably small so that there exists a positive number Λ_8 such that if $\operatorname{Re} \lambda > -\Lambda_8$, then

$$\begin{aligned}
& \left(\operatorname{Re} \lambda + \Lambda_8 \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_8 \right) (\|\partial_x u\|_2^2 + \|\partial_x^2 \phi\|_2^2 + \|\partial_{x_1}^2 w\|_2^2 + |\lambda|^2 \|u\|_2^2) \\
& + \|\partial_x^2 w\|_2^2 + \|\partial_x^2 \partial_{x_1} w\|_2^2 + |\lambda|^2 \|\nabla w\|_2^2 + \|\dot{\phi}_\lambda\|_{H^2}^2 \\
& \leq C \left\{ \|F\|_{H^2 \times H^1}^2 + |\lambda|^2 \|F\|_2^2 + \|\tilde{w}\|_{H^3} (1 + \|\tilde{w}\|_{H^3}) (\|\phi\|_{H^2}^2 + |\lambda|^2 \|\phi\|_2^2) \right. \\
& \quad \left. + \|w\|_2^2 + \|\phi\|_{H^{-1}}^2 \right\}.
\end{aligned} \tag{6.28}$$

Finally, consider (6.28) + $b_{10} \times \{(6.17)|_{j=0,k=1} + (6.18)|_{k=1}\}^2$. Taking b_{10} suitably small, we deduce that there exists a positive number Λ_9 such that if $\operatorname{Re} \lambda > -\Lambda_9$, then

$$\begin{aligned}
& \left(\operatorname{Re} \lambda + \Lambda_9 \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_9 \right) (\|\partial_x u\|_2^2 + \|\partial_x^2 \phi\|_2^2 + \|\partial_{x_1}^2 w\|_2^2 + |\lambda|^2 \|u\|_2^2) \\
& + \|\partial_x^2 w\|_2^2 + \|\partial_x^3 w\|_2^2 + |\lambda|^2 \|\nabla u\|_2^2 + \|\dot{\phi}_\lambda\|_{H^2}^2 \\
& \leq C \left\{ \|F\|_{H^2 \times H^1}^2 + |\lambda|^2 \|F\|_2^2 + \|\tilde{w}\|_{H^3} (1 + \|\tilde{w}\|_{H^3}) (\|\phi\|_{H^2}^2 + |\lambda|^2 \|\phi\|_2^2) \right. \\
& \quad \left. + \|w\|_2^2 + \|\phi\|_{H^{-1}}^2 \right\}.
\end{aligned} \tag{6.29}$$

We thus obtain (6.3). This completes the proof. $\square$

6.2 A priori estimates

We consider

$$\lambda u + \mathcal{L}(\varepsilon, \tilde{\sigma}, \tilde{V})u = F, \quad u \in QX^\ell \quad (\ell = 1, 2), \tag{6.30}$$

where $F \in Q(H_{per}^\ell \times H_{per}^{\ell-1})$ and

$$\mathcal{L}(\varepsilon, \tilde{\sigma}, \tilde{V}) = \mathcal{L}_{c_0,0} - \varepsilon \tilde{\sigma} Q \partial_{x_1} + \varepsilon Q N[V_1 + \varepsilon \tilde{V}]$$

with $\tilde{\sigma} \in \mathbb{R}$ and $\tilde{V} \in X^2$ satisfying $|\tilde{\sigma}| + \|\tilde{V}\|_{X^2} \leq M$. In this subsection we show the a priori estimates for solution u of (6.30).

We show the following a priori estimates.

Proposition 6.9. *Let $M > 0$ and assume that $|\tilde{\sigma}| + \|\tilde{V}\|_{X^2} \leq M$. Then there exist $\varepsilon_3 > 0$, $r_0 > 0$, $\Lambda > 0$ and $\{\lambda_j\}_{j=1}^K \subset \mathbb{C}$ with $|\lambda_j| \geq 2r_0$ such that if $|\varepsilon| \leq \varepsilon_3$ and*

$$\lambda \in \Sigma_0 = \{\lambda \in \mathbb{C}; \operatorname{Re} \lambda \geq -\Lambda, |\lambda - \lambda_j| \geq r_0, j = 1, \dots, K\},$$

the solution $u \in QX^1$ of (6.30) satisfies the estimate

$$\left(\operatorname{Re} \lambda + \Lambda\right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda\right) \|\partial_x u\|_2^2 + \|\partial_x^2 w\|_2^2 + |\lambda|^2 \|u\|_2^2 \leq C \|F\|_{H^1 \times L^2}^2 \quad (6.31)$$

uniformly for $\lambda \in \Sigma_0$. In addition, if $u \in QX^2$, then

$$\begin{aligned} & \left(\operatorname{Re} \lambda + \Lambda\right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda\right) (\|\partial_x u\|_2^2 + \|\partial_x^2 \phi\|_2^2 + |\lambda|^2 \|u\|_2^2) \\ & + \|\partial_x^2 w\|_{H^1}^2 + |\lambda|^2 \|\nabla u\|_2^2 \leq C \left\{ \|F\|_{H^2 \times H^1}^2 + |\lambda|^2 \|F\|_2^2 \right\} \end{aligned} \quad (6.32)$$

uniformly for $\lambda \in \Sigma_0$.

We note that $0 \in \Sigma_0$.

Proof. We first introduce frequency cut off operators. We expand $f \in L_{per}^2$ into the Fourier series $f = \sqrt{\frac{\alpha}{2\pi}} \sum_{k \in \mathbb{Z}} f_k(x_2) e^{i\alpha k x_1}$. We define $\Pi_{\leq N}$ and $\Pi_{\geq N}$ by

$$\Pi_{\leq N} f = \sqrt{\frac{\alpha}{2\pi}} \sum_{|k| \leq N} f_k(x_2) e^{i\alpha k x_1}$$

and

$$\Pi_{\geq N} f = \sqrt{\frac{\alpha}{2\pi}} \sum_{|k| \geq N} f_k(x_2) e^{i\alpha k x_1}$$

respectively. $\Pi_{< N}$ and $\Pi_{> N}$ are defined similarly. Observe that they are orthogonal projections on L_{per}^2 and

$$\|w\|_2 \leq \frac{1}{\alpha N} \|\nabla w\|_2, \quad \|\phi\|_{H^{-1}} \leq \frac{1}{\alpha N} \|\phi\|_2 \quad (6.33)$$

for $w \in \Pi_{\geq N} H_{per}^1$ and $\phi \in \Pi_{\geq N} L_{per}^2$ with $N \geq 1$.

We first prove (6.31). We write (6.30) as

$$\lambda u + \mathcal{L}_{c_0,0} u - \varepsilon \tilde{\sigma} Q J u + \varepsilon Q N[\tilde{w}] u = F. \quad (6.34)$$

Here $\tilde{w}$ is the function defined by

$$\tilde{w} = -\tilde{\sigma} e_1 + W_1 + \varepsilon \tilde{W}$$

with W_1 and $\tilde{W}$ being the velocity components of V_1 and $\tilde{V}$ respectively; and Ju and $N[\tilde{w}]u$ are defined by $Ju = {}^\top(0, \partial_{x_1} w)$ and $N[\tilde{w}]u = {}^\top(\operatorname{div}(\phi \tilde{w}), 0)$ for $u = {}^\top(\phi, w)$, respectively. Since $Q = I - P$, (6.34) is rewritten as

$$\lambda u + \mathcal{L}_{c_0,0} u + \varepsilon N[\tilde{w}]u = F + \varepsilon \tilde{\sigma} Q J u + \varepsilon P N[\tilde{w}]u. \quad (6.35)$$

Note that

$$\|QJu\|_{H^{\ell+1} \times H^\ell} \leq C \|\partial_{x_1} w\|_{H^\ell} \quad (\ell = 0, 1)$$

and

$$\|PN[\tilde{w}]u\|_{H^2 \times H^1} \leq C \|N[\tilde{w}]u\|_2 \leq C \|\tilde{w}\|_{H^3} \|\phi\|_{H^1}.$$

Applying (6.2) with v_s , $\tilde{w}$ and F replaced by $v_s - c_0 \mathbf{e}_1$, $\varepsilon \tilde{w}$ and $F + \varepsilon \tilde{\sigma} QJu + \varepsilon PN[\tilde{w}]u$, respectively, we see that

$$\begin{aligned} & \left(\operatorname{Re} \lambda + \Lambda \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda \right) \|\partial_x u\|_2^2 + \|\partial_x^2 w\|_2^2 + |\lambda|^2 \|u\|_2^2 \\ & \leq C \left\{ \|F\|_{H^1 \times L^2}^2 + |\varepsilon|^2 |\tilde{\sigma}|^2 \|QJu\|_{H^1 \times L^2}^2 + |\varepsilon|^2 \|PN[\tilde{w}]u\|_{H^1 \times L^2}^2 + \|w\|_2^2 \right. \\ & \quad \left. + \|\phi\|_{H^{-1}}^2 + |\varepsilon| \|w\|_{H^3} (1 + |\varepsilon| \|\tilde{w}\|_{H^3}) \|\phi\|_{H^1}^2 \right\} \\ & \leq C \left\{ \|F\|_{H^1 \times L^2}^2 + \|u_{<N}\|_2^2 + \frac{1}{\alpha^2 N^2} \left(\|\nabla w_{\geq N}\|_2^2 + \|\phi_{\geq N}\|_2^2 \right) \right. \\ & \quad \left. + |\varepsilon|^2 |\tilde{\sigma}|^2 \|\partial_{x_1} w\|_2 + |\varepsilon| \|\tilde{w}\|_{H^3} (1 + |\varepsilon| \|\tilde{w}\|_{H^3}) \|\phi\|_{H^1}^2 \right\}. \end{aligned}$$

It then follows that there exists $N_0 \in \mathbb{N}$ such that the inequality

$$\begin{aligned} & \left(\operatorname{Re} \lambda + \Lambda_{10} \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_{10} \right) \|\partial_x u\|_2^2 + \|\partial_x^2 w\|_2^2 + |\lambda|^2 \|u\|_2^2 \\ & \leq C \left\{ \|F\|_{H^1 \times L^2}^2 + \|u_{<N}\|_2^2 + |\varepsilon|^2 |\tilde{\sigma}|^2 \|\partial_{x_1} w\|_2^2 \right. \\ & \quad \left. + |\varepsilon| \|\tilde{w}\|_{H^3} (1 + |\varepsilon| \|\tilde{w}\|_{H^3}) \|\phi\|_{H^1}^2 \right\} \end{aligned} \tag{6.36}$$

holds with $\Lambda_{10} = \frac{1}{2}\Lambda$ uniformly for $N \geq N_0$.

To proceed further, we apply the following result on the spectral distribution proved by Iooss and Padula [6].

Lemma 6.10. ([6]) *There exists a constant $\tilde{\Lambda} > 0$ with $\tilde{\Lambda} \leq \Lambda_{10}$ such that*

$$\sigma(-\mathcal{L}_{c_0,0}) \cap \{\lambda; \operatorname{Re} \lambda \geq -\tilde{\Lambda}\} = \{\lambda_j\}_{j=0}^K,$$

where λ_j ($j = 0, 1, \dots, K$) are eigenvalues of $-\mathcal{L}_{c_0,0}$ with finite multiplicities.

We may assume $N_0 \geq 2$. Furthermore, by assumption (4.1), we may assume that $\lambda_0 = 0$ and $\lambda_j \neq 0$ for $j = 1, \dots, K$. By Lemma 6.10, we see that there is a positive number r_0 such that

$$|\lambda_j - \lambda_k| \geq 4r_0, \quad j \neq k, \quad j, k = 0, 1, \dots, K,$$

$$\rho(-\mathcal{L}_{c_0,0}|_{\Pi_{\leq N_0} QX}) \supset \Sigma_0 \equiv \{\lambda; \operatorname{Re} \lambda \geq -\tilde{\Lambda}, |\lambda - \lambda_j| \geq r_0, j = 1, \dots, K\}$$

and

$$(|\lambda| + 1) \|(\lambda + \mathcal{L}_{c_0,0}|_{\Pi_{\leq N_0} QX})^{-1} F\|_2 \leq C \|F\|_2 \tag{6.37}$$

uniformly for $\lambda \in \Sigma_0$. Note that $\Sigma_0 \ni 0$ since $\lambda_0 = 0$.

Let us estimate $\|u_{<N}\|_2$. Applying $\Pi_{<N_0}$ to (6.35), we have

$$\lambda u_{<N_0} + \mathcal{L}_{c_0,0} u_{<N_0} = F_{<N_0} - \varepsilon \Pi_{<N_0} N[\tilde{w}]u + \varepsilon \tilde{\sigma} QJu + \varepsilon PN[\tilde{w}]u.$$

Here we have used the fact $\Pi_{<N_0} P = P$. It then follows from (6.37) that

$$\begin{aligned} \|u_{<N_0}\|_2 &\leq C \left\{ \|F_{<N_0}\|_2 + |\varepsilon| \|\Pi_{<N_0} N[\tilde{w}]u\|_2 + |\varepsilon| |\tilde{\sigma}| \|QJu\|_2 + |\varepsilon| \|PN[\tilde{w}]u\|_2 \right\} \\ &\leq C \left\{ \|F\|_2 + |\varepsilon| |\tilde{\sigma}| \|\partial_{x_1} w\|_2 + |\varepsilon| \|\tilde{w}\|_\infty \|\nabla \phi\|_2 \right\} \\ &\leq C \left\{ \|F\|_2 + |\varepsilon| |\tilde{\sigma}| \|\partial_{x_1} w\|_2 + |\varepsilon| \|\tilde{w}\|_{H^3} \|\nabla \phi\|_2 \right\}. \end{aligned} \quad (6.38)$$

We see from (6.36) and (6.38) that

$$\begin{aligned} &\left(\operatorname{Re} \lambda + \tilde{\Lambda} \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \tilde{\Lambda} \right) \|\partial_x u\|_2^2 + \|\partial_x^2 w\|_2^2 + |\lambda|^2 \|u\|_2^2 \\ &\leq C \left\{ \|F\|_{H^1 \times L^2}^2 + |\varepsilon|^2 |\tilde{\sigma}|^2 \|\partial_{x_1} w\|_2^2 + |\varepsilon| \|\tilde{w}\|_{H^3} (1 + |\varepsilon| \|\tilde{w}\|_{H^3}) \|\phi\|_{H^1}^2 \right\} \end{aligned} \quad (6.39)$$

uniformly for $\lambda \in \Sigma_0$. Since $|\tilde{\sigma}| \leq M$ and $\|\tilde{w}\|_{H^3} \leq C(\|V_1\|_{H^3} + M)$, we conclude that there exists $\varepsilon_3 > 0$ such that if $|\varepsilon| \leq \varepsilon_3$, then

$$\left(\operatorname{Re} \lambda + \Lambda_{11} \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_{11} \right) \|\partial_x u\|_2^2 + \|\partial_x^2 w\|_2^2 + |\lambda|^2 \|u\|_2^2 \leq C \|F\|_{H^1 \times L^2}^2 \quad (6.40)$$

uniformly for $\lambda \in \Sigma_0$ with $\Lambda_{11} = \frac{1}{2} \tilde{\Lambda}$. This shows (6.31).

As for (6.32), by (6.3) and (6.40), we have

$$\begin{aligned} &\left(\operatorname{Re} \lambda + \Lambda_{11} \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_{11} \right) (\|\partial_x u\|_2^2 + \|\partial_x^2 \phi\|_2^2 + |\lambda|^2 \|u\|_2^2) \\ &\quad + \|\partial_x^2 w\|_{H^1}^2 + |\lambda|^2 \|\nabla u\|_2^2 \\ &\leq C \left\{ \|F\|_{H^2 \times H^1}^2 + |\lambda|^2 \|F\|_2^2 + |\varepsilon|^2 |\tilde{\sigma}|^2 (\|\partial_{x_1} w\|_{H^1}^2 + |\lambda|^2 \|\partial_{x_1} w\|_2^2) \right. \\ &\quad \left. + |\varepsilon| \|\tilde{w}\|_{H^3} (1 + |\varepsilon| \|\tilde{w}\|_{H^3}) (\|\nabla \phi\|_{H^1}^2 + |\lambda|^2 \|\phi\|_{H^1}^2) \right\} \end{aligned}$$

uniformly for $\lambda \in \Sigma_0$. Therefore, if $|\varepsilon| \leq \varepsilon_3$ (by taking ε_3 smaller if necessary), then

$$\begin{aligned} &\left(\operatorname{Re} \lambda + \Lambda_{12} \right)^2 \|u\|_2^2 + \left(\operatorname{Re} \lambda + \Lambda_{12} \right) (\|\partial_x u\|_2^2 + \|\partial_x^2 \phi\|_2^2 + |\lambda|^2 \|u\|_2^2) \\ &\quad + \|\partial_x^2 w\|_{H^1}^2 + |\lambda|^2 \|\nabla u\|_2^2 \\ &\leq C \left\{ \|F\|_{H^2 \times H^1}^2 + |\lambda|^2 \|F\|_2^2 \right\} \end{aligned} \quad (6.41)$$

uniformly for $\lambda \in \Sigma_0$ with $\Lambda_{12} = \frac{1}{2} \Lambda_{11}$. This completes the proof. $\square$

6.3 Invertibility

We finally prove the invertibility of $\mathcal{L}(\varepsilon, \tilde{\sigma}, \tilde{V})$. We first show the existence of solution of (6.30) in QX^ℓ ($\ell = 1, 2$) for sufficiently large $\lambda > 0$.

Proposition 6.11. *Let $\ell = 1, 2$ and assume that $|\varepsilon| \leq \varepsilon_1$. There exists $\mu_0 > 0$ such that if $\lambda \geq \mu_0$, then for any $F = {}^\top(f^0, f) \in Q(H_{per,*}^\ell \times H_{per}^{\ell-1})$, there exists a unique solution $u = {}^\top(\phi, w) \in QX^\ell$ of (6.30), and $u = {}^\top(\phi, w)$ satisfies*

$$\lambda \|\phi\|_{H^\ell} + \sum_{j=0}^{\ell+1} \lambda^{\frac{\ell+1-j}{2}} \|\partial_x^j w\|_2 \leq C \|f^0\|_{H^\ell} + C \sum_{j=0}^{\ell-1} \lambda^{\frac{\ell-1-j}{2}} \|\partial_x^j f\|_2.$$

Proof. We consider (6.35) instead of (6.30). Suppose that $u \in X^\ell$ is a solution of (6.35). Then

$$\lambda u + \mathcal{L}_{c_0,0} u = \varepsilon \tilde{\sigma} QJu - \varepsilon QN[\tilde{w}]u + F.$$

Applying P to both sides, we see that $\lambda Pu = 0$. Since $\lambda > 0$, we have $Pu = 0$, and hence $u \in QX^\ell$. Therefore, it suffices to show the existence of solution of (6.35) in X^ℓ .

Hereafter in the proof, we simply denote the density and velocity components of $PJu = P^\top(0, \partial_{x_1} w)$ ($u = {}^\top(\phi, w)$) by $P_d(\partial_{x_1} w)$ and $P_v(\partial_{x_1} w)$ respectively, i.e.,

$$PJ u = P^\top(0, \partial_{x_1} w) = {}^\top(P_d(\partial_{x_1} w), P_v(\partial_{x_1} w)),$$

and likewise, we denote the density and velocity components of $PN[\tilde{w}]u = P^\top(\operatorname{div}(\phi \tilde{w}), 0)$ with $u = {}^\top(\phi, w)$ by $P_d(\operatorname{div}(\phi \tilde{w}))$ and $P_v(\operatorname{div}(\phi \tilde{w}))$ respectively, i.e.,

$$PN[\tilde{w}]u = P^\top(\operatorname{div}(\phi \tilde{w}), 0) = {}^\top(P_d(\operatorname{div}(\phi \tilde{w})), P_v(\operatorname{div}(\phi \tilde{w}))).$$

We write (6.35) as

$$\lambda \phi + \operatorname{div}((\tilde{v}_s + \varepsilon \tilde{w})\phi) = -\varepsilon \tilde{\sigma} P_d(\partial_{x_1} w) + \varepsilon P_d(\operatorname{div}(\phi \tilde{w})) - \gamma^2 \operatorname{div} w + f^0, \quad (6.42)$$

$$\lambda w + Aw = B\phi - \varepsilon \tilde{\sigma} P_v(\partial_{x_1} w) + \varepsilon P_v(\operatorname{div}(\phi \tilde{w})) + f. \quad (6.43)$$

Here $\tilde{v}_s = v_s - (c_0 + \varepsilon \tilde{\sigma})\mathbf{e}_1$; A denotes the elliptic operator on L_{per}^2 defined by

$$Aw = -\nu \Delta w - \tilde{\nu} \nabla \operatorname{div} w + \tilde{v}_s^1 \partial_{x_1} w + (\partial_{x_2} \tilde{v}_s^1)(w \cdot \mathbf{e}_2)\mathbf{e}_1$$

with domain $D(A) = H_{per}^2 \cap H_{per,0}^1$; B is the operator on H_{per}^1 defined by

$$B\phi = -\nabla \phi + \frac{\nu}{\gamma^2} \phi \mathbf{e}_1.$$

By [4], there exists $\mu_1 > 0$ such that if $\lambda \geq \mu_1$, then, for any $f^0 \in H_{per,*}^\ell$, there exists a unique solution $\Phi \in H_{per,*}^\ell$ of

$$\lambda \Phi + \operatorname{div}(\Phi(\tilde{v}_s + \varepsilon \tilde{w})) = f^0, \quad (6.44)$$

and Φ satisfies the estimate

$$\|\Phi\|_{H^\ell} \leq \frac{C}{\lambda} \|f^0\|_{H^\ell}.$$

We denote by Φ_λ the solution map $f^0 \mapsto \Phi$ for (6.44). Then Φ_λ is a bounded linear operator on $H_{per,*}^\ell$ and

$$\|\Phi_\lambda f^0\|_{H^\ell} \leq \frac{C}{\lambda} \|f^0\|_{H^\ell}. \quad (6.45)$$

It then follows that (6.42) is equivalent to

$$\Psi_\lambda \phi = \Phi_\lambda(-\varepsilon \tilde{\sigma} P_d(\partial_{x_1} w) - \gamma^2 \operatorname{div} w + f^0), \quad (6.46)$$

where Ψ_λ is the operator defined by

$$\Psi_\lambda \phi = \phi - \varepsilon \Phi_\lambda P_d(\operatorname{div}(\phi \tilde{w})).$$

To solve (6.46), we show that the map $\Psi_\lambda : H_{per,*}^\ell \rightarrow H_{per,*}^\ell$ has a bounded inverse. By (6.45), we have

$$\|\varepsilon \Phi_\lambda P_d(\operatorname{div}(\phi \tilde{w}))\|_{H^\ell} \leq \frac{|\varepsilon|C}{\lambda} \|\operatorname{div}(\phi \tilde{w})\|_2 \leq \frac{\varepsilon_3 C}{\lambda} (\|V_1\|_{C^1} + M) \|\phi\|_{H^1}.$$

This implies that if $\lambda \geq \mu_2 = \max\{\mu_1, 2C\varepsilon_3(\|V_1\|_{C^1} + M)\}$, then $\|\varepsilon \Phi_\lambda P_d(\operatorname{div}(\phi \tilde{w}))\|_{H^\ell} \leq \frac{1}{2} \|\phi\|_{H^\ell}$ for $\ell = 1, 2$, and hence, $\Psi_\lambda : H_{per,*}^\ell \rightarrow H_{per,*}^\ell$ has a bounded inverse Ψ_λ^{-1} , and Ψ_λ^{-1} satisfies

$$\|\Psi_\lambda^{-1} \phi\|_{H^\ell} \leq 2 \|\phi\|_{H^\ell}. \quad (6.47)$$

In terms of Ψ_λ^{-1} , the solution ϕ of (6.46) is written as

$$\phi = \Psi_\lambda^{-1} \Phi_\lambda(-\varepsilon \tilde{\sigma} P_d(\partial_{x_1} w) - \gamma^2 \operatorname{div} w + f^0), \quad (6.48)$$

and, by (6.45) and (6.47), ϕ satisfies

$$\|\phi\|_{H^\ell} \leq \frac{C}{\lambda} \left\{ \|w\|_{H^{\ell+1}} + \|f^0\|_{H^\ell} \right\}. \quad (6.49)$$

From (6.43) and (6.48), we have

$$(\lambda + A)w = B_1[\tilde{w}] \Psi_\lambda^{-1} \Phi_\lambda(-\varepsilon \tilde{\sigma} P_d(\partial_{x_1} w) - \gamma^2 \operatorname{div} w + f^0) - \varepsilon \tilde{\sigma} P_v(\partial_{x_1} w) + f$$

with

$$B_1[\tilde{w}] \phi = B\phi + \varepsilon P_v(\operatorname{div}(\phi \tilde{w})).$$

This is equivalent to

$$(I - \Gamma_\lambda)w = (\lambda + A)^{-1} (B_1[\tilde{w}] \Psi_\lambda^{-1} \Phi_\lambda f^0 + f), \quad (6.50)$$

where Γ_λ is the operator defined by

$$\Gamma_\lambda w = (\lambda + A)^{-1} (B_1[\tilde{w}] \Psi_\lambda^{-1} \Phi_\lambda(-\varepsilon \tilde{\sigma} P_d(\partial_{x_1} w) - \gamma^2 \operatorname{div} w) - \varepsilon \tilde{\sigma} P_v(\partial_{x_1} w)).$$

Since A is strongly elliptic, there exists $\mu_3 > 0$ such that if $\lambda \geq \mu_3$, then $(\lambda + A)^{-1}f \in H_{per}^{\ell+1} \cap H_{per,0}^1$ for $f \in H^{\ell-1}$ and it holds that

$$\sum_{j=0}^{\ell+1} \lambda^{\frac{\ell+1-j}{2}} \|\partial_x^j (\lambda + A)^{-1}f\|_2 \leq C \sum_{j=0}^{\ell-1} \lambda^{\frac{\ell-1-j}{2}} \|\partial_x^j f\|_2. \quad (6.51)$$

Furthermore, for $j = 1, 2$, we have

$$\|B_1[\tilde{w}]\phi\|_{H^{j-1}} \leq C \left\{ \|\phi\|_{H^j} + |\varepsilon| \|\operatorname{div}(\phi \tilde{w})\|_2 \right\} \leq C \|\phi\|_{H^j}. \quad (6.52)$$

We now introduce the norm $|||w|||_{(\lambda)} = \sum_{j=0}^{\ell+1} \lambda^{\frac{\ell+1-j}{2}} \|\partial_x^j w\|_2$ of $H_{per}^{\ell+1}$ and show that the map $\Gamma_\lambda : H_{per}^{\ell+1} \cap H_{per,0}^1 \rightarrow H_{per}^{\ell+1} \cap H_{per,0}^1$ has a bounded inverse Γ_λ^{-1} . By (6.49) with $f^0 = 0$, (6.51) and (6.52), we see that if $\lambda \geq \max\{\mu_3, 1\}$, then

$$|||\Gamma_\lambda w|||_{(\lambda)} \leq C \sum_{j=0}^{\ell-1} \lambda^{\frac{\ell-1-j}{2}} \frac{1}{\lambda} \|w\|_{H^{j+2}} \leq \frac{C}{\lambda} \sum_{j=0}^{\ell+1} \lambda^{\frac{\ell+1-j}{2}} \|\partial_x^j w\|_2.$$

Therefore, there exists $\mu_4 > 0$ such that if $\lambda \geq \mu_4$, then

$$|||\Gamma_\lambda w|||_{(\lambda)} \leq \frac{1}{2} |||w|||_{(\lambda)},$$

and hence, $I - \Gamma_\lambda$ has a bounded inverse $(I - \Gamma_\lambda)^{-1}$, and $(I - \Gamma_\lambda)^{-1}$ satisfies the estimate

$$|||(I - \Gamma_\lambda)^{-1}f|||_{(\lambda)} \leq 2 |||f|||_{(\lambda)}.$$

In terms of $(I - \Gamma_\lambda)^{-1}$, the solution $w \in H^{\ell+1} \cap H_{per,0}^1$ of (6.50) is given by

$$w = (I - \Gamma_\lambda)^{-1}(\lambda + A)^{-1}(B_1[\tilde{w}]\Psi_\lambda^{-1}\Phi_\lambda f^0 + f)$$

and w satisfies the estimate

$$\sum_{j=0}^{\ell+1} \lambda^{\frac{\ell+1-j}{2}} \|\partial_x^j w\|_2 \leq C \|f^0\|_{H^\ell} + C \sum_{j=0}^{\ell-1} \lambda^{\frac{\ell-1-j}{2}} \|\partial_x^j f\|_2. \quad (6.53)$$

With this w , we define ϕ by (6.48). Then, by (6.49) and (6.53), we see that $\phi \in H_{per,*}^\ell$ and it holds that

$$\lambda \|\phi\|_{H^\ell} \leq C \left\{ \|f^0\|_{H^\ell} + \sum_{j=0}^{\ell-1} \lambda^{\frac{\ell-1-j}{2}} \|\partial_x^j f\|_2 \right\}.$$

This completes the proof. □

We are now in a position to prove Proposition 5.3 (ii).

Proof of Proposition 5.3 (ii). Let $|\varepsilon| \leq \varepsilon_3$ and $|\tilde{\sigma}| + \|\tilde{V}\|_{X^2} \leq M$. Define the operator $\mathcal{L}$ on $Q(H_{per,*}^\ell \times H_{per}^{\ell-1})$ ($\ell = 1, 2$) by

$$D(\mathcal{L}) = QX^\ell,$$

$$\mathcal{L} = \mathcal{L}(\varepsilon, \tilde{\sigma}, \tilde{V}) = \mathcal{L}_{c_0,0} - \varepsilon \tilde{\sigma} Q \partial_{x_1} + \varepsilon Q N[V_1 + \varepsilon \tilde{V}].$$

Set

$$\Sigma_1 = \Sigma_0 \cap \{\lambda; |\lambda| \leq \mu_0\}.$$

It follows from Proposition 6.9 that there exists a positive constant C_2 such that if $\lambda \in \rho(-\mathcal{L}) \cap \Sigma_1$, then

$$\|(\lambda + \mathcal{L})^{-1} F\|_{X^\ell} \leq C_2 \|F\|_{H^\ell \times H^{\ell-1}}. \quad (6.54)$$

Assume that $\mu \in \rho(-\mathcal{L}) \cap \Sigma_1$. Then, by (6.54), we have

$$\{\lambda; |\lambda - \mu| < \frac{1}{C_2}\} \cap \Sigma_1 \subset \rho(-\mathcal{L}) \quad (6.55)$$

and the estimate (6.54) holds for $\lambda \in \Sigma_1$ with $|\lambda - \mu| < \frac{1}{C_2}$.

Since Σ_1 is compact, there exists a finite number of balls B_j ($j = 1, \dots, N_1$) with radius $\frac{1}{2C_2}$ such that $\Sigma_1 \subset \cup_{j=1}^{N_1} B_j$. By Proposition 6.11, we have $\lambda_0 \in \rho(-\mathcal{L}) \cap \Sigma_1$, and hence, $\mu_0 \in B_j$ for some j . Since Σ_1 is connected, we see from (6.55) that $\Sigma_1 \subset \rho(-\mathcal{L})$. Since $0 \in \Sigma_1$, we conclude that $0 \in \rho(-\mathcal{L})$ and the estimate (6.54) holds for $\lambda = 0$. This completes the proof. $\square$

7 Proof of Lemma 6.2

In this section we give an outline of the proof of Lemma 6.2.

Proof of Lemma 6.2. Let $a = \frac{2\pi}{\alpha}$. In this section we write $\Omega_a = (0, a) \times (0, 1)$ instead of Ω_α . We set

$$G_1 = \left(-\frac{a}{4}, \frac{a}{4}\right), \quad G_2 = \left(\frac{a}{8}, \frac{7}{8}a\right),$$

and take $\psi_1, \psi_2 \in C^\infty$ satisfying

$$\begin{aligned} \psi_1 &\geq 0, \quad \left(-\frac{3}{16}a, \frac{3}{16}a\right) \subset \text{supp } \psi_1 \subset G_1, \\ \psi_2 &\geq 0, \quad \left(\frac{5}{32}a, \frac{27}{32}a\right) \subset \text{supp } \psi_2 \subset G_2. \end{aligned}$$

We define $\eta(x_1)$ by

$$\eta(x_1) = \sum_{j=1,2, k \in \mathbb{Z}} \psi_j(x_1 - ak).$$

Then $\eta \in C^\infty(\mathbb{R})$, $\eta(x_1 + a) = \eta(x_1)$ and $\eta(x_1) > 0$ for all $x_1 \in \mathbb{R}$. Setting

$$\phi_{j,k}(x_1) = \frac{\psi_j(x_1 - ak)}{\eta(x_1)},$$

we see that

$$\phi_{j,k} \in C_0^\infty(\mathbb{R}), \quad \text{supp } \phi_{j,k} \subset G_j + ak\mathbf{e}_1 \quad (j = 1, 2, k \in \mathbb{Z}),$$

$$\phi_{j,k}(x_1) = \frac{\psi_j(x_1 - ak)}{\eta(x_1 - ak)} = \phi_{j,0}(x_1 - ak) \quad (j = 1, 2, k \in \mathbb{Z}),$$

$$\sum_{j=1,2,k \in \mathbb{Z}} \phi_{j,k}(x_1) = 1 \quad (x_1 \in \mathbb{R}).$$

Let us consider the problem

$$\operatorname{div} v = f$$

for a given $f \in C_{per,0}^\infty(\Omega_a)$ with $\int_{\Omega_a} f(x) dx = 0$.

We set $Q_0 = G_1 \cup G_2$ and define f_0 by

$$f_0(x) = \phi_{1,0}(x_1)f(x) + \phi_{2,0}(x_1)f(x) \quad (x \in Q_0).$$

It then follows that $f_0 \in C_0^\infty(Q_0)$. Furthermore,

$$\begin{aligned} \int_{Q_0} f_0(x) dx &= \int_{G_1} \phi_{1,0}(x_1)f(x) dx + \int_{G_2} \phi_{2,0}(x_1)f(x) dx \\ &= \int_0^1 \left(\int_{-\frac{a}{4}}^0 \phi_{1,0}(x_1)f(x) dx_1 \right) dx_2 \\ &\quad + \int_0^1 \left(\int_0^{\frac{3}{4}a} (\phi_{1,0}(x_1) + \phi_{2,0}(x_1))f(x) dx_1 \right) dx_2 \\ &\quad + \int_0^1 \left(\int_{\frac{3}{4}a}^{\frac{7}{8}a} \phi_{2,0}(x_1)f(x) dx_1 \right) dx_2 \\ &= \int_0^1 \left(\int_{\frac{3}{4}a}^a \phi_{1,0}(x_1 - a)f(x - a\mathbf{e}_1) dx_1 \right) dx_2 + \int_0^1 \left(\int_0^{\frac{3}{4}a} f(x) dx_1 \right) dx_2 \\ &\quad + \int_0^1 \left(\int_{\frac{3}{4}a}^{\frac{7}{8}a} \phi_{2,0}(x_1)f(x) dx_1 \right) dx_2 \\ &= \int_0^1 \left(\int_{\frac{3}{4}a}^a (\phi_{1,1}(x_1) + \phi_{2,0}(x_1))f(x) dx_1 \right) dx_2 + \int_0^1 \left(\int_0^{\frac{3}{4}a} f(x) dx_1 \right) dx_2 \\ &= \int_{\Omega_a} f(x) dx = 0. \end{aligned}$$

Therefore, from [3, Theorem III. 3.2] and its proof, we see that there exist $v_j \in C_0^\infty(\mathbb{R})$ ($j = 1, 2$) such that $\operatorname{supp} v_j \subset G_j$ ($j = 1, 2$) and $v_0 = v_1 + v_2 \in C_0^\infty(Q_0)$ satisfies

$$\operatorname{div} v_0 = f_0,$$

$$\|\nabla v_0\|_{L^2(Q_0)} \leq C\|f_0\|_{L^2(Q_0)} \leq C\|f\|_{L^2(\Omega_a)}.$$

Let $\tilde{v}_0$ and $\tilde{f}_0$ be the zero extensions of v_0 and f_0 on $\mathbb{R}^2$, respectively, and define v by

$$v(x) = \sum_{k \in \mathbb{Z}} \tilde{v}_0(x - ak\mathbf{e}_1).$$

Then $v \in C_{per,0}^\infty$ and

$$\operatorname{div} v(x) = \sum_{k \in \mathbb{Z}} \operatorname{div} \tilde{v}_0(x - ak\mathbf{e}_1) = \sum_{k \in \mathbb{Z}} \tilde{f}_0(x - ak\mathbf{e}_1).$$

For $x \in \Omega_a \cap G_1$, we have

$$\sum_{k \in \mathbb{Z}} \tilde{f}_0(x - ak\mathbf{e}_1) = \sum_{j=1,2,k \in \mathbb{Z}} \phi_{j,k}(x_1) f_0(x - ak\mathbf{e}_1) = \sum_{j=1,2} \phi_{j,0}(x_1) f(x) = f(x).$$

Furthermore, for $x \in [\frac{a}{4}, \frac{3}{4}a) \times (0, 1)$, we have

$$\sum_{k \in \mathbb{Z}} \tilde{f}_0(x - ak\mathbf{e}_1) = \sum_{j=1,2,k \in \mathbb{Z}} \phi_{j,k}(x_1) f(x - ak\mathbf{e}_1) = \phi_{2,0}(x_1) f(x) = f(x),$$

and, for $x \in [\frac{3}{4}a, a) \times (0, 1)$, we have

$$\begin{aligned} \sum_{k \in \mathbb{Z}} \tilde{f}_0(x - ak\mathbf{e}_1) &= \sum_{j=1,2,k \in \mathbb{Z}} \phi_{j,k}(x_1) f_0(x - ak\mathbf{e}_1) \\ &= \phi_{1,1}(x_1) f(x - a\mathbf{e}_1) + \phi_{2,0}(x_1) f(x) \\ &= (\phi_{1,1}(x_1) + \phi_{2,0}(x_1)) f(x) = f(x). \end{aligned}$$

We thus conclude that $\operatorname{div} v(x) = f(x)$ for $x \in \Omega_a$. Moreover,

$$\|\nabla v\|_{L^2(\Omega_a)} \leq 2\|\nabla v_0\|_{L^2(Q_0)} \leq 2C\|f\|_{L^2(\Omega_a)}.$$

We next consider the case $f = \operatorname{div} \mathbf{g}$ with $\mathbf{g} = {}^\top(g_1, g_2)$, $g_j \in C_{per}^\infty(\overline{\Omega}_a)$ ($j = 1, 2$) and $\operatorname{div} \mathbf{g} \in C_{per,0}^\infty(\Omega_a)$. Following the proofs of [3, Lemma III. 3.5] and [3, Theorem III.3.3], one can show that v_0 satisfies

$$\|v_0\|_{L^2(Q_0)} \leq C\|\mathbf{g}\|_{L^2(Q_0)} \leq C\|\mathbf{g}\|_{L^2(\Omega_a)},$$

$$\|\nabla v_0\|_{L^2(Q_0)} \leq C\|f_0\|_{L^2(Q_0)} \leq C\|\operatorname{div} \mathbf{g}\|_{L^2(\Omega_a)}.$$

It then follows that

$$\|v\|_{L^2(\Omega_a)} \leq C\|\mathbf{g}\|_{L^2(\Omega_a)},$$

$$\|\nabla v\|_{L^2(\Omega_a)} \leq C\|\operatorname{div} \mathbf{g}\|_{L^2(\Omega_a)}.$$

This completes the proof. $\square$

8 Proof of Proposition 5.3 (i)

In this section we will give a proof of Proposition 5.3 (i). We denote $\tilde{L}_{\eta,k}$ and $\tilde{L}_{\eta,k}^*$ with $k = +1$ by $L(\alpha)$ and $L(\alpha)^*$. Then $L(\alpha)$ is expanded as

$$L(\alpha) = L^{(0)} + \alpha L^{(1)} + \alpha^2 L^{(2)},$$

where

$$\begin{aligned} L^{(0)} &= \begin{pmatrix} 0 & 0 & \gamma^2 \partial_{x_2} \\ -\frac{\nu}{\gamma^2} & -\nu \partial_{x_2}^2 & \partial_{x_2} v_s^1 \\ \partial_{x_2} & 0 & -(\nu + \tilde{\nu}) \partial_{x_2}^2 \end{pmatrix}, \\ L^{(1)} &= \begin{pmatrix} i v_s^1 & i \gamma^2 & 0 \\ i & i v_s^1 & -i \tilde{\nu} \partial_{x_2} \\ 0 & -i \tilde{\nu} \partial_{x_2} & i v_s^1 \end{pmatrix}, \\ L^{(2)} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & \nu + \tilde{\nu} & 0 \\ 0 & 0 & \nu \end{pmatrix}. \end{aligned}$$

Similarly, $L(\alpha)^*$ is expanded as

$$L(\alpha)^* = L^{(0)*} + \alpha L^{(1)*} + \alpha^2 L^{(2)*},$$

where

$$\begin{aligned} L^{(0)*} &= \begin{pmatrix} 0 & -\nu & -\gamma^2 \partial_{x_2} \\ 0 & -\nu \partial_{x_2}^2 & 0 \\ -\partial_{x_2} & \partial_{x_2} v_s^1 & -(\nu + \tilde{\nu}) \partial_{x_2}^2 \end{pmatrix}, \\ L^{(1)*} &= \begin{pmatrix} -i v_s^1 & -i \gamma^2 & 0 \\ -i & -i v_s^1 & -i \tilde{\nu} \partial_{x_2} \\ 0 & -i \tilde{\nu} \partial_{x_2} & -i v_s^1 \end{pmatrix}, \\ L^{(2)*} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & \nu + \tilde{\nu} & 0 \\ 0 & 0 & \nu \end{pmatrix}. \end{aligned}$$

Lemma 8.1. *There exists a positive number r_1 such that if $\alpha \leq r_1$, then $V_{\pm}$ and $V_{\pm}^*$ given in section 5.1 are represented as*

$$V_+(x) = (v^{(0)}(x_2) + \alpha v^{(1)}(x_2) + O(\alpha^2)) e^{i\alpha x_1}, \quad V_- = \overline{V_+},$$

$$V_+^*(x) = \frac{\alpha}{2\pi} (v^{(0)*}(x_2) + \alpha v^{(1)*}(x_2) + O(\alpha^2)) e^{i\alpha x_1}, \quad V_-^* = \overline{V_+^*},$$

where $v^{(0)} = {}^\top(\phi^{(0)}, w^{(0),1}, 0)$ with

$$\phi^{(0)} = 1, \quad w^{(0),1} = \frac{1}{2\gamma^2}(-x_2^2 + x_2),$$

$v^{(1)} = {}^\top(\phi^{(1)}, w^{(1),1}, w^{(1),2})$ with

$$\begin{aligned} \phi_1^{(1)}(x_2) &= -i \left(\frac{\nu}{\gamma^2} + \frac{\tilde{\nu}}{2\gamma^2} \right) (-x_2^2 + x_2 - \frac{1}{6}), \\ w_1^{(1),1}(x_2) &= -i \left(\frac{\nu}{\gamma^4} + \frac{\tilde{\nu}}{2\gamma^4} \right) \left(\frac{1}{12}x_2^4 - \frac{1}{6}x_2^3 + \frac{1}{12}x_2^2 \right) \\ &\quad - \frac{i}{12\nu\gamma^2} \left(\frac{1}{30}x_2^6 - \frac{1}{10}x_2^5 + \frac{1}{12}x_2^4 - \frac{1}{60}x_2 \right) - \frac{i}{2\nu}(-x_2^2 + x_2), \\ w_1^{(1),2}(x_2) &= -\frac{i}{\gamma^2} \left(-\frac{1}{3}x_2^3 + \frac{1}{2}x_2^2 - \frac{1}{6}x_2 \right), \end{aligned}$$

$v^{(0)*} = {}^\top(\phi^{(0)*}, 0, 0)$ with $\phi^{(0)*} = \gamma^2$, and $v^{(1)*} = {}^\top(\phi^{(1)*}, w^{(1),1*}, w^{(1),2*})$ with

$$w^{(1),1*} = \frac{i\gamma^2}{2\nu}(-x_2^2 + x_2).$$

Remark 8.2. Note that we will not use the explicit form of $\phi^{(1)*}$ and $w^{(1),2*}$.

Proof. We see from [8, Lemma 5.1] that $v^{(0)}$ and $v^{(0)*}$ are eigenfunctions for eigenvalue 0 of $-L^{(0)}$ and $-L^{(0)*}$, respectively, and the corresponding eigenprojections $\Pi^{(0)}$ and $\Pi^{(0)*}$ are given by

$$\Pi^{(0)}u = \langle \langle u, v^{(0)*} \rangle \rangle v^{(0)}, \quad \Pi^{(0)*}u = \langle \langle u, v^{(0)} \rangle \rangle v^{(0)*}.$$

Let P_α be the eigenprojection for λ_α . Then

$$P_\alpha = \Pi^{(0)} - \alpha(SL^{(1)}\Pi^{(0)} + \Pi^{(0)}L^{(1)}S) + O(\alpha^2),$$

where $S = [(I - \Pi^{(0)})L^{(0)}(I - \Pi^{(0)})]^{-1}$. Set $v_{+1} = P_\alpha v^{(0)}$. We see that v_{+1} is an eigenfunction for λ_α and

$$v_{+1} = v^{(0)} - \alpha SL^{(1)}v^{(0)} + O(\alpha^2).$$

Therefore, setting $v^{(1)} = -SL^{(1)}v^{(0)}$, we have the desired expression of $v^{(1)}$ from [8, Proposition 6.5], where $SL^{(1)}v^{(0)}$ is computed.

As for V_+^* , let P_α^* be the eigenprojection for $\lambda_\alpha^* = \overline{\lambda_\alpha}$. Then

$$P_\alpha^* = \Pi^{(0)*} - \alpha(S^*L^{(1)*}\Pi^{(0)*} + \Pi^{(0)*}L^{(1)*}S^*) + O(\alpha^2),$$

where $S^* = [(I - \Pi^{(0)*})L^{(0)*}(I - \Pi^{(0)*})]^{-1}$. Set $\tilde{v}_{+1}^* = P_\alpha^* v^{(0)*}$. Then $\tilde{v}_{+1}^*$ is an eigenfunction for λ_α^* and

$$\tilde{v}_{+1}^* = v^{(0)*} - \alpha S^*L^{(1)*}v^{(0)*} + O(\alpha^2).$$

Let us compute $\tilde{v}^{(1)*} = -S^*L^{(1)*}v^{(0)*}$ which is the solution of

$$L^{(0)*}u = -(I - \Pi^{(0)*})L^{(1)*}v^{(0)*}, \quad \langle \langle u, v^{(0)*} \rangle \rangle = 0.$$

By [8, Proposition 6.3], we have $\langle \langle L^{(1)}v^{(0)}, v^{(0)*} \rangle \rangle = \frac{i}{6}$, and hence,

$$\Pi^{(0)*}L^{(1)*}v^{(0)*} = \langle \langle L^{(1)*}v^{(0)*}, v^{(0)*} \rangle \rangle v^{(0)*} = \overline{\langle \langle L^{(1)}v^{(0)}, v^{(0)*} \rangle \rangle} v^{(0)*} = -\frac{i}{6}v^{(0)*}.$$

We set $f = {}^\top(f^{0*}, f^{1*}, f^{2*}) = -(I - \Pi^{(0)*})L^{(1)*}v^{(0)*}$. By a direct computation we have

$$f^{0*} = i\gamma^2 v_s^1 - \frac{i}{6}\gamma^2, \quad f^{1*} = i\gamma^2, \quad f^{2*} = 0.$$

It then follows that

$$\partial_{x_2}^2 w^1 = -\frac{i\gamma^2}{\nu}, \quad w^1|_{x_2=0,1} = 0. \quad (8.1)$$

This gives $w^1 = \frac{i\gamma^2}{2\nu}(-x_2^2 + x_2)$, and then w^2 and ϕ are given by

$$\partial_{x_2} w^2 = -\frac{1}{\gamma^2}(\nu w^1 + f^{0*}), \quad (8.2)$$

$$\partial_{x_2} \phi = (\partial_{x_2} v_s^1)w^1 - (\nu + \tilde{\nu})\partial_{x_2}^2 w^2, \quad \int_0^1 \phi dx_2 = -\gamma^2(w^1, w^{(0),1}).$$

Since $\langle \langle v^{(1)}, v^{(0)*} \rangle \rangle = \langle \langle \tilde{v}^{(1)*}, v^{(0)*} \rangle \rangle = 0$, we have $\langle \langle v_{+1}, \tilde{v}_{+1}^* \rangle \rangle = 1 + O(\alpha^2)$. Therefore, setting $v_{+1}^* = \tilde{v}_{+1}^* / \langle \langle v_{+1}, \tilde{v}_{+1}^* \rangle \rangle$, we have the desired result. This completes the proof. $\square$

We are now in a position to prove Proposition 5.3 (i).

Proof of Proposition 5.3 (i). By Lemma 8.1 and the relation that $-\partial_{x_2}^2 w^{(0),1} = \frac{1}{\gamma^2}\phi^{(0)}$, we have

$$\begin{aligned} \llbracket K_0 V_1 \rrbracket_1 &= \alpha^2 \left\{ (\partial_{x_2} w^{(0),1}, i w^{(1),2*}) + \left(-\frac{i}{\gamma^2} \phi^{(1)} - i \partial_{x_2}^2 w^{(1),1}, i w^{(1),1*} \right) \right. \\ &\quad \left. - 2 (i \partial_{x_2}^2 w^{(1),3}, i w^{(1),2*}) \right\} + O(\alpha^3) \\ &= \alpha^2 \left\{ - (w^{(0),1}, i \partial_{x_2} w^{(1),2*}) - \left(\frac{i}{\gamma^2} \phi^{(1)}, i w^{(1),1*} \right) - (i w^{(1),1}, i \partial_{x_2}^2 w^{(1),1*}) \right. \\ &\quad \left. + 2 (i \partial_{x_2} w^{(1),2}, i \partial_{x_2} w^{(1),2*}) \right\} + O(\alpha^3). \end{aligned}$$

By using (8.1), (8.2) and Lemma 8.1, we find that

$$\llbracket K_0 V_1 \rrbracket_1 = \frac{\alpha^2}{12\nu^2} \left\{ \left(\frac{1}{280} - \gamma^2 \right) + \frac{\nu^2}{10\gamma^2} \right\} + O(\alpha^3) > 0$$

for $\alpha \ll 1$. This completes the proof. $\square$

Acknowledgements. Y. Kagei was partly supported by JSPS KAKENHI Grant Number 24340028, 22244009, 24224003 15K13449.

References

- [1] M. Bause, J. G. Heywood, A. Novotny and M. Padula, An iterative scheme for steady compressible viscous flow, modified to treat large potential forces, in *Mathematical Fluid Mechanics, Recent results and open questions*, ed. by J. Neustupa, P. Penel, Birkhäuser, Basel (2001), pp. 27–46.
- [2] M. E. Bogovskii, Solution of the first boundary value problem for an equation of continuity of an incompressible medium, *Soviet Math. Dokl.*, **20** (1979), pp. 1094–1098.
- [3] G. P. Galdi, *An Introduction to the Mathematical Theory of the Navier-Stokes Equations*, Vol. 1, Springer-Verlag, New York (1994).
- [4] J. G. Heywood and M. Padula, On the steady transport equation, in *Fundamental Directions in Mathematical Fluid Mechanics*, ed. by G. P. Galdi, J. G. Heywood, R. Rannacher, Birkhäuser, Basel (2000), pp. 149–170.
- [5] J. G. Heywood and M. Padula, On the existence and uniqueness theory for steady compressible viscous flow, in *Fundamental Directions in Mathematical Fluid Mechanics*, ed. by G. P. Galdi, J. G. Heywood, R. Rannacher, Birkhäuser, Basel (2000), pp. 171–189.
- [6] G. Iooss and M. Padula, Structure of the linearized problem for compressible parallel fluid flows, *Ann. Univ. Ferrara, Sez. VII*, **43** (1998), pp. 157–171.
- [7] Y. Kagei, Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow, *Arch. Rational Mech. Anal.*, **205** (2012), pp. 585–650.
- [8] Y. Kagei and T. Nishida, Instability of plane Poiseuille flow in viscous compressible gas, *J. Math. Fluid Mech.*, **17** (2015), pp. 129–143.
- [9] S. A. Orszag, Accurate solution of the Orr-Sommerfeld stability equation, *J. Fluid Mech.*, **50** (1971), pp. 689–703.
- [10] T. Nishida, M. Padula and Y. Teramoto, Heat convection of compressible viscous fluids: I, *J. Math. Fluid Mech.*, **15** (2013), pp. 525–536.
- [11] T. Nishida, M. Padula and Y. Teramoto, Heat convection of compressible viscous fluids. II, *J. Math. Fluid Mech.*, **15** (2013), pp. 689–700.
- [12] A. Matsumura and T. Nishida, Initial boundary value problems for the equations of motion of compressible viscous and heat-conductive fluids. *Comm. Math. Phys.*, **89** (1983), pp. 445–464.
- [13] H. Sohr, *The Navier-Stokes equations: an elementary functional analytic approach*, Birkhäuser, Basel (2001).

List of MI Preprint Series, Kyushu University

The Global COE Program

Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p-adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials
- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials

- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings
- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric τ -functions of the q-Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications

- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain
- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection
- MI2009-26 Manabu YOSHIDA
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance

- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA
Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI
Finite element computation for scattering problems of micro-hologram using DtN map
- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO
Abstract collision systems on groups

- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA
An algebraic approach to underdetermined experiments
- MI2010-10 Kei HIROSE & Sadanori KONISHI
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE
Lagrangian approach to weakly nonlinear stability of an elliptical flow
- MI2010-21 Hiroki MASUDA
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiro FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra

- MI2010-25 Toshimitsu TAKAESU
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI
Spin-spin correlation functions of the q -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and spike detection via Gaussian basis expansions
- MI2010-32 Nobutaka NAKAZONO
Hypergeometric τ functions of the q -Painlevé systems of type $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA
CAP representations of inner forms of $Sp(4)$ with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER
Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide FUKUMOTO & Alexander B. SAMOKHIN
Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI
Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI
Anderson-Darling test and the Malliavin calculus
- MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling via Compressed Sensing

- MI2011-5 Hiroshi INOUE
Implications in Compressed Sensing and the Restricted Isometry Property
- MI2011-6 Daeju KIM & Sadanori KONISHI
Predictive information criterion for nonlinear regression model based on basis expansion methods
- MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI
Group variable selection via relevance vector machine
- MI2011-8 Jan BREZINA & Yoshiyuki KAGEI
Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow
Group variable selection via relevance vector machine
- MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC
Recursive structures in the multispecies TASEP
- MI2011-10 Kazunori YASUTAKE
On projective space bundle with nef normalized tautological line bundle
- MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA
An explicit formula for the discrete power function associated with circle patterns of Schramm type
- MI2011-12 Yoshiyuki KAGEI
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow
- MI2011-13 Vladimír CHALUPECKÝ & Adrian MUNTEAN
Semi-discrete finite difference multiscale scheme for a concrete corrosion model: approximation estimates and convergence
- MI2011-14 Jun-ichi INOBUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves
- MI2011-15 Hiroshi INOUE
A generalization of restricted isometry property and applications to compressed sensing
- MI2011-16 Yu KAWAKAMI
A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space
- MI2011-17 Naoyuki KAMIYAMA
Matroid intersection with priority constraints
- MI2012-1 Kazufumi KIMOTO & Masato WAKAYAMA
Spectrum of non-commutative harmonic oscillators and residual modular forms
- MI2012-2 Hiroki MASUDA
Mighty convergence of the Gaussian quasi-likelihood random fields for ergodic Levy driven SDE observed at high frequency

- MI2012-3 Hiroshi INOUE
A Weak RIP of theory of compressed sensing and LASSO
- MI2012-4 Yasuhide FUKUMOTO & Youich MIE
Hamiltonian bifurcation theory for a rotating flow subject to elliptic straining field
- MI2012-5 Yu KAWAKAMI
On the maximal number of exceptional values of Gauss maps for various classes of surfaces
- MI2012-6 Marcio GAMEIRO, Yasuaki HIRAOKA, Shunsuke IZUMI, Miroslav KRAMAR, Konstantin MISCHAIKOW & Vedit NANDA
Topological Measurement of Protein Compressibility via Persistence Diagrams
- MI2012-7 Nobutaka NAKAZONO & Seiji NISHIOKA
Solutions to a q -analog of Painlevé III equation of type $D_7^{(1)}$
- MI2012-8 Naoyuki KAMIYAMA
A new approach to the Pareto stable matching problem
- MI2012-9 Jan BREZINA & Yoshiyuki KAGEI
Spectral properties of the linearized compressible Navier-Stokes equation around time-periodic parallel flow
- MI2012-10 Jan BREZINA
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a time-periodic parallel flow
- MI2012-11 Daeju KIM, Shuichi KAWANO & Yoshiyuki NINOMIYA
Adaptive basis expansion via the extended fused lasso
- MI2012-12 Masato WAKAYAMA
On simplicity of the lowest eigenvalue of non-commutative harmonic oscillators
- MI2012-13 Masatoshi OKITA
On the convergence rates for the compressible Navier- Stokes equations with potential force
- MI2013-1 Abuduwaili PAERHATI & Yasuhide FUKUMOTO
A Counter-example to Thomson-Tait-Chetayev's Theorem
- MI2013-2 Yasuhide FUKUMOTO & Hirofumi SAKUMA
A unified view of topological invariants of barotropic and baroclinic fluids and their application to formal stability analysis of three-dimensional ideal gas flows
- MI2013-3 Hiroki MASUDA
Asymptotics for functionals of self-normalized residuals of discretely observed stochastic processes
- MI2013-4 Naoyuki KAMIYAMA
On Counting Output Patterns of Logic Circuits
- MI2013-5 Hiroshi INOUE
RIPless Theory for Compressed Sensing

- MI2013-6 Hiroshi INOUE
Improved bounds on Restricted isometry for compressed sensing
- MI2013-7 Hidetoshi MATSUI
Variable and boundary selection for functional data via multiclass logistic regression modeling
- MI2013-8 Hidetoshi MATSUI
Variable selection for varying coefficient models with the sparse regularization
- MI2013-9 Naoyuki KAMIYAMA
Packing Arborescences in Acyclic Temporal Networks
- MI2013-10 Masato WAKAYAMA
Equivalence between the eigenvalue problem of non-commutative harmonic oscillators and existence of holomorphic solutions of Heun's differential equations, eigenstates degeneration, and Rabi's model
- MI2013-11 Masatoshi OKITA
Optimal decay rate for strong solutions in critical spaces to the compressible Navier-Stokes equations
- MI2013-12 Shuichi KAWANO, Ibuki HOSHINA, Kazuki MATSUDA & Sadanori KONISHI
Predictive model selection criteria for Bayesian lasso
- MI2013-13 Hayato CHIBA
The First Painleve Equation on the Weighted Projective Space
- MI2013-14 Hidetoshi MATSUI
Variable selection for functional linear models with functional predictors and a functional response
- MI2013-15 Naoyuki KAMIYAMA
The Fault-Tolerant Facility Location Problem with Submodular Penalties
- MI2013-16 Hidetoshi MATSUI
Selection of classification boundaries using the logistic regression
- MI2014-1 Naoyuki KAMIYAMA
Popular Matchings under Matroid Constraints
- MI2014-2 Yasuhide FUKUMOTO & Youichi MIE
Lagrangian approach to weakly nonlinear interaction of Kelvin waves and a symmetry-breaking bifurcation of a rotating flow
- MI2014-3 Reika AOYAMA
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Parallel flow in a cylindrical domain
- MI2014-4 Naoyuki KAMIYAMA
The Popular Condensation Problem under Matroid Constraints

- MI2014-5 Yoshiyuki KAGEI & Kazuyuki TSUDA
Existence and stability of time periodic solution to the compressible Navier-Stokes equation for time periodic external force with symmetry
- MI2014-6 This paper was withdrawn by the authors.
- MI2014-7 Masatoshi OKITA
On decay estimate of strong solutions in critical spaces for the compressible Navier-Stokes equations
- MI2014-8 Rong ZOU & Yasuhide FUKUMOTO
Local stability analysis of azimuthal magnetorotational instability of ideal MHD flows
- MI2014-9 Yoshiyuki KAGEI & Naoki MAKIO
Spectral properties of the linearized semigroup of the compressible Navier-Stokes equation on a periodic layer
- MI2014-10 Kazuyuki TSUDA
On the existence and stability of time periodic solution to the compressible Navier-Stokes equation on the whole space
- MI2014-11 Yoshiyuki KAGEI & Takaaki NISHIDA
Instability of plane Poiseuille flow in viscous compressible gas
- MI2014-12 Chien-Chung HUANG, Naonori KAKIMURA & Naoyuki KAMIYAMA
Exact and approximation algorithms for weighted matroid intersection
- MI2014-13 Yusuke SHIMIZU
Moment convergence of regularized least-squares estimator for linear regression model
- MI2015-1 Hidetoshi MATSUI
Sparse regularization for multivariate linear models for functional data
- MI2015-2 Reika AOYAMA & Yoshiyuki KAGEI
Spectral properties of the semigroup for the linearized compressible Navier-Stokes equation around a parallel flow in a cylindrical domain
- MI2015-3 Naoyuki KAMIYAMA
Stable Matchings with Ties, Master Preference Lists, and Matroid Constraints
- MI2015-4 Reika AOYAMA & Yoshiyuki KAGEI
Large time behavior of solutions to the compressible Navier-Stokes equations around a parallel flow in a cylindrical domain
- MI2015-5 Kazuyuki TSUDA
Existence and stability of time periodic solution to the compressible Navier-Stokes-Korteweg system on R^3
- MI2015-6 Naoyuki KAMIYAMA
Popular Matchings with Ties and Matroid Constraints

- MI2015-7 Shoichi EGUCHI & Hiroki MASUDA
Quasi-Bayesian model comparison for LAQ models
- MI2015-8 Yoshiyuki KAGEI & Ryouta OOMACHI
Stability of time periodic solution of the Navier-Stokes equation on the half-space
under oscillatory moving boundary condition
- MI2015-9 Yoshiyuki KAGEI & Takaaki NISHIDA
Traveling waves bifurcating from plane Poiseuille flow of the compressible Navier-
Stokes equation