

## Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$

Kajiwara, Kenji  
Faculty of Mathematics, Kyushu University

Nakazono, Nobutaka  
Graduate School of Mathematics, Kyushu University

Tsuda, Teruhisa  
Faculty of Mathematics, Kyushu University

<https://hdl.handle.net/2324/15556>

---

出版情報 : MI Preprint Series. 2009-34, 2010-05-28. Oxford University Press  
バージョン :  
権利関係 :



# MI Preprint Series

Kyushu University  
The Global COE Program  
Math-for-Industry Education & Research Hub

## Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$

Kenji Kajiwara,  
Nobutaka Nakazono and  
Teruhisa Tsuda

MI 2009-34

( Received September 29, 2009 )

Faculty of Mathematics  
Kyushu University  
Fukuoka, JAPAN

# Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$

Kenji Kajiwara, Nobutaka Nakazono<sup>†</sup> and Teruhisa Tsuda

28 September. 2009

Faculty of Mathematics, Kyushu University,  
819-0395 Moto-oka, Fukuoka 819-0395, Japan

<sup>†</sup> Graduate School of Mathematics, Kyushu University,  
819-0395 Moto-oka, Fukuoka 819-0395, Japan

## Abstract

We consider the  $q$ -Painlevé III equation arising from the birational representation of the affine Weyl group of type  $(A_2 + A_1)^{(1)}$ . We study the reduction of the  $q$ -Painlevé III equation to the  $q$ -Painlevé II equation from the viewpoint of affine Weyl group symmetry. In particular, the mechanism of apparent inconsistency between the hypergeometric solutions to both equations is clarified by using factorization of difference operators and the  $\tau$  functions.

**2000 Mathematics Subject Classification:** 34M55, 39A13, 33D15, 33E17

**Keywords and Phrases:** affine Weyl group, discrete Painlevé equation, hypergeometric function

## 1 Introduction

The discrete Painlevé equations have been studied actively from various points of view. Together with the Painlevé equations, they are now regarded as one of the most important classes of equations in the theory of integrable systems (see, for example, [6]). Originally, the discrete Painlevé equations had been identified as single second-order equations [1–3, 33, 37] and then were generalized to simultaneous first-order equations. A typical example is the following equation known as a discrete Painlevé II equation [33, 37]:

$$x_{n+1} + x_{n-1} = \frac{(an + b)x_n + c}{1 - x_n^2}, \quad (1.1)$$

where  $x_n$  is the dependent variable,  $n$  is the independent variable, and  $a, b, c \in \mathbb{C}$  are parameters. By applying the singularity confinement criterion [7], (1.1) is generalized to

$$x_{n+1} + x_{n-1} = \frac{(an + b)x_n + c + (-1)^n d}{1 - x_n^2}, \quad (1.2)$$

where  $d$  is a parameter, with its integrability preserved. Introducing the dependent variables  $X_n$  and  $Y_n$  by

$$X_n = x_{2n}, \quad Y_n = x_{2n-1}, \quad (1.3)$$

then (1.2) can be rewritten as

$$Y_{n+1} + Y_n = \frac{(2an + b)X_n + c + d}{1 - X_n^2}, \quad X_{n+1} + X_n = \frac{(a(2n + 1) + b)Y_{n+1} + c - d}{1 - Y_{n+1}^2}. \quad (1.4)$$

Equation (1.4) is known as a discrete Painlevé III equation since it admits a continuous limit to the Painlevé III equation [5]. Conversely, (1.1) can be recovered from (1.4) by putting  $d = 0$  and (1.3). We call this procedure “symmetrization” of (1.4), which comes from the terminology of the Quispel–Roberts–Thompson (QRT) mapping [34, 35]. After this terminology, (1.4) is sometimes called the “asymmetric” discrete Painlevé II equation, and (1.1) is called the “symmetric” discrete Painlevé III equation [21].

It looks that the symmetrization is a simple specialization of parameters at the level of the equation, but some strange phenomena have been reported as to their particular solutions expressed in terms of hypergeometric functions (*hypergeometric solutions*). The hypergeometric solutions to (1.1) have been constructed as follows [9, 19]:

**Proposition 1.1** *For each  $N \in \mathbb{N}$ , let  $\tau_N^n$  be an  $N \times N$  determinant defined by*

$$\tau_N^n = \begin{vmatrix} H_n & H_{n+1} & \cdots & H_{n+N-1} \\ H_{n+2} & H_{n+3} & \cdots & H_{n+N+1} \\ \vdots & \vdots & \ddots & \vdots \\ H_{n+2N-2} & H_{n+2N-1} & \cdots & H_{n+3N-3} \end{vmatrix}, \quad (1.5)$$

where  $H_n$  is a function satisfying the three-term relation:

$$H_{n+1} - zH_n + nH_{n-1} = 0. \quad (1.6)$$

Then,

$$x_n = \frac{2}{z} \frac{\tau_{N+1}^{n+1} \tau_N^n}{\tau_{N+1}^n \tau_N^{n+1}} - 1, \quad (1.7)$$

satisfies (1.1) with the parameters

$$a = \frac{8}{z^2}, \quad b = \frac{4(1+2N)}{z^2}, \quad c = -\frac{4(1+2N)}{z^2}. \quad (1.8)$$

On the other hand, since (1.4) appears as the Bäcklund transformation of the Painlevé V equation [28, 38], its hypergeometric solutions are essentially the same as those to the Painlevé V equation [22, 31]. The explicit form of the hypergeometric solutions to (1.4) are given as follows:

**Proposition 1.2** *For each  $N \in \mathbb{N}$ , let  $\tau_N^{n,m}$  be an  $N \times N$  determinant defined by*

$$\tau_N^{n,m} = \begin{vmatrix} K_n^m & K_{n+1}^m & \cdots & K_{n+N-1}^m \\ K_{n+1}^m & K_{n+2}^m & \cdots & K_{n+N}^m \\ \vdots & \vdots & \ddots & \vdots \\ K_{n+N-1}^m & K_{n+N}^m & \cdots & K_{n+2N-2}^m \end{vmatrix}, \quad (1.9)$$

where  $K_n^m$  is a function satisfying

$$K_{n+1}^m - K_n^m - tK_{n+1}^{m+1} = 0, \quad nK_{n+1}^m - (n+t)K_n^m - (n-m)tK_n^{m+1} = 0. \quad (1.10)$$

Then,

$$X_n = 2(n+2N-1) \frac{\tau_{N+1}^{n,m} \tau_N^{n,m}}{\tau_{N+1}^{n-1,m-1} \tau_N^{n+1,m+1}} - 1, \quad Y_n = \frac{2}{t} \frac{\tau_{N+1}^{n-1,m-1} \tau_N^{n,m+1}}{\tau_{N+1}^{n-1,m} \tau_N^{n,m}} - 1, \quad (1.11)$$

satisfy (1.4) with the parameters

$$a = -\frac{4}{t}, \quad b = \frac{-4(-m + 2N - 1)}{t}, \quad c = \frac{2(1 + 2N)}{t}, \quad d = \frac{2(2m + 2N - 3)}{t}. \quad (1.12)$$

It is obvious that substituting  $d = 0$  into the hypergeometric solutions to (1.4) in Proposition 1.2 do not yield those to (1.1) in Proposition 1.1. In particular, we remark the following differences between the two solutions:

- (1) The hypergeometric functions are different. Equation (1.6) can be solved by considering the parabolic cylinder function (Weber function), while (1.10) can be solved by considering the confluent hypergeometric function. In fact, the former function is expressed as a specialization of the latter, but this specialization is not consistent with the symmetrization.
- (2) Structures of the determinant are different. The determinant (1.5) has asymmetry in the shift of index: the shift in the vertical direction is two while that in the horizontal direction is one. On the other hand, the determinant (1.9) is an ordinary Hankel determinant.

We note that similar phenomena have been reported also for some other discrete Painlevé equations [8, 18, 25]. Many integrable systems admit particular solutions expressed in terms of determinants, but such an asymmetric structure of the determinant solutions has been seen only in the hypergeometric solutions to the discrete Painlevé equations. Note here that these phenomena cannot be seen for the algebraic (or rational) solutions. For example, it is known that substituting  $d = 0$  into the determinant expression of the rational solutions to (1.4) yields those to (1.1); see [20, 23, 24].

The  $\tau$  function is one of the most important objects in the theory of integrable systems and is regarded as carrying the underlying fundamental mathematical structures. Concerning the discrete Painlevé equations, investigation of the  $\tau$  functions started [18, 19] through the search for the explicit formulae of the hypergeometric and algebraic solutions. In fact, the above mysterious asymmetric structure has been one motivation of further study.

It is now known that theory of birational representations of affine Weyl groups provides us with an algebraic tool to study the Painlevé systems [27, 29–32]. Moreover, a geometric framework of the two-dimensional Painlevé systems has been presented based on certain rational surfaces [15, 39]. Combining these results enables us to study the Painlevé systems effectively. For instance, it played a crucial role in the identification of hypergeometric functions that appear as the particular solutions to the Painlevé systems in Sakai’s classification [12–14].

The purpose of this paper is to clarify the mechanism of the phenomena of hypergeometric solutions from the viewpoint of the affine Weyl group symmetry. We shall take the  $q$ -Painlevé equation of type  $(A_2 + A_1)^{(1)}$  as an example, which is the simplest non-trivial discrete Painlevé system [39]. The key is to formulate the symmetrization in terms of the birational representation of the affine Weyl group, where the discrete Painlevé equation arises from the action of the translational subgroup. In fact, the discrete time evolution of the symmetric case can be regarded as a “half-step” of a translation of the affine Weyl group with restricted to a certain line in the parameter space. Conversely, we can derive various discrete Painlevé equations from elements of infinite order that are not only translations by taking a projection on a certain subspace of the parameters. We call such a procedure to obtain a “smaller” discrete time evolution of Painlevé type a *projective reduction*.

This paper is organized as follows: in Section 2, we introduce a  $q$ -Painlevé III equation and derive a  $q$ -Painlevé II equation by applying the symmetrization. Then we give a brief review on their hypergeometric solutions. In Section 3, we first introduce the family of Bäcklund transformations of the  $q$ -Painlevé III equation, which is a birational representation of the affine Weyl group of type  $(A_2 + A_1)^{(1)}$ . We next lift the representation on the level of  $\tau$  functions and derive various bilinear equations. We then clarify the mechanism of the inconsistency among the hypergeometric solutions by using this framework. Some concluding remarks are given in Section 4.

## 2 $q$ -P<sub>III</sub> and $q$ -P<sub>II</sub>

We consider the following system of  $q$ -difference equations [11, 17, 39]:

$$g_{n+1} = \frac{q^{2N+1}c^2}{f_n g_n} \frac{1 + a_0 q^n f_n}{a_0 q^n + f_n}, \quad f_{n+1} = \frac{q^{2N+1}c^2}{f_n g_{n+1}} \frac{1 + a_2 a_0 q^{n-m} g_{n+1}}{a_2 a_0 q^{n-m} + g_{n+1}}, \quad (2.1)$$

for the unknown functions  $f_n = f_n(m, N)$  and  $g_n = g_n(m, N)$  and the independent variable  $n \in \mathbb{Z}$ . Here  $m, N \in \mathbb{Z}$  and  $a_0, a_2, c, q \in \mathbb{C}^\times$  are parameters. Equation (2.1) has the (extended) affine Weyl group symmetry of type  $(A_2 + A_1)^{(1)}$  and is known as a  $q$ -Painlevé III equation ( $q$ -P<sub>III</sub>) since the continuous limit yields the Painlevé III equation. We also consider the following  $q$ -difference equation [25, 36]:

$$X_{k+1} = \frac{q^{2N+1}c^2}{X_k X_{k-1}} \frac{1 + a_0 q^{\frac{k}{2}} X_k}{a_0 q^{\frac{k}{2}} + X_k}, \quad (2.2)$$

for the unknown function  $X_k = X_k(N)$  and the independent variable  $k \in \mathbb{Z}$ . Equation (2.2) is a  $q$ -Painlevé II equation ( $q$ -P<sub>II</sub>) and actually it admits a continuous limit to the Painlevé II equation.

Note that substituting

$$m = 0, \quad a_2 = q^{\frac{1}{2}}, \quad (2.3)$$

and putting

$$f_k(0, N) = X_{2k}(N), \quad g_k(0, N) = X_{2k-1}(N), \quad (2.4)$$

in (2.1) yield (2.2).

We shall briefly review the hypergeometric solutions to  $q$ -P<sub>III</sub> and  $q$ -P<sub>II</sub> following [11, 25].

### 2.1 Hypergeometric solutions to $q$ -P<sub>III</sub>

First, we review the hypergeometric solutions to  $q$ -P<sub>III</sub>. For each  $N \in \mathbb{Z}_{\geq 0}$ , let  $\psi_N^{n,m}$  be an  $N \times N$  determinant defined by

$$\psi_N^{n,m} = \begin{vmatrix} F_{n,m} & F_{n+1,m} & \cdots & F_{n+N-1,m} \\ F_{n-1,m} & F_{n,m} & \cdots & F_{n+N-2,m} \\ \vdots & \vdots & \ddots & \vdots \\ F_{n-N+1,m} & F_{n-N+2,m} & \cdots & F_{n,m} \end{vmatrix}, \quad \psi_0^{n,m} = 1, \quad (2.5)$$

where  $F_{n,m}$  satisfies

$$\begin{aligned} F_{n+1,m} - F_{n,m} &= -a_0^2 q^{2n} F_{n,m-1}, \\ F_{n,m+1} - F_{n,m} &= -a_2^{-2} q^{2m+2} F_{n-1,m}. \end{aligned} \quad (2.6)$$

**Lemma 2.1** ([11])  $\psi_N^{n,m}$  satisfies the following bilinear difference equations:

$$a_0^2 q^{2n-2} \psi_{N+1}^{n-1,m-1} \psi_N^{n,m} - q^{2N} \psi_N^{n,m-1} \psi_{N+1}^{n-1,m} + \psi_N^{n-1,m-1} \psi_{N+1}^{n,m} = 0, \quad (2.7)$$

$$\psi_{N+1}^{n,m} \psi_N^{n,m-1} - q^{-2N} \psi_N^{n-1,m} \psi_{N+1}^{n+1,m+1} - a_0^2 q^{2n} \psi_N^{n,m} \psi_{N+1}^{n,m-1} = 0, \quad (2.8)$$

$$\psi_{N+1}^{n,m} \psi_N^{n-1,m-1} - \psi_{N+1}^{n,m-1} \psi_N^{n-1,m} + a_2^{-2} q^{2m} \psi_N^{n,m} \psi_{N+1}^{n-1,m-1} = 0, \quad (2.9)$$

$$\psi_{N+1}^{n,m-1} \psi_N^{n,m} - a_2^{-2} q^{2m} \psi_{N+1}^{n-1,m-1} \psi_N^{n+1,m} - \psi_N^{n,m-1} \psi_{N+1}^{n,m} = 0. \quad (2.10)$$

**Proposition 2.2** ([11]) The hypergeometric solutions to  $q$ - $P_{\text{III}}$ , (2.1), with  $c = 1$  are given by

$$f_n = -a_0 q^n \frac{\psi_{N+1}^{n,m-1} \psi_N^{n,m}}{\psi_{N+1}^{n,m} \psi_N^{n,m-1}}, \quad g_n = \frac{a_2}{a_0} q^{-n-m+1} \frac{\psi_{N+1}^{n,m} \psi_N^{n-1,m-1}}{\psi_{N+1}^{n-1,m-1} \psi_N^{n,m}}. \quad (2.11)$$

Proposition 2.2 follows from Lemma 2.1.

**Remark 2.3** (1)  $F_{n,m}$  satisfies the three-term relation:

$$F_{n+1,m} + (a_0^2 q^{2n} - a_2^{-2} q^{2m+2} - 1) F_{n,m} + a_2^{-2} q^{2m+2} F_{n-1,m} = 0, \quad (2.12)$$

whose general solution is given by

$$F_{n,m} = \frac{A}{(a_2^{-2} q^{2m+2}; q^2)_\infty} {}_1\varphi_1 \left( \begin{matrix} 0 \\ a_2^2 q^{-2m}; q^2, a_2^2 a_0^2 q^{2n-2m} \end{matrix} \right) + B (a_2^{-2} q^{2m+4}; q^2)_\infty (a_2^{-1} q^{m+1})^{2n} {}_1\varphi_1 \left( \begin{matrix} 0 \\ a_2^{-2} q^{2m+4}; q^2, a_0^2 q^{2n+2} \end{matrix} \right). \quad (2.13)$$

Here,  $A$  and  $B$  are arbitrary constants, and  ${}_1\varphi_1$  is the basic hypergeometric function defined by [4]

$${}_1\varphi_1 \left( \begin{matrix} a \\ b \end{matrix}; q, z \right) = \sum_{k=0}^{\infty} \frac{(a; q)_k}{(b; q)_k (q; q)_k} (-1)^k q^{\frac{k(k-1)}{2}} z^k, \quad (a; q)_k = \prod_{i=1}^k (1 - a q^{i-1}). \quad (2.14)$$

(2)  $\psi_N^{n,m}$  satisfies the discrete Toda equation:

$$\psi_{N+1}^{n,m} \psi_{N-1}^{n,m} - (\psi_N^{n,m})^2 + \psi_N^{n+1,m} \psi_N^{n-1,m} = 0. \quad (2.15)$$

In general, (2.15) admits a solution expressed in terms of the Toeplitz type determinant

$$\psi_N^{n,m} = \det (c_{n-i+j,m})_{i,j=1,\dots,N} \quad (N > 0), \quad (2.16)$$

for an arbitrary function  $c_{n,m}$  under the boundary conditions

$$\psi_0^{n,m} = 1, \quad \psi_N^{n,m} = 0 \quad (N < 0). \quad (2.17)$$

Since the hypergeometric solutions to  $q$ - $P_{\text{III}}$  satisfy the conditions (2.17), the bilinear equation (2.15) is regarded as to fix the determinant structure of the solutions.

## 2.2 Hypergeometric solutions to $q$ -P<sub>II</sub>

Next, we review the hypergeometric solutions to  $q$ -P<sub>II</sub>. For each  $N \in \mathbb{Z}_{\geq 0}$ , let  $\phi_N^k$  be an  $N \times N$  determinant defined by

$$\phi_N^k = \begin{vmatrix} G_k & G_{k-1} & \cdots & G_{k-N+1} \\ G_{k+2} & G_{k+1} & \cdots & G_{k-N+3} \\ \vdots & \vdots & \ddots & \vdots \\ G_{k+2N-2} & G_{k+2N-3} & \cdots & G_{k+N-1} \end{vmatrix}, \quad \phi_0^k = 1, \quad (2.18)$$

where  $G_k$  satisfies

$$G_{k+1} - G_k + \frac{1}{a_0^2 q^k} G_{k-1} = 0. \quad (2.19)$$

**Lemma 2.4** ([25])  $\phi_N^k$  satisfies the following bilinear difference equations:

$$a_0^{-2} q^{-k+1} \phi_{N+1}^{k-2} \phi_N^{k+1} + \phi_{N+1}^k \phi_N^{k-1} - q^{-N} \phi_{N+1}^{k-1} \phi_N^k = 0, \quad (2.20)$$

$$q^N \phi_{N+1}^{k+1} \phi_N^{k-2} + a_0^{-2} q^{-k-N} \phi_{N+1}^{k-1} \phi_N^k - \phi_{N+1}^k \phi_N^{k-1} = 0. \quad (2.21)$$

**Proposition 2.5** ([25]) The hypergeometric solutions to  $q$ -P<sub>II</sub>, (2.2), with  $c = 1$  are given by

$$X_k = -a_0 q^{\frac{k}{2}+N} \frac{\phi_{N+1}^k \phi_N^{k-1}}{\phi_{N+1}^{k-1} \phi_N^k}. \quad (2.22)$$

Proposition 2.5 follows from Lemma 2.4.

**Remark 2.6** (1) The general solution to (2.19) is given by

$$G_k = \Theta(-a_0 q^{\frac{2k+1}{4}}; q^{\frac{1}{2}}) \times \left\{ A e^{\frac{k\pi i}{2}} {}_1\varphi_1 \left( \begin{matrix} 0 \\ -q^{\frac{1}{2}} \end{matrix}; q^{\frac{1}{2}}, -ia_0 q^{\frac{3+2k}{4}} \right) + B e^{-\frac{k\pi i}{2}} {}_1\varphi_1 \left( \begin{matrix} 0 \\ -q^{\frac{1}{2}} \end{matrix}; q^{\frac{1}{2}}, ia_0 q^{\frac{3+2k}{4}} \right) \right\}. \quad (2.23)$$

Here,  $\Theta(a; q)$  denotes the Jacobi theta function, which is defined by

$$\Theta(a; q) = (a; q)_{\infty} (qa^{-1}; q)_{\infty}, \quad (2.24)$$

and satisfies

$$\Theta(qa; q) = -a^{-1} \Theta(a; q). \quad (2.25)$$

(2)  $\phi_N^k$  also satisfies the bilinear equation

$$\phi_{N+1}^k \phi_{N-1}^{k+1} - \phi_N^k \phi_N^{k+1} + \phi_N^{k+2} \phi_N^{k-1} = 0, \quad (2.26)$$

which is a variant of the discrete Toda equation. Under the conditions

$$\phi_0^k = 1, \quad \phi_N^k = 0 \quad (N < 0), \quad (2.27)$$

(2.26) admits a solution expressed by

$$\phi_N^k = \det \left( c_{k+2i-j-1} \right)_{i,j=1,\dots,N} \quad (N > 0), \quad (2.28)$$

for an arbitrary function  $c_k$ . Hence, (2.26) can be regarded as the bilinear equation that fixes the determinant structure of the hypergeometric solutions to  $q$ -P<sub>II</sub>.



## 2.3 Comparing the hypergeometric solutions

By comparing the hypergeometric solutions to  $q$ -P<sub>III</sub> and  $q$ -P<sub>II</sub> (see Propositions 2.2 and 2.5, respectively) one may immediately notice that a naïve application of the specialization, (2.3), to the former does not yield the latter. As analogous to the phenomena seen in Section 1, we find the following differences between the two solutions:

- (i) the hypergeometric functions are different. In fact, substituting  $a_2 = q^{\frac{1}{2}}$  into (2.12) and (2.13) do not yield (2.19) and (2.23), respectively;
- (ii) the determinant structures are different.

**Remark 2.7** The correspondence between the rational solutions to  $q$ -P<sub>III</sub> (see [10]) and that to  $q$ -P<sub>II</sub> (see [25]) are straightforward. It is easily verified that substituting  $a_2 = q^{\frac{1}{2}}$  into the former yields the latter.

## 3 Projective reduction from $q$ -P<sub>III</sub> to $q$ -P<sub>II</sub>

### 3.1 Birational representation of $\widetilde{W}((A_2 + A_1)^{(1)})$

We formulate the family of Bäcklund transformations of  $q$ -P<sub>III</sub> as a birational representation of the extended affine Weyl group of type  $(A_2 + A_1)^{(1)}$  [11, 17]. We refer to [27] for basic ideas of this formulation.

We define the transformations  $s_i$  ( $i = 0, 1, 2$ ) and  $\pi$  on the variables  $f_j$  ( $j = 0, 1, 2$ ) and parameters  $a_k$  ( $k = 0, 1, 2$ ) by

$$s_i(a_j) = a_j a_i^{-a_{ij}}, \quad s_i(f_j) = f_j \left( \frac{a_i + f_i}{1 + a_i f_i} \right)^{u_{ij}}, \quad (3.1)$$

$$\pi(a_i) = a_{i+1}, \quad \pi(f_i) = f_{i+1}, \quad (3.2)$$

for  $i, j \in \mathbb{Z}/3\mathbb{Z}$ . Here the symmetric  $3 \times 3$  matrix

$$A = (a_{ij})_{i,j=0}^2 = \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix}, \quad (3.3)$$

is the Cartan matrix of type  $A_2^{(1)}$ , and the skew-symmetric one

$$U = (u_{ij})_{i,j=0}^2 = \begin{pmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{pmatrix}, \quad (3.4)$$

represents an orientation of the corresponding Dynkin diagram. We also define the transformations

$w_j$  ( $j = 0, 1$ ) and  $r$  by

$$w_0(f_i) = \frac{a_i a_{i+1} (a_{i-1} a_i + a_{i-1} f_i + f_{i-1} f_i)}{f_{i-1} (a_i a_{i+1} + a_i f_{i+1} + f_i f_{i+1})}, \quad w_0(a_i) = a_i, \quad (3.5)$$

$$w_1(f_i) = \frac{1 + a_i f_i + a_i a_{i+1} f_i f_{i+1}}{a_i a_{i+1} f_{i+1} (1 + a_{i-1} f_{i-1} + a_{i-1} a_i f_{i-1} f_i)}, \quad w_1(a_i) = a_i, \quad (3.6)$$

$$r(f_i) = \frac{1}{f_i}, \quad r(a_i) = a_i, \quad (3.7)$$

for  $i \in \mathbb{Z}/3\mathbb{Z}$ .

**Proposition 3.1 ([17])**  $\langle s_0, s_1, s_2, \pi, w_0, w_1, r \rangle$  forms the extended affine Weyl group of type  $(A_2 + A_1)^{(1)}$ . Namely, the transformations satisfy the fundamental relations

$$s_i^2 = (s_i s_{i+1})^3 = \pi^3 = 1, \quad \pi s_i = s_{i+1} \pi \quad (i \in \mathbb{Z}/3\mathbb{Z}), \quad w_0^2 = w_1^2 = r^2 = 1, \quad r w_0 = w_1 r, \quad (3.8)$$

and the actions of  $\widetilde{W}(A_2^{(1)}) = \langle s_0, s_1, s_2, \pi \rangle$  and  $\widetilde{W}(A_1^{(1)}) = \langle w_0, w_1, r \rangle$  commute with each other.

In general, for a function  $F = F(a_i, f_j)$  we let an element  $w \in \widetilde{W}((A_2 + A_1)^{(1)})$  act as  $w.F(a_i, f_j) = F(a_i.w, f_j.w)$ , that is,  $w$  acts on the arguments from the right. Note that  $a_0 a_1 a_2 = q$  and  $f_0 f_1 f_2 = qc^2$  are invariant under the actions of  $\widetilde{W}((A_2 + A_1)^{(1)})$  and  $\widetilde{W}(A_2^{(1)})$ , respectively. We define the translations  $T_i$  ( $i = 1, 2, 3, 4$ ) by

$$T_1 = \pi s_2 s_1, \quad T_2 = s_1 \pi s_2, \quad T_3 = s_2 s_1 \pi, \quad T_4 = r w_0, \quad (3.9)$$

whose actions on parameters  $a_i$  ( $i = 0, 1, 2$ ) and  $c$  are given by

$$\begin{aligned} T_1 &: (a_0, a_1, a_2, c) \mapsto (q a_0, q^{-1} a_1, a_2, c), \\ T_2 &: (a_0, a_1, a_2, c) \mapsto (a_0, q a_1, q^{-1} a_2, c), \\ T_3 &: (a_0, a_1, a_2, c) \mapsto (q^{-1} a_0, a_1, q a_2, c), \\ T_4 &: (a_0, a_1, a_2, c) \mapsto (a_0, a_1, a_2, qc). \end{aligned} \quad (3.10)$$

Note that  $T_i$  ( $i = 1, 2, 3, 4$ ) commute with each other and  $T_1 T_2 T_3 = 1$ . The action of  $T_1$  on  $f$ -variables can be expressed as

$$T_1(f_1) = \frac{qc^2}{f_1 f_0} \frac{1 + a_0 f_0}{a_0 + f_0}, \quad T_1(f_0) = \frac{qc^2}{f_0 T_1(f_1)} \frac{1 + a_2 a_0 T_1(f_1)}{a_2 a_0 + T_1(f_1)}. \quad (3.11)$$

Or, applying  $T_1^n T_2^m T_4^N$  ( $n, m, N \in \mathbb{Z}$ ) on (3.11) and putting

$$f_{i,N}^{n,m} = T_1^n T_2^m T_4^N(f_i) \quad (i = 0, 1, 2), \quad (3.12)$$

we obtain

$$f_{1,N}^{n+1,m} = \frac{q^{2N+1} c^2}{f_{1,N}^{n,m} f_{0,N}^{n,m}} \frac{1 + a_0 q^n f_{0,N}^{n,m}}{a_0 q^n + f_{0,N}^{n,m}}, \quad f_{0,N}^{n+1,m} = \frac{q^{2N+1} c^2}{f_{0,N}^{n,m} f_{1,N}^{n+1}} \frac{1 + a_2 a_0 q^{n-m} f_{1,N}^{n+1,m}}{a_2 a_0 q^{n-m} + f_{1,N}^{n+1,m}}, \quad (3.13)$$

which is equivalent to  $q$ -P<sub>III</sub>, (2.1). Then,  $T_i$  ( $i = 1, 2, 3, 4$ ) are regarded as Bäcklund transformations of  $q$ -P<sub>III</sub>.

In order to formulate the symmetrization to  $q$ -P<sub>II</sub>, it is crucial to introduce the transformation  $R_1$  defined by

$$R_1 = \pi^2 s_1, \quad (3.14)$$

which satisfies

$$R_1^2 = T_1. \quad (3.15)$$

The actions of  $R_1$  are given by

$$R_1 : (a_0, a_1, a_2, c) \mapsto (a_2 a_0, a_0^{-1}, a_1 a_0, c), \quad (3.16)$$

$$R_1(f_0) = \frac{qc^2}{f_0 f_1} \frac{1 + a_0 f_0}{a_0 + f_0}, \quad R_1(f_1) = f_0, \quad (3.17)$$

which describe the zig-zag motion around the line  $a_2 = q^{\frac{1}{2}}$  on the parameter space. However, if we put  $a_2 = q^{\frac{1}{2}}$ , then  $R_1$  becomes the translation on the line  $a_2 = q^{\frac{1}{2}}$  with the step  $q^{\frac{1}{2}}$  (see Figure 1). In fact, the actions of  $R_1$  are now given by

$$R_1 : (a_0, a_1, c) \mapsto (q^{\frac{1}{2}} a_0, q^{-\frac{1}{2}} a_1, c), \quad (3.18)$$

$$R_1(f_0) = \frac{qc^2}{f_0 f_1} \frac{1 + a_0 f_0}{a_0 + f_0}, \quad R_1(f_1) = f_0. \quad (3.19)$$

Applying  $R_1^k T_4^N$  on (3.19) and putting

$$f_{i,N}^k = R_1^k T_4^N(f_i) \quad (i = 0, 1, 2), \quad (3.20)$$

we have

$$f_{0,N}^{k+1} = \frac{q^{2N+1} c^2}{f_{0,N}^k f_{0,N}^{k-1}} \frac{1 + a_0 q^{\frac{k}{2}} f_{0,N}^k}{a_0 q^{\frac{k}{2}} + f_{0,N}^k}, \quad (3.21)$$

which is equivalent to  $q$ -P<sub>II</sub>, (2.2). Then,  $R_1$  and  $T_4$  are regarded as Bäcklund transformations of  $q$ -P<sub>II</sub>.




Figure 1. Action of  $R_1$  on the parameter space  $\mathbf{a} = (a_0, a_1, a_2) \in (\mathbb{C}^\times)^3$  with  $a_0 a_1 a_2 = q$ . Left: generic case. Right:  $a_2 = q^{\frac{1}{2}}$ .

In general, it is possible to obtain various discrete dynamical systems of Painlevé type from elements of infinite order that are not necessarily translations in the affine Weyl group by taking a projection on an appropriate sublattice of corresponding root lattice. We call such a procedure a *projective reduction*.

By using the above formulation, we can now explain why the difference of hypergeometric solutions to  $q$ -P<sub>III</sub> and that to  $q$ -P<sub>II</sub> occurs.

### 3.2 Hypergeometric functions

First, we explain about the difference of hypergeometric functions. For convenience, we define the function  $H_{n,m}$  by

$$H_{n,m} = \Theta(a_0^{-2} q^{2n+1}; q^2) F_{n+\frac{1}{2},m}, \quad (3.22)$$

where  $F_{n,m}$  is given in Remark 2.3. Then, we obtain from (2.12) with  $a_2 = q^{\frac{1}{2}}$  the three-term relation for  $H_{n,0}$ :

$$\left[ T_1^2 + \left( q^{-3-2n} a_0^{-2} + q^{-2-2n} a_0^{-2} - 1 \right) T_1 + q^{-3-4n} a_0^{-4} \right] H_{n,0} = 0. \quad (3.23)$$

Let  $n = 0$ . Since  $R_1^2 = T_1$ , the linear difference operator in (3.23) is fourth order with respect to  $R_1$ . Moreover, it admits the following factorization into the second order linear difference operators:

$$T_1^2 + \left( a_0^{-2} q^{-3} + a_0^{-2} q^{-2} - 1 \right) T_1 + a_0^{-4} q^{-3} = \left( R_1^2 + R_1 + a_0^{-2} q^{-2} \right) \left( R_1^2 - R_1 + a_0^{-2} q^{-1} \right). \quad (3.24)$$

On the other hand, the three-term relation for  $G_0$  (see (2.19)) can be expressed as

$$\left( R_1^2 - R_1 + a_0^{-2} q^{-1} \right) G_0 = 0. \quad (3.25)$$

Note that the second factor in the right-hand side of (3.24) is exactly the operator in (3.25), thus,  $G_0$  also satisfies (3.23) with  $n = 0$ . This factorization (3.24) implies that  $G_k$  can not be obtained simply from  $F_{n,m}$  by a specialization of parameters (2.3).

### 3.3 Determinant structure

Next, in order to discuss the difference of determinant structures, we need to introduce the  $\tau$  functions and lift the representation to the Weyl group on the level of  $\tau$  functions [17, 40]. We introduce  $\tau_i$  and  $\bar{\tau}_i$  ( $i \in \mathbb{Z}/3\mathbb{Z}$ ) with

$$f_i = q^{\frac{1}{3}} c^{\frac{2}{3}} \frac{\bar{\tau}_{i+1} \tau_{i-1}}{\tau_{i+1} \bar{\tau}_{i-1}}. \quad (3.26)$$

**Proposition 3.2 ([40])** *We define the action of  $s_i$  ( $i = 0, 1, 2$ ),  $\pi$ ,  $w_j$  ( $j = 0, 1$ ), and  $r$  on  $\tau_k$  and  $\bar{\tau}_k$  ( $k = 0, 1, 2$ ) by the following formulae:*

$$\begin{cases} s_i(\tau_i) = \frac{u_i \tau_{i+1} \bar{\tau}_{i-1} + \bar{\tau}_{i+1} \tau_{i-1}}{u_i^{\frac{1}{2}} \bar{\tau}_i}, & s_i(\tau_j) = \tau_j \quad (i \neq j), \\ s_i(\bar{\tau}_i) = \frac{v_i \bar{\tau}_{i+1} \tau_{i-1} + \tau_{i+1} \bar{\tau}_{i-1}}{v_i^{\frac{1}{2}} \tau_i}, & s_i(\bar{\tau}_j) = \bar{\tau}_j \quad (i \neq j), \end{cases} \quad (3.27)$$

$$\pi(\tau_i) = \tau_{i+1}, \quad \pi(\bar{\tau}_i) = \bar{\tau}_{i+1}, \quad (3.28)$$

$$\begin{cases} w_0(\bar{\tau}_i) = \frac{a_{i+1}^{\frac{1}{3}} (\bar{\tau}_i \tau_{i+1} \tau_{i+2} + u_{i-1} \tau_i \bar{\tau}_{i+1} \tau_{i+2} + u_{i+1}^{-1} \tau_i \tau_{i+1} \bar{\tau}_{i+2})}{a_{i+2}^{\frac{1}{3}} \bar{\tau}_{i+1} \bar{\tau}_{i+2}}, \\ w_0(\tau_i) = \tau_i, \end{cases} \quad (3.29)$$

$$\begin{cases} w_1(\tau_i) = \frac{a_{i+1}^{\frac{1}{3}}(\tau_i \bar{\tau}_{i+1} \bar{\tau}_{i+2} + v_{i-1} \bar{\tau}_i \tau_{i+1} \bar{\tau}_{i+2} + v_{i+1}^{-1} \bar{\tau}_i \bar{\tau}_{i+1} \tau_{i+2})}{a_{i+2}^{\frac{1}{3}} \tau_{i+1} \tau_{i+2}}, \\ w_1(\bar{\tau}_i) = \bar{\tau}_i, \end{cases} \quad (3.30)$$

$$r(\tau_i) = \bar{\tau}_i, \quad r(\bar{\tau}_i) = \tau_i, \quad (3.31)$$

with

$$u_i = q^{-\frac{1}{3}} c^{-\frac{2}{3}} a_i, \quad v_i = q^{\frac{1}{3}} c^{\frac{2}{3}} a_i, \quad (3.32)$$

where  $i, j \in \mathbb{Z}/3\mathbb{Z}$ . Then,  $\langle s_0, s_1, s_2, \pi, w_0, w_1, r \rangle$  realizes the affine Weyl group  $\widetilde{W}((A_2 + A_1)^{(1)})$ .




Figure 2. Configuration of the  $\tau$  functions on the lattice with  $N = 0$ .

Then, we define the  $\tau$  functions  $\tau_N^{n,m}$  ( $n, m, N \in \mathbb{Z}$ ) by

$$\tau_N^{n,m} = T_1^n T_2^m T_4^N(\tau_1). \quad (3.33)$$

We note that  $\tau_0 = \tau_0^{-1,0}$ ,  $\tau_1 = \tau_0^{0,0}$ ,  $\tau_2 = \tau_0^{0,1}$ ,  $\bar{\tau}_0 = \tau_1^{-1,0}$ ,  $\bar{\tau}_1 = \tau_1^{0,0}$ , and  $\bar{\tau}_2 = \tau_1^{0,1}$ .

**Proposition 3.3** *The action of  $\widetilde{W}((A_2 + A_1)^{(1)})$  on  $\tau_N^{n,m}$  is*

$$s_0(\tau_N^{n,m}) = \tau_N^{-n,m-n}, \quad s_1(\tau_N^{n,m}) = \tau_N^{m-1,n+1}, \quad s_2(\tau_N^{n,m}) = \tau_N^{n-m,-m}, \quad \pi(\tau_N^{n,m}) = \tau_N^{-m,n-m+1}, \quad (3.34)$$

$$w_0(\tau_N^{n,m}) = \tau_{-N}^{n,m}, \quad w_1(\tau_N^{n,m}) = \tau_{2-N}^{n,m}, \quad r(\tau_N^{n,m}) = \tau_{1-N}^{n,m}. \quad (3.35)$$

For convenience, we put

$$\alpha_i = a_i^{\frac{1}{6}}, \quad \gamma = c^{\frac{1}{6}}, \quad Q = q^{\frac{1}{6}}. \quad (3.36)$$

Though it is possible to derive more various bilinear difference equations from Proposition 3.2, we present here only the equations that are directly relevant to  $q$ -P<sub>III</sub>, (3.11).

**Proposition 3.4** *The following bilinear equations hold:*

$$\tau_{N+1}^{n,m} \tau_N^{n+1,m+1} - Q^{-3n+3m+2N-2} \gamma^2 \alpha_1^3 \tau_N^{n+1,m} \tau_{N+1}^{n,m+1} + Q^{-6n+6m+4N-4} \gamma^4 \alpha_1^6 \tau_N^{n,m} \tau_{N+1}^{n+1,m+1} = 0, \quad (3.37)$$

$$\tau_{N+1}^{n+1,m+1} \tau_N^{n+1,m} - Q^{3n+2N+4} \gamma^2 \alpha_0^3 \tau_N^{n,m} \tau_{N+1}^{n+2,m+1} + Q^{6n+4N+8} \gamma^4 \alpha_0^6 \tau_N^{n+1,m+1} \tau_{N+1}^{n+1,m} = 0, \quad (3.38)$$

$$\tau_{N+1}^{n+1,m+1} \tau_N^{n,m} - Q^{-3n+3m-2N-4} \gamma^{-2} \alpha_1^3 \tau_{N+1}^{n+1,m} \tau_N^{n,m+1} + Q^{-6n+6m-4N-8} \gamma^{-4} \alpha_1^6 \tau_N^{n+1,m+1} \tau_{N+1}^{n,m} = 0, \quad (3.39)$$

$$\tau_{N+1}^{n+1,m} \tau_N^{n+1,m+1} - Q^{3n-2N+2} \gamma^{-2} \alpha_0^3 \tau_{N+1}^{n,m} \tau_N^{n+2,m+1} + Q^{6n-4N+4} \gamma^{-4} \alpha_0^6 \tau_N^{n+1,m} \tau_{N+1}^{n+1,m+1} = 0, \quad (3.40)$$

$$\tau_{N+1}^{n,m} \tau_{N-1}^{n,m} + Q^{-8n+4m-4} \alpha_0^{-4} \alpha_1^4 (\tau_N^{n,m})^2 - Q^{-2n+m-1} \alpha_0^{-1} \alpha_1 \tau_N^{n+1,m} \tau_N^{n-1,m} = 0. \quad (3.41)$$

The proof of Proposition 3.4 will be given in the appendix A.1.

As seen below  $q$ -P<sub>III</sub>, (3.11) or (3.13), can be obtained from the bilinear equations. Noticing that

$$f_{0,N}^{n,m} = Q^{4N+2} \gamma^4 \frac{\tau_{N+1}^{n,m} \tau_N^{n,m+1}}{\tau_N^{n,m} \tau_{N+1}^{n,m+1}}, \quad f_{1,N}^{n,m} = Q^{4N+2} \gamma^4 \frac{\tau_{N+1}^{n,m+1} \tau_N^{n-1,m}}{\tau_N^{n,m+1} \tau_{N+1}^{n-1,m}}, \quad f_{2,N}^{n,m} = Q^{4N+2} \gamma^4 \frac{\tau_{N+1}^{n-1,m} \tau_N^{n,m}}{\tau_N^{n-1,m} \tau_{N+1}^{n,m}}, \quad (3.42)$$

we can rewrite (3.37) and (3.39) as

$$1 + Q^{-6n+6m-6} \alpha_1^6 f_{1,N}^{n+1,m} = Q^{-3n+3m+2N-2} \gamma^2 \alpha_1^3 \frac{\tau_N^{n+1,m} \tau_{N+1}^{n,m+1}}{\tau_{N+1}^{n,m} \tau_N^{n+1,m+1}}, \quad (3.43)$$

$$1 + Q^{6n-6m+6} \alpha_1^{-6} f_{1,N}^{n+1,m} = Q^{3n-3m+2N+4} \gamma^2 \alpha_1^{-3} \frac{\tau_{N+1}^{n+1,m} \tau_N^{n,m+1}}{\tau_{N+1}^{n,m} \tau_N^{n+1,m+1}}, \quad (3.44)$$

respectively. Dividing (3.44) by (3.43), we have

$$\frac{1 + Q^{6n-6m+6} \alpha_1^{-6} f_{1,N}^{n+1,m}}{1 + Q^{-6n+6m-6} \alpha_1^6 f_{1,N}^{n+1,m}} = Q^{6n-6m+6} \alpha_1^{-6} \frac{\tau_{N+1}^{n+1,m} \tau_N^{n,m+1}}{\tau_N^{n+1,m} \tau_{N+1}^{n,m+1}} = Q^{6n-6m+6} \alpha_1^{-6} \frac{f_{0,N}^{n,m}}{f_{2,N}^{n,m}}, \quad (3.45)$$

which is equivalent to the second equation of (3.13). Similarly, (3.38) and (3.40) yield the first equation of (3.13).

For the hypergeometric solutions, we relate the  $\tau$  functions to the determinants  $\psi_N^{n,m}$ , (2.5), by multiplication of appropriate “gauge” factor. Set

$$\tau_N^{n,m} = (-1)^{\frac{N(N+1)}{2}} Q^{-2(2n-m)N^2+6Nn} \alpha_0^{-4N^2+6N} \alpha_2^{-2N^2} \left( \frac{\Theta(-Q^{-6n} \alpha_0^{-6}, Q^6) \Theta(-Q^{6m} \alpha_2^{-6}, Q^6)}{\Theta(Q^{-6(n-m)} \alpha_0^{-6} \alpha_2^{-6}, Q^6)} \right)^N \\ \times \Gamma(Q^{2n-m+1} \alpha_0^2 \alpha_2; Q, Q) \Gamma(Q^{-n+2m-1} \alpha_1^2 \alpha_0; Q, Q) \Gamma(Q^{-n-m} \alpha_2^2 \alpha_1; Q, Q) \psi_N^{n,m-1}, \quad (3.46)$$

where  $\Gamma(a; p, q)$  denotes the Elliptic gamma function, which is defined by

$$\Gamma(a; p, q) = \frac{(q^2 a^{-1}; p, q)_\infty}{(a; p, q)_\infty}, \quad (3.47)$$

and satisfies

$$\Gamma(qa; q, q) = \Theta(a, q) \Gamma(a; q, q). \quad (3.48)$$

Let  $\gamma = 1$ . Then the bilinear equations (3.37)–(3.41) can be rewritten as

$$\psi_{N+1}^{n,m} \psi_N^{n+1,m+1} - Q^{-12n+12N} \alpha_0^{-12} \psi_N^{n+1,m} \psi_{N+1}^{n,m+1} + Q^{-12n} \alpha_0^{-12} \psi_N^{n,m} \psi_{N+1}^{n+1,m+1} = 0, \quad (3.49)$$

$$\psi_{N+1}^{n+1,m+1} \psi_N^{n+1,m} - Q^{-12N} \psi_N^{n,m} \psi_{N+1}^{n+2,m+1} - Q^{12n+12} \alpha_0^{12} \psi_N^{n+1,m+1} \psi_{N+1}^{n+1,m} = 0, \quad (3.50)$$

$$\psi_{N+1}^{n+1,m+1} \psi_N^{n,m} - \psi_{N+1}^{n+1,m} \psi_N^{n,m+1} + Q^{12m+12} \alpha_2^{-12} \psi_N^{n+1,m+1} \psi_{N+1}^{n,m} = 0, \quad (3.51)$$

$$\psi_{N+1}^{n+1,m} \psi_N^{n+1,m+1} - Q^{12m+12} \alpha_2^{-12} \psi_{N+1}^{n,m} \psi_N^{n+2,m+1} - \psi_N^{n+1,m} \psi_{N+1}^{n+1,m+1} = 0, \quad (3.52)$$

$$\psi_{N+1}^{n,m} \psi_{N-1}^{n,m} - (\psi_N^{n,m})^2 + \psi_N^{n+1,m} \psi_N^{n-1,m} = 0, \quad (3.53)$$

respectively. Equations (3.49)–(3.52) are equivalent to (2.7)–(2.10). Note that (3.53) is exactly the discrete Toda equation, (2.15), which fixes the determinant structure of the hypergeometric solutions as mentioned in Remark 2.3.

**Remark 3.5** The gauge factor  $\tau_N^{n,m}/\psi_N^{n,m-1}$  in (3.46) is obtained by solving the overdetermined system of the bilinear difference equations with  $\gamma = 1$  under the boundary conditions  $\tau_N^{n,m} = 0$  ( $N \in \mathbb{Z}_{<0}$ ) [26].

Let us consider the bilinear equations for  $q$ -P<sub>II</sub>. Since we need  $R_1$ ,  $\tau_i$ , and  $\bar{\tau}_i$  ( $i \in \mathbb{Z}/3\mathbb{Z}$ ), the lattice is restricted to the “unit-strip” (see Figure 3). Therefore, we have only to consider  $\tau_N^{n,0}$  and  $\tau_N^{n,1}$  ( $n, N \in \mathbb{Z}$ ). We set

$$\tau_N^k = R_1^k T_4^N(\tau_1). \quad (3.54)$$

Note that

$$\tau_0 = \tau_0^{-2}, \quad \tau_1 = \tau_0^0, \quad \tau_2 = \tau_0^{-1}, \quad \bar{\tau}_0 = \tau_1^{-2}, \quad \bar{\tau}_1 = \tau_1^0, \quad \bar{\tau}_2 = \tau_1^{-1}. \quad (3.55)$$

In general, it follows that

$$\tau_N^{n,0} = \tau_N^{2n}, \quad \tau_N^{n,1} = \tau_N^{2n-1}, \quad (3.56)$$

and

$$f_{0,N}^k = Q^{4N+2} \gamma^4 \frac{\tau_{N+1}^k \tau_N^{k-1}}{\tau_N^k \tau_{N+1}^{k-1}}. \quad (3.57)$$




Figure 3. The actions of  $R_1$  on  $\tau_i$  ( $i = 0, 1, 2$ ).

**Proposition 3.6** *The following bilinear equations hold:*

$$Q^{\frac{-3k+4N+2}{2}} \gamma^2 \alpha_0^{-3} \tau_N^{k+1} \tau_{N+1}^{k-2} - Q^{-3k+4N+2} \gamma^4 \alpha_0^{-6} \tau_N^{k-1} \tau_{N+1}^k - \tau_{N+1}^{k-1} \tau_N^k = 0, \quad (3.58)$$

$$Q^{\frac{-3k-4N-2}{2}} \gamma^{-2} \alpha_0^{-3} \tau_{N+1}^{k+1} \tau_N^{k-2} - Q^{-3k-4N-2} \gamma^{-4} \alpha_0^{-6} \tau_N^k \tau_{N+1}^{k-1} - \tau_{N+1}^k \tau_N^{k-1} = 0, \quad (3.59)$$

$$\tau_{N+1}^k \tau_{N-1}^{k+1} - Q^{\frac{k-4N+1}{2}} \gamma^{-2} \alpha_0 \tau_N^{k+2} \tau_N^{k-1} - Q^{-k+4N-1} \gamma^4 \alpha_0^{-2} \tau_N^k \tau_N^{k+1} = 0. \quad (3.60)$$

The proof of Proposition 3.6 will be given in the appendix A.2.

One can obtain  $q$ -P<sub>II</sub>, (3.19), from Proposition 3.6 as follows. Equations (3.58) and (3.59) can be rewritten as

$$1 + Q^{-3k} \alpha_0^{-6} f_{0,N}^k = Q^{\frac{-3k+4N+2}{2}} \gamma^2 \alpha_0^{-3} \frac{\tau_N^{k+1} \tau_{N+1}^{k-2}}{\tau_{N+1}^{k-1} \tau_N^k}, \quad (3.61)$$

$$1 + Q^{3k} \alpha_0^6 f_{0,N}^k = Q^{\frac{3k+4N+2}{2}} \gamma^2 \alpha_0^3 \frac{\tau_{N+1}^{k+1} \tau_N^{k-2}}{\tau_{N+1}^{k-1} \tau_N^k}. \quad (3.62)$$

Dividing (3.62) by (3.61), we have

$$\frac{1 + Q^{3k} \alpha_0^6 f_{0,N}^k}{1 + Q^{-3k} \alpha_0^{-6} f_{0,N}^k} = Q^{3k} \alpha_0^6 \frac{\tau_{N+1}^{k+1} \tau_N^{k-2}}{\tau_N^{k+1} \tau_{N+1}^{k-2}} = Q^{3k-12N-6} \gamma^{12} \alpha_0^6 f_{0,N}^{k+1} f_{0,N}^k f_{0,N}^{k-1}, \quad (3.63)$$

which is equivalent to (3.21).

For hypergeometric solutions, by putting  $\gamma = 1$  and

$$\tau_N^k = (-1)^{\frac{N(N-1)}{2}} Q^{N(N-1)(k+n)} \alpha_0^{2N(N-1)} \frac{\Gamma(Q^{\frac{2k+3}{2}} \alpha_0^2; Q, Q) \Gamma(Q^{-\frac{k}{2}} \alpha_0^{-1}; Q, Q) \Gamma(Q^{-\frac{k+3}{2}} \alpha_0^{-1}; Q, Q)}{\Theta(Q^{3k+1} \alpha_0^6, Q^3)^N} \phi_n^k, \quad (3.64)$$

we can rewrite the bilinear equations (3.58), (3.59), and (3.60) as

$$Q^{6N-6k+6} \alpha_0^{-12} \phi_N^{k+1} \phi_{N+1}^{k-2} + Q^{6N} \phi_N^{k-1} \phi_{N+1}^k - \phi_{N+1}^{k-1} \phi_N^k = 0, \quad (3.65)$$

$$Q^{6N} \phi_{N+1}^{k+1} \phi_N^{k-2} + Q^{-6N-6k} \alpha_0^{-12} \phi_N^k \phi_{N+1}^{k-1} - \phi_{N+1}^k \phi_N^{k-1} = 0, \quad (3.66)$$

$$\phi_{N+1}^k \phi_{N-1}^{k+1} - \phi_N^k \phi_N^{k+1} + \phi_N^{k+2} \phi_N^{k-1} = 0, \quad (3.67)$$

which are equivalent to (2.20), (2.21), and (2.26), respectively. The determinant structure of the hypergeometric solutions is fixed by (3.67) as was explained in Remark 2.6.

Therefore, the difference of the determinant structures of the hypergeometric solutions to  $q$ -P<sub>III</sub> and that to  $q$ -P<sub>II</sub> originates from the following procedures:

- (i) the specialization  $a_2 = q^{\frac{1}{2}}$  and the restriction of  $\tau$  functions on the “unit-strip”;
- (ii) taking the half-step translation  $R_1$  instead of  $T_1$  as a time evolution.

These result in the difference of the bilinear equations (3.41) (or (3.53)) and (3.60) (or (3.67)), which fix the determinant structure of the hypergeometric solutions.

## 4 Concluding remarks

In this paper, we have clarified the mechanism that gives rise to the apparent “inconsistency” in the hypergeometric solutions to  $q$ -P<sub>III</sub> and that to  $q$ -P<sub>II</sub> by using their underlying affine Weyl group symmetry. In general, it is also possible to explain the inconsistency among the hypergeometric solutions to other symmetric and asymmetric discrete Painlevé equations (see, for example, Propositions 1.1 and 1.2).



We shall make a slightly technical remark on the solutions to  $q$ -P<sub>II</sub>. Besides the hypergeometric solution to  $q$ -P<sub>II</sub> in Proposition 2.5, one can also obtain another solution to  $q$ -P<sub>II</sub> from that to  $q$ -P<sub>III</sub> in Proposition 2.2 through a naïve specialization (2.3). This solution, however, takes different expressions according to the parity of the time variable  $k$  of  $q$ -P<sub>II</sub>, (2.2). On the other hand, the solution in Proposition 2.5 forms a smooth function in  $k$ . In this sense it is more natural as a solution to  $q$ -P<sub>II</sub>.

Before closing, we demonstrate another example of the projective reductions. Let us consider the following system of difference equations [28]:

$$Z_n + X_n = \frac{3na + b_1}{Y_n} + t, \quad X_{n+1} + Y_n = \frac{(3n+1)a + b_2}{Z_n} + t, \quad Y_{n+1} + Z_n = \frac{(3n+2)a + b_3}{Z_n} + t, \quad (4.1)$$

where  $X_n$ ,  $Y_n$ , and  $Z_n$  are the dependent variables,  $n \in \mathbb{Z}$  is the independent variable, and  $a, b_1, b_2, b_3, t \in \mathbb{C}$  are parameters. Equation (4.1) is one of the discrete Painlevé systems of type  $A_3^{(1)}$ . Namely, it arises from a Bäcklund transformation of the Painlevé V equation, which describes a translation in a different direction from (1.4). Putting  $b_1 = b_2 = b_3 = b$ ,  $X_n = x_{3n-1}$ ,  $Y_n = x_{3n}$ , and  $Z_n = x_{3n+1}$ , we can reduce (4.1) to

$$x_{n+1} + x_{n-1} = \frac{an + b}{x_n} + t, \quad (4.2)$$

which is known as a discrete Painlevé I equation [36]. This reduction from (4.1) to (4.2) is a typical example of the projective reductions other than a symmetrization.

It seems that various projective reductions of the discrete Painlevé systems change the underlying symmetry and yield a number of intriguing problems. One interesting project is to make a list of the hypergeometric functions that appear as the solutions to all the symmetric discrete Painlevé equations in Sakai's classification [13, 14, 39]. These will be discussed in forthcoming papers [16].

**Acknowledgement.** The authors would like to express their sincere thanks to Prof. M. Noumi for fruitful discussions and valuable suggestions. They also acknowledge continuous encouragement by Prof. T. Masuda, Prof. H. Sakai, and Prof. Y. Yamada. This work has been partially supported by the JSPS Grant-in-Aid for Scientific Research No. 19340039.

## A Derivation of bilinear equations

In this appendix, we derive various bilinear equations for  $\tau$  functions from the birational representations of  $\widetilde{W}\left((A_2 + A_1)^{(1)}\right)$  given in Proposition 3.2.

### A.1 Bilinear equations for $q$ -P<sub>III</sub>

We use the notations introduced in (3.33) and (3.36). For convenience, we classify the bilinear equations into six types so that any equations which belong to the same type can be transformed into each other by the action of  $\widetilde{W}\left((A_2 + A_1)^{(1)}\right)$ .

**Proposition A.1 (Type I: Discrete Toda type)** *The following bilinear equations hold:*

$$\tau_{N+1}^{n,m} \tau_{N-1}^{n,m} + Q^{4n-8m+4} \alpha_1^{-4} \alpha_2^4 (\tau_N^{n,m})^2 - Q^{n-2m+1} \alpha_1^{-1} \alpha_2 \tau_N^{n,m+1} \tau_N^{n,m-1} = 0, \quad (\text{A.1})$$

$$\tau_{N+1}^{n,m} \tau_{N-1}^{n,m} + Q^{4n+4m} \alpha_0^4 \alpha_2^{-4} (\tau_N^{n,m})^2 - Q^{n+m} \alpha_0 \alpha_2^{-1} \tau_N^{n+1,m+1} \tau_N^{n-1,m-1} = 0, \quad (\text{A.2})$$

$$\tau_{N+1}^{n,m} \tau_{N-1}^{n,m} + Q^{-8n+4m-4} \alpha_0^{-4} \alpha_1^4 (\tau_N^{n,m})^2 - Q^{-2n+m-1} \alpha_0^{-1} \alpha_1 \tau_N^{n+1,m} \tau_N^{n-1,m} = 0. \quad (\text{A.3})$$




Figure 4. Configuration of  $\tau$  functions for the bilinear equations of type I. Left: (A.1), center: (A.2), right: (A.3).

**Proof.** Application of  $T_4 = rw_0$  on  $\bar{\tau}_0$  yields

$$T_4(\bar{\tau}_0) = c^{-\frac{2}{3}} a_0^{-\frac{1}{3}} a_1^{-1} a_2^{-\frac{2}{3}} \frac{\bar{\tau}_0 \bar{\tau}_1}{\tau_1} + c^{\frac{2}{3}} a_0^{\frac{1}{3}} a_1^{\frac{2}{3}} a_2 \frac{\bar{\tau}_0 \bar{\tau}_2}{\tau_2} + a_1^{\frac{1}{3}} a_2^{-\frac{1}{3}} \frac{\tau_0 \bar{\tau}_1 \bar{\tau}_2}{\tau_1 \tau_2}, \quad (\text{A.4})$$

which is rearranged as

$$T_4(\bar{\tau}_0) - c^{-\frac{2}{3}} a_0^{-\frac{1}{3}} a_1^{-1} a_2^{-\frac{2}{3}} \frac{\bar{\tau}_0 \bar{\tau}_1}{\tau_1} \left( \frac{q^{\frac{1}{3}} c^{\frac{2}{3}} a_1 \tau_0 \bar{\tau}_2 + \bar{\tau}_0 \tau_2}{\bar{\tau}_0 \tau_2} \right) \left( \frac{q^{\frac{1}{3}} c^{\frac{2}{3}} a_2 \tau_1 \bar{\tau}_0 + \bar{\tau}_1 \tau_0}{\bar{\tau}_1 \tau_0} \right) + a_1^{-\frac{2}{3}} a_2^{\frac{2}{3}} \frac{\bar{\tau}_0^2}{\tau_0} = 0. \quad (\text{A.5})$$

Applying  $T_2 = s_2 \pi s_1$  and  $T_3 = s_2 s_1 \pi$  on  $\bar{\tau}_0$  and  $\bar{\tau}_1$ , respectively, we obtain

$$q^{\frac{1}{6}} c^{\frac{1}{3}} a_1^{\frac{1}{2}} \tau_1 T_2(\bar{\tau}_0) = q^{\frac{1}{3}} c^{\frac{2}{3}} a_1 \tau_0 \bar{\tau}_2 + \bar{\tau}_0 \tau_2, \quad (\text{A.6})$$

$$q^{\frac{1}{6}} c^{\frac{1}{3}} a_2^{\frac{1}{2}} \tau_2 T_3(\bar{\tau}_1) = q^{\frac{1}{3}} c^{\frac{2}{3}} a_2 \tau_1 \bar{\tau}_0 + \bar{\tau}_1 \tau_0. \quad (\text{A.7})$$

Using (A.6) and (A.7), we can rewrite (A.5) as

$$T_4^2(\tau_0) \tau_0 + a_1^{-\frac{2}{3}} a_2^{\frac{2}{3}} T_4(\tau_0)^2 - a_1^{-\frac{1}{6}} a_2^{\frac{1}{6}} T_2 T_4(\tau_0) T_3 T_4(\tau_1) = 0. \quad (\text{A.8})$$

Then by applying  $T_1^{l+1} T_2^m T_4^{n-1}$ ,  $T_1^l T_2^m T_4^{n-1} \pi$ , and  $T_1^l T_2^{m-1} T_4^{n-1} \pi^2$  on (A.8), we obtain (A.1), (A.2), and (A.3), respectively.  $\square$

Figure 4 shows the configuration of  $\tau$  functions in the bilinear equations. Each bilinear equation takes the form of a linear combination of the three quadratic terms in  $\tau$  functions. In the left figure, we mark the first, the second, and the third multiplication of  $\tau$  functions of (A.1) with the square, the circle, and the triangle, respectively. In the rest of this paper, we use similar representations as above.

**Proposition A.2 (Type II: Discrete 2d-Toda type)** *The following bilinear difference equations hold:*

$$(1 - Q^{-12m} \alpha_2^{12}) \tau_{N+1}^{n,m} \tau_{N-1}^{n,m} + Q^{n-11m} \alpha_0 \alpha_2^{11} \tau_N^{n+1,m+1} \tau_N^{n-1,m-1} - Q^{n-2m} \alpha_0 \alpha_2^2 \tau_N^{n,m+1} \tau_N^{n,m-1} = 0, \quad (\text{A.9})$$

$$(1 - Q^{12n} \alpha_0^{12}) \tau_{N+1}^{n,m} \tau_{N-1}^{n,m} + Q^{10n+m} \alpha_0^{10} \alpha_2^{-1} \tau_N^{n+1,m} \tau_N^{n-1,m} - Q^{n+m} \alpha_0 \alpha_2^{-1} \tau_N^{n+1,m+1} \tau_N^{n-1,m-1} = 0, \quad (\text{A.10})$$

$$(1 - Q^{12n-12m} \alpha_0^{12} \alpha_2^{12}) \tau_{N+1}^{n,m} \tau_{N-1}^{n,m} + Q^{10n-11m} \alpha_0^{10} \alpha_2^{11} \tau_N^{n+1,m} \tau_N^{n-1,m} - Q^{n-2m} \alpha_0 \alpha_2^2 \tau_N^{n,m+1} \tau_N^{n,m-1} = 0. \quad (\text{A.11})$$




Figure 5. Configuration of  $\tau$  functions for the bilinear equations of type II. Left: (A.9), center: (A.10), right: (A.11).

**Proof.** Equation (A.9) is derived by eliminating  $\tau_{l,m,n}$  from (A.1) and (A.2). We obtain (A.10) and (A.11) in a similar manner.  $\square$

**Proposition A.3 (Type III)** *The following bilinear equations hold:*

$$(Q^{4n-8m+4} \alpha_1^{-4} \alpha_2^4 - Q^{4l+4m} \alpha_0^4 \alpha_2^{-4}) (\tau_N^{n,m})^2 + Q^{n+m} \alpha_0 \alpha_2^{-1} \tau_N^{n+1,m+1} \tau_N^{n-1,m-1} - Q^{n-2m+1} \alpha_1^{-1} \alpha_2 \tau_N^{n,m+1} \tau_N^{n,m-1} = 0, \quad (\text{A.12})$$

$$(Q^{4n+4m} \alpha_0^4 \alpha_2^{-4} - Q^{-8n+4m-4} \alpha_0^{-4} \alpha_1^4) (\tau_N^{n,m})^2 + Q^{-2n+m-1} \alpha_0^{-1} \alpha_1 \tau_N^{n+1,m} \tau_N^{n-1,m} - Q^{n+m} \alpha_0 \alpha_2^{-1} \tau_N^{n+1,m+1} \tau_N^{n-1,m-1} = 0, \quad (\text{A.13})$$

$$(Q^{-8n+4m-4} \alpha_0^{-4} \alpha_1^4 - Q^{4n-8m+4} \alpha_1^{-4} \alpha_2^4) (\tau_N^{n,m})^2 - Q^{-2n+m-1} \alpha_0^{-1} \alpha_1 \tau_N^{n+1,m} \tau_N^{n-1,m} + Q^{n-2m+1} \alpha_1^{-1} \alpha_2 \tau_N^{n,m+1} \tau_N^{n,m-1} = 0. \quad (\text{A.14})$$

**Proof.** We obtain (A.12) by eliminating  $\tau_{l,m,n+1} \tau_{l,m,n-1}$  from (A.1) and (A.2). Other equations can be derived in a similar manner.  $\square$

**Proposition A.4 (Type IV)** *The following bilinear equation holds:*

$$Q^{-3n} \alpha_0^{-3} (1 - Q^{-12m} \alpha_2^{-12}) \tau_N^{n+1,m} \tau_N^{n-1,m} - Q^{-3m} \alpha_2^3 (1 - Q^{-12l} \alpha_0^{-12}) \tau_N^{n,m+1} \tau_N^{n,m-1} + (Q^{-12m} \alpha_2^{-12} - Q^{-12l} \alpha_0^{-12}) \tau_N^{n+1,m+1} \tau_N^{n-1,m-1} = 0. \quad (\text{A.15})$$

**Proof.** Equation (A.15) can be derived by eliminating  $\tau_N^{n,m}$  from (A.12) and (A.13).  $\square$




Figure 6. Configuration of  $\tau$  functions for the bilinear equations of type III. Left: (A.12), center: (A.13), right: (A.14).




Figure 7. Configuration of  $\tau$  functions for the bilinear equations of type IV.

**Proposition A.5 (Type V)** *The following bilinear equations hold:*

$$\tau_{N+1}^{n,m} \tau_{N-1}^{n+1,m+1} - Q^{n+m-2N} \gamma^{-2} \alpha_0^2 \alpha_1^2 \tau_N^{n+1,m} \tau_N^{n,m+1} - Q^{-2n+2m-4N} \gamma^4 \alpha_0^{-4} \alpha_1^{-2} \tau_N^{n,m} \tau_N^{n+1,m+1} = 0, \quad (\text{A.16})$$

$$\tau_{N+1}^{n+1,m} \tau_{N-1}^{n,m} - Q^{-2n+m-2N} \gamma^{-2} \alpha_0^{-3} \alpha_1^{-1} \alpha_2^{-2} \tau_N^{n+1,m+1} \tau_N^{n,m-1} - Q^{4n-2m+4N} \gamma^4 \alpha_0^6 \alpha_1^2 \alpha_2^4 \tau_N^{n,m} \tau_N^{n+1,m} = 0, \quad (\text{A.17})$$

$$\tau_{N+1}^{n+1,m+1} \tau_{N-1}^{n+1,m} - Q^{n-2m-2N+1} \gamma^{-2} \alpha_1^{-1} \alpha_2 \tau_N^{n,m} \tau_N^{n+2,m+1} - Q^{-2n+4m+4N-2} \gamma^4 \alpha_1^2 \alpha_2^{-2} \tau_N^{n+1,m+1} \tau_N^{n+1,m} = 0, \quad (\text{A.18})$$

$$\tau_{N+1}^{n+1,m+1} \tau_{N-1}^{n,m} - Q^{n+m+2N} \gamma^2 \alpha_0^2 \alpha_1^2 \tau_N^{n,m+1} \tau_N^{n+1,m} - Q^{-2n-2m-4N} \gamma^{-4} \alpha_0^{-4} \alpha_1^{-2} \tau_N^{n+1,m+1} \tau_N^{n,m} = 0, \quad (\text{A.19})$$

$$\tau_{N+1}^{n,m} \tau_{N-1}^{n+1,m} - Q^{-2n+m+2N} \gamma^2 \alpha_0^{-3} \alpha_1^{-1} \alpha_2^{-2} \tau_N^{n,m-1} \tau_N^{n+1,m+1} - Q^{4n-2m-4N} \gamma^{-4} \alpha_0^6 \alpha_1^2 \alpha_2^4 \tau_N^{n,m} \tau_N^{n+1,m} = 0, \quad (\text{A.20})$$

$$\tau_{N+1}^{n+1,m} \tau_{N-1}^{n+1,m+1} - Q^{n-2m+2N} \gamma^2 \alpha_0 \alpha_2^2 \tau_N^{n+2,m+1} \tau_N^{n,m} - Q^{-2n+4m-4N} \gamma^{-4} \alpha_0^{-2} \alpha_2^{-4} \tau_N^{n+1,m} \tau_N^{n+1,m+1} = 0. \quad (\text{A.21})$$

**Proof.** First, we prove (A.16)–(A.18). We rewrite (A.4) as

$$T_4(\bar{\tau}_0) - c^{-\frac{2}{3}} a_0^{-\frac{1}{3}} a_1^{-1} a_2^{-\frac{2}{3}} \frac{\bar{\tau}_1}{\tau_1 \tau_2} \left( q^{\frac{1}{3}} c^{\frac{2}{3}} a_1 \tau_0 \bar{\tau}_2 + \bar{\tau}_0 \tau_2 \right) - c^{\frac{2}{3}} a_0^{\frac{1}{3}} a_1^{\frac{2}{3}} a_2 \frac{\bar{\tau}_0 \bar{\tau}_2}{\tau_2} = 0. \quad (\text{A.22})$$

By using (A.6), we have from (A.22) that

$$T_4(\bar{\tau}_0) \tau_2 - c^{-\frac{1}{3}} a_0^{-\frac{1}{6}} a_1^{-\frac{1}{3}} a_2^{-\frac{1}{2}} \bar{\tau}_1 T_2(\bar{\tau}_0) - c^{\frac{2}{3}} a_0^{\frac{1}{3}} a_1^{\frac{2}{3}} a_2 \bar{\tau}_0 \bar{\tau}_2 = 0, \quad (\text{A.23})$$

which is equivalent to

$$T_1^{-1} T_4^2(\tau_1) T_2(\tau_1) - c^{-\frac{1}{3}} a_0^{-\frac{1}{6}} a_1^{-\frac{1}{3}} a_2^{-\frac{1}{2}} T_4(\tau_1) T_1^{-1} T_2 T_4(\tau_1) - c^{\frac{2}{3}} a_0^{\frac{1}{3}} a_1^{\frac{2}{3}} a_2 T_1^{-1} T_4(\tau_1) T_2 T_4(\tau_1) = 0. \quad (\text{A.24})$$




Figure 8. Configuration of  $\tau$  functions for the bilinear equations of type V. Upper left: (A.16), upper center: (A.17), upper right: (A.18), lower left: (A.19), lower center: (A.20), lower right: (A.21).

We obtain (A.16), (A.17), and (A.18) by applying  $T_1^{l+1}T_2^mT_4^{n-1}$ ,  $T_1^{l+1}T_2^mT_4^{n-1}\pi$ , and  $T_1^{l+1}T_2^mT_4^{n-1}\pi^2$  on (A.24), respectively.

Next, we prove (A.19)–(A.21). We rewrite (A.4) as

$$T_4(\bar{\tau}_0) - a_1^{\frac{1}{3}}a_2^{-\frac{1}{3}}\frac{\bar{\tau}_2}{\tau_1\tau_2}\left(q^{\frac{1}{3}}c^{\frac{2}{3}}a_2\tau_1\bar{\tau}_0 + \bar{\tau}_1\tau_0\right) - c^{-\frac{2}{3}}a_0^{-\frac{1}{3}}a_1^{-1}a_2^{-\frac{2}{3}}\frac{\bar{\tau}_0\bar{\tau}_1}{\tau_1} = 0. \quad (\text{A.25})$$

By using (A.7), we have from (A.25) that

$$T_4^2(\tau_0)\tau_1 - c^{\frac{1}{3}}a_0^{\frac{1}{6}}a_1^{\frac{1}{2}}a_2^{\frac{1}{3}}T_3T_4(\tau_1)T_4(\tau_2) - c^{-\frac{2}{3}}a_0^{-\frac{1}{3}}a_1^{-1}a_2^{-\frac{2}{3}}T_4(\tau_0)T_4(\tau_1) = 0. \quad (\text{A.26})$$

We obtain (A.19), (A.20), and (A.21) by applying  $T_1^{l+1}T_2^mT_4^{n-1}\pi^2$ ,  $T_1^{l+1}T_2^mT_4^{n-1}$ , and  $T_1^{l+1}T_2^mT_4^{n-1}\pi$  on (A.26), respectively.  $\square$

**Proposition A.6 (Type VI)** *The following bilinear equations hold:*

$$\tau_{N+1}^{n,m}\tau_N^{n+1,m+1} - Q^{-3n+3m+2N-2}\gamma^2\alpha_1^3\tau_N^{n+1,m}\tau_{N+1}^{n,m+1} + Q^{-6n+6m+4N-4}\gamma^4\alpha_1^6\tau_N^{n,m}\tau_{N+1}^{n+1,m+1} = 0, \quad (\text{A.27})$$

$$\tau_{N+1}^{n+1,m}\tau_N^{n,m} - Q^{-3m+2N+1}\gamma^2\alpha_2^3\tau_{N+1}^{n+1,m+1}\tau_{N+1}^{n,m-1} + Q^{-6m+4N+2}\gamma^4\alpha_2^6\tau_N^{n+1,m}\tau_{N+1}^{n,m} = 0, \quad (\text{A.28})$$

$$\tau_{N+1}^{n+1,m+1}\tau_N^{n+1,m} - Q^{3n+2N+4}\gamma^2\alpha_0^3\tau_N^{n,m}\tau_{N+1}^{n+2,m+1} + Q^{6n+4N+8}\gamma^4\alpha_0^6\tau_N^{n+1,m+1}\tau_{N+1}^{n+1,m} = 0, \quad (\text{A.29})$$

$$\tau_{N+1}^{n+1,m+1}\tau_N^{n,m} - Q^{-3n+3m-2N-4}\gamma^{-2}\alpha_1^3\tau_{N+1}^{n+1,m}\tau_N^{n,m+1} + Q^{-6n+6m-4N-8}\gamma^{-4}\alpha_1^6\tau_N^{n+1,m+1}\tau_{N+1}^{n,m} = 0, \quad (\text{A.30})$$

$$\tau_{N+1}^{n,m}\tau_N^{n+1,m} - Q^{-3m-2N-1}\gamma^{-2}\alpha_2^3\tau_{N+1}^{n+1,m+1}\tau_N^{n,m-1} + Q^{-6m-4N-2}\gamma^{-4}\alpha_2^6\tau_N^{n,m}\tau_{N+1}^{n+1,m} = 0, \quad (\text{A.31})$$

$$\tau_{N+1}^{n+1,m}\tau_N^{n+1,m+1} - Q^{3n-2N+2}\gamma^{-2}\alpha_0^3\tau_{N+1}^{n,m}\tau_N^{n+2,m+1} + Q^{6n-4N+4}\gamma^{-4}\alpha_0^6\tau_N^{n+1,m}\tau_{N+1}^{n+1,m+1} = 0. \quad (\text{A.32})$$




Figure 9. Configuration of  $\tau$  functions for the bilinear equations of type VI. Upper left: (A.27), upper center: (A.28), upper right: (A.29) lower left: (A.30), lower center: (A.31), lower right: (A.32).

**Proof.** First, we prove (A.27)–(A.29). Equations (A.27), (A.28), and (A.29) can be derived by applying  $T_1^{l+1}T_2^mT_4^n$ ,  $T_1^{l+1}T_2^mT_4^n\pi$ , and  $T_1^{l+1}T_2^mT_4^n\pi^2$  on (A.7), respectively.

Next, we prove (A.30)–(A.32). By applying  $T_2$  on  $\tau_0$ , we obtain

$$q^{-\frac{1}{6}}c^{-\frac{1}{3}}a_1^{\frac{1}{2}}\bar{\tau}_1T_2(\tau_0) - q^{-\frac{1}{3}}c^{-\frac{2}{3}}a_1\tau_2\bar{\tau}_0 - \bar{\tau}_2\tau_0 = 0. \quad (\text{A.33})$$

Equations (A.30), (A.31), and (A.32) can be derived by applying  $T_1^{l+1}T_2^mT_4^n$ ,  $T_1^{l+1}T_2^mT_4^n\pi$ , and  $T_1^{l+1}T_2^mT_4^n\pi^2$  on (A.33), respectively.  $\square$

**Remark A.7** The bilinear equations in Proposition 3.4 correspond to (A.27), (A.29), (A.30), (A.32), and (A.3).

## A.2 Bilinear equations for $q$ -P<sub>II</sub>

The bilinear equations for  $q$ -P<sub>II</sub> are derived from the equations in Section A.1. Since the parameter space and  $\tau$  functions are restricted, we only have to pick up the bilinear equations that consist of the  $\tau$  functions on the “unit-strip,” and to rewrite them in terms of  $R_1$  instead of  $T_1$  (see Figure 3). Therefore, only the bilinear equations of type V and VI are relevant. We use the notation in (3.54).

**Proposition A.8** *The following bilinear equations hold:*

$$\tau_{N+1}^{k+1}\tau_{N-1}^{k+2} - Q^{\frac{k-4N+2}{2}}\gamma^{-2}\alpha_0\tau_N^{k+3}\tau_N^k - Q^{-k+4N-2}\gamma^4\alpha_0^{-2}\tau_N^{k+1}\tau_N^{k+2} = 0, \quad (\text{A.34})$$

$$\tau_{N+1}^{k+2}\tau_{N-1}^{k+1} - Q^{\frac{k+4N+2}{2}}\gamma^2\alpha_0\tau_N^{k+3}\tau_N^k - Q^{-k-4N-2}\gamma^{-4}\alpha_0^{-2}\tau_N^{k+2}\tau_N^{k+1} = 0, \quad (\text{A.35})$$

$$Q^{-\frac{3k-4N+4}{2}}\gamma^2\alpha_0^{-3}\tau_N^{k+3}\tau_{N+1}^k - Q^{-3k+4N-4}\gamma^4\alpha_0^{-6}\tau_{N+1}^{k+1}\tau_{N+1}^{k+2} - \tau_{N+1}^{k+1}\tau_N^{k+2} = 0, \quad (\text{A.36})$$

$$Q^{-\frac{3k+4N+8}{2}}\gamma^{-2}\alpha_0^{-3}\tau_{N+1}^{k+3}\tau_N^k - Q^{-3k-4N-8}\gamma^{-4}\alpha_0^{-6}\tau_N^{k+2}\tau_{N+1}^{k+1} - \tau_{N+1}^{k+2}\tau_N^{k+1} = 0. \quad (\text{A.37})$$

**Proof.** Noticing (3.55), we obtain from (A.23)

$$R_1^{-2}T_4^2(\tau_1)R_1^{-1}(\tau_1) - q^{-\frac{5}{12}}c^{-\frac{1}{3}}a_0^{\frac{1}{6}}T_4(\tau_1)R_1^{-3}T_4(\tau_1) - q^{\frac{5}{6}}c^{\frac{2}{3}}a_0^{-\frac{1}{3}}R_1^{-2}T_4(\tau_1)R_1^{-1}T_4(\tau_1) = 0, \quad (\text{A.38})$$

from which (A.34) is derived by applying  $R_1^{m+3}T_4^{n-1}$ . Similarly, we have

$$T_4^2(\tau_1)R_1^{-1}(\tau_1) - q^{\frac{1}{3}}c^{\frac{1}{3}}a_0^{\frac{1}{6}}R_1T_4(\tau_1)R_1^{-2}T_4(\tau_1) - q^{-\frac{2}{3}}c^{-\frac{2}{3}}a_0^{-\frac{1}{3}}T_4(\tau_1)R_1^{-1}T_4(\tau_1) = 0. \quad (\text{A.39})$$

by applying  $\pi$  on (A.26). Then we obtain (A.35) by applying  $R_1^{m+2}T_4^{n-1}$  on (A.39). Equation (A.36) is derived by applying  $R_1^{m+3}T_4^n$  on

$$q^{\frac{1}{6}}c^{\frac{1}{3}}a_0^{-\frac{1}{2}}\tau_1R_1^{-3}T_4(\tau_1) - q^{\frac{1}{3}}c^{\frac{2}{3}}a_0^{-1}R_1^{-2}(\tau_1)R_1^{-1}T_4(\tau_1) - R_1^{-2}T_4(\tau_1)R_1^{-1}(\tau_1) = 0, \quad (\text{A.40})$$

which follows from (A.6). Finally, we obtain (A.37) by applying  $R_1^{m+3}T_4^n$  on

$$q^{-\frac{1}{6}}c^{-\frac{1}{3}}a_0^{-\frac{1}{2}}T_4(\tau_1)R_1^{-3}(\tau_1) - q^{-\frac{1}{3}}c^{-\frac{2}{3}}a_0^{-1}R_1^{-1}(\tau_1)R_1^{-2}T_4(\tau_1) - R_1^{-1}T_4(\tau_1)R_1^{-2}(\tau_1) = 0, \quad (\text{A.41})$$

which is follows from (A.33).  $\square$




Figure 10. Configuration of  $\tau$  functions for the bilinear equations in Proposition A.8. The figures correspond to (A.34), (A.35), (A.36), and (A.37), respectively, from the left to the right.

**Remark A.9** The bilinear equations in Proposition 3.6 correspond to (A.36), (A.37), and (A.34).

## References

- [1] E. Brézin and V.A. Kazakov, Exactly solvable theories of closed strings, Phys. Lett. B **236** (1990) 144–150.
- [2] M.R. Douglas and S.H. Shenker, Strings in less than one dimension, Nucl. Phys. B **335** (1990) 635–654.
- [3] A.S. Fokas, A.R. Its and A.V. Kitaev, The isomonodromy approach to matrix models in 2D quantum gravity, Comm. Math. Phys. **147** (1992) 395–430.
- [4] G. Gasper and M. Rahman, Basic Hypergeometric Series, Encyclopedia of Mathematics and Its Applications **35** (Cambridge University Press, Cambridge, 1990).
- [5] B. Grammaticos, F.W. Nijhoff, V. Papageorgiou, A. Ramani and J. Satsuma, Linearization and solutions of the discrete Painlevé III equation, Phys. Lett. A **185** (1994) 446–452.

- [6] B. Grammaticos and A. Ramani, Discrete Painlevé equations: a review, *Lect. Notes Phys.* **644** (2004) 245–321.
- [7] B. Grammaticos, A. Ramani and V. Papageorgiou, Do integrable mappings have the Painlevé property?, *Phys. Rev. Lett.* **67** (1991) 1825–1828.
- [8] T. Hamamoto, K. Kajiwara and N.S. Witte, Hypergeometric solutions to the  $q$ -Painlevé equation of type  $(A_1 + A'_1)^{(1)}$ , *Int. Math. Res. Not.* **2006** (2006) Article ID 84619.
- [9] K. Kajiwara, The discrete Painlevé II equation and the classical special functions, in *Symmetries and integrability of difference equations*, eds. by P. Clarkson and F.W. Nijhoff, London Math. Soc. Lecture Note Ser. 255 (Cambridge University Press, Cambridge, 1999) 217–227.
- [10] K. Kajiwara, On a  $q$ -difference Painlevé III equation. II. Rational solutions, *J. Nonlin. Math. Phys.* **10** (2003) 282–303.
- [11] K. Kajiwara and K. Kimura, On a  $q$ -difference Painlevé III equation. I. Derivation, symmetry and Riccati type solutions, *J. Nonlin. Math. Phys.* **10** (2003) 86–102.
- [12] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada,  ${}_{10}E_9$  solution to the elliptic Painlevé equation, *J. Phys. A: Math. Gen.* **36** (2003) L263–L272.
- [13] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, Hypergeometric solutions to the  $q$ -Painlevé equations, *Int. Math. Res. Not.* **2004** (2004) 2497–2521.
- [14] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, Construction of hypergeometric solutions to the  $q$ -Painlevé equations, *Int. Math. Res. Not.* **2005** (2005) 1441–1463.
- [15] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, Point configurations, Cremona transformations and the elliptic difference Painlevé equation, *Sémin. Congr.* **14** (2006) 169–198.
- [16] K. Kajiwara and N. Nakazono, In preparation.
- [17] K. Kajiwara, M. Noumi and Y. Yamada, A study on the fourth  $q$ -Painlevé equation, *J. Phys. A: Math. Gen.* **34** (2001) 8563–8581.
- [18] K. Kajiwara, Y. Ohta and J. Satsuma, Casorati determinant solutions for the discrete Painlevé III equation, *J. Math. Phys.* **36** (1995) 4162–4174.
- [19] K. Kajiwara, Y. Ohta, J. Satsuma, B. Grammaticos and A. Ramani, Casorati determinant solutions for the discrete Painlevé-II equation, *J. Phys. A: Math. Gen.* **27** (1994) 915–922.
- [20] K. Kajiwara, K. Yamamoto and Y. Ohta, Rational solutions for the discrete Painlevé II equation, *Phys. Lett. A* **232** (1997) 189–199.
- [21] M.D. Kruskal, K.M. Tamizhmani, B. Grammaticos and A. Ramani, Asymmetric discrete Painlevé equations, *Regul. Chaotic Dyn.* **5** (2000) 273–280.
- [22] T. Masuda, Classical transcendental solutions of the Painlevé equations and their degeneration, *Tohoku Math. J.* **56** (2004) 467–490.



- [23] T. Masuda, Y. Ohta and K. Kajiwara, Rational solutions to the Painlevé V equation and the universal characters, *RIMS Kokyuroku* **1203** (2001) 97–108 (in Japanese).
- [24] T. Masuda, Y. Ohta and K. Kajiwara, A determinant formula for a class of rational solutions of Painlevé V equation, *Nagoya J. Math.* **168** (2002) 1–25.
- [25] S. Nakao, K. Kajiwara and D. Takahashi, Multiplicative  $dP_{II}$  and its ultradiscretization, *Reports of RIAM Symposium No. 9ME-S2*, Kyushu University (1998) 125–130 (in Japanese).
- [26] N. Nakazono, In preparation.
- [27] M. Noumi, *Painlevé equations through symmetry* (American Mathematical Society, Providence, 2004).
- [28] Y. Ohta, Self-dual structure of the discrete Painlevé equations, *RIMS Kokyuroku* **1098** (1999) 130–137 (in Japanese).
- [29] K. Okamoto, Studies on the Painlevé equations. III. Second and Fourth Painlevé equation,  $P_{II}$  and  $P_{IV}$ , *Math. Ann.* **275** (1986) 221–255.
- [30] K. Okamoto, Studies on the Painlevé equations. I. Sixth Painlevé equation  $P_{VI}$ , *Ann. Mat. Pura Appl.* **146** (1987) 337–381.
- [31] K. Okamoto, Studies on the Painlevé equations. II. Fifth Painlevé equation  $P_V$ , *Japan. J. Math.* **13** (1987) 47–76.
- [32] K. Okamoto, Studies on the Painlevé equations. IV. Third Painlevé equation  $P_{III}$ , *Funct. Ekvac.* **30** (1987) 305–332.
- [33] V. Periwal and D. Shevitz, Unitary-matrix models as exactly solvable string theories, *Phys. Rev. Lett.* **64** (1990) 1326–1329.
- [34] G.R.W. Quispel, J.A.G Roberts and C.J. Thompson, Integrable mappings and soliton equations, *Phys. Lett. A* **126** (1988) 419–421.
- [35] G.R.W. Quispel, J.A.G Roberts and C.J. Thompson, Integrable mappings and soliton equations II, *Physica D* **34** (1989) 183–192.
- [36] A. Ramani and B. Grammaticos, Discrete Painlevé equations: coalescences, limits and degeneracies, *Physica A* **228** (1996) 150–159.
- [37] A. Ramani, B. Grammaticos and J. Hietarinta, Discrete versions of the Painlevé equations, *Phys. Rev. Lett.* **67** (1991) 1829–1832.
- [38] A. Ramani, Y. Ohta, J. Satsuma and B. Grammaticos, Self-duality and Schlesinger chains for the asymmetric  $dP_{II}$  and  $qP_{III}$  equations, *Comm. Math. Phys.* **192** (1998) 67–76,
- [39] H. Sakai, Rational surfaces associated with affine root systems and geometry of the Painlevé equations, *Comm. Math. Phys.* **220** (2001) 165–229.
- [40] T. Tsuda, Tau functions of  $q$ -Painlevé III and IV equations, *Lett. Math. Phys.* **75** (2006) 39–47.

# List of MI Preprint Series, Kyushu University

The Global COE Program  
Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI  
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA  
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO  
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU  
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI  
Torsion points of abelian varieties with values in infinite extensions over a  $p$ -adic field
- MI2008-6 Yoshiyuki TOMIYAMA  
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI  
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA  
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA  
Alpha-determinant cyclic modules and Jacobi polynomials

- MI2008-10 Sangyeol LEE & Hiroki MASUDA  
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA  
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO  
On the  $L^2$  a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA  
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials
- MI2008-14 Takashi NAKAMURA  
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA  
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO  
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI  
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI  
Variable selection for functional regression model via the  $L_1$  regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI  
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI  
Flat modules and Groebner bases over truncated discrete valuation rings

- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA  
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI  
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI  
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA  
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO  
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA  
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA  
Hypergeometric  $q$ -functions of the  $q$ -Painlevé system of type  $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA  
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA  
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications
- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI  
Large time behavior of the semigroup on  $L^p$  spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain

- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE  
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO  
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO  
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI  
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI  
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO  
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA  
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO  
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI  
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO  
On very accurate enclosure of the optimal constant in the a priori error estimates for  $H_0^2$ -projection

- MI2009-26 Manabu YOSHIDA  
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI  
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA  
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA  
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance
- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA  
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI  
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA  
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA  
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO and Teruhisa TSUDA  
Projective reduction of the discrete Painlevé system of type  $(A_2 + A_1)^{(1)}$