

Mineral Uptake and Soluble Carbohydrates of Lettuce : Effects of Air Temperatures and Mineral Supply Levels

Sung, Jwakyung

Soil and Fertilizer Division, NAAS, RDA

Cho, Minji

Soil and Fertilizer Division, NAAS, RDA

Sonn, Yeonkyu

Soil and Fertilizer Division, NAAS, RDA

Yun, Hejin

Soil and Fertilizer Division, NAAS, RDA

他

<https://doi.org/10.5109/1564080>

出版情報 : 九州大学大学院農学研究院紀要. 61 (1), pp.41-46, 2016-02-29. Faculty of Agriculture, Kyushu University

バージョン :

権利関係 :

Mineral Uptake and Soluble Carbohydrates of Lettuce : Effects of Air Temperatures and Mineral Supply Levels

Jwakyung SUNG¹, Minji CHO^{1,†}, Yeonkyu SONN^{1,†}, Hejin YUN¹, Seongsoo KANG¹,
Yoshiyuki SHINOBI and Taek-Keun OH^{2,*}

Science for Bioproduction Environment, Faculty of Agriculture, Kyushu University, Hakozaki 6–10–1,
Higashi-ku, Fukuoka city 812–8581, Japan

(Received November 11, 2015 and accepted November 19, 2015)

Both low and high temperatures affect plant growth and development at whole plant level, tissue and even cell level through a variety of metabolic changes. In this paper we investigated the extent of the inhibition of growth, macro-element uptake and soluble carbohydrate production, and the effect of extra-supply of minerals as a means of the recovery from the damage. Lettuce plants were grown four different growth temperatures (15/8, 22/15, 25/18 and 28/21°C), and extra-supply of minerals was composed of 1.5- and 2.0-fold stronger than the standard nutrition (1/2 strength of Hoagland's solution). Low and high temperatures significantly adversely affected lettuce growth, mineral uptake and soluble carbohydrate production, and the extra-supply of minerals didn't play any crucial roles to promote and recover them. Merely, P uptake was slightly increased by an extra-supply of nutrients in all temperature conditions. From this study, it was strongly suggested that further research should include the effect of extra-supply of minerals when plants are recovered from low or high temperature stress, and metabolic interaction such as mineral uptake and carbohydrates between shoot and root.

Key words: Lettuce, Mineral uptake, Carbohydrates, Temperature stress, Mineral supply

INTRODUCTION

The production of vegetable crops has shown significant yield increases in South Korea since over the last several decades. Transitory or constantly high temperatures cause a variety of morpho-anatomical, physiological and biochemical changes in plants, which affect plant growth and development and may result in a significant reduction in economic yield. At moderately high temperatures, injuries or death to crop plants may occur only after long-term exposure, and it may be derived from a limited photosynthesis, assimilate partitioning, and water and nutrient use efficiency (Kuiper, 1964; Walker, 1969; Ruter and Ingram, 1990). A well-known consequence of elevated temperature in plants is the damage caused by heat-induced imbalance in photosynthesis and respiration; in general, the rate of photosynthesis decreases whereas dark- and photo-respiration rates increase considerably under high temperatures (Nakamoto and Hiyama, 1999). Assimilate partitioning, taking place via apoplastic and symplastic pathways under high temperatures, has significant effects on transport and transfer processes in plants although it strongly depends genotypes (Yang *et al.*, 2002; Taiz and Zeiger, 2006). Mineral uptake by plants from the rhizosphere is directly influenced by soil temperature, and the lower (chilling) and higher (heat) temperatures compared to the optimum

lead to a considerable reduction in mineral uptake (Cumbus and Nye, 1984; Raju *et al.*, 1990; Tindall *et al.*, 1990).

Lettuce (*Lactuca sativa* L.) is the second largest leafy crops after Chinese cabbage, and the scale of greenhouses-based cultivation throughout South Korea was approximately 2,968 ha (MAFRA statistics, 2013). Furthermore, a year round cultivation of lettuce often causes the unexpected heat (high in summer season) and chilly (low in winter season) stresses, and thus results in deleterious effects on the growth and yield. Although the literature on plant responses to temperatures such as mineral uptake and carbohydrate production is abundant, the information about the effects of extra-supply of minerals there is only a very few. The objective of this study was to know the extent of a damage of growth, mineral uptake and carbohydrate production under low or high temperature conditions, and to know whether there is any effect of extra-supply of minerals to recover and promote those in lettuce plant.

MATERIALS AND METHODS

Plant materials and growth conditions

This study was performed in an environment-controlled growth chamber, NAAS, RDA, South Korea in 2014. The uniformly growing seedlings of lettuce were transplanted into 1 L plastic box filled with pure sand soil, and then fed with 1/2 strength of Hoagland's solution which is composed as follows; 5 mM Ca(NO₃)₂, 5 mM KNO₃, 2 mM MgSO₄, 0.5 mM KH₂PO₄, 1.5 mM Fe-EDTA, 1 mM NH₄NO₃, 2 μM H₃BO₃, 0.2 μM MnCl₂, 0.01 μM ZnSO₄, 0.01 μM CuSO₄, and 0.03 μM H₂MoO₄. The 1/2 strength of Hoagland's solution, 100 mL per box, was

¹ Soil and Fertilizer Division, NAAS, RDA, Wanju, Jeollabuk-do, 55365, Korea

² Department of Bio-Environmental Chemistry, Chungnam National University, Daejeon 305–764, Korea

[†] The authors contributed equally to this paper as a co-first author.

* Corresponding author (E-mail: ok5382@cnu.ac.kr) (T.K. OH)

supplied every day during the experiment. Plants were grown in four different temperature conditions, low (15/8°C, day/night), optimal (22/15°C), and high temperature (25/18 and 28/21°C). To investigate the effect of an extra-supply of minerals to mitigate the damage from temperature stress, lettuce plants were grown in three different nutrient conditions, standard (1/2 strength of Hoagland's solution), 1.5- and 2.0-fold stronger nutrient solutions. Lettuce plants were assigned with the completely randomized two factor factorial design (temperature and mineral supply) in an environment-controlled chamber and were taken to determine the contents of carbohydrates and mineral elements at 5, 10 and 15 days after treatment (DAT).

Measurement of nutrients

The oven-dried samples (0.3 g) which were at 80°C for 48 h were soaked in 5 mL of 368 mmol L⁻¹ salicylic acid in 84.7% sulfuric acid (H₂SO₄) for 24 h then digested in a digestion system, heated to 300°C for 3 h, followed by several drops of hydrogen peroxide (H₂O₂). The extracted solution was transferred to 100 mL volumetric flasks and then diluted to 100 mL with deionized water for mineral assays. The N concentration was colorimetrically determined using the automatic flow injection analyzer (BRAN LUBBE, Germany). The P concentration was measured using the molybdate-blue colorimetric method (UV-2450, Shimadzu, Japan) and cation concentrations were determined with ICP-OES (INTEGRA XMP, GBC, Australia).

Measurement of soluble carbohydrates

Soluble sugar from dried shoots and roots was determined by the reaction of 1.0 mL of the alcoholic extract with 2.0 mL fresh 0.2% anthrone in sulfuric acid (w/v); the absorbance was read at 630 nm. After the extraction of the soluble fractions, the solid fraction was used for starch analysis. Starch was firstly extracted with 9.3 N (normal concentration) of perchloric acid and followed by 4.6 N. The extracts were combined and starch concentration was determined after reaction with the anthrone reagent. Glucose was used as the standard for soluble sugar.

Statistical analysis

This experiment was performed with the completely randomized two factor factorial design (temperature and nutrient level) with three repeats. The analysis of variance (ANOVA) was conducted to find effects of treatments. Least significant difference (LSD) was performed to determine the significance of the difference between the means of treatments. An α error value of 0.05 was chosen to indicate statistical significance. All statistical analysis was performed using version 9.01 of SAS (SAS Institute Inc, Cary, NC).

RESULTS

Effects of temperature and extra-supplied nutrient on lettuce growth

The shoot growth of lettuce (dry weight-based) from each temperature condition was shown in Fig. 1, and the growth rates between temperature conditions represented significant differences ($p < 0.01$ and 0.001) at three time points. The growth was the highest at the optimal (22/15°C) and moderately high (25/18°C) temperature conditions during the whole experiment period whereas the low (15/8°C) and high (28/21°C) temperature conditions represented the marked reduction in the growth from Day 10. The effect of extra-supply of minerals (1.5 and 2.0-fold stronger) as a means of recovering the damage from temperature stresses was not observed (Fig. 2), and especially the extra-supply made the growth worse at high temperature condition.




Fig. 1. Effect of temperature stresses on tomato plants grown under 1/2 strength Hoagland's nutrient solution ($n=3$). Star marks, *, ** and ***, mean significant difference at $p < 0.05$, 0.01 and 0.001 , respectively.




Fig. 2. Effect of an extra-supply of nutrient solution on lettuce growth grown under four different temperature regimes at 15 DAT ($n=3$). The concentration, $\times 1.0$, means the 1/2 strength Hoagland's nutrient solution. The letters above vertical bars mean significant differences from LSD test.

Effects of temperature and extra-supplied minerals on concentration and uptake of macro-elements

The concentrations of macro-nutrients in temperature-affected lettuce resulted in a tendency of the overall increase except N (67% compared to the optimal) and K (55%) concentration, which represented a marked reduction, at low temperature (Table 1). An increase in the concentration was obvious Ca (1.9 fold as high compared to the optimal) at low temperature and P (1.9 fold as high) at high temperature. By contrast, the uptake of macro-nutrients was greatly affected by temperature stresses (Table 2), and the uptake rates were ranged from 26 to 71% at the low temperature to optimal and from 48 to 66% in the high temperature. The uptake rate under moderately high temperature (25/18°C) resulted in slight decrease of N, P and K, and in great increase of Ca and Mg. The extra supply of minerals was non-effective to promote the nutrient uptake as lettuce was exposed extremely adverse temperature conditions although an effect was partially observed at 2.0-fold stronger mineral supply (Table 3). The supply of 1.5-fold stronger minerals led to the significant reduction in most elements which ranged from 18 to 56% at low temperature (15/8°C) to optimal temperature (22/15°C) and from 30 to 71% at high temperature (28/21°C). The supply of 2.0-fold stronger minerals promoted widely the nutrient uptake

at low temperature (15/8°C) compared to control ($\times 1.0$ of low temperature), whereas there was no effect at high temperature. Unlike other macro-elements, P uptake was slightly increased by an extra-supply of nutrients at all temperature conditions.

Effects of temperature and extra-supplied nutrients on soluble carbohydrates production

The contents (glucose equi.) of soluble sugars and starch greatly differed with both temperature conditions and the period of temperature treatment (Fig. 3). The levels of soluble sugars and starch were the highest in optimal temperature at 5 and 10 DAT and in moderately high temperature at 15 DAT, and the relative rates of soluble sugars and starch in low and high temperatures represented 39 ($21.1 \pm 1.5 \text{ mg g}^{-1} \text{ DW}$), 37 ($13.0 \pm 1.1 \text{ mg g}^{-1} \text{ DW}$), 26 ($13.9 \pm 2.1 \text{ mg g}^{-1} \text{ DW}$) and 43 ($15.4 \pm 0.7 \text{ mg g}^{-1} \text{ DW}$) %, respectively, at 15 DAT. The extra supply of nutrients to enhance the accumulation of soluble sugars and starch didn't show any effect in all temperature conditions (Fig. 4). The contents of soluble sugars and starch were the highest in moderately high temperature and followed by optimal, low and high temperatures. Moreover, the level of soluble sugars was significantly reduced by low temperature, and starch was markedly decreased in both low and high temperatures.

Table 1. The range of macro-element concentrations of temperature-affected lettuce

Temperature (day/night, °C)	N	P ₂ O ₅	K ₂ O	CaO	MgO
	----- %, DW -----				
15/8	1.64 ~ 3.91 (67)	0.38 ~ 0.49 (123)	2.57 ~ 5.02 (55)	1.21 ~ 3.04 (190)	0.32 ~ 0.65 (112)
22/15	3.32 ~ 4.00 (100)	0.34 ~ 0.39 (100)	6.08 ~ 6.66 (100)	1.19 ~ 1.43 (100)	0.39 ~ 0.51 (100)
25/18	3.34 ~ 4.64 (106)	0.30 ~ 0.50 (110)	5.74 ~ 7.48 (104)	1.23 ~ 2.34 (138)	0.42 ~ 0.80 (143)
28/21	4.92 ~ 5.10 (137)	0.58 ~ 0.78 (193)	6.87 ~ 7.67 (115)	1.11 ~ 1.83 (114)	0.43 ~ 0.62 (117)

[†] Data indicate the range of the concentrations measured at three sampling points, 5, 10 and 15 DAT, and the data within the parenthesis represent a concentration index of macro-elements when the concentrations are defined as 100 in optimal temperature condition (22/15°C)

Table 2. The uptake of macro-elements of temperature-affected lettuce at 15 DAT



Temperature (day/night, °C)	N	P ₂ O ₅	K ₂ O	CaO	MgO
	----- mg plant ⁻¹ , DW -----				
15/8	26.5 ± 9.0b	3.5 ± 0.5c	35.6 ± 7.6c	22.2 ± 6.3bc	5.5 ± 1.2c
22/15	87.2 ± 5.3a	7.7 ± 0.5a	137.6 ± 7.2a	31.1 ± 2.0b	11.0 ± 0.8b
25/18	73.3 ± 14.3a	6.5 ± 1.2a	126.3 ± 25.6a	50.7 ± 9.8a	17.1 ± 3.1a
28/21	42.5 ± 5.0b	5.1 ± 0.6b	65.3 ± 8.0b	15.1 ± 1.5c	5.3 ± 0.6c
F-value (p<0.0001)	27.53	17.59	34.53	20.03	30.48

[†] Data indicate the uptake of macro-elements of tomato shoots at three different temperature conditions at 15 days after treatment, and tomato plants were grown under the 1/2 strength Hoagland's nutrient solution. The letters mean significant differences from LSD test (n=3)

Table 3. The effect of the extra-supplied nutrients on the uptake rates of macro-elements

Temperature (day/night, °C)	Nutrient supply	N	P ₂ O ₅	K ₂ O	CaO	MgO
15/8	× 1.0	30	46	26	71	50
	× 1.5	19	50	18	56	34
	× 2.0	31	71	28	108	58
22/15	× 1.0	100	100	100	100	100
	× 1.5	98	103	113	109	101
	× 2.0	98	100	119	144	139
25/18	× 1.0	84	85	92	163	156
	× 1.5	94	96	108	139	134
	× 2.0	78	96	105	160	171
28/21	× 1.0	49	66	47	49	48
	× 1.5	34	71	32	30	33
	× 2.0	39	78	38	39	41
F-value						
Temperature		82.42***	26.98***	136.02***	47.13***	90.00***
Nutrient		0.61	3.24	0.95	5.53*	7.29**
Temperature × Nutrient		1.08	0.84	1.68	1.44	1.11

[†] The data indicate a percentage of macro-element uptake when the uptake is defined as 100 in optimal supply (1/2 strength Hoagland nutrient solution) at optimal temperature condition (22/15°C). The concentration, × 1.0, means the 1/2 strength Hoagland's nutrient solution

**Fig. 3.** Effect of temperature stresses on the accumulation of soluble sugars and starch in the leaves of lettuce plants grown under 1/2 strength Hoagland's nutrient solution at 15 DAT (n=3). The letters above vertical bars mean significant differences from LSD test.**Fig. 4.** Effect of an extra-supply of nutrient solution on the accumulation of soluble sugars and starch in the leaves of lettuce plants grown under three different temperature conditions at 15 DAT (n=3). The concentration, × 1.0, means the 1/2 strength Hoagland's nutrient solution. The letters above vertical bars mean significant differences from LSD test.

DISCUSSION

Temperature stresses (low and high temperature) are the major environmental factors affecting plant growth and development, and also induce morphological, physiological and biochemical changes in plants. It is well known that crop growth rates are significantly reduced by temperature stresses as results of a decrease in metabolic processes and photosynthesis (Sharkova, 2001; Wise *et al.*, 2004; Waraich *et al.*, 2012), shoot and root growth inhibition, and leaf senescence (Vollenweider and Gunthardt-Goerg, 2005), and our result also represented only one-third in low or high temperature conditions compared to the optimal. An extra-supply of minerals to promote lettuce growth didn't have any effect in low or high temperature stress (Fig. 2), the growth was not differed in the optimal or moderately high temperature conditions, and even high temperature led to a marked reduction in the growth under 1.5- and 2.0-fold stronger mineral conditions. Waraich *et al.* (2011) reported that N fertilization mitigated the adverse effects of temperature stresses which induced the reduction in photosynthesis and growth by photo-oxidative damage (Waraich *et al.*, 2011). Increasing K supply also alleviated the damage of leaf and stem damage and decrease in crop yield (Grewal and Singh, 1980; Hakerlerler *et al.*, 1997), however the effect mentioned above was not observed in our present work. The concentration and uptake of macro-elements was strongly different with temperature conditions. Certainly, high temperature led to substantial accumulation of macro-elements, while low temperature represented lower N and K concentrations or higher P, Ca and Mg concentrations. This suggests the existence of a controlling mechanism in plant, particularly root, which is mediated by physiological responses such as transpiration, mineral absorption, and leaf and root growth. Marked reduction in mineral concentration has been considered as a factor responsible for temperature stresses (Ali *et al.*, 1998; Tindall *et al.*, 1990; Engels and Marschner, 1996; Du and Tachibana, 1994). However, mineral concentrations except N and K at low temperature in lettuce shoots increased at both low and high temperatures, and it might be interpreted to be the mineral accumulation as the result of restricted utilization by growth cessation. Undoubtedly, a remarkable growth inhibition which was caused by low and high temperature stresses adversely affected macro-element uptake (Table 2), this means that mineral uptake greatly depends on ambient temperature conditions, similar to the observation by Ali *et al.* (1998) and Tindall *et al.* (1990). The extra-supply of minerals as a means of promoting mineral uptake wasn't noticeable in all temperature conditions although two-fold stronger minerals affected a slight increase in macro-element uptake at low temperature stress (Table 3). Interestingly, an extra-supply of P led to the significant increase in P uptake as lettuce plant was suffered from temperature stresses. Low and high temperatures markedly reduced soluble sugars and starch levels in lettuce shoots (Fig. 3). More carbon losses due to increased respiration and the shortage of non-structural

carbohydrate have long been considered as factors responsible for the growth inhibition under various temperature stresses (Youngner and Nudge, 1968; Canmore-Neumann and Kafkafi, 1983), consistent with our observation. Starch metabolism (diurnal fluctuation) is very sensitive to changes in the environment, and soluble sugars that accumulate in response to stress can function as osmolytes to maintain cell turgor against adverse environments, particularly temperature stresses (Madden *et al.*, 1985; Todaka *et al.*, 2000; Kaplan and Guy, 2004; Basu *et al.*, 2007; Kempa *et al.*, 2008). In this study, a decline of soluble sugars and starch levels is possibly considered as a result of restricted water and nutrient absorption and limited photosynthesis although further study is needed for the clarification of carbohydrate metabolism. The extra-supply of minerals didn't also have any effect to enhance soluble sugars and starch levels, namely photosynthesis. In conclusion, temperature stresses caused significant reduction in crop growth, mineral uptake and carbohydrate production while macro-element concentrations in lettuce shoots seemed likely to be a temperature-dependent response. Moreover, extra-supply of minerals didn't play crucial roles to promote an adverse damage caused by temperature stresses. Further research should include the effect of extra-supply of minerals when plants are recovered from low or high temperature stress, and metabolic interaction such as mineral uptake and carbohydrates between shoot and root.

ACKNOWLEDGMENT

This work was carried out with the support of "Cooperative Research Program for Agriculture Science & Technology Development (Project No. PJ010899)" Rural Development Administration, Republic of Korea

REFERENCES

- Ali I. E. A., U. Kafkafi, I. Yamaguchi, Y. Sugimoto and S. Inanaga 1998 Response of oilseed rape plant to low root temperature and nitrate: ammonium ratios. *J. Plant Nutr.*, **21**: 1463-1481
- Basu P. S., M. Ali and S. K. Chaturvedi 2007 Osmotic adjustment increases water uptake, remobilization of assimilates and maintains photosynthesis in chickpea under drought. *Indian J. Exp. Bot.*, **45**: 261-267
- Canmore-Neumann F. and U. Kafkafi 1983 The effect of root temperature and $\text{NO}_3^-/\text{NH}_4^+$ ratio on strawberry plants. I. Growth, flowering, and root development. *Agric. J.*, **75**: 941-947
- Cumby I. P. and P. H. Nye 1984 Root zone temperature effects on growth and phosphate absorption in rape *Brassica napus* cv. Emerald. *J. Exp. Bot.*, **36**: 219-227
- Du Y. C. and S. Tachibana 1994 Effect of supraoptimal root temperature on the growth, root respiration and sugar content of cucumber plants. *Scientia Horticulturae*, **58**: 289-301
- Engels C. and H. Marschner 1996 Effects of suboptimal root zone temperature and shoot demand on net translocation of micro-nutrients from the roots to the shoot of maize. *Plant and Soil*, **186**: 311-320
- Grewal J. S. and S. N. Singh 1980 Effect of potassium nutrition on frost damage and yield of potato plants on alluvial soils of the Punjab (India). *Plant Soil*, **57**: 105-110
- Hakerlerler H., M. Oktay, N. Eryüce and B. Yagmur 1997 Effect of potassium sources on the chilling tolerance of some vege-

- table seedlings grown in hotbeds, in Johnston, A. E.: Food Security in the WANA Region, the Essential Need for Balanced Fertilization. International Potash Institute, Basel, pp. 317–327
- Kaplan F., J. Kopka, D. W. Haskell, W. Zhao, K. C. Schiller, N. Gatzke, D. Y. Sung and C. L. Guy 2004 Exploring the temperature-stress metabolome of Arabidopsis. *Plant Physiol.*, **136**: 4159–4168
- Kempa S., J. Krasensky, S. Dal Santo, J. Kopka and C. Jonak 2008 A central role of synthesizing fructans and enhancing chilling tolerance. *J. Exp. Bot.*, **59**: 793–802
- Kuiper P. J. C 1964 Water uptake of higher plants as affected by root temperature. Meded. Land-bouwhogesch. *Wageningen*, **64**: 1–11
- Madden T. D. M. B. Bally, M. J. Hope, P. R. Cullis, H. P. Schieren, and A. S. Janoff 1985 Protection of large unilamellar vesicles by trehalose during dehydration: retention of vesicle contents. *Biochimica et Biophysica Acta.*, **817**: 67–74
- Ministry of Agriculture, Food and Rural Affairs (MAFRA) statistics. 2013 p. 15
- Nakamoto H. and T. Hiyama 1999 Heat-shock proteins and temperature stress. In “Handbook of Plant and Crop Stress” ed. by M. Pessarakli, Marcel Dekker, New York, pp. 399–416
- Raju P. S., R. B. Clark, J. R. Ellis and J. W. Maranville 1990 Effects of species of VA-mycorrhizal fungi on growth and mineral uptake of sorghum at different temperatures. *Plant and Soil*, **121**: 165–170
- Ruter J. M. and D. L. Ingram 1990 ¹⁴Carbon-labeled photosynthate partitioning in *Ilex crenata* ‘Rotundifolia’ at supraoptimal root-zone temperatures. *J. Am. Soc. Hortic. Sci.*, **115**: 1008–1013
- Sharkova V. E 2001 The effect of heat shock on the capacity of wheat plants to restore their photosynthetic electron transport after photoinhibition or repeated heating. *Russ. J. Plant Physiol.*, **48**: 793–797
- Taiz L. and E. Zeiger 2006 Plant physiology. Sinauer Associates Inc. Publishers, Massachusetts
- Tindall J. A., H. A. Mills and D. E. Radcliffe 1990 The effect of root zone temperature on nutrient uptake of tomato. *J. Plant Nutr.*, **13**: 939–956
- Todaka D., H. Matsushima and Y. Morohashi 2000 Water stress enhances beta-amylase activity in cucumber cotyledons. *J. Exp. Bot.*, **51**: 739–745
- Vollenweider P. and M. S. Gunthardt-Goerg 2005 Diagnosis of abiotic and biotic stress factors using the visible symptoms in foliage. *Environ. Pollut.*, **137**: 455–465
- Walker J. M 1969 One-degree increments in soil temperature affect maize seedling behavior. *Soil Sci. Soc. Am. Proc.*, **33**: 729–736
- Waraich E. A., R. Ahmad, A. Halim and T. Aziz 2012 Alleviation of temperature stress by nutrient management in crop plants : a review. *J. Soil Sci. Plant Nurt.*, **12**: 221–244
- Waraich E. A., R. Ahmad, M. Ashraf, Y. Saifullah and M. Ahmad 2011 Improving agricultural water use efficiency by nutrient management in crop plants. *Acta Agriculturae Scandinavica, Section B – Plant Soil Sci.*, **61**: 291–304
- Wise R. R., A. J. Olson, S. M. Chrader and T. D. Sharkey 2004 Electron transport is the functional limitation of photosynthesis in field-grown Pima cotton plants at high temperature. *Plant Cell Environ.*, **27**: 717–724
- Yang J., R. G. Sears, B. S. Gill and G. M. Paulsen 2002 Genotypic differences in utilization of assimilate sources during maturation of wheat under chronic heat and heat shock stresses. *Euphytica*, **125**: 179–188
- Youngner V. B. and F. G. Nudge 1968 Growth and carbohydrate storage of three *Poa pratensis* L. strains as influenced by temperatures. *Crop Science*, **8**: 455–456