The It^o-Nisio theorem, quadratic Wiener functionals, and 1-solitons

Ikeda, Nobuyuki

Taniguchi, Setsuo Faculty of Mathematics, Kyushu University

https://hdl.handle.net/2324/16258

出版情報:MI Preprint Series. 2010-4, 2010-01-19. 九州大学大学院数理学研究院 バージョン: 権利関係: **MI Preprint Series**

Kyushu University The Global COE Program Math-for-Industry Education & Research Hub

The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons

Nobuyuki Ikeda & Setsuo Taniguchi

MI 2010-4

(Received January 19, 2010)

Faculty of Mathematics Kyushu University Fukuoka, JAPAN

The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons

Nobuyuki Ikeda^a

^aSuzukakedai 2-40-3, Sanda, Hyogo, 669-1322, Japan

Setsuo Taniguchi ^{b,1,*}

^bFaculty of Mathematics, Kyushu University, Fukuoka 819-0395, Japan

Abstract

Among Professor Kiyosi Itô's achievements, there is the Itô-Nisio theorem, a completely general theorem relative to the Fourier series decomposition of the Brownian motion. In this paper, some of its applications will be reviewed, and new applications to 1-soliton solutions to the Korteweg-de Vries (KdV in short) equation and Eulerian polynomials will be given.

Key words: Itô-Nisio theorem, quadratic Wiener functional, stochastic area, 1-soliton, Eulerian polynomial *1991 MSC:* 60H05, 60H07

1 Introduction

Professor Kiyosi Itô's careful study about the work by N. Wiener on Brownian motion started around 1943, after he moved to Nagoya University; in the foreword to "Kiyosi Itô Selected papers", he said ([21, pp.xiv-xv])

"Although I had heard much of N. Wiener's great contribution to probability theory, I had not read his work carefully until I went to Nagoya. Even his

^{*} Corresponding author. Tel: +81-92-802-4424; Fax: +81-92-802-4405.

Email addresses: ikenobu@pearl.ocn.ne.jp (Nobuyuki Ikeda),

taniguch@math.kyushu-u.ac.jp (Setsuo Taniguchi).

URL: http://www2.math.kyushu-u.ac.jp/~taniguch/ (Setsuo Taniguchi).

 $^{^1}$ Research supported partially by Grant-in-Aid for Scientific Research (B)(1) 18340038

theory of Brownian motion I learned from Lévy's book and Doob's papers. Reading some of his papers I was impressed by the originality with which he initiated not only measure-theoretic probability theory but also paththeoretic process theory as early as the 1920's."

He then learned the Fourier series expansion of Brownian motion by N. Wiener in 1924 ([41]) and R. E. A. C. Paley-N. Wiener in 1934 ([34]). Itô was particularly interested in the latter paper, where the construction of the path of Brownian motion was used to construct the Wiener measure. As Itô wrote in the Commentary in [42, pp.513-519], it is not easy to read Wiener's paper of year 1923 ([40]), in which the existence of the Wiener measure was first proved, because of the heuristic nature of its presentation. In contrast, the Fourier series expansion given in Paley-Wiener [34] is more satisfactory from the logical point of view. Being interested in the method, Itô introduced the Fourier series expansion in detailed and clearly understandable manner in his Japanese book [20, §§38-39] published in 1953. This is an example of Itô's attitudes to devote extraordinary efforts to approaching central problems in probability theory with prior knowledge as little as possible. After the publication of his book, the Japanese mathematicians with interests in probability theory easily understood Chapters IX and X in Paley-Wiener [34].

It should be also mentioned that the sections from 64 to 66 in Itô's book are devoted to the theory of stochastic differential equations, which was created by him in 1942 ([16]). In 1950's, in Japan, many probabilists started the study of probability theory with this book, and then read together the books by J. Doob [3] and W. Feller [5].

Another Fourier series expansion of Brownian motion is the one by P. Lévy in 1940 ([27]), where the Haar wavelet expansion was used. The wavelet expansion is now used widely. For example, see Z. Ciesielski [1], M. Kac [24], and M. Pinsky [36].

Retaining an interest in the Fourier series expansion of Brownian motion, Itô established a theorem in 1968 in the joint paper with Makiko Nisio ([23]), the Itô-Nisio theorem, which is a complete generalization of Wiener's construction of the Brownian motion; they formulated the expansion as a problem on convergence of sums of independent random variables with values in a Banach space, and applied the concept of tightness due to Y. Prohorov [37] to see the convergence. Their result in the most simple case asserts that

Theorem 1 (Itô-Nisio[23], p.45, Theorem 5.2) Denote by H the Cameron-Martin space of real valued functions over [0,1], that is, the real separable Hilbert space of all real, absolutely continuous functions h on [0,1] with h(0) = 0, possessing square integrable derivatives. Let $\{u_n\}_{n=1}^{\infty}$ be an orthonormal basis of H, and $\{\xi_n\}_{n=1}^{\infty}$ be a sequence of independent and identically distributed

random variables, each of which obeys N(0,1), the normal distribution of mean 0 and variance 1. Then, the sequence

$$\sum_{n=1}^{\infty} \xi_n u_n(t), \quad t \in [0,1]$$

converges uniformly in t a.s., and the limit process $\{S(t)\}_{t\in[0,1]}$ is a 1-dimensional Brownian motion.

In addition to this, we would like to remark that Itô was very much interested in Prohorov's work [37]. For example, in 1964, 4 years before the above article on the Fourier series expansion appeared, he wrote another paper with M. Nisio ([22]) using Prohorov's result. Itô had been interested in the metric space structure of the space of probability measures. It should be recalled that the metric space structure of $\mathcal{P}(\mathbb{R})$ (\equiv the space of probability measures on \mathbb{R}) was already studied in 1930's by Lévy ([26]), and, in 1956, Prohorov introduced a distance between two measures on a complete separable metric space, which is analogous to the Lévy distance on $\mathcal{P}(\mathbb{R})$. Being very much interested in the results of 1930's, in 1943, about two decades before the joint articles with Nisio, Itô wrote an introductory book to probability theory in Japanese, where he gave a concise account of the metric space structure of $\mathcal{P}(\mathbb{R})$ ([17]).

Now returning to our subject, we recall the first section of Lévy [28, pp.171-172], where he discusses the importance of using the Fourier series expansion. After giving two definitions of the Brownian motion: the first one in the fourth line on page 171:

"Let X(t) be Wiener's well known random function, defined up to an additive constant by the condition

(1.1.1)
$$X(t) - X(t_0) = \xi \sqrt{t - t_0}, \quad t > t_0,$$

 ξ being a real and normalized Laplacian (often called Gaussian) random variable"

and the second one in the fourteenth line:

"(1.1.5)
$$X(t) = \frac{\xi' t}{\sqrt{2\pi}} + \sum_{n=1}^{\infty} \frac{1}{n\sqrt{\pi}} [\xi_n(\cos nt - 1) + \xi'_n \sin nt],$$

the Greek letters indicating normalized Laplacian random variables, all independent of each other"

he wrote

"Thus, the same random function may be defined by (1.1.1) or by (1.1.5). This theorem was proved by N. Wiener [9] in 1924, and ten years later, formula (1.1.5) was used as a definition by Paley and Wiener. Starting from one or the other point of view, it is easy to prove that X(t) is almost surely a well defined and continuous function; $\delta X(t)$ is generally $O(\sqrt{dt}) (dt > 0)$, and not O(dt). Thus X(t) is not differentiable.

The explicit representation of X(t), given by (1.1.5), is often very useful. Yet, during more than twenty years, the author and other mathematicians did not have the idea of using it. We shall now apply it to the study of the Brownian plane curve."

Lévy's method has been well known recently, and used widely by many people, for example, [24] and so on. In particular, it is indispensable in the study of quadratic Wiener functionals, which plays an essential role in stochastic analysis. Moreover, as for the Malliavin calculus, another important research field in stochastic analysis, P. Malliavin has taken the advantage of the Itô-Nisio theorem in many situations.

As N. Wiener wrote in his autobiography [43, pp.37-39], when the study on Brownian motion, his first major mathematical work, started, he was strongly influenced by the book of J. Perrin [35]. Perrin noted similarities between on one hand the non-differentiable function, which was constructed by K. Weierstrass in 1872 with the use of sums of trigonometric functions, and on the other hand the irregular movement of physical Brownian motion. The result by Weierstrass was refined by G. Hardy in 1916 ([9]). Having close exchanges with Hardy and Littlewood, Wiener must have had a detailed knowledge of these preceding results. He began to study the Brownian motion in early 1920. There is a mention that makes one guess that, from the very beginning, he had in his mind an idea of using Fourier series expansions. Actually he tackled straightforwardly the problems of Fourier series expansion in the paper of year 1924, and in the joint paper with Paley [34], he discussed systematically about the construction of Wiener measure along this line. See also [28].

As was mentioned before, the studies of the Fourier series expansion of Brownian motion bore as a fruit Lévy's Haar function expansion, and then came up to the Itô-Nisio theorem, the complete generalization. On one hand, if one approaches the study of diffusion processes with heat kernels, then the Fourier series for regular functions play a key role. On the other hand, studying Brownian motion from the point of view of path behaviour (particle movement), then the Itô-Nisio theorem and the Fourier series of irregular functions play a fundamental role. In this paper, we shall see several more concrete examples where Itô-Nisio's Fourier series expansions are indispensable, and we shall give some applications of them. As shown in his study attitude of deepening the understanding of the Wiener measure by constructing the paths of Brownian motion, Itô had a tendency to dwell on essentials of problems. This can be also seen as he developed the understanding of Kolmogorov's work [25] on diffusion processes by stochastic differential equations. Apart from stochastic topics, we would like to exhibit more episodes of Itô's interests in fundamental subjects. Itô, who returned again and again to logical structures of research subjects, was interested in extensive range of mathematics, not only probability theory. For example, in the late 1940's, he was fascinated by Gödel's theory, and his review [18] of the Japanese translation of Gödel's book received attention of Japanese mathematicians who were interested in Gödel's work. In 1960's, through the lecture by Martin Löf at Aarhus University, he was also very much interested in the concept of a random sequence by Kolmogorov.

All of Itô's mathematical works, including the Itô-Nisio theorem, consistently gave motivations and directions for our researches. Not only in his lectures, but also Itô was willing to tell young mathematicians privately what was behind his research topics. By such private conversations, we learned a lot of things and enriched our understandings over many topics. It is a great honor for us to write a paper for this Tribute to Professor Kiyosi Itô, and on this occasion we would like to express our deepest respect and gratitude to him for all he taught us directly and indirectly.

2 A generalization of Lévy's stochastic area formula

Let W be the space of all continuous functions w on $[0, \infty)$ taking values in \mathbb{R}^2 with w(0) = 0, and let P stand for the Wiener measure on W. For $x \in \mathbb{R}^2$, we set

$$w_x(s) = x + w(s), \quad s \in [0, \infty), \ w \in W.$$

To avoid any confusion, under P, we continue to write w(s) and do not use $w_0(s)$. Take the differential 1-form $\theta = (1/2)(x^1dx^2 - x^2dx^1), x = (x^1, x^2) \in \mathbb{R}^2$, on \mathbb{R}^2 . Its exterior derivative $d\theta$ is the area element $dx^1 \wedge dx^2$. We define $S(t, w_x)$ to be the stochastic line integral of θ along the curve $[0, t] \ni s \mapsto w_x(s) \in \mathbb{R}^2$;

$$S(t, w_x) = \int_{w_x[0,t]} \theta.$$

For stochastic line integrals, see [12,15]. S(t, w), i.e. $S(t, w_x)$ with x = 0, is called *Lévy's stochastic area*. It holds:

$$S(t, w_x) = \frac{1}{2} \int_0^t \langle Jw_x(s), dw_x(s) \rangle,$$

where $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $\langle \cdot, \cdot \rangle$ denotes the standard inner product in \mathbb{R}^2 ; $\langle x, y \rangle = x^1 y^1 + x^2 y^2, \ x = (x^1, x^2), \ y = (y^1, y^2) \in \mathbb{R}^2.$

From this expression with Itô integral, we see that the stochastic area is a quadratic Wiener functional, i.e., an element of the homogeneous chaos of order 2 of Wiener-Itô, which plays a key role in stochastic analysis. For details, see the second proof of Proposition 2.

While Lévy defined the stochastic area in his own manner in [27], it is now standard to use Itô's stochastic integral to define the area. The discontinuity of S(t, w) in w requires special care in defining the stochastic area ([15,29]).

Let $\alpha, \beta \in \mathbb{R}$, and define

$$p(t, x, y; \alpha, \beta) = \int_{W} \exp\left(\sqrt{-1}\,\alpha S(t, w_x) - \frac{\beta^2}{2} \int_{0}^{t} |w_x(s)|^2 ds\right) \delta_y(w_x(t)) P(dw), \tag{1}$$

where $\delta_y(w_x(t))$ stands for Watanabe's pull-back of the Dirac measure δ_y concentrating at $y \in \mathbb{R}^2$ through $w_x(t)$. For the pull-back, see [15]. Then $p(t, x, y; \alpha, \beta)$ is the fundamental solution to the partial differential equation

$$\frac{\partial u}{\partial t} = \left(\mathcal{L}_{\alpha} - \frac{\beta^2}{2}|x|^2\right)u,$$

where

$$\mathcal{L}_{\alpha} = \frac{1}{2}\Delta + \frac{\alpha\sqrt{-1}}{2} \left(x^1 \frac{\partial}{\partial x^2} - x^2 \frac{\partial}{\partial x^1} \right) - \frac{\alpha^2}{8} |x|^2, \tag{2}$$

 Δ being the Laplacian. See [15]. The heat equation was deeply studied by Gaveau [7].

It follows from the result by H. Matsumoto ([31]) that

Proposition 2 Let $\alpha, \beta \in \mathbb{R}$, and set $m_1 = (\alpha^2 + 4\beta^2)^{1/2}$. Then it holds that

$$\int_{W} \exp\left(\sqrt{-1}\,\alpha S(t,w) - \frac{\beta^2}{2} \int_{0}^{t} |w(s)|^2 ds\right) \delta_y(w(t)) P(dw)$$

= $\frac{1}{2\pi t} \frac{m_1 t/2}{\sinh(m_1 t/2)} \exp\left(-\frac{1}{2t} \frac{m_1 t/2}{\tanh(m_1 t/2)} |y|^2\right), \quad y \in \mathbb{R}^2.$ (3)

Matsumoto's result is based on the Van Vleck formula, the formula established in [11], which is a Wiener integral counterpart to the application in the Feynman path integral theory of the result due to Van Vleck [39] on fundamental solutions.

PROOF. The assertion is a special case of the observation made by Matsumoto in [31, pp.172-173]. We shall give the proof after briefly revisiting his result.

Let $k_1, k_2 > 0$ and $B \in \mathbb{R}$. Denote by $q(t, a, b), t \ge 0, a, b \in \mathbb{R}^2$, the heat kernel associated with the differential operator

$$-\left\{\frac{1}{2}\sum_{j=1}^{2} \left(\sqrt{-1}\frac{\partial}{\partial x^{j}} - B\theta_{j}\right)^{2} + \frac{1}{2}\left\{k_{1}^{2}(x^{1})^{2} + k_{2}^{2}(x^{2})^{2}\right\}\right\}$$
$$= \mathcal{L}_{B} - \frac{1}{2}\left\{k_{1}^{2}(x^{1})^{2} + k_{2}^{2}(x^{2})^{2}\right\},$$

where $\theta_1 = x_2/2$ and $\theta_2 = -x_1/2$, and \mathcal{L}_B is the operator defined in (2). Setting

$$m_1 = \{(k_1 + k_2)^2 + B^2\}, \quad m_2 = \{(k_1 - k_2)^2 + B^2\},\$$
$$s_1 = \frac{m_1 + m_2}{2}, \quad s_2 = \frac{-m_1 + m_2}{2},\$$

$$K(t) = 2k_1k_2B^2(\cosh(s_1t)\cosh(s_2t) - 1) -\{B^2(k_1^2 + k_2^2) + (k_1 - k_2)^2\}\sinh(s_1t)\sinh(s_2t),$$

$$\alpha_i(t) = s_1(s_2^2 - k_i^2) \cosh(s_1 t) \sinh(s_2 t) -s_2(s_1^2 - k_i^2) \sinh(s_1 t) \cosh(s_2 t),$$

$$\beta_i(t) = s_2(s_1^2 - k_i^2)\sinh(s_1t) - s_1(s_2^2 - k_i^2)\sinh(s_2t), \quad i = 1, 2,$$

$$\gamma(t) = 2k_1k_2\{\cosh(s_1t)\cosh(s_2t) - 1\} + (m_1^2 - 2k_1k_2)\sinh(s_1t)\sinh(s_2t),$$

$$\begin{split} \widetilde{S}_{cl}(t,a,b) &= \frac{m_1 m_2}{K(t)} \alpha_1(t) \{ (a^1)^2 + (b_1)^2 \} + \frac{m_1 m_2}{K(t)} \beta_1(t) a^1 b^1 \\ &+ \frac{m_1 m_2}{K(t)} \alpha_2(t) \{ (a^2)^2 + (b^2)^2 \} + \frac{m_1 m_2}{K(t)} \beta_2(t) a^2 b^2 \\ &+ \frac{\sqrt{-1} B(k_1^2 - k_2^2)}{2K(t)} \gamma(t) (a^1 a^2 - b^1 b^2) \\ &- \frac{\sqrt{-1} k_1 k_2 m_1 m_2 B}{K(t)} \{ \cosh(s_1 t) - \cosh(s_2 t) \} (a^1 b^2 - a^2 b^1), \end{split}$$

Matsumoto [31] showed that

$$q(t,a,b) = \frac{1}{2\pi} \left(\frac{k_1 k_2 m_1^2 m_2^2}{K(t)}\right)^{1/2} \exp(-\tilde{S}_{cl}(t,a,b)).$$
(4)

In our situation, $B = \alpha$, $k_1 = k_2 = \beta$, a = 0, and b = y. Hence $m_1 = (\alpha^2 + 4\beta^2)^{1/2}$ and $m_2 = |\alpha|$. Using the identities: $\cosh x \cosh y - \sinh x \sinh y = \cosh(x-y)$ and $\cosh x - 1 = 2\sinh^2(x/2)$, we see that

$$K(t) = 4\alpha^2 \beta^2 \sinh^2(m_1 t/2),$$

which implies:

$$\frac{k_1 k_2 m_1^2 m_2^2}{K(t)} = \frac{(m_1/2)^2}{\sinh^2(m_1 t/2)}.$$

Since

$$s_1(s_2^2 - \beta^2) = s_2(s_1^2 - \beta^2) = -\beta^2 |\alpha|$$

and $\sinh x \cosh y - \cosh x \sinh y = \sinh(x - y)$, it holds that

$$\alpha_1(t) = \alpha_2(t) = |\alpha|\beta^2 \sinh(m_1 t).$$

Then

$$\widetilde{S}_{cl}(t,0,y) = \frac{m_1 m_2}{2K(t)} \alpha_1(t) |y|^2 = \frac{1}{2t} \frac{m_1 t/2}{\tanh(m_1 t/2)} |y|^2.$$

Plugging these into (4), we obtain the desired identity. \Box

Remark 3 The identity in the case when $\alpha = 0$ can be shown as an application of the Feynman-Kac formula. If $\beta = 0$ and y = 0, then the identity is well-known Lévy's formula ([28]).

Remark 4 Observe that

$$\alpha \int_{0}^{t} w^{1}(s) dw^{2}(s) + \beta \int_{0}^{t} w^{2}(s) dw^{1}(s) = \frac{\alpha - \beta}{2} S(t, w) + \frac{\alpha + \beta}{2} w^{1}(t) w^{2}(t).$$

This identity was used by M. Yor [44] to study the joint distribution

$$\left(\int_{0}^{1} w^{1}(s)dw^{2}(s), \int_{0}^{1} w^{2}(s)dw^{1}(s)\right).$$

Using this expression, we obtain a variant of Matsumoto's result to the operator

$$\left(\sqrt{-1}\frac{\partial}{\partial x^1} + \frac{B_1 x^2}{2}\right)^2 + \left(\sqrt{-1}\frac{\partial}{\partial x^2} + \frac{B_2 x^1}{2}\right)^2 + \frac{1}{2}\left\{k_1^2 (x^1)^2 + k_2^2 (x^2)^2\right\}$$

with $B_1, B_2 \in \mathbb{R}$.

In the remainder of this section, we shall give an alternative proof of Proposition 2 with $\beta = 0$ by using the general Fourier series expansion of Brownian motion due to Itô-Nisio. The method was first used by Lévy [28] in the case when y = 0. The following observation was essentially made by Ikeda-Watanabe [15, pp.476-478], and we are aiming to make clear which specified trigonometric functions are involved in this computation.

PROOF. (An alternative proof of Proposition 2 for $\beta = 0$)

Suppose $\beta = 0$. Due to the scaling property of Brownian motion, it suffices to show (3) for t = 1. Hence we work on the space $\{w|_{[0,1]} | w \in W\}$ of restrictions of elements in W on [0, 1]. For the sake of simplicity, we use the same letter W to indicate the space of restrictions. Let H be the corresponding Cameron-Martin space. Then:

$$L^2(W;P) = \bigoplus_{n=0}^{\infty} \mathcal{C}_n$$

denotes the decomposition of $L^2(W; P)$, the Hilbert space of square integrable variables with respect to P, in terms of the homogeneous chaos of Wiener-Itô ([19]). A quadratic Wiener functional is an element of C_2 . By the result of Itô on multiple Wiener integrals ([19]), each quadratic Wiener functional $F \in C_2$ admits a kernel representation;

$$F(w) = \sum_{i,j=1}^{2} \left\{ \int_{0}^{1} \int_{0}^{t} F^{ij}(t,s) dw^{j}(s) dw^{i}(t) + \int_{0}^{1} \int_{0}^{s} F^{ij}(t,s) dw^{i}(t) dw^{j}(s) \right\},$$

where F^{ij} is square integrable on $[0, 1]^2$ with respect to the Lebesgue measure and $F^{ij}(t, s) = F^{ji}(s, t), t, s \in [0, 1], i, j = 1, 2$. Setting the 2×2 matrix F(t, s)to be $(F^{ij}(t, s))_{1 \le i,j \le 2}$, we define a symmetric Hilbert-Schmidt operator B of H into H by

$$(Bh)'(t) = \int_0^1 F(t,s)h'(s)ds, \quad h \in H,$$

where h' denotes the derivative of h. Such a correspondence between quadratic Wiener functionals and symmetric Hilbert-Schmidt operators of H into H is bijective ([13]).

It is easily seen that S(1, w) is a quadratic Wiener functional. The corresponding kernel $F(t, s) = (F^{ij}(t, s))_{1 \le i, j \le 2}$ is given by

$$F^{11}(t,s) = F^{22}(t,s) = 0,$$

$$F^{12}(t,s) = F^{21}(s,t) = \begin{cases} \frac{1}{4}, & 0 \le t \le s \le 1, \\ \\ -\frac{1}{4}, & 0 \le s < t \le 1. \end{cases}$$

Moreover, if we set

$$\mathcal{I}[h] = \binom{\mathcal{I}[h^1]}{\mathcal{I}[h^2]}, \quad \mathcal{I}[h^i](s) = \int_0^s h^i(u) du, \quad h \in H,$$

then the Hilbert-Schmidt operator B associated with S(1, w) is represented as

$$B = B_V + B_F,\tag{5}$$

where

$$(B_V h)(s) = \frac{1}{2}\mathcal{I}[Jh](s), \quad (B_F h)(s) = -\frac{1}{4}(Jh)(1)s, \quad s \in [0,1].$$

Let $H_0 = \{h \in H | h(1) = 0\}$ and set $B^{\#} = \pi_0 B \pi_0$, π_0 being the orthogonal projection of H onto H_0 . Then the above decomposition (5) implies that

$$(B^{\#}h)(s) = \frac{1}{2}\mathcal{I}[Jh](s) - \frac{1}{2}\mathcal{I}[Jh](1)s, \quad s \in [0,1].$$

It is then easily checked that $B^{\#}$ admits the eigenfunction expansion:

$$B^{\#} = \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{1}{2} \frac{1}{2n\pi} \{ k_n \otimes k_n + \widetilde{k}_n \otimes \widetilde{k}_n \},\$$

where $k \otimes k : H \to H$ is defined by $(k \otimes k)(h) = \langle k, h \rangle_H k, h \in H$, and

$$k_n(s) = \frac{1}{2n\pi} \binom{\cos(2n\pi s) - 1}{\sin(2n\pi s)}, \ s \in [0, 1], \quad \tilde{k}_n = Jk_n.$$

See [13,32]. This leads us to the orthonormal basis $\{\phi_n^1, \psi_{n+1}^1, \phi_n^2, \psi_{n+1}^2\}_{n=0}^{\infty}$ of H defined by

$$\begin{split} \phi_0^1(s) &= \binom{s}{0}, \quad \phi_0^2(s) = \binom{0}{s}, \\ \phi_n^1(s) &= \frac{\sqrt{2}}{2n\pi} \binom{\cos(2n\pi s) - 1}{0}, \quad \phi_n^2(s) = \frac{\sqrt{2}}{2n\pi} \binom{0}{\cos(2n\pi s) - 1}, \\ \psi_n^1(s) &= \frac{\sqrt{2}}{2n\pi} \binom{\sin(2n\pi s)}{0}, \quad \psi_n^2(s) = \frac{\sqrt{2}}{2n\pi} \binom{0}{\sin(2n\pi s)}. \end{split}$$

Applying the Itô-Nisio theorem, we obtain the expansion:

$$w(s) = \sum_{i=1}^{2} \left\{ \sum_{n=0}^{\infty} \xi_n^{(i)}(w) \phi_n^i(s) + \sum_{m=1}^{\infty} \eta_m^{(i)}(w) \psi_m^i(s) \right\}$$

where

$$\xi_n^{(i)}(w) = \int_0^1 \langle (\phi_n^i)'(s), dw(s) \rangle, \quad \eta_m^{(i)}(w) = \int_0^1 \langle (\psi_m^i)'(s), dw(s) \rangle.$$

Since $X_0^i(w) = w^i(1)$, in terms of components, this can be rewritten as

$$w^{i}(s) = w^{i}(1)\phi_{0}^{i,i} + \sum_{n=1}^{\infty} \xi_{n}^{(i)}(w)\phi_{n}^{i,i}(s) + \sum_{n=1}^{\infty} \eta_{n}^{(i)}(w)\psi_{n}^{i,i}(s), \quad i = 1, 2,$$

where $\phi_n^i = \begin{pmatrix} \phi_n^{i,1} \\ \phi_n^{i,2} \end{pmatrix}$ and $\psi_n^i = \begin{pmatrix} \psi_n^{i,1} \\ \psi_n^{i,2} \end{pmatrix}$. It is easily seen that $\{w^i(1), \xi_n^{(i)}, \eta_n^{(i)}; i = \psi_n^{i,1}\}$.

 $1, 2, n = 1, 2, \ldots$ are independent random variables, each of which obeys N(0, 1). These expansions are exactly the same as the ones used in [15, pp.476-477]. In particular, we have that, *P*-a.s.,

$$S(1,w) = \sum_{n=1}^{\infty} \frac{1}{2\pi n} \Big\{ (\eta_n^{(1)}(w) - \sqrt{2} w^1(1)) \xi_n^{(2)} - (\eta_n^{(2)}(w) - \sqrt{2} w^2(1)) \xi_n^{(1)} \Big\}.$$

As was seen by Ikeda-Watanabe, it holds that

$$E[e^{\sqrt{-1}\alpha S(1)}|w(1) = y] = \prod_{n=1}^{\infty} \left(1 + \frac{\alpha^2}{(2n\pi)^2}\right)^{-1} \exp\left(-\sum_{n=1}^{\infty} \frac{(\alpha/2\pi n)^2}{1 + (\alpha/2\pi n)^2}|y|^2\right),\tag{6}$$

where $E[\cdot|w(1) = y]$ stands for the conditional expectation given w(1) = y. In conjunction with the well-known formulas

$$\sinh x = x \prod_{n=1}^{\infty} \left(1 + \frac{x^2}{\pi^2 n^2} \right), \quad \coth x = \frac{1}{x} + 2x \sum_{n=1}^{\infty} \frac{1}{x^2 + \pi^2 n^2} \tag{7}$$

and the identity

$$\int_{W} e^{\sqrt{-1}\alpha S(1,w)} \delta_y(w(1)) P(dw) = \frac{1}{2\pi} e^{-|y|^2/2} E[e^{\sqrt{-1}\alpha S(1)} | w(1) = y]$$

we arrive at the desired identity. 2 $\hfill\square$

 $[\]overline{^2$ In [15, p.478]}, the multiplication by 1/2 is missing when the infinite series expression of cot is applied.

Remark 5 The first product in (6) can be rewritten as

$$\prod_{n=1}^{\infty} \left(1 + \frac{\alpha^2}{(2n\pi)^2} \right)^{-1} = \left\{ \det_2(I - 2\sqrt{-1}\,\alpha B^{\#}) \right\}^{-1/2}$$

Remark 6 It is well known ([13,28,32]) that the Hilbert-Schmidt operator B is decomposed as

$$B = \sum_{n \in \mathbb{Z}} \frac{1}{2} \frac{1}{(2n+1)\pi} \{ h_n^B \otimes h_n^B + \tilde{h}_n^B \otimes \tilde{h}_n^B \}$$

where

$$h_n^B(s) = \frac{1}{(2n+1)\pi} \left(\frac{\cos((2n+1)\pi t) - 1}{\sin((2n+1)\pi t)} \right), \quad \tilde{h}_n^B = Jh_n^B$$

In particular, B has the eigenvalues $1/\{2(2n\pi)\pi\}$, $n \in \mathbb{Z}$, the multiplicity of each being two, and:

$$\int_{W} e^{\sqrt{-1}\,\alpha S(1)} dP = \frac{1}{\cosh(\alpha/2)} = \{\det_2(I - 2\sqrt{-1}\alpha B)\}^{-1/2}.$$

Remark 7 The formulas in (7) are the well known Euler formulas. The latter proof relies upon these formulas. Conversely, the Euler formulas can be shown by combining Proposition 2 and the identity (6).

In [24, Sections 2 and 7], M. Kac also gave two such directions to approach Lévy's formula in the case when $\alpha = 0$, i.e., the case of harmonic oscillator. The importance of proving Lévy's formula from two directions is discussed there. Two such types of computation are widely known in the theory of Feynman path integrals. See for example [6, Problems 2-2, 3-8, pp.71-73].

3 Lévy's stochastic area formula for Ornstein-Uhlenbeck processes

In [14,38], the authors gave a probabilistic approach to reflectionless potentials, soliton solutions and the τ -function to the KdV equation. Their observation is based on quadratic Wiener functionals obtained as the square norms on time interval for Ornstein-Uhlenbeck processes. In this section, we investigate stochastic areas determined by Ornstein-Uhlenbeck processes, and establish a similar result to the one in [14]. In the case of the Brownian motion, i.e., the Ornstein-Uhlenbeck process with p = 0, the result is well known ([8,28]). We continue to work on the Wiener space (W, P) of 2-dimensional Brownian

motion. To apply the KdV equation, real numbers x, y, z instead of s, t, u are used to indicate time parameters.

For $p \in \mathbb{R}$, define the 2-dimensional Ornstein-Uhlenbeck process $\{\xi^p(z)\}_{z\geq 0}$ to be the solution to the stochastic differential equation

$$d\xi^{p}(y) = dw(y) + p\xi^{p}(y)dy, \quad \xi^{p}(0) = 0,$$

that is, if we represent: $\xi^p(y)=(\xi^{p,1}(y),\xi^{p,2}(y))$ and $w(y)=(w^1(y),w^2(y)),$ then

$$d\xi^{p,i}(y) = dw^{i}(y) + p\xi^{p,i}(y)dy, \quad \xi^{p,i}(0) = 0, \quad i = 1, 2.$$

In the sequel, let x > 0. By the Maruyama-Girsanov theorem [30,15], the process $\left\{w(z) - \int_0^z pw(y)dy\right\}_{z\in[0,x]}$ is a 2-dimensional Brownian motion under the probability measure $\exp(p\int_0^x \langle w(y), dw(y) \rangle - (p^2/2)\int_0^x |w(y)|^2 dy) dP$. Since $\int_0^x \langle w(y), dw(y) \rangle = \frac{1}{2}|w(x)|^2 - x$, we have:

$$\int_{W} \Phi(\{\xi^{p}(z)\}_{z \in [0,x]}) dP$$

= $\int_{W} \Phi(\{w(z)\}_{z \in [0,x]}) \exp\left(\frac{p}{2}|w(x)|^{2} - \frac{p^{2}}{2}\int_{0}^{x}|w(y)|^{2}dy\right) P(dw)e^{-px}$

for every $\Phi \in C_b(W_x)$, where $W_x = \{w|_{[0,x]} | w \in W\}$. In particular, for $\Psi \in C_b(W_x)$ such that $\Psi(\{\xi^p(z)\}_{z\in[0,x]})$ is smooth in the sense of the Malliavin calculus, it also holds that, for any $b \in \mathbb{R}^2$,

$$\int_{W} \Psi(\{\xi^{p}(z)\}_{z\in[0,x]}) \delta_{b}(\xi^{p}(x)) dP$$

= $\int_{W} \Psi(\{w(z)\}_{z\in[0,x]}) \exp\left(-\frac{p^{2}}{2} \int_{0}^{x} |w(y)|^{2} dy\right) \delta_{b}(w(x)) P(dw) e^{p\{(|b|^{2}/2)-x\}}.$

Thus we arrive at

Proposition 8 For $b \in \mathbb{R}^2$, $\alpha, C \in \mathbb{R}$, it holds that

$$\int_{W} \exp\left(\frac{\sqrt{-1}\alpha}{2} \int_{0}^{x} \langle J\xi^{p}(y), d\xi^{p}(y) \rangle - C|\xi^{p}(x)|^{2}\right) \delta_{b}(\xi^{p}(x)) dP$$
$$= \int_{W} \exp\left(\sqrt{-1}\alpha S(x, w) - \frac{p^{2}}{2} \int_{0}^{x} |w(y)|^{2} dy\right) \delta_{b}(w(x)) P(dw)$$

$$\times \exp\left(\left\{\frac{p}{2} - C\right\}|b|^2 - px\right).\tag{8}$$

It should be mentioned that the above expression is meaningful for any C, while the integrand $\exp(\cdots)$ on the left hand side is not smooth in the sense of the Malliavin calculus for C < 0 with large absolute value. Namely, by the localization via $\delta_b(\xi^p(x))$, one may replace $|\xi^p(x)|^2$ by $\psi(|\xi^p(x)|^2)$ with smooth $\psi : \mathbb{R} \to \mathbb{R}$ which is compactly supported and equal to the identity function on the interval $[0, 2|b|^2)$. After this replacement, one can take the pairing with Watanabe's pull-back.

In conjunction with Proposition 2, this proposition implies that

Theorem 9 Let $\alpha, C \in \mathbb{R}$ and set $m_1 = (\alpha^2 + 4p^2)^{1/2}$. Then it holds that

$$\int_{W} \exp\left(\frac{\sqrt{-1}\alpha}{2} \int_{0}^{x} \langle J\xi^{p}(y), d\xi^{p}(y) \rangle - C |\xi^{p}(y)|^{2}\right) \delta_{b}(\xi^{p}(y)) dP$$
$$= \frac{1}{2\pi x} \frac{m_{1}x/2}{\sinh(m_{1}x/2)} \exp\left(-\left\{\frac{1}{2x} \frac{m_{1}x/2}{\tanh(m_{1}x/2)} + \left(C - \frac{p}{2}\right)\right\} |b|^{2} - px\right).$$
(9)

Moreover, if $C \ge p/2$, then it holds that

$$\int_{W} \exp\left(\frac{\sqrt{-1} \alpha}{2} \int_{0}^{x} \langle J\xi^{p}(y), d\xi^{p}(y) \rangle - C |\xi^{p}(x)|^{2}\right) dP$$
$$= \frac{m_{1} e^{-px}}{m_{1} \cosh(m_{1} x/2) + (4C - 2p) \sinh(m_{1} x/2)}.$$
(10)

PROOF. The identity (9) follows immediately from Propositions 2 and 8. If $C \ge p/2$, then

$$\frac{1}{2x}\frac{m_1x/2}{\tanh(m_1x/2)} + \left(C - \frac{p}{2}\right) > 0.$$

Integrating (9) in b over \mathbb{R}^2 , we come to the identity:

$$\int_{W} \exp\left(\frac{\sqrt{-1} \alpha}{2} \int_{0}^{x} \langle J\xi^{p}(y), d\xi^{p}(y) \rangle - C |\xi_{x}^{p}|^{2}\right) dP$$

= $\frac{m_{1}/2}{\sinh(m_{1}x/2)} \left\{ 2 \left(\frac{1}{2x} \frac{m_{1}x/2}{\tanh(m_{1}x/2)} + \left(C - \frac{p}{2}\right)\right) \right\}^{-1} e^{-px},$

which implies (10). \Box

We first apply Theorem 9 to reflectionless potentials. For $\eta, m > 0$, the reflectionless potential with scattering data (η, m) is the function on \mathbb{R} of the form

$$-2(d/dx)^2 \log\left(1 + \frac{m}{2\eta} e^{-2\eta x}\right).$$

See [33]. We have that

Corollary 10 We continue to use the same notation as in Theorem 9. Let $C \in [p/2, (p/2) + (m_1/4))$, and put

$$Q(x) = \log\left(\int_{W} \exp\left(\frac{\sqrt{-1}\,\alpha}{2}\int_{0}^{x} \langle J\xi^{p}(y), d\xi^{p}(y) \rangle - C|\xi^{p}(x)|^{2}\right) dP\right).$$

Then the function $q = 2(d/dx)^2Q$ is the reflectionless potential with scattering data

$$\left(\frac{m_1}{2}, \frac{m_1(m_1 - 4C + 2p)}{m_1 + 4C - 2p}\right).$$

If $p \leq 0$, then we can take C = 0 in the above equation and

$$2(d/dx)^2 \log\left(\int_{W} \exp\left(\frac{\sqrt{-1}\,\alpha}{2}\int_{0}^{x} \langle J\xi^p(y), d\xi^p(y) \rangle\right) dP\right)$$

is a reflectionless potential.

PROOF. Define $\gamma \geq 0$ so that

$$m_1 \tanh \gamma = 4C - 2p$$
, i.e., $e^{-2\gamma} = \frac{m_1 - 4C + 2p}{m_1 + 4C - 2p}$. (11)

Since $\cosh(t + s) = \cosh t \cosh s + \sinh t \sinh s$, we have:

$$m_1 \cosh(m_1 x/2) + (4C - 2p) \sinh(m_1 x/2)$$
$$= \frac{m_1}{\cosh \gamma} \frac{e^{(m_1 x/2) + \gamma}}{2} (1 + e^{-2\gamma} e^{-m_1 x}).$$

By Theorem 9, it then holds:

$$q(x) = -2(d/dx)^2 \log(1 + e^{-2\gamma} e^{-m_1 x}).$$

Thus q is the reflectionless potential with scattering data $(m_1/2, m_1 e^{-2\gamma})$. \Box

As an application of the corollary, we consider a stochastic representation of 1-soliton solutions to the KdV equation. After deforming the Brownian motion into Ornstein-Uhlenbeck processes with parameter p, we shall obtain a 1-parameter family of soliton solutions of the KdV equation which varies according to the initial condition.

To see this, for $p \in \mathbb{R}, t \ge 0$, we set

$$V^{p}(x,t) = \log\left(\int_{W} \exp\left(\frac{\sqrt{-1}\alpha}{2} \int_{0}^{x} \langle J\xi^{p}(y), d\xi^{p}(y) \rangle - \left\{\frac{p}{2} + \frac{m_{1}}{4} \tanh(m_{1}^{3}t/8)\right\} |\xi^{p}(x)|^{2}\right) dP\right)$$

and

$$v^p(x,t) = 2(\partial/\partial x)^2 V(x,t).$$

The 1-parameter family $\{v^p; p \in \mathbb{R}\}$ satisfies:

Corollary 11 $\{v^p; p \in \mathbb{R}\}$ is a 1-parameter family of 1-soliton solutions v^p to the KdV equation

$$\frac{\partial u}{\partial t} = \frac{3}{2} u \frac{\partial u}{\partial x} + \frac{1}{4} \frac{\partial^3 u}{\partial x^3} \quad \text{with } u(x,0) = -\frac{(\alpha^2 + 4p^2)^{1/2}}{2\cosh^2((\alpha^2 + 4p^2)^{1/2}x)}.$$

PROOF. Due to Corollary 10 and (11), for each $t \ge 0$, $v^p(\cdot, t)$ is the reflectionless potential with scattering data $(m_1/2, m_1 e^{-2(m_1/2)^3 t})$. Then it is well known that v^p is a 1-soliton solution (cf. [33]). The initial value $v^p(x, 0)$ is easily computed. \Box

4 Eulerian polynomials and stochastic area

Recently F. Hirzebruch [10] and A. Cohen [2] began to study several types of Eulerian polynomials from a new point of view. In [28], Lévy pointed out that

the stochastic area is related to Euler and Bernoulli numbers. In this section, we shall see that the stochastic area is also related to Eulerian polynomials introduced by Euler [4] (also see [2,10]).

Define the Eulerian polynomials $P_n(\xi)$, n = 0, 1, ..., by

$$\sum_{k=0}^{\infty} (k+1)^n \xi^k = \frac{P_n(\xi)}{(1-\xi)^{n+1}}, \quad |\xi| < 1.$$

For the convenience of the reader, let us recall that

$$\frac{1}{1-\xi} = \sum_{k=0}^{\infty} \xi^k, \quad |\xi| < 1,$$

so that $P_0(\xi) = 1$. Differentiating both sides of the previous identity, we get:

$$P_1(\xi) = 1.$$

Further differentiating both sides of the identity, thus defining P_n , by induction, we see that P_n is of degree n - 1 (cf. [10]).

From the definition, we easily obtain the exponential generating function for the Eulerian polynomials ([10]):

$$\sum_{n=0}^{\infty} P_n(\xi) \frac{\lambda^n}{n!} = \frac{(1-\xi)e^{(1-\xi)\lambda}}{1-\xi e^{(1-\xi)\lambda}} \quad \text{for } \lambda \in \mathbb{R} \text{ with } |\xi|e^{(1-\xi)\lambda} < 1.$$
(12)

We shall show that

Proposition 12 For $-1 < \xi \leq 0$ and $\lambda \in \mathbb{R}$ with $|\xi|e^{(1-\xi)\lambda} < 1$ and $|\lambda| < 1$, *it holds:*

$$\sum_{n=0}^{\infty} P_n(\xi) \frac{\lambda^n}{n!} = \int_{W} e^{\sqrt{-1}(1-\xi)\lambda S(1,w) + (1-\xi)\lambda|w(1)|^2/4} P(dw) \\ \times \int_{W} e^{\sqrt{-1}(1-\xi)\lambda S(1,w) + (1+\xi)\lambda|w(1)|^2/4} P(dw).$$
(13)

The above product may be gathered into one integration with respect to 4dimensional Brownian motion. Before proceeding to the proof, we see that exact representations of Eulerian polynomials follow from (13). Namely, the identity (13) yields:

$$P_n(\xi) = \sum_{k=0}^n \frac{n!}{k!(n-k)!} \int_W \left\{ \sqrt{-1} \left(1-\xi\right) S(1,w) + \frac{(1-\xi)|w(1)|^2}{4} \right\}^k P(dw) \\ \times \int_W \left\{ \sqrt{-1} \left(1-\xi\right) S(1,w) + \frac{(1+\xi)|w(1)|^2}{4} \right\}^{n-k} P(dw).$$
(14)

For example, we can compute $P_1(\xi)$, $P_2(\xi)$, and $P_3(\xi)$ as follows. Let

$$C_{j,k} = \int_{W} S(1,w)^{j} |w(1)|^{2k} P(dw).$$

Applying Itô's formula, we see:

$$C_{1,0} = C_{1,1} = C_{1,2} = C_{3,0} = 0, (15)$$

$$C_{0,k} = 2^k k!, \quad k = 1, 2, 3, \quad C_{2,0} = \frac{1}{4}, \quad C_{2,1} = \frac{5}{6}.$$
 (16)

Then it follows from (14) and (15) that

$$P_{1}(\xi) = \frac{1}{2}C_{0,1},$$

$$P_{2}(\xi) = \left\{-2C_{2,0} + \frac{C_{0,2}}{2^{3}} + \frac{C_{0,1}^{2}}{2^{3}}\right\} + 4C_{2,0}\xi + \left\{-2C_{2,0} + \frac{C_{0,2}}{2^{3}} - \frac{C_{0,1}^{2}}{2^{3}}\right\}\xi^{2},$$

$$P_{3}(\xi) = \left\{-\frac{3}{2}C_{2,1} + \frac{1}{2^{5}}C_{0,3} - \frac{3}{2}C_{0,1}C_{2,0} + \frac{3}{2^{5}}C_{0,1}C_{0,2}\right\} + \left\{3C_{2,1} + 3C_{0,1}C_{2,0}\right\}\xi + \left\{-\frac{3}{2}C_{2,1} + \frac{3}{2^{5}}C_{0,3} - \frac{3}{2}C_{0,1}C_{2,0} - \frac{3}{2^{5}}C_{0,1}C_{0,2}\right\}\xi^{2}.$$

Substituting (16) into these, we arrive at the well known expressions ([10]):

$$P_1(\xi) = 1, \quad P_2(\xi) = 1 + \xi, \quad P_3(\xi) = 1 + 4\xi + \xi^2.$$

For the proof of Proposition 12, we prepare a lemma.

Lemma 13 For $a \in \mathbb{R}$ and $\beta \in \mathbb{R}$ with $|a| \leq 1/2$ and $-a\beta < 1$, it holds:

$$\int_{W} e^{\sqrt{-1}\beta\{S(1,w) + \sqrt{-1}a|w(1)|^{2}/2\}} P(dw) = \left\{ \left(\frac{1}{2} + a\right)e^{\beta/2} + \left(\frac{1}{2} - a\right)e^{-\beta/2} \right\}^{-1}.$$

PROOF. Since $-a\beta < 1$, $e^{\sqrt{-1}\beta\{S(1,w)+\sqrt{-1}a|w(1)|^2/2\}}$ is smooth in the sense of the Malliavin calculus. Then we have:

$$\int_{W} e^{\sqrt{-1}\beta\{S(1,w)+\sqrt{-1}a|w(1)|^{2}/2\}} P(dw)$$

$$= \int_{\mathbb{R}^{2}} dy \int_{W} e^{\sqrt{-1}\beta\{S(1,w)+\sqrt{-1}a|w(1)|^{2}/2\}} \delta_{y}(w(1)) P(dw).$$

By Proposition 2, this is equal to

$$\frac{1}{2\pi} \frac{\beta/2}{\sinh(\beta/2)} \int_{\mathbb{R}^2} dy \exp\left(-\frac{\beta/2}{\sinh(\beta/2)} \left\{\cosh(\beta/2) + 2a\sinh(\beta/2)\right\} \frac{|y|^2}{2}\right).$$

Notice that

$$\cosh(\beta/2) + 2a\sinh(\beta/2) = \left(\frac{1}{2} + a\right)e^{\beta/2} + \left(\frac{1}{2} - a\right)e^{-\beta/2} > 0.$$

Then, by an elementary change of variables, we obtain the desired identity. \Box

We now proceed to the:

PROOF of Proposition 12. Suppose that $-1 < \xi \leq 0$, $|\xi|e^{(1-\xi)\lambda} < 1$, and $|\lambda| < 1$. Observe that

$$\sum_{n=0}^{\infty} P_n(\xi) \frac{\lambda^n}{n!} = e^{(1-\xi)\lambda/2} \left\{ \frac{1}{1-\xi} e^{-(1-\xi)\lambda/2} - \frac{\xi}{1-\xi} e^{(1-\xi)\lambda/2} \right\}^{-1}.$$
 (17)

Applying Lemma 13 with a = -1/2 and $\beta = (1 - \xi)\lambda$, we have:

$$\int_{W} e^{\sqrt{-1}(1-\xi)\lambda S(1,w) + (1-\xi)\lambda|w(1)|^2/4} P(dw) = e^{(1-\xi)\lambda/2}.$$
(18)

If $a = -(1 + \xi)/\{2(1 - \xi)\}$ and $\beta = (1 - \xi)\lambda$, then $|a| \le 1/2$ and $-a\beta < 1$. Applying Lemma 13 again, since $(1/2) + a = -\xi/(1 - \xi)$ and $(1/2) - a = 1/(1 - \xi)$, we obtain:

$$\int_{W} e^{\sqrt{-1}(1-\xi)\lambda S(1,w) + (1+\xi)\lambda|w(1)|^{2}/4} P(dw)$$

= $\left\{ \frac{1}{1-\xi} e^{-(1-\xi)\lambda/2} - \frac{\xi}{1-\xi} e^{(1-\xi)\lambda/2} \right\}^{-1}.$

Plugging this and (18) into (17), we obtain the desired identity. \Box

Let $P(B_n; \xi)$ be the Eulerian polynomial of type B_n , where we have borrowed the notation from Cohen [2]. Then it holds ([2,10]):

$$\sum_{k=0}^{\infty} (2k+1)^n \xi^k = \frac{P(B_n;\xi)}{(1-\xi)^{n+1}}.$$

A similar result as above can be shown for $P(B_n; \xi)$, n = 0, 1, ... Namely, the exponential generating function satisfies:

$$\sum_{n=0}^{\infty} P(B_n;\xi) \frac{\lambda^n}{n!} = \frac{(1-\xi)e^{(1-\xi)\lambda}}{1-\xi e^{2(1-\xi)\lambda}} \quad \text{for } |\xi|e^{2(1-\xi)\lambda} < 1.$$

It follows from this expression that

$$P(B_n; -1) = 2^n E_n,$$

where the Euler number E_n is the *n*-th derivative of $1/\cosh x$ at x = 0 ([10]). Suppose that $-1 < \xi \le 0$, $|\xi|e^{2(1-\xi)\lambda} < 1$, and $|\lambda| < 1$. Applying Lemma 13 with $a = -(1+\xi)/\{2(1-\xi)\}$ and $\beta = 2(1-\xi)\lambda$, we obtain:

$$\sum_{n=0}^{\infty} P(B_n;\xi) \frac{\lambda^n}{n!} = \int_{W} e^{2\sqrt{-1}(1-\xi)\lambda S(1,w) + (1+\xi)\lambda|w(1)|^2/2} P(dw),$$

and

$$P(B_n;\xi) = \iint_W \left\{ 2\sqrt{-1} \left(1-\xi\right) S(1,w) + \frac{(1+\xi)|w(1)|^2}{2} \right\}^n P(dw).$$

References

- Z. Ciesielski, Hölder conditions for realizations of Gaussian processes, Trans. Amer. Math. Soc. 99 (1961) 403-413.
- [2] A. Cohen, Eulerian polynomials of spherical type, Münster J. of Math. 1 (2008) 1–8.
- [3] J.L. Doob, Stochastic processes, John Wiley, 1953.
- [4] L. Euler, Remarques sur un beau rapport entre les séries des puissances tant directes que réciproques, Académie des sciences de Berlin, Lu en 1749, Opera Omnia Serie I, 15, 70–90.
- [5] W. Feller, An introduction to probability theory and its applications, vol.1, 2, John Wiley, 1950, 1966.

- [6] R. P. Feynman and A. R. Hibbs, Quantum Mechanics and Path Integrals, McGraw-Hill, 1965.
- [7] B. Gaveau, Principe de moindre action, propagation de la chaleur et estimées sous elliptiques sur certains groupes nilpotents, Acta Math. 139 (1977) 95–153.
- [8] K. Hara and N. Ikeda, Quadratic Wiener functionals and dynamics on Grassmannians, Bull. Sci. math. 125 (2001) 481–528
- [9] G. H. Hardy, Weierstrass's Non-Differentiable Function, Trans. Amer. Math. Soc. 17 (1916) 301–325.
- [10] F. Hirzebruch, Eulerian polynomials, Münster J. of Math. 1 (2008) 9–14.
- [11] N. Ikeda, S. Kusuoka, S. Manabe, Lévy's stochastic area formula and related problems, in "Stochastic analysis (Ithaca, NY, 1993)", 281–305, Proc. Sympos. Pure Math., 57, Amer. Math. Soc., 1995.
- [12] N. Ikeda and S. Manabe, Integral of differential forms along the path of diffusion processes, Publ. Res. Inst. Math. Sci. 15 (1979) 827–852.
- [13] N. Ikeda and S. Manabe, Van Vleck-Pauli formula for Wiener integrals and Jacobi fields, in "Itô's stochastic calculus and probability theory", pp. 141–156, Springer, Tokyo, 1996.
- [14] N. Ikeda and S. Taniguchi, Quadratic Wiener functionals, Kalman-Bucy filters, and the KdV equation, in: H. Kunita, S. Watanabe, Y. Takahashi (Eds.) Stochastic Analysis and Related Topics in Kyoto, In honor of Kiyosi Itô, Adv. Studies Pure Math. 41, Math. Soc. Japan, Tokyo, 2004, pp. 167–187.
- [15] N. Ikeda and S. Watanabe, Stochastic differential equations and diffusion processes, Second Edition, North-Holland/Kodansha, 1989.
- [16] K. Itô, Differential equations determining a Markov process (in Japanese), Jour. Pan-Japan Math. Coll. 244 (1942), 1352–1400 (English translation: [21] pp. 42– 75).
- [17] K. Itô, Foundation of Probability Theory (in Japanese), Iwanami, 1943. (The reference Books[a] in the Bibliography of Kiyosi Itô, Kiyosi Itô Selected Papers [21])
- [18] K. Itô, A Review on the Japanese translation of "Kurt Gödel: The Consistency Continuum Hypothesis" (in Japanese), Sugaku 1 (1947) 47–48.
- [19] K. Itô, Multiple Wiener integral, Jour. Math. Soc. Japan 3 (1951) 157–169
- [20] K. Itô, Probability Theory (in Japanese), Iwanami, 1953. (The reference Books[b] in the Bibliography of Kiyosi Itô, Kiyosi Itô Selected Papers [21])
- [21] K. Itô, Kiyosi Itô Selected Papers, D. W. Stroock and S. R. S. Varadhan eds, Springer, 1986.
- [22] K. Itô and M. Nisio, On stationary solutions of a stochastic differential equation, Jour. Math. Kyoto Univ. 4 (1964), 1–75.

- [23] K. Itô and M. Nisio, On the convergence of sums of independent Banach space valued random variables, Osaka Jour. Math. 5 (1968) 35–48.
- [24] M. Kac, Integration in function space and some of its applications, Lezioni Fermiane, Accademia Nazionale del Lincei Scuola Norm. Sup. Pisa, 1980.
- [25] A. N. Kolmogorov, Über die analytischen Methoden in der Wahrscheinlichkeitsrechnung, Math. Ann. 104 (1931) 415–458.
- [26] P. Lévy, Théorie de l'addition des variables aléatoires, Gauthier-Villars, 1937.
- [27] P. Lévy, Le mouvement brownien plan, Amer. J. Math. 62 (1940) 487–550.
- [28] P. Lévy, Wiener's random function, and other Laplacian random functions, Proc. 2nd Berkeley Symp. Math. Statistics Probab. 1950, pp. 171–187, University of California Press, 1951.
- [29] T. Lyons and Z. Qian, System control and rough paths, Oxford Math. Monographs, Oxford University Press, 2002.
- [30] G. Maruyama, On the transition probability functions of the Markov process, Nat. Sci. Rep. Ochanomizu Univ. 5 (1954) 10–20.
- [31] H. Matsumoto, Semiclassical asymptotics of eigenvalues for Schrödinger operators with magnetic fields, Jour. Funct. Anal. 129 (1995) 168–190.
- [32] H. Matsumoto and S. Taniguchi, Wiener functionals of second order and their Lévy measures, Elect. Jour. Probab. 7 No.14 (2002) 1–30.
- [33] T. Miwa, M. Jimbo and E. Date, Solitons, Cambridge Univ. Press, Cambridge, 2000.
- [34] R. E. A. C. Paley and N. Wiener, Fourier Transforms in the Complex Domain. Amer. Math. Soc. Colloq. Publ. 19. Providence, R.I., AMS, 1934.
- [35] J. Perrin, Les Atomes, Félix Alcan, 1913.
- [36] M. Pinsky, Introduction to Fourier analysis and wavelets, Brooks/Cole, Pacific Grove, 2002.
- [37] Y. Prohorov, Convergence of random processes and limit theorems in probability theory, Theor. Prob. Appl. 1 (1956), 157–214.
- [38] S. Taniguchi, Brownian sheet and reflectionless potentials, Stoch. Proc. Appl. 116 (2006) 293–309
- [39] J. H. Van Vleck, The correspondence principle in the statistical interpretation of quantum mechanics, Proc. Nat. Acad. Sci. U.S.A. 14 (1928), 178–188.
- [40] N. Wiener, Differential space, Math. Phys. 2 (1923) 131-174.
- [41] N. Wiener, Un problème de probabilités dénombrables, Bull. Soc. Math. France 52 (1924) 569–578.

- [42] N. Wiener, Collected works with commentaries / Norbert Wiener ; vol. 1, P. Masani ed., MIT Press, 1976.
- [43] N. Wiener, I am a mathematician, The later life of a prodigy, Doubleday & Co., 1956
- [44] M. Yor, Remarques sur une formule de Paul Lévy, Séminaire Prob. XIV, Lect. Notes Math. 784 (1980) 343–346.

List of MI Preprint Series, Kyushu University

The Global COE Program Math-for-Industry Education & Research Hub

\mathbf{MI}

MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI Abstract collision systems simulated by cellular automata

MI2008-2 Eiji ONODERA

The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds

- MI2008-3 Hiroaki KIDO On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU Numerical computations of cavity flow problems by a pressure stabilized characteristiccurve finite element scheme
- MI2008-5 Yoshiyasu OZEKI Torsion points of abelian varieties with values in nfinite extensions over a padic field
- MI2008-6 Yoshiyuki TOMIYAMA Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA Alpha-determinant cyclic modules and Jacobi polynomials

- MI2008-10 Sangyeol LEE & Hiroki MASUDA Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials
- MI2008-14 Takashi NAKAMURA Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCHI & Yuichiro TAGUCHII Flat modules and Groebner bases over truncated discrete valuation rings

- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA Hypergeometric -functions of the q-Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA On Gaussian decay estimates of solutions to some linear elliptic equations and its applications
- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain

- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection

- MI2009-26 Manabu YOSHIDA Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance
- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI Finite element computation for scattering problems of micro-hologram using DtN map

- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIKE & Sadanori KONISHI Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons