

Gender Disparities in Quality of Life and Psychological Disturbance in Patients With Implantable Cardioverter-Defibrillators

ラハマワティ, アニタ

<https://hdl.handle.net/2324/1654708>

出版情報：九州大学, 2015, 博士（医学）, 課程博士
バージョン：
権利関係：やむを得ない事由により本文ファイル非公開（2）



氏 名 : Anita Rahmawati
ア ニ タ ラ ハ マ ワ テ ィ

論 文 名 : Gender Disparities in Quality of Life and Psychological Disturbance in Patients
With Implantable Cardioverter-Defibrillators

(植込み型除細動器(ICD)患者の QOL と心理的反応における性差)

区 分 : 甲

論 文 内 容 の 要 旨

Background: Implantable cardioverter-defibrillator (ICD) has improved prognosis in fatal arrhythmia and the number of ICD implantations has increased. ICD-related psychological problems and impaired quality of life (QOL), however, have been observed. This study examined whether gender differences exist in QOL and psychological disturbances in ICD patients.

Methods and Results: Consecutive outpatients (n=179; mean age, 60.5±15.9 years; 81% male) with ICD implantations completed questionnaires consisting of the Short Form-8 (SF-8), Beck Depression Inventory, Impact of Event Scale-Revised (IES-R), State-Trait Anxiety Inventory, and Worries about ICD. One-way multivariate analysis of variance (MANOVA) showed women to have impaired QOL on the role physical functioning (F15,157=4.57, P<0.05) and bodily pain (F15,157=5.26, P<0.05) subscales of the SF-8. More women reported depression (F15,157=5.37, P<0.05) and worry about ICD than men (F15,157=6.62, P<0.05). Moreover, women also had higher IES-R scores indicating post-traumatic stress disorder (PTSD) than men (F15,157=5.87, P<0.05).

Conclusions: Women reported poorer QOL on 2 subscales: role physical functioning and bodily pain. There was a significant relationship between gender and depression, worry about ICD, and PTSD, but not for anxiety. Female patients need more psychological interventions following ICD implantation.