九州大学学術情報リポジトリ
Kyushu University Institutional Repository

Asymptotic profiles for the compressible Navier-Stokes equations on the whole space

Kagei, Yoshiyuki
Faculty of Mathematics, Kyushu University

Okita, Masatoshi
National Institute of Technology, Kurume Colloge

https://hdl.handle.net/2324/1655023

出版情報：MI Preprint Series. 2016-2, 2016-04-04. 九州大学大学院数理学研究院
バージョン：
権利関係：

KYUSHU UNIVERSITY

MI Preprint Series

Mathematics for Industry
Kyushu University

Asymptotic profiles for the compressible Navier-Stokes equations on the whole space

Yoshiyuki Kagei
& Masatoshi Okita

MI 2016-2

(Received April 4, 2016)

Institute of Mathematics for Industry
Graduate School of Mathematics
Kyushu University
Fukuoka, JAPAN

Asymptotic profiles for the compressible Navier-Stokes equations on the whole space

Yoshiyuki Kagei[1] and Masatoshi Okita[2]

[1] Faculty of Mathematics,
Kyushu University,
Nishi-ku, Motooka 744,
Fukuoka 819-0395, Japan

[2] National Institute of Technology,
Kurume Colloge,
Kurume-shi, Komorino 1-1-1,
Fukuoka 830-8555, Japan

Abstract: This paper is concerned with large time behavior of the strong solutions of the compressible Navier-Stokes equation in the whole space around the motionless state. It was shown by Kawashima-Matsumura-Nishida (1979) and Hoff-Zumbrun (1995) that the perturbation of the motionless state is time-asymptotic to the solution of the linearized problem. In this paper we will give the second-order asymptotics of strong solutions.

Key Words: compressible Navier-Stokes equations; asymptotic profiles ; convergence rate.

2010 Mathematics Subject Classification Numbers. 35Q30, 76N15.

1 Introduction

This paper studies the initial value problem for the compressible Navier-Stokes equation in $\mathbb{R}^n$, $n \geq 3$:

$$\begin{cases} \partial_t \rho + \operatorname{div} m = 0, \\ \partial_t m + \operatorname{div}\left(\frac{m \bigotimes m}{\rho}\right) + \nabla P(\rho) = \mu \Delta\left(\frac{m}{\rho}\right) + (\mu + \mu')\nabla \operatorname{div}\left(\frac{m}{\rho}\right), \\ (\rho, m)(0, x) = (\rho_0, m_0)(x). \end{cases} \tag{1}$$

Here $t > 0$, $x = {}^T(x_1, x_2, \cdots, x_n) \in \mathbb{R}^n$, and the superscript ${}^T\cdot$ stands for the transposition; the unknown functions $\rho = \rho(t, x) > 0$ and $m = m(t, x) = {}^T(m_1(t, x), m_2(t, x), \cdots, m_n(t, x))$ denote the density and momentum, respectively; $P = P(\rho)$ is the pressure that is assumed to be a function of the density ρ; μ and μ' are the viscosity coefficients satisfying the conditions $\mu > 0$ and $\frac{2}{n}\mu + \mu' \geq 0$; and div , ∇ and Δ denote the usual divergence, gradient and Laplacian with respect to x, respectively. The notation $\operatorname{div}\left(\frac{m \bigotimes m}{\rho}\right)$ means that its j-th component is given by $\operatorname{div}\left(\frac{m_j m}{\rho}\right)$.

We assume that $P(\rho)$ is smooth in a neighborhood of $\bar{\rho}$ with $P'(\bar{\rho}) > 0$, where $\bar{\rho}$ is a given positive constant.

In this paper we investigate asymptotic properties of strong solutions of problem (1) around the constant stationary solution $(\bar{\rho}, 0)$.

Matsumura and Nishida [6] showed the global in time existence of the solution of (1) for $n = 3$, provided that the initial perturbation $u_0 = {}^T(\gamma(\rho_0 - \bar{\rho}), {}^T m_0)$, with $\gamma = \sqrt{P'(\bar{\rho})}$, is sufficiently small in $H^3(\mathbb{R}^3) \cap L^1(\mathbb{R}^3)$. Furthermore, the following decay estimates were obtained in [6]

$$\|\nabla^k u(t)\|_{L^2} \leq C(1 + t)^{-\frac{3}{4} - \frac{k}{2}}, \quad k = 0, 1, \tag{2}$$

where $u(t) = {}^T(\gamma(\rho - \bar{\rho}), {}^T m)$. (See also [7].) Kawashima, Matsumura and Nishida [4] proved that the solution is time asymptotic to the one of the linearized problem. It was shown in [4] that

$$\|u(t) - G(t) * u_0\|_{L^2} \leq CL(t)(1+t)^{-\frac{3}{4}-\frac{1}{2}},$$

if u_0 is sufficiently small in $H^3(\mathbb{R}^3) \cap L^1(\mathbb{R}^n)$ for $n = 3$. Here $G(t) * u_0 = G(t, \cdot) * u_0$, where $G(t, x)$ is Green's matrix for the linearized system at ${}^T(\bar{\rho}, 0)$ for (1) and $*$ denotes the convolution with respect to x. Hoff and Zumbrun [3] derived a more detailed description of the large-time behavior of $u(t)$ in L^p for all $1 \leq p \leq \infty$; the following estimates were established:

$$\|u(t)\|_{L^p} \leq C \begin{cases} (1+t)^{-\frac{n}{2}(1-\frac{1}{p})}, & 2 \leq p \leq \infty, \\ (1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{n-1}{4}(1-\frac{2}{p})} L(t), & 1 \leq p < 2, \end{cases}$$

and

$$\|u(t) - G(t) * u_0\|_{L^p} \leq CL(t) \begin{cases} (1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{1}{2}}, & 2 \leq p \leq \infty, \\ (1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{n}{4}(1-\frac{2}{p})-\frac{1}{2}+\eta}, & 1 \leq p < 2, \end{cases}$$

where η is any positive number. Moreover, it was proved in [3] that $u(t)$ is time-asymptotic to the solution $\tilde{u}(t) = {}^T(\tilde{\sigma}, {}^T\tilde{m})$ of the following linear effective artificial viscosity system:

$$\begin{cases} \partial_t \tilde{\sigma} + \gamma \operatorname{div} \tilde{m} - \frac{1}{2}(\mu_2 + \mu_1)\Delta\tilde{\sigma} = 0, \\ \partial_t \tilde{m} - \mu_1 \Delta \tilde{m} - \frac{1}{2}(\mu_2 - \mu_1)\nabla \operatorname{div} \tilde{m} + \gamma \nabla \tilde{\sigma} = 0, \end{cases} \tag{3}$$

where $\mu_1 = \frac{\mu}{\bar{\rho}}$ and $\mu_2 = \frac{\mu+\mu'}{\bar{\rho}}$. More precisely,

$$\begin{aligned} & \|u(t) - \tilde{G}(t) * u_0\|_{L^p}, \ \|u(t) - \tilde{G}(t, \cdot) \int u_0 dx\|_{L^p} \\ \leq \quad & CL(t) \begin{cases} (1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{1}{2}}, & 2 \leq p \leq \infty, \\ (1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{n}{4}(1-\frac{2}{p})-\frac{1}{2}+\eta}, & 1 \leq p < 2, \end{cases} \end{aligned}$$

for any positive constant η, where $\tilde{G}$ is Green's matrix for (3). Here $L(t) = \log(1+t)$ when $n = 2$, and is otherwise identically one. We also mention that Kobayashi and Shibata [5] proved the following estimates

$$\|\partial_t^j \partial_x^\alpha G_1(t, x)\|_{L^p(\mathbb{R}^n_x)} \leq C(1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{j+|\alpha|}{2}}$$

for $2 \leq p \leq \infty$, and

$$\|\partial_t^j \partial_x^\alpha G_1(t, x)\|_{L^p(\mathbb{R}^n_x)} \leq C \begin{cases} (1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{n-1}{4}(1-\frac{2}{p})-\frac{j+|\alpha|}{2}}, & n \geq 3 \text{ and } n\text{: odd}, \\ (1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{n}{4}(1-\frac{2}{p})-\frac{j+|\alpha|}{2}}, & n \geq 2 \text{ and } n\text{: even}, \end{cases}$$

for $1 \leq p \leq \infty$, any positive integer j and any multi-index α. Here G_1 is a low frequency part of G, i.e., $G_1 = \Phi * G$ with $\mathfrak{F}[\Phi] \in C^\infty$, $\operatorname{Supp} \mathfrak{F}[\Phi] \subset \{\xi \in \mathbb{R}^n | |\xi| \leq \frac{A}{\sqrt{2}}\}$ and $\mathfrak{F}[\Phi](\xi) = 1$ for $|\xi| \leq \frac{A}{2}$. Here, $\mathfrak{F}[\Phi]$ denotes the Fourier transform of Φ, $A = \frac{2\gamma}{\mu_1+\mu_2}$ with $\mu_1 = \frac{\mu}{\bar{\rho}}$ and $\mu_2 = \frac{\mu+\mu'}{\bar{\rho}}$.

In this paper we give the second order term in the asymptotic expansion of $u(t)$ as $t \to \infty$. The main result of this paper is stated as follows

Theorem 1.1. *Assume that $n \geq 3$. Then there exists $\epsilon > 0$ such that if*

$$u_0 \in H^{s_0+1} \cap L^1_l \cap L^2_l$$

with $s_0 = [\frac{n}{2}] + 1$ and

$$\|u_0\|_{H^{s_0+1} \cap L^1} \leq \epsilon,$$

for $l=\frac{1}{2}$ when $n=3$, and $l=1$ when $n\geq 4$, then

$$\begin{aligned}
&\big\|u(t)-G(t)*u_0-\sum_{i=1}^n \partial_i G_1(t.\cdot)\int_0^\infty\int_{\mathbb{R}^n}\mathcal{F}_i^0 dyds\big\|_{L^2},\\
&\big\|u(t)-G(t)*u_0-\sum_{i=1}^n \partial_i \tilde{G}(t.\cdot)\int_0^\infty\int_{\mathbb{R}^n}\mathcal{F}_i^0 dyds\big\|_{L^2}\\
\leq\ & C(1+t)^{-\frac{n}{4}-\frac{3}{4}}\begin{cases}\log(1+t), & n=3,\\ (1+t)^{-\frac{1}{4}}\log(1+t), & n=4,\\ (1+t)^{-\frac{1}{4}}, & n\geq 5,\end{cases}
\end{aligned}$$

for $t\geq 0$, where

$$\mathcal{F}_i^0=\begin{pmatrix}0\\ F_i^0\end{pmatrix}$$

with $F_i^0=\left(F_{ij}^0\right)_{i,j=1}^n$ being the matrix defined by

$$\begin{aligned}
F_{ij}^0 &= -\delta_{ji}\{\frac{1}{2}(\rho-\bar{\rho})^2 P_{\rho\rho}(\bar{\rho})\}-\frac{m_j m_i}{\bar{\rho}}\\
&\quad +\delta_{ji}\{\frac{1}{2\gamma^3}\sigma^3\int_0^1(1-\theta)^2\partial_\rho^3 P(\frac{1}{\gamma}\sigma\theta+\bar{\rho})d\theta\}+\frac{\sigma m_j m_i}{\bar{\rho}(\sigma+\gamma\bar{\rho})}.
\end{aligned}$$

Remark 1.2. *Let $n\geq 3$. One can also obtain*

$$\big\|u(t)-G(t)*u_0-\sum_{i=1}^n \partial_i G_1(t,\cdot)\int_0^\infty\int_{\mathbb{R}^n}\mathcal{F}_i^0 dyds\big\|_{L^p}\leq CK(t)(1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{3}{4}}$$

for $2\leq p\leq\infty$, and

$$\begin{aligned}
&\big\|u(t)-G(t)*u_0-\sum_{i=1}^n \partial_i G_1(t,\cdot)\int_0^\infty\int_{\mathbb{R}^n}\mathcal{F}_i^0 dyds\big\|_{L^p}\\
\leq\ & CK(t)\begin{cases}(1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{n-1}{4}(1-\frac{2}{p})-\frac{3}{4}}, & n\geq 3 \text{ and } n\text{: odd},\\ (1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{n}{4}(1-\frac{2}{p})-\frac{3}{4}}, & n\geq 2 \text{ and } n\text{: even}\end{cases}
\end{aligned}$$

for $1\leq p<2$, and initial perturbation u_0 sufficiently small in some Sobolev spaces and weighted Lebesgue space. Here $K(t)=\begin{cases}\log(1+t), & n=3,\\ (1+t)^{-\frac{1}{4}}\log(1+t), & n=4,\\ (1+t)^{-\frac{1}{4}}, & n\geq 5.\end{cases}$

Remark 1.3. *Asymptotic expansion of the solution of the linearized equation is given as follows:*

$$\|G(t)*u_0-G_1(t,\cdot)\int u_0dy+\sum_{|\alpha|=1}\partial_x^\alpha G_1(t,\cdot)\int y^\alpha u_0dy\|_{L^2}\leq C(1+t)^{-\frac{n}{4}-1},$$

for $t\geq 0$ and $u_0\in L_2^1\cap L^2$. Hence we have

$$\begin{aligned}
&\big\|u(t)-G_1(t,\cdot)\int u_0dy+\sum_{|\alpha|=1}\partial_x^\alpha G_1(t,\cdot)\int y^\alpha u_0dy-\sum_{i=1}^n\partial_i G_1(t.\cdot)\int_0^\infty\int_{\mathbb{R}^n}\mathcal{F}_i^0 dyds\big\|_{L^2}\\
\leq\ & C(1+t)^{-\frac{n}{4}-\frac{3}{4}}\begin{cases}\log(1+t), & n=3,\\ (1+t)^{-\frac{1}{4}}\log(1+t), & n=4,\\ (1+t)^{-\frac{1}{4}}, & n\geq 5,\end{cases}
\end{aligned}$$

for $t\geq 0$ and sufficiently small $u_0\in H^{s_0+1}\cap L_2^1\cap L_l^2$.

This paper is organized as follows. In section 2 we introduce notation and an auxiliary lemma. In section 3 we reformulate the problem; we rewrite the equation into a symmetric one and formulate the problem for a system of low and high frequency parts. Section 4 is devoted to the proof of the main theorem. In section 5 we establish a decay estimate of L^2-moments.

2 Preliminaries

In this section we first introduce the notation which will be used throughout this paper. We then introduce some auxiliary lemmas which will be useful in the proof of the main result.

Let $L^p(1 \le p \le \infty)$ denote the usual L^p-Lebesgue space on $\mathbb{R}^n$. For a nonnegative integer m, we denote by H^m the usual L^2-Sobolev space of order m. For $1 \le p \le \infty$ and $0 \le l$, $L^p\big((1+|x|^l)^p dx\big)$ stands for the weighted L^p spaces over $\mathbb{R}^n$ defined by

$$L^p_l := L^p\big((1+|x|^l)^p dx\big) = \{u : \int_{\mathbb{R}^n} |u(x)|^p (1+|x|^l)^p dx < \infty\}.$$

The inner-product of L^2 is denoted by $(\cdot,\cdot)$.

For any integer $k \ge 0, \nabla^k f$ denotes all of k-th derivatives of f.

For a function f, we denote its Fourier transform by $\mathfrak{F}[f] = \hat{f}$:

$$\mathfrak{F}[f](\xi) = \hat{f}(\xi) = \int_{\mathbb{R}^n} f(x) e^{-ix\cdot\xi} dx \quad (\xi \in \mathbb{R}).$$

The inverse Fourier transform is denoted by $\mathfrak{F}^{-1}[f] = \check{f}$,

$$\mathfrak{F}^{-1}[f](x) = \check{f}(x) = (2\pi)^{-n} \int_{\mathbb{R}^n} f(\xi) e^{i\xi\cdot x} d\xi \quad (x \in \mathbb{R}).$$

We close this section with the Gagliardo-Nirenberg-Sobolev inquality.

Lemma 2.1. *Assume that $u(x) \in L^q(\mathbb{R}^n)$, $\nabla^m u(x) \in L^r(\mathbb{R}^n)$ with $1 \le q, r \le \infty$. Then for each integer $j \in [0, m]$, we have*

$$\|\nabla^j u\|_{L^p} \le C \|\nabla^m u\|^{\alpha}_{L^r} \|u\|^{1-\alpha}_{L^q}.$$

Here

$$\frac{1}{p} = \frac{j}{n} + \alpha\big(\frac{1}{r} - \frac{m}{n}\big) + (1-\alpha)\frac{1}{q}, \qquad \frac{j}{m} \le \alpha \le 1.$$

See, e.g., [2], for the proof.

3 Reformulation of the problem

In this section we first rewrite system (1) into the one for the perturbation. We next state the existence of strong solutions. We then introduce some auxiliary lemmas which will be useful in the proof of the main result.

Let us rewrite the problem (1) in a symmetric form. We define μ_1, μ_2 and γ by

$$\mu_1 = \frac{\mu}{\bar{\rho}},\ \mu_2 = \frac{\mu + \mu'}{\bar{\rho}},\ \gamma = \sqrt{P'(\bar{\rho})}.$$

We introduce the new unknown functions

$$\sigma(t,x) = \gamma\big(\rho(t,x) - \bar{\rho}\big), \quad m(t,x) = m(t,x).$$

The initial value problem (1) is then reformulated as

$$\begin{cases} \partial_t \sigma + \gamma \mathrm{div}\, m = 0, \\ \partial_t m - \mu_1 \Delta m - \mu_2 \nabla \mathrm{div}\, m + \gamma \nabla \sigma = \mathrm{div}\, F(u, \partial_x u), \\ (\sigma, m)(0,x) = (\sigma_0, m_0)(x), \end{cases} \tag{4}$$

where, $u = \begin{pmatrix} \sigma \\ m \end{pmatrix}$, and $F(u, \partial_x u)$ is the $n \times n$ matrix $(F_{jk}(u, \partial_x u))$ defined by

$$\begin{aligned} F_{jk}(u, \partial_x u) \quad = \quad & -\delta_{jk} \frac{1}{2\gamma^2} \sigma^2 P_{\rho\rho}(\bar{\rho}) - \frac{m_j m_k}{\bar{\rho}} \\ & + \delta_{jk} \big\{ \frac{1}{2\gamma^3} \sigma^3 \int_0^1 (1-\theta)^2 \partial_\rho^3 P(\frac{1}{\gamma}\sigma\theta + \bar{\rho}) d\theta \big\} + \frac{\sigma m_j m_k}{\bar{\rho}(\sigma + \gamma\bar{\rho})} \\ & - \delta_{jk} \mu_2 \mathrm{div}\, \big(\frac{\sigma m}{\sigma + \gamma\bar{\rho}}\big) - \mu_1 \partial_{x_k} \big(\frac{\sigma m_j}{\sigma + \gamma\bar{\rho}}\big), \end{aligned}$$

and the j-th component of $\operatorname{div} F$ is given by $\sum_{k=1}^{n} \partial_{x_k} F_{jk}$.

The global existence of strong solutions was proved by Matsumura and Nishida [6] by a combination of the spectral analysis and the energy method for the density-velocity formulation for (4). We here restate the global existence in terms of the density and momentum.

Proposition 3.1 (Matsumura-Nishida [6])**.** *Let $n \geq 2$ and let $u_0 \in H^{s_0+1}$. There exist a positive constant ϵ_1 such that if*

$$\|u_0\|_{H^{s_0+1}} \leq \epsilon_1,$$

then problem (4) *has a unique global solution $u \in C([0,\infty); H^{s_0+1})$.*

Proposition 3.1 were proved for the case $n = 3$ in [6]. In a similar manner one can see that Proposition 3.1 holds for $n \geq 2$.

We set

$$A = \begin{pmatrix} 0 & -\gamma\nabla\cdot \\ -\gamma\nabla & \mu_1\Delta + \mu_2\nabla\nabla\cdot \end{pmatrix}.$$

By using operator A, problem (4) is written as

$$\partial_t u - Au = \operatorname{div}\mathcal{F}, \quad u|_{t=0} = u_0, \tag{5}$$

where

$$\mathcal{F} = \mathcal{F}(u, \partial_x u) = \begin{pmatrix} 0 \\ F(u, \partial_x u) \end{pmatrix}, \quad u_0 = \begin{pmatrix} \sigma_0 \\ m_0 \end{pmatrix}.$$

We introduce a semigroup generated by A. We set

$$E(t)u := \mathfrak{F}^{-1}[e^{\hat{A}(\xi)t}\hat{u}] = G(t) * u \quad \text{for } u \in L^2,$$

where

$$\hat{A}(\xi) = \begin{pmatrix} 0 & -i\gamma^T\xi \\ -i\gamma\xi & -\mu_1|\xi|^2 I_n - \mu_2\xi^T\xi \end{pmatrix} \quad \text{and} \quad G(t,x) = \mathfrak{F}^{-1}[e^{\hat{A}(\xi)t}](x).$$

Lemma 3.2. *(i) The set of all eigenvalues of $\hat{A}(\xi)$ consists of $\lambda_i(\xi)$ $(i = 1, 2, 3)$, where*

$$\begin{cases} \lambda_1(\xi) = -\frac{1}{2}(\mu_1+\mu_2)|\xi|^2 + i\gamma|\xi|\sqrt{1 - \frac{(\mu_1+\mu_2)^2}{4\gamma^2}|\xi|^2}, \\ \lambda_2(\xi) = -\frac{1}{2}(\mu_1+\mu_2)^2|\xi|^2 - i\gamma|\xi|\sqrt{1 - \frac{(\mu_1+\mu_2)}{4\gamma^2}|\xi|^2}, \\ \lambda_3(\xi) = -\mu_1|\xi|^2, \end{cases}$$

for all $\xi \in \mathbb{R}^n$.

(ii) $e^{t\hat{A}(\xi)}$ has the spectral resolution

$$e^{t\hat{A}(\xi)} = \sum_{j=1}^{3} e^{t\lambda_j(\xi)} P_j(\xi)$$

for all $|\xi| \neq \frac{2\gamma}{\sqrt{\mu_1+\mu_2}}$, where $P_j(\xi)$ is the eigenprojection for $\lambda_j(\xi)$.

For $|\xi| = \frac{2\gamma}{\sqrt{\mu_1+\mu_2}}$, we have $\lambda_1(\xi) = \lambda_2(\xi) = -\frac{\mu_1+\mu_2}{2}|\xi|^2$ and

$$e^{t\hat{A}(\xi)} = e^{t\lambda_1(\xi)}\big(I + t(\hat{A}(\xi) - \lambda_1 I)\big)P_1(\xi) + e^{t\lambda_3(\xi)}P_3(\xi).$$

We see from Lemma 3.2 that

$$\lambda_1(\xi) \sim -\frac{1}{2}(\mu_1+\mu_2)|\xi|^2 + i\gamma|\xi|, \quad \lambda_2(\xi) \sim -\frac{1}{2}(\mu_1+\mu_2)|\xi|^2 - i\gamma|\xi| \qquad (|\xi| \to 0)$$

and

$$\lambda_1(\xi) \sim -(\mu_1+\mu_2)|\xi|^2, \quad \lambda_2(\xi) \sim -\frac{\gamma^2}{\mu_1+\mu_2} \qquad (|\xi| \to \infty),$$

and hence, $G(t) * u$ has different characters in its low and high frequency parts.

We next decompose the solution u of (5) into its low and high frequency parts. Let $\hat{\Phi}$ be a function in $C^\infty(\mathbb{R}^n)$ such that $\hat{\Phi}(\xi) = \begin{cases} 1 & |\xi| \le \frac{A}{2}, \\ 0 & |\xi| \ge \frac{A}{\sqrt{2}}, \end{cases}$ where $A = \frac{2\gamma}{\sqrt{\mu_1+\mu_2}}$. We set

$$G_1(t,x) = \begin{pmatrix} L_{11}(t,x) & L_{12}(t,x) \\ L_{21}(t,x) & L_{22}(t,x) \end{pmatrix},$$

where

$$\begin{aligned}
L_{11}(t,x) &= \mathfrak{F}^{-1}\left[\frac{\lambda_1(\xi)e^{\lambda_2(\xi)t} - \lambda_2(\xi)e^{\lambda_1(\xi)t}}{\lambda_1(\xi)-\lambda_2(\xi)}\hat{\Phi}(\xi)\right](x), \\
L_{12}(t,x) &= -i\gamma\mathfrak{F}^{-1}\left[{}^T\xi\frac{e^{\lambda_1(\xi)t} - e^{\lambda_2(\xi)t}}{\lambda_1(\xi)-\lambda_2(\xi)}\hat{\Phi}(\xi)\right](x), \\
L_{21}(t,x) &= {}^TL_{12}(t,x), \\
L_{22}(t,x) &= K_1(t,x) + K_2(t,x) - K_3(t,x), \\
K_1(t,x) &= \mathfrak{F}^{-1}\left[e^{\lambda_3(\xi)t}\hat{\Phi}(\xi)\right](x)I, \quad I \text{ is unit matrix}, \\
K_2(t,x) &= \mathfrak{F}^{-1}\left[\frac{\lambda_1(\xi)e^{\lambda_1(\xi)t} - \lambda_2(\xi)e^{\lambda_2(\xi)t}}{\lambda_1(\xi)-\lambda_2(\xi)}\frac{\xi_j\xi_k}{|\xi|^2}\hat{\Phi}(\xi)\right](x), \\
K_3(t,x) &= \mathfrak{F}^{-1}\left[e^{\lambda_3(\xi)t}\frac{\xi_j\xi_k}{|\xi|^2}\hat{\Phi}(\xi)\right](x).
\end{aligned}$$

We define the operators S_1 and $\tilde{S}_\infty$ on L^2 by

$$S_1u := \mathfrak{F}^{-1}\left[\hat{\Phi}\hat{u}\right] \quad \tilde{S}_\infty u := \mathfrak{F}^{-1}\left[(1-\hat{\Phi})\hat{u}\right], \quad u \in L^2.$$

In terms of S_1 and $\tilde{S}_\infty$, we decompose the solution $u(t)$ of (5) as

$$u(t) = u_1(t) + u_\infty(t), \quad u_1(t) = S_1u(t),\ u_\infty(t) = \tilde{S}_\infty u(t).$$

It then follows that $u_1(t)$ and $u_\infty(t)$ are governed by systems (6) and (7) − (8) given in the following proposition.

Proposition 3.3. *Let $u = {}^T(\sigma, {}^Tm)$ be a solution of problem (5) on $[0,\infty)\times\mathbb{R}^n$. Then, $u_1(t)$ and $u_\infty(t)$ satisfy*

$$u_1(t) = G_1(t) * u_0 + \int_0^t G_1(t-s) * \operatorname{div}\mathcal{F}(u, \partial_x u)(s)ds \tag{6}$$

and

$$\partial_t u_\infty - Au_\infty = \tilde{S}_\infty \operatorname{div}\mathcal{F}(u, \partial_x u), \tag{7}$$

$$u_\infty|_{t=0} = u_{0\infty}, \tag{8}$$

where $u_{0\infty} = \tilde{S}_\infty u_0$.

See, e.g., [8], for the proof.

4 Proof of main result

In this section we prove Theorem 1.1. We have the following decay estimate for high frequency part.

Proposition 4.1 ([8]). *There exists $\epsilon > 0$ such that if*

$$u_0 \in H^{s_0} \cap L^1$$

and

$$\|u_0\|_{H^{s_0}\cap L^1} \le \epsilon,$$

then we have

$$\|\nabla^k u(t)\|_{L^2} \le C\epsilon(1+t)^{-\frac{n}{4}-\frac{k}{2}}$$

for $k = 1, 2, \cdots, s_0$, and

$$\|u_\infty(t)\|_{L^2} \le C\epsilon(1+t)^{-\frac{n}{4}-1}.$$

In view of Proposition 4.1 we expect that the large-time behavior of solution is described by the behavior of the low frequency part. Hence, we investigate asymptotic properties of the low frequency part of solution in the form of the integral equation (6). We first consider the linear term $G_1(t) * u_0$ in (6). We have the following estimates for $G_1(t) * u_0$.

Lemma 4.2 (Kobayashi-Shibata [5]). *Let $n \ge 2$ and let j and α be any positive integer and any multi-index, respectively. Then, for any $t \ge 0$, we have*

$$\|\partial_t^j \partial_x^\alpha G_1(t,x)\|_{L^p(\mathbb{R}^n_x)} \le C(1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{j+|\alpha|}{2}}$$

for $2 \le p \le \infty$, and

$$\|\partial_t^j \partial_x^\alpha G_1(t,x)\|_{L^p(\mathbb{R}^n_x)} \le C \begin{cases} (1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{n-1}{4}(1-\frac{2}{p})-\frac{j+|\alpha|}{2}}, & n \ge 3 \text{ and } n\text{: odd}, \\ (1+t)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{n}{4}(1-\frac{2}{p})-\frac{j+|\alpha|}{2}}, & n \ge 2 \text{ and } n\text{: even}, \end{cases}$$

for $1 \le p < 2$.

On the other hand, the high frequency part of $G(t) * u_0$ decays exponentially. In fact, we have the following estimate.

Proposition 4.3. *There exists constants $C, c_0 > 0$ independent of t such that*

$$\|G_1(t) * u_0 - G(t) * u_0\|_{L^2} \le Ce^{-c_0 t}\|u_0\|_{L^2}.$$

Proof. By Plancherel's theorem, we have

$$\begin{aligned} \|G_1(t) * u_0 - G(t) * u_0\|_{L^2} &\le C\Big(\int_{|\xi|\ge\frac{A}{2}} \big|e^{\hat{A}(\xi)t}\hat{u_0}\big|^2 d\xi\Big)^{\frac{1}{2}} \\ &\le Ce^{-c_0 t}\Big(\int_{\mathbb{R}^n} |\hat{u_0}(\xi)|^2 d\xi\Big)^{\frac{1}{2}} \\ &\le Ce^{-c_0 t}\|u_0\|_{L^2}. \end{aligned}$$

□

The following estimates show that $\tilde{G}(t) * u_0$ is time-asymptotic to $G_1(t) * u_0$.

Proposition 4.4 (Hoff-Zumbrun [3]). *Let $\tilde{G}$ be Green's matrix for the linear artificial viscosity system (3). Then for each nonnegative integer k, $u_0 \in L^1 \cap H^k$, and $|\alpha| \le k$, there is a positive constant C such that*

$$\|\partial_x^\alpha \tilde{G}(t) * u_0\|_{L^2} \le C(1+t)^{\frac{n}{4}-\frac{|\alpha|}{2}}\|u_0\|_{L^1\cap H^k},$$

$$\|\partial_x^\alpha [G_1(t) * u_0 - \tilde{G}(t) * u_0]\|_{L^2} \le C(1+t)^{-\frac{n}{4}-\frac{|\alpha|}{2}-\frac{1}{2}}\|u_0\|_{L^1\cap H^k}.$$

The linear term $G_1(t) * u_0$ has the following expansion.

Proposition 4.5. *Suppose $u_0 \in L^1_2$. Then we obtain*

$$\|G_1(t) * u_0 - G_1(t,\cdot)\int u_0(y)dy + \sum_{|\alpha|=1} \partial_x^\alpha G_1(t,\cdot)\int y^\alpha u_0 dy\|_{L^2} \le C(1+t)^{-\frac{n}{4}-1}$$

Proof. Applying Taylor's formula we have

$$\begin{aligned} G_1(t) * u_0 &= \int G_1(t, x-y)u_0(y)dy \\ &= G_1(t,x)\int u_0(y)dy - \sum_{|\alpha|=1} \partial_x^\alpha G_1(t,x)\int y^\alpha u_0(y)dy \\ &\quad + \sum_{|\alpha|=2}\int\int_0^1 \big(\partial_x^\alpha G\big)(x-\theta y)y^\alpha u_0(y)d\theta dy. \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} \|G_1(t) * u_0 - G_1(t,\cdot)\int u_0(y)dy + \sum_{|\alpha|=1} \partial_x^\alpha G_1(t,\cdot)\int y^\alpha u_0 dy\|_{L^2} &\le \|\nabla^2 G_1(t)\|_{L^2}\|u_0\|_{L^1_2} \\ &\le C(1+t)^{-\frac{n}{4}-1}\|u_0\|_{L^1_2}. \end{aligned}$$

□

We next give the asymptotic leading term for the nonlinearity of (6).

Theorem 4.6. *Assume that $n \ge 3$. Let $l = \frac{1}{2}$ when $n = 3$ or $l = 1$ when $n \ge 4$. There exists $\epsilon > 0$ such that if*

$$u_0 \in H^{s_0+1} \cap L^1_l \cap L^2_l$$

and

$$\|u_0\|_{H^{s_0+1}\cap L^1} \le \epsilon,$$

then we have

$$\begin{aligned} &\Big\|\int_0^t G_1(t-s) * \operatorname{div}\mathcal{F}(u,\partial_x u)(s)ds - \sum_{i=1}^n \partial_i G_1(t.\cdot)\int_0^\infty\int_{\mathbb{R}^n}\mathcal{F}_i^0 dyds\Big\|_{L^2} \\ \le\ & C(1+t)^{-\frac{n}{4}-\frac{3}{4}}\begin{cases} \log(1+t), & n=3, \\ (1+t)^{-\frac{1}{4}}\log(1+t), & n=4, \\ (1+t)^{-\frac{1}{4}}, & n\ge 5 \end{cases} \end{aligned}$$

for $t \ge 0$.

To prove Theorem 4.6 we introduce the following decay estimates of L^2-moments of solutions of problem (4).

Proposition 4.7. *Let $l = \frac{1}{2}$ or $l = 1$ for $n \ge 3$. There exists $\epsilon > 0$ such that if*

$$u_0 \in H^{s_0+1} \cap L^1_l \cap L^2_l$$

and

$$\|u_0\|_{H^{s_0}} \le \epsilon,$$

then the solution $u(t)$ of (4) satisfies

$$\big\||x|^l u(t)\big\|_{L^2} \le C(1+t)^{-\frac{n}{4}+l}.$$

We will give a proof of Proposition 4.7 in section 5.

Proof of Theorem 4.6. Direct calculation gives

$$\begin{aligned}
&\int_0^t G_1(t-s)*\operatorname{div}\mathcal{F}(u,\partial_x u)(s)ds-\sum_{i=1}^n \partial_i G_1(t.\cdot)\int_0^\infty\int_{\mathbb{R}^n}\mathcal{F}_i^0 dyds\\
=\;&\int_0^{\frac{t}{2}}\int\sum_{i=1}^n\big(\partial_{x_i}G_1(t-s,x-y)-\partial_{x_i}G_1(t,x)\big)\mathcal{F}_i^0 dyds\\
&+\int_0^{\frac{t}{2}}\int\sum_{i=1}^n\partial_{x_i}G_1(t-s,x-y)\mathcal{F}_i^1 dyds\\
&+\int_{\frac{t}{2}}^t G_1(t-s)*\operatorname{div}\mathcal{F}(u,\partial_x u)(s)ds+\sum_{i=1}^n\partial_{x_i}G_1(t,\cdot)\int_{\frac{t}{2}}^\infty\int\mathcal{F}_i^0 dyds\\
=:\;&J_1+J_2+J_3+J_4,
\end{aligned}$$

where $\mathcal{F}_i^1=\begin{pmatrix}0\\ F_i^1(u,\partial_x u)\end{pmatrix}$ with $F_i^1(u,\partial_x u)=\big(F_{ij}^1(u,\partial_x u)\big)_j^n$, $F_{ij}^1(u,\partial_x u)=-\delta_{ji}\mu_2\operatorname{div}\big(\frac{\sigma m}{\sigma+\gamma\bar\rho}\big)-\mu_1\partial_{x_i}\big(\frac{\sigma m_j}{\sigma+\gamma\bar\rho}\big)$.

We first estimate the L^2 norms of J_3 and J_4:

$$\begin{aligned}
\|J_3\|_{L^2}=\Big\|\int_{\frac{t}{2}}^t G_1(t-s)*\operatorname{div}\mathcal{F}ds\Big\|_{L^2}&\le C\int_{\frac{t}{2}}^t\|u\|_{L^\infty}\|\nabla u\|_{H^1}\\
&\le C\int_{\frac{t}{2}}^t(1+s)^{-\frac{3n}{4}-\frac{1}{2}}\le C(1+t)^{-\frac{n}{4}-1},
\end{aligned}$$

$$\begin{aligned}
\|J_4\|_{L^2}=\Big\|\sum_{i=1}^n\partial_{x_i}G_1(t,\cdot)\Big\|_{L^2}\int_{\frac{t}{2}}^\infty\int\mathcal{F}_i^0 dyds&\le C(1+t)^{-\frac{n}{4}-\frac{1}{2}}\int_{\frac{t}{2}}^t\|u\|_{L^2}^2 ds\\
&\le C(1+t)^{-\frac{n}{4}-1}.
\end{aligned}$$

We next estimate J_2. We put $F_i^2(u,\partial_x u)=\big(F_{i,jk}^2(u,\partial_x u)\big)_{j,k=1}^n$ with $F_{i,jk}^2(u,\partial_x u)=-\delta_{ji}\mu_2\big(\frac{\sigma m_k}{\sigma+\gamma\bar\rho}\big)$ for $1\le i\le n$ and $F^3(u,\partial_x u)=\big(F_j^3(u,\partial_x u)\big)_{j=1}^n$ with $F_j^3(u,\partial_x u)=-\mu_1\big(\frac{\sigma m_j}{\sigma+\gamma\bar\rho}\big)$; and we set $\mathcal{F}_i^2=\begin{pmatrix}0\\ F_i^2(u,\partial_x u)\end{pmatrix}$ and $\mathcal{F}^3=\begin{pmatrix}0\\ F^3(u,\partial_x u)\end{pmatrix}$. By using $\mathcal{F}_i^2$ and $\mathcal{F}^3$, we can write $\mathcal{F}_i^1=\sum_{k=1}^n\partial_{x_k}\mathcal{F}_{i,k}^2+\partial_{x_i}\mathcal{F}^3$, where $\mathcal{F}_{i,k}^2=\begin{pmatrix}0\\ F_{ik}^2(u,\partial_x u)\end{pmatrix}$ with $F_{ik}^2(u,\partial_x u)=\big(F_{i,jk}^2(u,\partial_x u)\big)_{j=1}^n$. Hence we obtain

$$\begin{aligned}
\|J_2\|_{L^2}&=\Big\|\int_0^{\frac{t}{2}}\int\sum_{i=1}^n\partial_{x_i}G_1(t-s,x-y)\mathcal{F}_i^1 dyds\Big\|_{L^2}\\
&=\Big\|\int_0^{\frac{t}{2}}\int\sum_{i=1}^n\partial_{x_i}G_1(t-s,x-y)\{\sum_{k=1}^n\partial_{x_k}\mathcal{F}_{i,k}^2+\partial_{x_i}\mathcal{F}^3\}dyds\Big\|_{L^2}\\
&=\Big\|\int_0^{\frac{t}{2}}\int\{\sum_{i=1}^n\sum_{k=1}^n\partial_{x_i}\partial_{x_k}G_1(t-s,x-y)\mathcal{F}_{i,k}^2\}\\
&\qquad+\{\sum_{i=1}^n\partial_{x_i}^2G_1(t-s,x-y)\mathcal{F}^3\}dyds\Big\|_{L^2}\\
&\le C\int_0^{\frac{t}{2}}\|\nabla^2G_1(t-s)\|_{L^2}\|u(s)\|_{L^2}^2 ds\\
&\le C\int_0^{\frac{t}{2}}(1+t-s)^{-\frac{n}{4}-1}(1+s)^{-\frac{n}{2}}ds\\
&\le C(1+t)^{-\frac{n}{4}-1}.
\end{aligned}$$

Let us next consider J_1. We decompose J_1 as

$$\begin{aligned} J_1 &= \int_0^{\frac{t}{2}} \int \sum_{i=1}^n \big(\partial_{x_i} G_1(t-s,x-y) - \partial_{x_i} G_1(t,x)\big)\mathcal{F}_i^4 dyds \\ &\quad + \int_0^{\frac{t}{2}} \int \sum_{i=1}^n \big(\partial_{x_i} G_1(t-s,x-y) - \partial_{x_i} G_1(t,x)\big)\mathcal{F}_i^5 dyds \\ &=: I_1 + I_2, \end{aligned}$$

where $\mathcal{F}_i^4 = \begin{pmatrix} 0 \\ F_i^4(u) \end{pmatrix}$, and $\mathcal{F}_i^5 = \begin{pmatrix} 0 \\ F_i^5(u) \end{pmatrix}$ with $F_i^4(u) = \big(F_{ij}^4(u)\big)_{j=1}^n$, $F_{ij}^4(u) = -\delta_{ji}\big\{\frac{1}{2\gamma^2}\sigma^2 P_{\rho\rho}(\bar{\rho})\big\} - \frac{m_j m_i}{\bar{\rho}}$, and $F_i^5(u) = \big(F_{ij}^5(u)\big)_{j=1}^n$ with $F_{ij}^5(u) = \delta_{ji}\big\{\frac{1}{2\gamma^3}\sigma^3 \int_0^1 (1-\theta)^2 \partial_\rho^3 P(\frac{1}{\gamma}\sigma\theta + \bar{\rho})d\theta\big\} + \frac{\sigma m_j m_i}{\bar{\rho}(\sigma+\gamma\bar{\rho})}$.

Moreover, we decompose I_1 as

$$\begin{aligned} I_1 &= \int_0^{\frac{t}{2}} \int \sum_{i=1}^n \big(\partial_{x_i} G_1(t-s,x-y) - \partial_{x_i} G_1(t,x)\big)\mathcal{F}_i^4 dyds \\ &= \int_0^{\frac{t}{2}} \int \sum_{i=1}^n \big(\partial_{x_i} G_1(t-s,x-y) - \partial_{x_i} G_1(t-s,x)\big)\mathcal{F}_i^4 dyds \\ &\quad + \int_0^{\frac{t}{2}} \int \sum_{i=1}^n \big(\partial_{x_i} G_1(t-s,x) - \partial_{x_i} G_1(t,x)\big)\mathcal{F}_i^4 dyds \\ &=: I_{11} + I_{12}. \end{aligned}$$

We first estimate I_{11} and I_{12} when $n = 3$. By using Lemma 2.1 and Lemma 4.2, we have

$$\begin{aligned} \|I_{11}\|_{L^2} &= \Big\| \int_0^{\frac{t}{2}} \int \sum_{i=1}^3 \big(\partial_{x_i} G_1(t-s,x-y) - \partial_{x_i} G_1(t-s,x)\big)\mathcal{F}_i^4(s,y) dyds \Big\|_{L^2} \\ &\le C \int_0^{\frac{t}{2}} \int \big\|\nabla\big\{G_1(t-s,x-y) - G_1(t-s,x)\big\}\big\|_{L^2} \sum_{i=1}^3 \big|\mathcal{F}_i^4(s,y)\big| dyds \\ &\le C \int_0^{\frac{t}{2}} \int \big\|G_1(t-s,x-y) - G_1(t-s,x)\big\|_{L^2}^{\frac{1}{2}} \big\|\nabla^2\big\{G_1(t-s,x-y) - G_1(t-s,x)\big\}\big\|_{L^2}^{\frac{1}{2}} \\ &\quad \times \sum_{i=1}^3 \big|\mathcal{F}_i^4(s,y)\big| dyds \\ &\le C \int_0^{\frac{t}{2}} \int \Big\| \int_0^1 \big|\nabla G_1(t-s,x-\tau y)\big| d\tau \Big\|_{L^2}^{\frac{1}{2}} \big\|\nabla^2\big\{G_1(t-s,x-y) - G_1(t-s,x)\big\}\big\|_{L^2}^{\frac{1}{2}} |y|^{\frac{1}{2}} \\ &\quad \times \sum_{i=1}^3 \big|\mathcal{F}_i^4(s,y)\big| dyds \\ &\le C \int_0^{\frac{t}{2}} (1+t-s)^{-\frac{1}{2}(\frac{3}{4}+\frac{1}{2})} (1+t-s)^{-\frac{1}{2}(\frac{3}{4}+1)} \sum_{i=1}^3 \big\||x|^{\frac{1}{2}} \mathcal{F}_i^4(s,x)\big\|_{L^1} ds \\ &\le C \int_0^{\frac{t}{2}} (1+t-s)^{-\frac{3}{2}} \||x|^{\frac{1}{2}} u\|_{L^2} \|u\|_{L^2} ds \\ &\le C(1+t)^{-\frac{3}{2}} \int_0^{\frac{t}{2}} (1+s)^{-1} ds \le C(1+t)^{-\frac{3}{2}} \log(1+t), \end{aligned}$$

and

$$
\begin{aligned}
\|I_{12}\|_{L^2} &= \Big\| \int_0^{\frac{t}{2}} \int \sum_{i=1}^{3} \big(\partial_{x_i} G_1(t-s,x) - \partial_{x_i} G_1(t,x)\big) \mathcal{F}_i^4(s,y) dyds \Big\|_{L^2} \\
&\le C \int_0^{\frac{t}{2}} \int \big\| G_1(t-s,x) - G_1(t,x) \big\|_{L^2}^{\frac{1}{2}} \big\| \nabla^2 \big\{ G_1(t-s,x) - G_1(t,x) \big\} \big\|_{L^2}^{\frac{1}{2}} \\
&\qquad \times \sum_{i=1}^{3} \big| \mathcal{F}_i^4(s,y) \big| dyds \\
&\le C \int_0^{\frac{t}{2}} \int \Big\| \int_0^1 \big| \partial_t G_1(t-\tau s,x) \big| d\tau \Big\|_{L^2}^{\frac{1}{2}} \\
&\quad \times \big\| \nabla^2 \big\{ G_1(t-s,x) - G_1(t,x) \big\} \big\|_{L^2}^{\frac{1}{2}} s^{\frac{1}{2}} \sum_{i=1}^{3} \big| \mathcal{F}_i^4(s,y) \big| dyds \\
&\le C(1+t)^{-\frac{3}{2}} \int_0^{\frac{t}{2}} s^{\frac{1}{2}} \|u\|_{L^2} \|u\|_{L^2} ds \\
&\le C(1+t)^{-\frac{3}{2}} \int_0^{\frac{t}{2}} (1+s)^{-1} ds \le C(1+t)^{-\frac{3}{2}} \log(1+t).
\end{aligned}
$$

Hence we obtain the estimate of $\|I_1\|_{L^2}$ for $n = 3$.

We next consider the case $n \ge 4$. We have

$$
\begin{aligned}
\|I_{11}\|_{L^2} &= \Big\| \int_0^{\frac{t}{2}} \int \sum_{i=1}^{n} \big(\partial_{x_i} G_1(t-s,x-y) - \partial_{x_i} G_1(t-s,x)\big) \mathcal{F}_i^4(s,y) dyds \Big\|_{L^2} \\
&= \Big\| \int_0^{\frac{t}{2}} \int \sum_{i=1}^{n} \int_0^1 \partial_{x_i} \nabla G_1(t-s,x-\tau y) d\tau y \mathcal{F}_i^4(s,y) dyds \Big\|_{L^2} \\
&\le C \int_0^{\frac{t}{2}} (1+t-s)^{-\frac{n}{4}-1} \| |x| u \|_{L^2} \|u\|_{L^2} ds \\
&\le C(1+t)^{-\frac{n}{4}-1} \int_0^{\frac{t}{2}} (1+s)^{-\frac{n}{2}+1} ds \\
&\le C(1+t)^{-\frac{n}{4}-1} \begin{cases} \log(1+t), & n = 4, \\ 1, & n \ge 5. \end{cases}
\end{aligned}
$$

Similarly, we obtain

$$
\begin{aligned}
\|I_{12}\|_{L^2} &= \Big\| \int_0^{\frac{t}{2}} \int \sum_{i=1}^{n} \big(\partial_{x_i} G_1(t-s,x) - \partial_{x_i} G_1(t,x)\big) \mathcal{F}_i^4(s,y) dyds \Big\|_{L^2} \\
&= \Big\| \int_0^{\frac{t}{2}} \int \sum_{i=1}^{n} \int_0^1 \partial_{x_i} \partial_t G_1(t-\tau s,x) d\tau \, s \mathcal{F}_i^4(s,y) dyds \Big\|_{L^2} \\
&\le C(1+t)^{-\frac{n}{4}-1} \int_0^{\frac{t}{2}} (1+s)^{-\frac{n}{2}+1} ds \\
&\le C(1+t)^{-\frac{n}{4}-1} \begin{cases} \log(1+t), & n = 4, \\ 1, & n \ge 5, \end{cases}
\end{aligned}
$$

and the estimate of I_1 is obtained.

As for I_2, we can estimate similarly to the proof of the estimate of I_1:

$$\begin{aligned}
\|I_2\|_{L^2} &\leq C\Big\| \int_0^{\frac{t}{2}} \int \sum_{i=1}^n \big(\partial_{x_i} G_1(t-s,x-y) - \partial_{x_i} G_1(t-s,x)\big)\mathcal{F}_i^5 dyds\Big\|_{L^2} \\
&\quad + C\Big\| \int_0^{\frac{t}{2}} \int \sum_{i=1}^n \big(\partial_{x_i} G_1(t-s,x) - \partial_{x_i} G_1(t,x)\big)\mathcal{F}_i^5 dyds\Big\|_{L^2} \\
&\leq C\int_0^{\frac{t}{2}} (1+t-s)^{-\frac{n}{4}-1} \||x|^{\frac{1}{2}} u\|_{L^2}^2 \|\sigma\|_{L^\infty} ds \\
&\quad + C\int_0^{\frac{t}{2}} (1+t-s)^{-\frac{n}{4}-1} s\|u\|_{L^2}^2 \|\sigma\|_{L^\infty} ds \\
&\leq C(1+t)^{-\frac{n}{4}-1} \int_0^{\frac{t}{2}} (1+s)^{-n+1} ds \leq C(1+t)^{-\frac{n}{4}-1}.
\end{aligned}$$

This completes the proof. □

Proof of Theorem 1.1. It follows from Proposition 4.1, Proposition 4.3 and Theorem 4.6 that

$$\begin{aligned}
&\Big\|u(t) - G(t) * u_0 - \sum_{i=1}^n \partial_i G_1(t,\cdot) \int_0^\infty \int_{\mathbb{R}^n} \mathcal{F}_i^0 dyds\Big\|_{L^2} \\
= &\Big\|u_1(t) + u_\infty(t) - G(t) * u_0 - \sum_{i=1}^n \partial_i G_1(t,\cdot) \int_0^\infty \int_{\mathbb{R}^n} \mathcal{F}_i^0 dyds\Big\|_{L^2} \\
\leq &\Big\|G_1(t) * u - G(t) * u_0\Big\|_{L^2} \\
&+\Big\| \int_0^t G_1(t-s) * \operatorname{div} \mathcal{F}(u,\partial_x u)(s) ds - \sum_{i=1}^n \partial_i G_1(t,\cdot) \int_0^\infty \int_{\mathbb{R}^n} \mathcal{F}_i^0 dyds\Big\|_{L^2} \\
&+\Big\|u_\infty(t)\Big\|_{L^2} \\
\leq &\ C(1+t)^{-\frac{n}{4}-\frac{3}{4}} \begin{cases} \log(1+t), & n=3, \\ (1+t)^{-\frac{1}{4}}\log(1+t), & n=4, \\ (1+t)^{-\frac{1}{4}}, & n\geq 5. \end{cases}
\end{aligned}$$

Moreover, in a similar manner to the proof of Theorem 4.6, we have

$$\begin{aligned}
&\Big\| \int_0^t G_1(t-s) * \operatorname{div} \mathcal{F}(u,\partial_x u)(s) ds - \sum_{i=1}^n \partial_i \tilde{G}(t,\cdot) \int_0^\infty \int_{\mathbb{R}^n} \mathcal{F}_i^0 dyds\Big\|_{L^2} \\
\leq &\Big\| \int_0^t G_1(t-s) * \operatorname{div} \mathcal{F}(u,\partial_x u)(s) ds - \int_0^t \tilde{G}(t-s) * \operatorname{div} \mathcal{F}(u,\partial_x u)(s) ds\Big\|_{L^2} \\
&+\Big\| \int_0^t \tilde{G}(t-s) * \operatorname{div} \mathcal{F}(u,\partial_x u)(s) ds - \sum_{i=1}^n \partial_i \tilde{G}(t,\cdot) \int_0^\infty \int_{\mathbb{R}^n} \mathcal{F}_i^0 dyds\Big\|_{L^2} \\
\leq &\ C(1+t)^{-\frac{n}{4}-\frac{3}{4}} \begin{cases} \log(1+t), & n=3, \\ (1+t)^{-\frac{1}{4}}\log(1+t), & n=4, \\ (1+t)^{-\frac{1}{4}}, & n\geq 5. \end{cases}
\end{aligned}$$

Here we used proposition 4.4. We thus obtain the desired decay estimate in Theorem 1.1. □

5 Proof of Proposition 4.7

In this section we will give a proof of Proposition 4.7. Let us reformulate the problem (1) by using the velocity. We set $v = \frac{m}{\rho}$ and $v_0 = \frac{m_0}{\rho_0}$. Then (1) is rewritten as

$$\begin{cases} \partial_t \rho + \mathrm{div}\,(\rho v) = 0, \\ \partial_t v + (v \cdot \nabla) v + \frac{\nabla P(\rho)}{\rho} = \frac{\mu}{\rho}\Delta v + \frac{\mu+\mu'}{\rho}\nabla(\mathrm{div}\, v), \\ (\rho, v)(0,x) = (\rho_0, v_0)(x). \end{cases} \tag{9}$$

Moreover we rewrite the system (9) into the one for perturbation. By using the new unknown function

$$\phi(t,x) = \frac{\rho(t,x) - \bar{\rho}}{\bar{\rho}}, \quad w(t,x) = \frac{1}{\gamma} u(t,x),$$

the initial value problem (9) is reformulated as

$$\begin{cases} \partial_t \phi + \gamma \mathrm{div}\, w = F_1(U), \\ \partial_t w - \mu_1 \triangle w - \mu_2 \nabla(\mathrm{div}\, w) + \gamma \nabla \phi = F_2(U), \\ (\phi, w)(0,x) = (\phi_0, w_0)(x), \end{cases} \tag{10}$$

where $U = \begin{pmatrix} \phi \\ w \end{pmatrix}$,

$$F_1(U) = -\gamma(w \cdot \nabla \phi + \phi \mathrm{div}\, w),$$

$$\begin{aligned} F_2(U) &= -\gamma(w \cdot \nabla) w - \mu_1 \frac{\phi}{\phi+1} \triangle w - \mu_2 \frac{\phi}{\phi+1} \nabla(\mathrm{div}\, w) \\ &\quad + \Big(\frac{\bar{\rho}\gamma}{\phi+1} - \frac{\bar{\rho}}{\gamma} \frac{\int_0^1 P''(s\bar{\rho}\phi + \bar{\rho}) ds}{\phi+1} \Big) \phi \nabla \phi. \end{aligned}$$

We have the following estimates for the solution U of (10) which were obtaind by Matsumura and Nishida [6, 7].

Proposition 5.1. *There exists a $\epsilon > 0$ such that if*

$$\|U_0\|_{H^{s_0+1}} \le \epsilon.$$

Then we obtain

$$\Big(\int_0^\infty \|\nabla U\|_{H^{s_0}}^2 ds \Big)^{\frac{1}{2}} + \Big(\int_0^\infty \|\nabla w\|_{H^{s_0+1}}^2 ds \Big)^{\frac{1}{2}} \le C\epsilon$$

See, for instance, [6, 7] for a proof of Proposition 5.1.

By using operator A, problem (10) is written as

$$\partial_t U - AU = F(U), \quad U|_{t=0} = U_0, \tag{11}$$

where

$$F(U) = \begin{pmatrix} F_1(U) \\ F_2(U) \end{pmatrix}, \quad U_0 = \begin{pmatrix} \phi_0 \\ w_0 \end{pmatrix}.$$

In terms of G, solution U is written as

$$\begin{aligned} U(t) &= G(t) * U_0 + \int_0^t G(t-s) * F(U) ds \\ &= G_1(t) * U_0 + G_\infty(t) * U_0 + \int_0^t G_1(t-s) * F(U) ds + \int_0^t G_\infty(t-s) * F(U) ds \end{aligned} \tag{12}$$

where $G_\infty(t)$ is the high frequency part of G, i.e., $G_\infty(t)$ is given by $G_\infty(t) = G(t) - G_1(t)$.

To prove Proposition 4.7, we introduce some notation. We set

$$M_l(t) := \sup_{0\le\tau\le t} (1+\tau)^{\frac{n}{4}-l} \big\| |x|^l U(\tau)\big\|_2.$$

Let us introduce the dyadic partition of unity. We take $\phi \in C_0^\infty$ satisfying

$$\mathrm{Supp}\,\phi \subset \{x \in \mathbb{R}^n | \frac{3}{4} \le |x| \le \frac{8}{3}\},$$

$$\sum_{j\in\mathbb{Z}} \phi(2^{-j}x) = 1 \quad \textit{for } x \in \mathbb{R}^n\backslash\{0\},$$

$$\mathrm{Supp}\,\phi(2^{-j}\cdot) \cap \mathrm{Supp}\,\phi(2^{-j'}\cdot) = \emptyset \quad \textit{for } |j-j'| \ge 2.$$

We set $h = \mathfrak{F}[\phi]$ and define the dyadic blocks by

$$\dot{\triangle}_j \hat{u} = 2^{jn} \int_{\mathbb{R}^n} h(2^j\eta)\hat{u}(\xi-\eta)d\eta \quad \text{if } j \in \mathbb{Z}.$$

The cut-off operator $\dot{S}_j$ is defined by

$$\dot{S}_j \hat{u} = \sum_{k\le j-1} \dot{\triangle}_k \hat{u}.$$

The paraproducts of $\hat{u}$ and $\hat{v}$ are defined by

$$\dot{T}_{\hat{u}}\hat{v} := \sum_j \dot{S}_{j-1}\hat{u}\dot{\triangle}_j\hat{v}, \qquad \dot{R}(\hat{u},\hat{v}) = \sum_{|k-j|\le 1} \dot{\triangle}_k\hat{u}\dot{\triangle}_j\hat{v}.$$

The Bony decomposition

$$\hat{u}\hat{v} = \dot{T}_{\hat{u}}\hat{v} + \dot{T}_{\hat{v}}\hat{u} + \dot{R}(\hat{u},\hat{v})$$

is well known. See, for instance, [1]

We have the following estimate for $|x|^{\frac{1}{2}} u * v$.

Lemma 5.2. *Let $\nabla^k \hat{u} \in L^\infty$ for $k = 0, 1,$ and $(1+|x|^{\frac{1}{2}})v \in L^2$. We have*

$$\big\||x|^{\frac{1}{2}} u * v\big\|_{L^2} \le C\big\{\|\nabla_\xi \hat{u}\|_{L^\infty} + \|\hat{u}\|_{L^\infty}\big\}\big\{\|v\|_{L^2} + \||x|^{\frac{1}{2}} v\|_{L^2}\big\}.$$

Proof. We obtain

$$\begin{aligned}
\big\||x|^{\frac{1}{2}} u * v\big\|_{L^2}^2 &= \big\|\sum_j \phi(2^{-j}x)|x|^{\frac{1}{2}} u * v\big\|_{L^2}^2 \\
&\le C\sum_j \{2^{\frac{1}{2}j}\big\|\phi(2^{-j}x)u * v\big\|_{L^2}\}^2 \\
&\le C\sum_j \{2^{\frac{1}{2}j}\big\|\dot{\triangle}_k \hat{u}\hat{v}\big\|_{L^2}\}^2 \\
&\le C\sum_j \{2^j\|\dot{\triangle}_j \dot{T}_{\hat{u}}\hat{v}\|_{L^2}^2 + 2^j\|\dot{\triangle}_j \dot{T}_{\hat{v}}\hat{u}\|_{L^2}^2 + 2^j\|\dot{\triangle}_j \dot{R}(\hat{u},\hat{v})\|_{L^2}^2\}.
\end{aligned}$$

Each term on the right-hand side is estimated as

$$\begin{aligned}
\sum_j 2^j\|\dot{\triangle}_k \dot{T}_{\hat{u}}\hat{v}\|_{L^2}^2 &= \sum_j 2^j\|\dot{\triangle}_j \sum_k \dot{S}_{k-1}\hat{u}\dot{\triangle}_k\hat{v}\|_{L^2}^2 \\
&= \sum_j 2^j\|\dot{\triangle}_j \sum_{|j-k|\le 5} \dot{S}_{k-1}\hat{u}\dot{\triangle}_k\hat{v}\|_{L^2}^2 \\
&\le C\sum_j 2^j \sum_{|j-k|\le 5} \|\dot{S}_{k-1}\hat{u}\|_{L^\infty}^2 \|\dot{\triangle}_k\hat{v}\|_{L^2}^2 \\
&\le C\|\hat{u}\|_{L^\infty}^2 \sum_j 2^j\|\dot{\triangle}_j\hat{v}\|_{L^2}^2 \\
&\le C\|\hat{u}\|_{L^\infty}^2 \||x|^{\frac{1}{2}} v\|_{L^2}^2,
\end{aligned}$$

$$
\begin{aligned}
\sum_j 2^j \|\dot{\triangle}_k \dot{T}_{\hat{v}} \hat{u}\|_{L^2}^2 &\le C \sum_j 2^j \sum_{|j-k|\le 5} \|\dot{S}_{k-1}\hat{v}\|_{L^2}^2 \|\dot{\triangle}_k \hat{u}\|_{L^\infty}^2 \\
&\le C\|v\|_{L^2}^2 \sum_j 2^j \|\triangle_j \hat{u}\|_{L^\infty}^2 \\
&\le C\|v\|_{L^2}^2 \big\{ \|\hat{u}\|_{L^\infty}^2 \sum_{j<0} 2^j + \sup_j (2^j \|\triangle_j \hat{u}\|_{L^\infty})^2 \sum_{j\ge 0} 2^{-j} \big\} \\
&\le \|v\|_{L^2}^2 \big\{ \|\nabla_\xi \hat{u}\|_{L^\infty}^2 + \|\hat{u}\|_{L^\infty}^2 \big\},
\end{aligned}
$$

and

$$
\begin{aligned}
\sum_j 2^j \|\dot{\triangle}_j \dot{R}(\hat{u},\hat{v})\|_{L^2}^2 &\le C \sum_j 2^j \sum_{k>j-2} \|\tilde{\triangle}_k \hat{u} \dot{\triangle}_k \hat{v}\|_{L^2}^2 \\
&\le C \sum_j \sum_{k>j-2} 2^{j-k} \|\tilde{\triangle}_k \hat{u}\|_{L^\infty}^2 2^k \|\dot{\triangle}_k \hat{v}\|_{L^2}^2 \\
&\le C\|\hat{u}\|_{L^\infty}^2 \sum_k 2^k \|\dot{\triangle}_k \hat{v}\|_{L^2}^2 \sum_{j<k+2} 2^{j-k} \\
&\le C\|\hat{u}\|_{L^\infty}^2 \||x|^{\frac{1}{2}} v\|_{L^2}^2,
\end{aligned}
$$

where $\tilde{\triangle}_k = \triangle_{k-1} + \triangle_k + \triangle_{k+1}$. This completes the proof. □

$G_1(t)$ and $G_\infty(t)$ satisfy the following estimate.

Lemma 5.3. *Let k be a nonnegative integer. Then there hold*

$$
\big\| |x|^l \nabla^k G_1(t) * U_0(t) \big\|_{L^2} \le C(1+t)^{-\frac{n}{4}-\frac{k}{2}+l} \|U_0\|_{L^1} + C(1+t)^{-\frac{n}{4}-\frac{k}{2}} \big\| |x|^l U_0 \big\|_{L^1},
$$

$$
\big\| |x|^l \nabla^k G_\infty(t) * U_0(t) \big\|_{L^2} \le C e^{-c_0 t} \big\{ \|\nabla^k U_0\|_{L^2} + \big\| |x|^l \nabla^k U_0 \big\|_{L^2} \big\}
$$

for $l = 0, \frac{1}{2}, 1$ and $t \ge 0$, where c_0 is a positive constant independent of t.

Proof. The proof for the case of $l = 0$ is well known. When $l = 1$, we have, for any α with $|\alpha| = k$,

$$
\begin{aligned}
\big\| |x| \partial_x^\alpha G_1(t) \big\|_{L^2} &\le C \Big\{ \int_{|\xi|\le 2A} \big| \nabla_\xi \big((i\xi)^\alpha e^{\hat{A}(\xi)t} \hat{\Phi} \big) \big|^2 d\xi \Big\}^{\frac{1}{2}} \\
&\le C \Big\{ \int_{|\xi|\le 2A} \big(t|\xi|^k e^{-\beta|\xi|^2 t} \big)^2 d\xi \Big\}^{\frac{1}{2}} + C \Big\{ \int_{|\xi|\le 2A} \big(|\xi|^{k-1} e^{-\beta|\xi|^2 t} \big)^2 d\xi \Big\}^{\frac{1}{2}} \\
&\quad + C \Big\{ \int_{|\xi|\le 2A} \big(|\xi|^k e^{-\beta|\xi|^2 t} \big)^2 d\xi \Big\}^{\frac{1}{2}} \\
&\le C(1+t)^{-\frac{n}{4}-\frac{k}{2}+1},
\end{aligned}
$$

where β is positive constant. When $l = \frac{1}{2}$ we obtain

$$
\begin{aligned}
\big\| |x|^{\frac{1}{2}} \nabla^k G_1(t) \big\|_{L^2} &\le C \big\| |x| \nabla^k G_1(t) \big\|_{L^2}^{\frac{1}{2}} \big\| \nabla^k G_1(t) \big\|_{L^2}^{\frac{1}{2}} \\
&\le C(1+t)^{-\frac{n}{4}-\frac{k}{2}+\frac{1}{2}}.
\end{aligned}
$$

Hence we have

$$
\begin{aligned}
\big\| |x|^l \nabla^k G_1(t) * U_0 \big\|_{L^2} &\le \big\| |x|^l \nabla^k G_1(t) \big\|_{L^2} \|U_0\|_{L^1} + \big\| \nabla^k G_1(t) \big\|_{L^2} \big\| |x|^l U_0 \big\|_{L^1} \\
&\le C(1+t)^{-\frac{n}{4}-\frac{k}{2}+l} \|U_0\|_{L^1} + C(1+t)^{-\frac{n}{4}-\frac{k}{2}} \big\| |x|^l U_0 \big\|_{L^1}
\end{aligned}
$$

for $l = \frac{1}{2}, 1$.

We next consider the estimate of the high frequency part. A direct computation gives the following estimates for $\hat{G}_\infty(t,\xi)$:

$$|\nabla_\xi^k \hat{G}_\infty(t,\xi)| \le C(k)\big(e^{-\theta t}|\xi|^{-k-1} + e^{-\theta|\xi|^2 t}|\xi|^{-k}\big), \tag{13}$$

where θ is a positive constant. By using Lemma 5.2 and (13), we have

$$\begin{aligned} & \big\||x|^l \nabla^k G_\infty(t) * U_0\big\|_{L^2} \\ \le\ & \big\{\|\nabla_\xi \hat{G}_\infty\|_{L^\infty} + \|\hat{G}_\infty\|_{L^\infty}\big\}\big\{\||x|^l U_0\|_{L^2}\|U_0\|_{L^2}\big\} \\ \le\ & Ce^{-c_0 t}\|\nabla^k U_0\|_{L^2} + Ce^{-c_0 t}\big\||x|^l \nabla^k U_0\big\|_{L^2} \end{aligned}$$

for $l = \frac{1}{2}$, 1, where c_0 is a positive constant. This completes the proof. □

We are in a position to complete the proof of Proposition 4.7.

Proposition 5.4. *Let $n \ge 3$ and let $l = \frac{1}{2}$ or $l = 1$. There exists $\epsilon > 0$ such that if*

$$U_0 \in H^{s_0+1} \cap L^1_l \cap L^2_l$$

and

$$\|U_0\|_{H^{s_0+1}\cap L^1} \le \epsilon,$$

then we have

$$M_l(t) \le C\|U_0\|_{L^1_l \cap L^2_l} + C\epsilon$$

for $t \in [0,\infty)$.

Proof. By Proposition 4.1, Proposition 4.3, Lemma 5.3 and (12), we see that

$$\begin{aligned} \big\||x|^l U(\tau)\big\|_{L^2} \le\ & C\big\||x|^l G_1(\tau) * U_0\big\|_{L^2} + C\big\||x|^l G_\infty(\tau) * U_0\big\|_{L^2} \\ & + C\int_0^\tau \big\||x|^l G_1(\tau-s) * F(U)\|_{L^2} ds + C\int_0^\tau \big\||x|^l G_\infty(\tau-s) * F(U)\|_{L^2} ds \\ \le\ & C(1+\tau)^{-\frac{n}{4}+l}\|U_0\|_{L^1} + C(1+\tau)^{-\frac{n}{4}}\big\||x|^l U_0\big\|_{L^1} + Ce^{-c_0\tau}\big\|(1+|x|^l)U_0\|_{L^2} \\ & + C\int_0^\tau \big\{(1+\tau-s)^{-\frac{n}{4}+l}\|F(U)\|_{L^1} + (1+\tau-s)^{-\frac{n}{4}}\big\||x|^l F(U)\big\|_{L^1}\big\} ds \\ & + C\int_0^\tau e^{-c_0(\tau-s)}\|(1+|x|^l)F(U)\|_{L^2} ds \\ \le\ & C(1+\tau)^{-\frac{n}{4}+l}\|U_0\|_{L^1_l\cap L^2_l} + C\int_0^\tau (1+\tau-s)^{-\frac{n}{4}+l}\|U\|_{L^2}\|\nabla U\|_{H^1} ds \\ & + C\int_0^\tau (1+\tau-s)^{-\frac{n}{4}}\big\||x|^l U\big\|_{L^2}\|\nabla U\|_{H^1} ds \\ & + C\int_0^\tau e^{-c_o(\tau-s)}\big\{\big\||x|^l U\big\|_{L^2} + \big\|U\big\|_{L^2}\big\}\big\{\|\nabla U\|_{H^{s_0}} + \|\nabla w\|_{H^{s_0+1}}\big\} ds \\ \le\ & C(1+\tau)^{-\frac{n}{4}+l}\|U_0\|_{L^1_l\cap L^2_l} + C\epsilon^2\int_0^\tau (1+\tau-s)^{-\frac{n}{4}+l}(1+s)^{-\frac{n}{2}-\frac{1}{2}} ds \\ & + C\epsilon M_l(t)\int_0^\tau (1+\tau-s)^{-\frac{n}{4}}(1+s)^{-\frac{n}{2}-\frac{1}{2}+l} ds \\ & + C\big\{M_l(t)+\epsilon\big\}\big\{\big(\int_0^\tau e^{-2c_0(\tau-s)}(1+s)^{-\frac{n}{2}+2l} ds\big)^{\frac{1}{2}}\big(\int_0^\tau \|\nabla U\|^2_{H^{s_0}} + \|\nabla w\|^2_{H^{s_0+1}} ds\big)^{\frac{1}{2}}\big\} \\ \le\ & C(1+\tau)^{-\frac{n}{4}+l}\|U_0\|_{L^1_l\cap L^2_l} + C\epsilon(1+\tau)^{-\frac{n}{4}+l} + C\epsilon(1+\tau)^{-\frac{n}{4}+l}M_l(t) \end{aligned}$$

for $k = 0, 1$ and $0 \le \tau \le t$. Hence we have

$$(1+\tau)^{\frac{n}{4}-l}\big\||x|^l U\big\|_{L^2} \le C\|U_0\|_{L^1_l\cap L^2_l} + C\epsilon + C\epsilon M_l(t).$$

Taking the supremum in $\tau \in [0,t]$, we obtain the desired estimate if ϵ is sufficiently small. □

The desired decay estimate in Proposition 4.7 now follows from Proposition 5.4.

Acknowledgements. Y. Kagei was partly supported by JSPS KAKENHI Grant Number 24340028, 15K13449, 24224003.

References

[1] H. Bahouri, J. Y. Chemin, R. Danchin *Fourier analysis and nonlinear partial differential equation*, Springer, Heidelberg, 2011.

[2] A. Friedman, *Partial Differential Equations*, Holt, Rinehert and Winston, New York, 1969.

[3] D. Hoff, K. Zumbrun, Multi-dimensional Diffusion Waves for the Navier-Stokes Equations of Compressible Flow, Indiana Univ. Math. J. **44** (1995), 604-676.

[4] S. Kawashima, A. Matsumura, T. Nishida, On the Fluid Dynamical Approximation to the Boltzmann Equation at the Level of the Navier-Stokes Equation, Comm. Math.Phys. **70** (1979), 97-124.

[5] T. Kobayashi, Y. Shibata, Remark on the rate of decay of solutions to linearized Compressible Navier-Stokes Equation, Pacific J. Math. **207** (2002), 199-234.

[6] A. Matsumura, T. Nishida, The initial value problem for the equation of motion of compressible viscous and heat-conductive fluids, Proc. Japan Acad. Ser. A **55** (1979), 337-342.

[7] A. Matsumura, T. Nishida, The initial value problems for the equation of motion of compressible viscous and heat-conductive gases, J. Math. Kyoto Univ. **20** (1980), 67-104.

[8] M. Okita, On the convergence rates for the compressible Navier- Stokes equations with potential force, Kyushu J. Math. **68** (2014), 261-286

List of MI Preprint Series, Kyushu University

The Global COE Program
Math-for-Industry Education & Research Hub

MI

MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata

MI2008-2 Eiji ONODERA
The intial value problem for a third-order dispersive flow into compact almost Hermitian manifolds

MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space

MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme

MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in nfinite extensions over a p-adic field

MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields

MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited

MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition

MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials

MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE

MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds

MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator

MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials

MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality

MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions

MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings

MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors

MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization

MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions

MI2009-5 Toshiro HIRANOUCHI & Yuichiro TAGUCHII
Flat modules and Groebner bases over truncated discrete valuation rings

MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations

MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow

MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization

MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity

MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption

MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve

MI2009-12 Tetsu MASUDA
Hypergeometric τ-functions of the q-Painlevé system of type $E_8^{(1)}$

MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination

MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications

MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain

MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring

MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions

MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force

MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application

MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions

MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations

MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces

MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions

MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map

MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2-projection

MI2009-26 Manabu YOSHIDA
Ramification of local fields and Fontaine's property (Pm)

MI2009-27 Yu KAWAKAMI
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space

MI2009-28 Masahisa TABATA
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme

MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance

MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis

MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI
Hecke's zeros and higher depth determinants

MI2009-32 Olivier PIRONNEAU & Masahisa TABATA
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type

MI2009-33 Chikashi ARITA
Queueing process with excluded-volume effect

MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA
Projective reduction of the discrete Painlevé system of type$(A_2 + A_1)^{(1)}$

MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI
Finite element computation for scattering problems of micro-hologram using DtN map

MI2009-36 Reiichiro KAWAI & Hiroki MASUDA
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes

MI2009-37 Hiroki MASUDA
On statistical aspects in calibrating a geometric skewed stable asset price model

MI2010-1 Hiroki MASUDA
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes

MI2010-2 Reiichiro KAWAI & Hiroki MASUDA
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations

MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI
Hyper-parameter selection in Bayesian structural equation models

MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons

MI2010-5 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and detecting change point via the relevance vector machine

MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI
Semi-supervised logistic discrimination via graph-based regularization

MI2010-7 Teruhisa TSUDA
UC hierarchy and monodromy preserving deformation

MI2010-8 Takahiro ITO
Abstract collision systems on groups

MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA
An algebraic approach to underdetermined experiments

MI2010-10 Kei HIROSE & Sadanori KONISHI
Variable selection via the grouped weighted lasso for factor analysis models

MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA
Derivation of specific conditions with Comprehensive Groebner Systems

MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow

MI2010-13 Reiichiro KAWAI & Hiroki MASUDA
On simulation of tempered stable random variates

MI2010-14 Yoshiyasu OZEKI
Non-existence of certain Galois representations with a uniform tame inertia weight

MI2010-15 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency

MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO
The value distribution of the Gauss map of improper affine spheres

MI2010-17 Kazunori YASUTAKE
On the classification of rank 2 almost Fano bundles on projective space

MI2010-18 Toshimitsu TAKAESU
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field

MI2010-19 Reiichiro KAWAI & Hiroki MASUDA
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling

MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE
Lagrangian approach to weakly nonlinear stability of an elliptical flow

MI2010-21 Hiroki MASUDA
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test

MI2010-22 Toshimitsu TAKAESU
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs

MI2010-23 Takahiro ITO, Mitsuhiko FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Composition, union and division of cellular automata on groups

MI2010-24 Toshimitsu TAKAESU
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra

MI2010-25 Toshimitsu TAKAESU
On the Essential Self-Adjointness of Anti-Commutative Operators

MI2010-26 Reiichiro KAWAI & Hiroki MASUDA
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling

MI2010-27 Chikashi ARITA & Daichi YANAGISAWA
Exclusive Queueing Process with Discrete Time

MI2010-28 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Motion and Bäcklund transformations of discrete plane curves

MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE
On the Number of the Pairing-friendly Curves

MI2010-30 Chikashi ARITA & Kohei MOTEGI
Spin-spin correlation functions of the q-VBS state of an integer spin model

MI2010-31 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and spike detection via Gaussian basis expansions

MI2010-32 Nobutaka NAKAZONO
Hypergeometric τ functions of the q-Painlevé systems of type $(A_2 + A_1)^{(1)}$

MI2010-33 Yoshiyuki KAGEI
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows

MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI
Milnor-Selberg zeta functions and zeta regularizations

MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms

MI2010-36 Takanori YASUDA
CAP representations of inner forms of $Sp(4)$ with respect to Klingen parabolic subgroup

MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER
Dynamical analysis of the exclusive queueing process

MI2011-1 Yasuhide FUKUMOTO& Alexander B. SAMOKHIN
Singular electromagnetic modes in an anisotropic medium

MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI
Asymptotic tail dependence of the normal copula

MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI
Anderson-Darling test and the Malliavin calculus

MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling via Compressed Sensing

MI2011-5 Hiroshi INOUE
Implications in Compressed Sensing and the Restricted Isometry Property

MI2011-6 Daeju KIM & Sadanori KONISHI
Predictive information criterion for nonlinear regression model based on basis expansion methods

MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI
Group variable selection via relevance vector machine

MI2011-8 Jan BREZINA & Yoshiyuki KAGEI
Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow
Group variable selection via relevance vector machine

MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC
Recursive structures in the multispecies TASEP

MI2011-10 Kazunori YASUTAKE
On projective space bundle with nef normalized tautological line bundle

MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA
An explicit formula for the discrete power function associated with circle patterns of Schramm type

MI2011-12 Yoshiyuki KAGEI
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow

MI2011-13 Vladimír CHALUPECKÝ & Adrian MUNTEAN
Semi-discrete finite difference multiscale scheme for a concrete corrosion model: approximation estimates and convergence

MI2011-14 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves

MI2011-15 Hiroshi INOUE
A generalization of restricted isometry property and applications to compressed sensing

MI2011-16 Yu KAWAKAMI
A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space

MI2011-17 Naoyuki KAMIYAMA
Matroid intersection with priority constraints

MI2012-1 Kazufumi KIMOTO & Masato WAKAYAMA
Spectrum of non-commutative harmonic oscillators and residual modular forms

MI2012-2 Hiroki MASUDA
Mighty convergence of the Gaussian quasi-likelihood random fields for ergodic Levy driven SDE observed at high frequency

MI2012-3 Hiroshi INOUE
A Weak RIP of theory of compressed sensing and LASSO

MI2012-4 Yasuhide FUKUMOTO & Youich MIE
Hamiltonian bifurcation theory for a rotating flow subject to elliptic straining field

MI2012-5 Yu KAWAKAMI
On the maximal number of exceptional values of Gauss maps for various classes of surfaces

MI2012-6 Marcio GAMEIRO, Yasuaki HIRAOKA, Shunsuke IZUMI, Miroslav KRAMAR, Konstantin MISCHAIKOW & Vidit NANDA
Topological Measurement of Protein Compressibility via Persistence Diagrams

MI2012-7 Nobutaka NAKAZONO & Seiji NISHIOKA
Solutions to a q-analog of Painlevé III equation of type $D_7^{(1)}$

MI2012-8 Naoyuki KAMIYAMA
A new approach to the Pareto stable matching problem

MI2012-9 Jan BREZINA & Yoshiyuki KAGEI
Spectral properties of the linearized compressible Navier-Stokes equation around time-periodic parallel flow

MI2012-10 Jan BREZINA
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a time-periodic parallel flow

MI2012-11 Daeju KIM, Shuichi KAWANO & Yoshiyuki NINOMIYA
Adaptive basis expansion via the extended fused lasso

MI2012-12 Masato WAKAYAMA
On simplicity of the lowest eigenvalue of non-commutative harmonic oscillators

MI2012-13 Masatoshi OKITA
On the convergence rates for the compressible
Navier- Stokes equations with potential force

MI2013-1 Abuduwaili PAERHATI & Yasuhide FUKUMOTO
A Counter-example to Thomson-Tait-Chetayev's Theorem

MI2013-2 Yasuhide FUKUMOTO & Hirofumi SAKUMA
A unified view of topological invariants of barotropic and baroclinic fluids and their application to formal stability analysis of three-dimensional ideal gas flows

MI2013-3 Hiroki MASUDA
Asymptotics for functionals of self-normalized residuals of discretely observed stochastic processes

MI2013-4 Naoyuki KAMIYAMA
On Counting Output Patterns of Logic Circuits

MI2013-5 Hiroshi INOUE
RIPless Theory for Compressed Sensing

MI2013-6 Hiroshi INOUE
Improved bounds on Restricted isometry for compressed sensing

MI2013-7 Hidetoshi MATSUI
Variable and boundary selection for functional data via multiclass logistic regression modeling

MI2013-8 Hidetoshi MATSUI
Variable selection for varying coefficient models with the sparse regularization

MI2013-9 Naoyuki KAMIYAMA
Packing Arborescences in Acyclic Temporal Networks

MI2013-10 Masato WAKAYAMA
Equivalence between the eigenvalue problem of non-commutative harmonic oscillators and existence of holomorphic solutions of Heun's differential equations, eigenstates degeneration, and Rabi's model

MI2013-11 Masatoshi OKITA
Optimal decay rate for strong solutions in critical spaces to the compressible Navier-Stokes equations

MI2013-12 Shuichi KAWANO, Ibuki HOSHINA, Kazuki MATSUDA & Sadanori KONISHI
Predictive model selection criteria for Bayesian lasso

MI2013-13 Hayato CHIBA
The First Painleve Equation on the Weighted Projective Space

MI2013-14 Hidetoshi MATSUI
Variable selection for functional linear models with functional predictors and a functional response

MI2013-15 Naoyuki KAMIYAMA
The Fault-Tolerant Facility Location Problem with Submodular Penalties

MI2013-16 Hidetoshi MATSUI
Selection of classification boundaries using the logistic regression

MI2014-1 Naoyuki KAMIYAMA
Popular Matchings under Matroid Constraints

MI2014-2 Yasuhide FUKUMOTO & Youichi MIE
Lagrangian approach to weakly nonlinear interaction of Kelvin waves and a symmetry-breaking bifurcation of a rotating flow

MI2014-3 Reika AOYAMA
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Parallel flow in a cylindrical domain

MI2014-4 Naoyuki KAMIYAMA
The Popular Condensation Problem under Matroid Constraints

MI2014-5 Yoshiyuki KAGEI & Kazuyuki TSUDA
Existence and stability of time periodic solution to the compressible Navier-Stokes equation for time periodic external force with symmetry

MI2014-6 This paper was withdrawn by the authors.

MI2014-7 Masatoshi OKITA
On decay estimate of strong solutions in critical spaces for the compressible Navier-Stokes equations

MI2014-8 Rong ZOU & Yasuhide FUKUMOTO
Local stability analysis of azimuthal magnetorotational instability of ideal MHD flows

MI2014-9 Yoshiyuki KAGEI & Naoki MAKIO
Spectral properties of the linearized semigroup of the compressible Navier-Stokes equation on a periodic layer

MI2014-10 Kazuyuki TSUDA
On the existence and stability of time periodic solution to the compressible Navier-Stokes equation on the whole space

MI2014-11 Yoshiyuki KAGEI & Takaaki NISHIDA
Instability of plane Poiseuille flow in viscous compressible gas

MI2014-12 Chien-Chung HUANG, Naonori KAKIMURA & Naoyuki KAMIYAMA
Exact and approximation algorithms for weighted matroid intersection

MI2014-13 Yusuke SHIMIZU
Moment convergence of regularized least-squares estimator for linear regression model

MI2015-1 Hidetoshi MATSUI & Yuta UMEZU
Sparse regularization for multivariate linear models for functional data

MI2015-2 Reika AOYAMA & Yoshiyuki KAGEI
Spectral properties of the semigroup for the linearized compressible Navier-Stokes equation around a parallel flow in a cylindrical domain

MI2015-3 Naoyuki KAMIYAMA
Stable Matchings with Ties, Master Preference Lists, and Matroid Constraints

MI2015-4 Reika AOYAMA & Yoshiyuki KAGEI
Large time behavior of solutions to the compressible Navier-Stokes equations around a parallel flow in a cylindrical domain

MI2015-5 Kazuyuki TSUDA
Existence and stability of time periodic solution to the compressible Navier-Stokes-Korteweg system on R^3

MI2015-6 Naoyuki KAMIYAMA
Popular Matchings with Ties and Matroid Constraints

MI2015-7 Shoichi EGUCHI & Hiroki MASUDA
Quasi-Bayesian model comparison for LAQ models

MI2015-8 Yoshiyuki KAGEI & Ryouta OOMACHI
Stability of time periodic solution of the Navier-Stokes equation on the half-space under oscillatory moving boundary condition

MI2016-1 Momonari KUDO
Analysis of an algorithm to compute the cohomology groups of coherent sheaves and its applications

MI2016-2 Yoshiyuki KAGEI & Masatoshi OKITA
Asymptotic profiles for the compressible Navier-Stokes equations on the whole space