Asymptotic behavior of the linearized semigroup at space-periodic stationary solution of the compressible Navier-Stokes equation

Enomoto, Shota
Graduate School of Mathematics, Kyushu University

Kagei, Yoshiyuki
Faculty of Mathematics, Kyushu University

<http://hdl.handle.net/2324/1655024>

出版情報 : MI Preprint Series. 2016-3, 2016-04-14. 九州大学大学院数理学研究院
バージョン :
権利関係 :

MI Preprint Series

Mathematics for Industry
Kyushu University

**Asymptotic behavior of the
linearized semigroup at
space-periodic stationary
solution of the compressible
Navier-Stokes equation**

**Shota Enomoto
& Yoshiyuki Kagei**

MI 2016-3

(Received April 14, 2016)

Institute of Mathematics for Industry
Graduate School of Mathematics
Kyushu University
Fukuoka, JAPAN

Asymptotic behavior of the linearized semigroup at space-periodic stationary solution of the compressible Navier-Stokes equation

Shota Enomoto¹ and Yoshiyuki Kagei²

¹ Graduate School of Mathematics,
Kyushu University,
Nishi-ku, Motooka 744,
Fukuoka, 819-0395, JAPAN

² Faculty of Mathematics,
Kyushu University,
Nishi-ku, Motooka 744,
Fukuoka 819-0395, Japan

Abstract

The asymptotic behavior of the linearized semigroup at spatially periodic stationary solution of the compressible Navier-Stokes equation in a periodic layer of $\mathbb{R}^n$ ($n = 2, 3$) is investigated. It is shown that if the Reynolds and Mach numbers are sufficiently small, then the linearized semigroup is decomposed into two parts; one behaves like a solution of an $n - 1$ dimensional linear heat equation as time goes to infinity and the other one decays exponentially.

Keywords: Compressible Navier-Stokes equation, space-periodic stationary solution, linearized semigroup, spectrum.

1 Introduction

This paper studies the stability of stationary solution to the compressible Navier-Stokes equation

$$\partial_t \rho + \operatorname{div}(\rho v) = 0, \quad (1.1)$$

$$\rho(\partial_t v + (v \cdot \nabla)v) - \mu \Delta v - (\mu + \mu') \nabla \operatorname{div} v + \nabla p(\rho) = \rho g \quad (1.2)$$

in a periodic layer Ω_* of $\mathbb{R}^n$ with $n = 2, 3$:

$$\Omega_* = \{x = (x', x_n); x' = (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1}, \omega_{*,1}(x') < x_n < \omega_{*,2}(x')\},$$

where $\omega_{*,1}$ and $\omega_{*,2}$ are smooth Q_* -periodic functions in x' with the periodic cell $Q_* = \prod_{j=1}^{n-1} [-\frac{\pi}{\alpha_{*,j}}, \frac{\pi}{\alpha_{*,j}})$ for positive constants $\alpha_{*,j}$ ($j = 1, \dots, n-1$), namely, $\omega_{*,1}$ and $\omega_{*,2}$ are smooth functions satisfying $\omega_{*,j}(x' + \frac{2\pi}{\alpha_i} \mathbf{e}'_i) = \omega_{*,j}(x')$ ($i = 1, \dots, n-1, j = 1, 2$) with $\mathbf{e}'_i = {}^\top(0, \dots, \overset{i}{1}, \dots, 0) \in \mathbb{R}^{n-1}$; $\rho = \rho(x, t)$ and $v = {}^\top(v^1(x, t), \dots, v^n(x, t))$ denote the unknown density and velocity, respectively, at $x \in \Omega_*$ and $t \geq 0$; $p = p(\rho)$ is the pressure that is assumed to be a smooth function of ρ ; $g = {}^\top(g^1(x), \dots, g^n(x))$ is a given external force; and, μ and μ' are the viscosity coefficients that are assumed to be constants. Here and in what follows the superscript $^\top$ stands for the transposition.

We assume that μ and μ' satisfy $\mu > 0$ and $\frac{2}{n}\mu + \mu' \geq 0$. Furthermore we assume that $\frac{\mu'}{\mu}$ satisfies

$$\frac{\mu'}{\mu} \leq \mu_1 \quad (1.3)$$

for a certain constant $\mu_1 > 0$. We also assume that p satisfies $p'(\rho_*) > 0$ for a given constant $\rho_* > 0$, and that $g = {}^\top(g^1(x), \dots, g^n(x))$ is Q_* -periodic in x' .

We consider (1.1)-(1.2) under the no-slip boundary condition on $\{x_n = \omega_{*,j}(x')\}$ ($j = 1, 2$) for v :

$$v|_{x_n=\omega_{*,j}(x')} = 0 \quad (x' \in \mathbb{R}^{n-1}, j = 1, 2). \quad (1.4)$$

One can see that if g is sufficiently small, the system (1.1)-(1.2) with (1.4) has a Q_* -periodic stationary solution $\bar{u}_s = {}^\top(\bar{\rho}_s, \bar{v}_s)$, whose components $\bar{\rho}_s$ and $\bar{v}_s$ are in general non-uniform in x' and x_n .

The purpose of this paper is to study the large time behavior of solutions around $\bar{u}_s$. As a first step of the analysis, we here investigate spectral properties of the linearized semigroup around $\bar{u}_s = {}^\top(\bar{\rho}_s, \bar{v}_s)$.

Large time behavior of solutions of compressible Navier-Stokes equations in unbounded domains exhibit interesting phenomenon, and asymptotic behavior of solutions in large time have been obtained in detail. See, e.g., [7, 9, 14, 17, 18, 19, 20, 21, 22, 24] for the cases of multi-dimensional whole space, half space and exterior domains. In addition to these domains, the stability of flows in infinite layers and cylindrical domains has been paid much attention, e.g., the stability of parallel flows and periodic (in space or time) flow patterns, and so on, whose mathematical analysis is still under development in the case of compressible flows. A difficulty in the mathematical analysis appears due to the non-uniform velocity field of parallel flows and periodic flow patterns, which makes the hyperbolic aspect of the equations (1.1)-(1.2) stronger, and thus, the stability analysis is getting more difficult

compared with that of the motionless state (i.e., the flow with zero velocity field). As for the stability of stationary flow with non-zero velocity field on the whole space, we note that an interesting analysis was given by Shibata and Tanaka [24] to establish the decay rate of perturbations.

Concerning the large time behavior of solutions in the case of an n dimensional flat layer $\{x = (x', x_n); x' \in \mathbb{R}^{n-1}, 0 < x_n < 1\}$, (1.1)-(1.4) has a simple stationary solution $\bar{u}_s = {}^\top(\bar{\rho}_s(x_n), \bar{v}_s(x_n))$, called parallel flow, when g takes the form $g = {}^\top(g'(x_n), 0, \dots, 0, g^n(x_n))$. It was proved in [12, 13] that parallel flow is asymptotically stable for small initial perturbation if the Reynolds and Mach numbers are sufficiently small. (See also [5, 6] for time-periodic parallel flows.) Furthermore, the asymptotic behavior of the perturbation is described by an $n - 1$ dimensional linear heat equation in the case $n \geq 3$, and by a one-dimensional viscous Burgers equation in the case $n = 2$. The proof is based on the energy method and the spectral analysis of the linearized semigroup which is investigated by using the Fourier transform in x' variable. Similar results hold for the parallel flow in the case of cylindrical domain [1, 2, 3].

The analysis in [2, 6, 12] heavily depends on the structure of parallel flow that is uniform in x' variable. Due to this structure, the Fourier transform in x' is useful to investigate the stability of parallel flow, and one of the points in the analysis in [2, 6, 12] is that the *low frequency part* of the linearized operator can be regarded as a perturbation of the linearized operator around the motionless state which is sectorial. As for the Q_* -periodic stationary solution $\bar{u}_s = {}^\top(\bar{\rho}_s, \bar{v}_s)$ under consideration, the argument in [2, 6, 12] is formally applicable if one uses the Bloch transform instead of the Fourier transform. However, since the velocity field $\bar{v}_s$ is in general non-uniform in both x' and x_n , due to the term $\bar{v}_s \cdot \nabla \phi$ in the linearized equation of (1.1), the *low frequency part* of the linearized operator is no longer regarded as a simple perturbation of the linearized one around the motionless state. Therefore, the argument in [2, 6, 12] cannot be directly applied to the problem under consideration. To overcome this difficulty we will make use of the solvability result on the stationary transport equation given by Heywood and Padula [10], which makes the formal argument a rigorous one.

Our main results of this paper is summarized as follows. After introducing suitable non-dimensional variables, the equations for the perturbation $u = {}^\top(\phi, w) = {}^\top(\gamma^2(\rho - \rho_s), v - v_s)$ takes the following form:

$$\partial_t \phi + \operatorname{div}(\phi v_s) + \gamma^2 \operatorname{div}(\rho_s w) = f^0, \quad (1.5)$$

$$\begin{aligned} \partial_t w - \frac{\nu}{\rho_s} \Delta w - \frac{\tilde{\nu}}{\rho_s} \nabla \operatorname{div} w + \nabla \left(\frac{P'(\rho_s)}{\gamma^2 \rho_s} \phi \right) \\ + \frac{1}{\gamma^2 \rho_s^2} (\nu \Delta v_s + \tilde{\nu} \nabla \operatorname{div} v_s) \phi + v_s \cdot \nabla w + w \cdot \nabla v_s = \tilde{f}, \end{aligned} \quad (1.6)$$

$$w|_{\partial\Omega} = 0, \quad (1.7)$$

$$u|_{t=0} = u_0 = {}^\top(\phi_0, w_0). \quad (1.8)$$

Here $u_s = {}^\top(\rho_s, v_s)$ denotes the non-dimensionalization of $\bar{u}_s = {}^\top(\bar{\rho}_s, \bar{v}_s)$, and ν , $\tilde{\nu}$ and γ are non-dimensional parameters. The Reynolds number Re and Mach number Ma are given by $\operatorname{Re} = \frac{1}{\nu}$ and $\operatorname{Ma} = \frac{1}{\gamma}$, respectively. The terms f_0 and $\tilde{f}$ on the right of (1.5) and (1.6) are nonlinear terms given by

$$\begin{aligned} f_0 &= -\operatorname{div}(\phi w), \\ \tilde{f} &= -w \cdot \nabla w - \frac{\phi}{\rho_s(\gamma^2 \rho_s + \phi)} \left(\nu \Delta w + \tilde{\nu} \nabla \operatorname{div} w - \frac{\nu \phi}{\gamma^2 \rho_s} \Delta v_s - \frac{\tilde{\nu} \phi}{\gamma^2 \rho_s} \nabla \operatorname{div} v_s \right) \\ &\quad + \frac{\phi}{\gamma^2 \rho_s^2} \nabla \left(P^{(1)} \left(\rho_s, \frac{\phi}{\gamma^2} \right) \frac{\phi}{\gamma^2} \right) + \frac{\phi^2}{\gamma^2 \rho_s^2 (\gamma^2 \rho_s + \phi)} \nabla \left(P \left(\rho_s + \frac{\phi}{\gamma^2} \right) \right) \\ &\quad + \frac{1}{\rho_s} \nabla \left(P^{(2)} \left(\rho_s, \frac{\phi}{\gamma^2} \right) \frac{\phi^2}{\gamma^4} \right) \end{aligned}$$

with

$$P^{(1)}(\rho_s, \phi) = \int_0^1 P'(\rho_s + \theta \phi) d\theta, \quad P^{(2)}(\rho_s, \phi) = \int_0^1 (1 - \theta) P''(\rho_s + \theta \phi) d\theta.$$

The linearized problem for (1.5)-(1.8) is then formulated as

$$\partial_t u + Lu = 0, \quad u|_{t=0} = u_0, \quad (1.9)$$

where L is the operator of the form

$$\begin{aligned} L &= \begin{pmatrix} \operatorname{div}(\cdot v_s) & \gamma^2 \operatorname{div}(\rho_s \cdot) \\ \nabla \left(\frac{P'(\rho)}{\gamma^2 \rho_s} \cdot \right) & -\frac{\nu}{\rho_s} \Delta - \frac{\tilde{\nu}}{\rho_s} \nabla \operatorname{div} \end{pmatrix} \\ &\quad + \begin{pmatrix} 0 & 0 \\ \frac{1}{\gamma^2 \rho_s^2} (\nu \Delta v_s + \tilde{\nu} \nabla \operatorname{div} v_s) & v_s \cdot \nabla + {}^\top \nabla v_s \end{pmatrix}. \end{aligned}$$

We consider the operator L as an operator on $L^2(\Omega)$ with domain $D(L) = \{u = {}^\top(\phi, w) \in L^2(\Omega); w \in H_0^1(\Omega), Lu \in L^2(\Omega)\}$. We will prove that there exist positive constants ν_0 and $\tilde{\gamma}_0$ such that if ν and $\frac{\gamma^2}{\nu}$ are sufficiently large, then the linearized semigroup e^{-tL} satisfies

$$\|\Pi e^{-tL} u_0\|_{L^2(\Omega)} \leq C(1+t)^{-\frac{n-1}{4}} \|u_0\|_{L^1(\Omega)}, \quad (1.10)$$

$$\|\Pi e^{-tL}u_0 - [e^{-tH}\sigma_0]u^{(0)}\|_{L^2(\Omega)} \leq Ct^{-\frac{n-1}{4}-\frac{1}{2}}\|u_0\|_{L^1(\Omega)}, \quad (1.11)$$

where Π is a bounded projection on $L^2(\Omega)$ with $\Pi L \subset L\Pi$, and the operator $H : L^2(\mathbb{R}^{n-1}) \rightarrow L^2(\mathbb{R}^{n-1})$ with domain $D(H) = H^2(\mathbb{R}^{n-1})$ is given by

$$H\sigma = -\frac{\gamma^2}{\nu} \sum_{j,k=1}^{n-1} a_{jk} \partial_{x_j} \partial_{x_k} \sigma + \sum_{j=1}^{n-1} a_j \partial_{x_j} \sigma \quad (\sigma \in D(H))$$

with $a_j \in \mathbb{R}$ and $a_{jk} \in \mathbb{R}$ satisfying $\sum_{j,k=1}^{n-1} a_{jk} \eta_j \eta_k \geq C|\eta'|^2$ ($\eta' \in \mathbb{R}^{n-1}$) and σ_0 is given by $\sigma_0(x') = \frac{|Q|}{|\Omega_{per}|} \int_{\omega_1(x')}^{\omega_2(x')} \phi_0(x', x_n) dx_n$. Here $Q = \Pi_{j=1}^{n-1} [-\frac{\pi}{\alpha_j}, \frac{\pi}{\alpha_j})$ is the non-dimensional periodic cell and Ω_{per} is the non-dimensional basic period domain defined by

$$\Omega_{per} = \{x = (x', x_n); |x_i| < \frac{\pi}{\alpha_i} \ (i = 1, \dots, n-1), \ \omega_1(x') < x_n < \omega_2(x')\}.$$

Moreover, $u^{(0)} = {}^\top(\phi^{(0)}, w^{(0)})$ is Q -periodic function in x' and it satisfies the following problem on the basic period domain Ω_{per} :

$$\begin{cases} \operatorname{div}(\phi^{(0)}v_s) + \gamma^2 \operatorname{div}(\rho_s w^{(0)}) = 0 & \text{in } \Omega_{per}, \\ \nabla \left(\frac{P'(\rho_s)}{\gamma^2 \rho_s} \phi^{(0)} \right) + \frac{1}{\gamma^2 \rho_s^2} (\nu \Delta v_s + \tilde{\nu} \nabla \operatorname{div} v_s) \phi^{(0)} \\ \quad - \frac{\nu}{\rho_s} \Delta w^{(0)} - \frac{\tilde{\nu}}{\rho_s} \nabla \operatorname{div} w^{(0)} + v_s \cdot \nabla w^{(0)} + w^{(0)} \cdot \nabla v_s = 0 & \text{in } \Omega_{per}, \\ w^{(0)}|_{\{x_n=\omega_j(x')\}} = 0 \ (j = 1, 2), \\ \int_{\Omega_{per}} \phi^{(0)} = 1. \end{cases}$$

Concerning the $(I - \Pi)$ -part, as in [1, 2, 15], one can prove the exponential decay estimate:

$$\|e^{-tL}(I - \Pi)u_0\|_{H^1(\Omega)} \leq C\{e^{-at}\|u_0\|_{H^1 \times \tilde{H}^1} + t^{-\frac{1}{2}}\|w_0\|_{L^2}\} \quad (1.12)$$

for $u_0 = {}^\top(\phi_0, w_0) \in H^1(\Omega) \times \tilde{H}^1(\Omega)$ with some positive number a . Here $\tilde{H}^1(\Omega)$ denotes the set of all locally H^1 functions in Ω whose tangential derivatives near $\partial\Omega$ belong to $L^2(\Omega)$. See Theorem 3.3 for the details.

The proof of the main results are based on the method of the Bloch transformation as in [15] where the stability of the motionless state on a periodic layer was studied. By using the Bloch transformation one can reduce the linearized problem (1.9) on Ω to the problem $\partial_t u + L_{\eta'} u = 0$ on Ω_{per} under Q -periodic boundary conditions in x' . Here $L_{\eta'}$ is the linear operator obtained by replacing the partial derivatives ∂_{x_j} ($j = 1, \dots, n-1$) in L by $\partial_{x_j} + i\eta_j$ with parameter $\eta' = (\eta_1, \dots, \eta_{n-1}) \in Q'$, where Q' is the dual

cell defined by $Q' = \Pi_{j=1}^{n-1}[-\frac{\alpha_j}{2}, \frac{\alpha_j}{2}]$. In the case $|\eta'| \ll 1$ the operator $L_{\eta'}$ can be regarded as a perturbation of L_0 . By using the solvability results on stationary transport equations by Heywood-Padula [10], we investigate the eigenspace for the eigenvalue 0 of L_0 , which enables us to apply the analytic perturbation theory to $L_{\eta'}$ with $|\eta'| \ll 1$. We show that

$$\rho(-L_{\eta'}) \supset \{\lambda \in \mathbb{C}; \operatorname{Re} \lambda > -\beta_0, |\lambda| \geq \frac{\beta_0}{2}\},$$

$$\sigma(-L_{\eta'}) \cap \{\lambda \in \mathbb{C}; |\lambda| < \frac{\beta_0}{2}\} = \{\lambda_{\eta'}\}$$

for some $\beta_0 > 0$, where $\lambda_{\eta'} = i \sum_{j=1}^{n-1} a_j \eta_j - \frac{\gamma^2}{\nu} \sum_{j,k=1}^{n-1} a_{jk} \eta_j \eta_k + O(|\eta'|^3)$ as $\eta' \rightarrow 0$. It then follows that this part of e^{-tL} behaves as in (1.10) and (1.11). As for the remaining part of η' , one can establish the exponential decay estimate (1.12) as in the arguments given in [1, 2, 15].

This paper is organized as follows. In section 2 we introduce notations, function spaces and operators. In section 3 we state the main results of this paper. Section 4 is devoted to the proof of (1.10) and (1.11). In section 4.1 we formulate the linearized problem and reduce it to the ones on the basic period domain Ω_{per} by the Bloch transform. In section 4.2 we investigate the null space of L_0 . The spectrum of $L_{\eta'}$ with $|\eta'| \ll 1$ is then investigated in section 4.3. Section 4.4 is devoted to estimating the eigenprojection, and in section 4.5 the estimates (1.10) and (1.11) are established. In section 5 we give resolvent estimates for the principal part of L_0 by the Matsumura-Nishida energy method [22] and the result of Heywood-Padula [10].

2 Preliminaries

In this section we introduce notation used in this paper.

Let D be a domain. $L^p(D)$ ($1 \leq p \leq \infty$) denotes Lebesgue space on D and its norm is denoted by $\|\cdot\|_{L^p(D)}$. Similarly, $H^m(D)$ denotes the m -th order L^2 -Sobolev space on D and its norm is denoted by $\|\cdot\|_{H^m(D)}$. $C_0^m(D)$ stands for the set of C^m -functions having compact support on D . Furthermore, we denote the completion of $C_0^\infty(D)$ in $H^m(D)$ by $H_0^m(D)$, and the dual space of $H_0^m(D)$ is denoted by $H^{-m}(D)$. $\mathcal{S}(\mathbb{R}^k)$ denotes the Schwartz space on $\mathbb{R}^k$.

We simply write the set of all vector fields $w = {}^\top(w^1, \dots, w^n)$ on D with $w^j \in L^p(D)$ (resp. $H^m(D)$) ($j = 1, \dots, n$) as $L^p(D)$ (resp. $H^m(D)$), and its norm is denoted by $\|\cdot\|_{L^p(D)}$ (resp. $\|\cdot\|_{H^m(D)}$). We define $\|\cdot\|_{H^k(D) \times H^m(D)}$ by $\|u\|_{H^k(D) \times H^m(D)} = (\|\phi\|_{H^k(D)}^2 + \|w\|_{H^m(D)}^2)^{\frac{1}{2}}$ for $u = {}^\top(\phi, w)$ satisfying $\phi \in$

$H^k(D)$ and $w \in H^m(D)$. In particular, we write $\|u\|_{H^k(D) \times H^k(D)} = \|u\|_{H^k(D)}$ in the case $k = m$.

We next rewrite the problem (1.1)-(1.2) with (1.4) into the one in a non-dimensional form. We introduce the following non-dimensional variables:

$$x = l\tilde{x}, \quad t = \frac{l}{V}\tilde{t}, \quad v = V\tilde{v}, \quad \rho = \rho_*\tilde{\rho}, \quad p = \rho_*V^2P, \quad g = \frac{V^2}{l}\tilde{g}.$$

Here

$$l = |\Omega_{*,per}|^{\frac{1}{n}}, \quad V = \sqrt{l[g]_{H^2(\Omega_{*,per})}}, \quad [g]_{H^2(\Omega_{*,per})} = \sqrt{\sum_{s=0}^m l^{-n+2s} \|\partial_{\tilde{x}}^s g\|_{L^2(\Omega_{*,per})}^2}.$$

The problem (1.1)-(1.2) with (1.4) is transformed into the following non-dimensional problem in $\tilde{\Omega}$:

$$\partial_{\tilde{t}}\tilde{\rho} + \operatorname{div}_{\tilde{x}}(\tilde{\rho}\tilde{v}) = 0, \quad (2.1)$$

$$\tilde{\rho}(\partial_{\tilde{t}}\tilde{v} + (\tilde{v} \cdot \nabla_{\tilde{x}})\tilde{v}) - \nu\Delta_{\tilde{x}}\tilde{v} - \tilde{\nu}\nabla_{\tilde{x}}\operatorname{div}_{\tilde{x}}\tilde{v} + \nabla_{\tilde{x}}P(\tilde{\rho}) = \tilde{\rho}\tilde{g}, \quad (2.2)$$

$$\tilde{v}|_{x_n=\omega_j(x')} = 0 \quad (j = 1, 2). \quad (2.3)$$

Here Ω is the non-dimensional periodic layer:

$$\Omega = \{x = (x', x_n); x' = (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1}, \omega_1(x') < x_n < \omega_2(x')\}$$

with ω_1 and ω_2 being smooth Q -periodic functions of x' with the period cell

$$Q = \prod_{j=1}^{n-1} [-\frac{\pi}{\alpha_j}, \frac{\pi}{\alpha_j}),$$

where $\alpha_j = l\alpha_{*,j}$; and ν , $\tilde{\nu}$ and γ are non-dimensional parameters defined by

$$\nu = \frac{\mu}{\rho_*lV}, \quad \tilde{\nu} = \nu + \nu', \quad \nu' = \frac{\mu'}{\rho_*lV}, \quad \gamma = \frac{\sqrt{p'(\rho_*)}}{V}.$$

The assumption (1.3) is written as

$$\frac{\nu'}{\nu} \leq \nu_* \quad (2.4)$$

for some positive number ν_* . Note that under (2.4) we have

$$\nu \leq \nu + \tilde{\nu} \leq (\nu_* + 2)\nu. \quad (2.5)$$

In what follows, we omit tildes $\sim$ of $\tilde{\rho}$, $\tilde{v}$, $\tilde{x}$, $\tilde{t}$, $\tilde{p}$ and $\tilde{g}$ for simplicity in notation.

We next define Q' , $\Sigma_{j,\pm}(j = 1, \dots, n-1)$ and Σ_n by

$$Q' := \Pi_{i=1}^{n-1} \left[-\frac{\alpha_i}{2}, \frac{\alpha_i}{2} \right), \quad \Sigma_{j,\pm} := \{x \in \bar{\Omega}; x_j = \pm \frac{\pi}{\alpha_j}\} \quad (j = 1, \dots, n-1)$$

and

$$\Sigma_n := \{x \in \partial\Omega; x' \in Q, x_n = \omega_j(x'), j = 1, 2\}.$$

The inner product of $L^2(\Omega_{per})$ is defined by

$$(f, g) = \int_{\Omega_{per}} f(x) \overline{g(x)} dx$$

for $f, g \in L^2(\Omega_{per})$. Furthermore the mean value of $f \in L^2(\Omega_{per})$ over Ω_{per} is represented as $\langle \cdot \rangle$, i.e.,

$$\langle f \rangle = \int_{\Omega_{per}} f(x) dx.$$

Note that $|\Omega_{per}| = 1$ due to the non-dimensionalization.

We denote the $k \times k$ identity matrix by I_k . We define $(n+1) \times (n+1)$ diagonal matrices Q_0 and $\tilde{Q}$ is defined by

$$Q_0 = \text{diag}(1, 0, \dots, 0), \quad \tilde{Q} = \text{diag}(0, 1, \dots, 1).$$

Note that $Q_0 u = {}^\top(\phi, 0)$ and $\tilde{Q} u = {}^\top(0, w)$ for $u = {}^\top(\phi, w)$.

The Fourier transform of $f = f(x')$ ($x' \in \mathbb{R}^{n-1}$) is denoted by $\hat{f}$ or $\mathcal{F}[f]$:

$$\hat{f}(\xi') = \mathcal{F}[f](\xi') = \int_{\mathbb{R}^{n-1}} f(x') e^{-i\xi' \cdot x'} dx', \quad \xi' \in \mathbb{R}^{n-1}.$$

We denote the inverse Fourier transform by $\mathcal{F}^{-1}$:

$$\mathcal{F}^{-1}[f](x') = \frac{1}{(2\pi)^{n-1}} \int_{\mathbb{R}^{n-1}} f(\xi') e^{i\xi' \cdot x'} d\xi', \quad x' \in \mathbb{R}^{n-1}.$$

For an operator A , we denote by $\text{Ker}(A)$ and $\text{R}(A)$ the null space and the range of A , respectively. The resolvent set and the spectrum of A are denoted by $\rho(A)$ and $\sigma(A)$, respectively. We denote the commutator of operators A and B by $[A, B]$, i.e., $[A, B]f = A(Bf) - B(Af)$.

We introduce function spaces of periodic functions as follows:

$$\begin{aligned} L_{per}^2(\Omega_{per}) &:= \{u|_{\Omega_{per}}; u \in L_{loc}^2(\bar{\Omega}), u(x', x_n) \text{ is } Q\text{-periodic in } x'\}, \\ C_{per}^\infty(\bar{\Omega}_{per}) &:= \{u|_{\Omega_{per}}; u \in C^\infty(\bar{\Omega}), u(x', x_n) \text{ is } Q\text{-periodic in } x'\}, \end{aligned}$$

$$\begin{aligned}
C_{0,per}^\infty(\Omega_{per}) &:= \{u \in C_{per}^\infty(\overline{\Omega_{per}}); u = 0 \text{ in a neighborhood of } \partial\Omega\}, \\
H_{per}^l(\Omega_{per}) &:= \text{the closure of } C_{per}^\infty(\overline{\Omega_{per}}) \text{ in } H^l(\Omega_{per}), \\
H_{0,per}^l(\Omega_{per}) &:= \text{the closure of } C_{0,per}^\infty(\overline{\Omega_{per}}) \text{ in } H^l(\Omega_{per}).
\end{aligned}$$

We note that $L_{per}^2(\Omega_{per})$ can be identified with $L^2(\Omega_{per})$, and that $H_{per}^l(\Omega_{per})$ and $H_{0,per}^1(\Omega_{per})$ are given by

$$\begin{aligned}
H_{per}^l(\Omega_{per}) &= \{u \in H^l(\Omega_{per}); \partial_{x'}^\beta u|_{\Sigma_{j,-}} = \partial_{x'}^\beta u|_{\Sigma_{j,+}}, 1 \leq j \leq n-1, |\beta| \leq l-1\}, \\
H_{0,per}^1(\Omega_{per}) &= \{u \in H_{per}^1(\Omega_{per}); u|_{\Sigma_n} = 0\}.
\end{aligned}$$

Moreover, we define $L_{per,*}^2(\Omega_{per})$ by

$$L_{per,*}^2(\Omega_{per}) := \{f \in L_{per}^2(\Omega_{per}); \langle f \rangle = 0\},$$

and $H_{per,*}^l(\Omega_{per})$ by $H_{per,*}^l(\Omega_{per}) = H_{per}^l(\Omega_{per}) \cap L_{per,*}^2(\Omega_{per})$. We will abbreviate $L_{per}^2(\Omega_{per})$ to L_{per}^2 , and likewise, $H_{per}^l(\Omega_{per})$ to H_{per}^l , $\dots$, and so on. The norm $\|\cdot\|_{L^p(\Omega_{per})}$ is written as $\|\cdot\|_p$, and similarly, $\|\cdot\|_{H^l(\Omega_{per})}$ as $\|\cdot\|_{H^l}$.

We next introduce the Bloch transformation.

Definition 2.1. We define the operator T by $(T\varphi)(x', \eta')$ ($x' \in \mathbb{R}^{n-1}, \eta' \in \mathbb{R}^{n-1}$) for $\varphi \in \mathcal{S}(\mathbb{R}^{n-1})$, where

$$\begin{aligned}
(T\varphi)(x', \eta') &= \frac{1}{(2\pi)^{\frac{n-1}{2}}|Q|^{\frac{1}{2}}} \sum_{(k_1, \dots, k_{n-1}) \in \mathbb{Z}^{n-1}} \hat{\varphi}(\eta' + \sum_{j=1}^{n-1} k_j \alpha_j \mathbf{e}'_j) e^{i \sum_{j=1}^{n-1} k_j \alpha_j x_j} \\
&= \frac{1}{|Q'|^{\frac{1}{2}}} \sum_{(l_1, \dots, l_{n-1}) \in \mathbb{Z}^{n-1}} \varphi(x' + \sum_{j=1}^{n-1} l_j \frac{2\pi}{\alpha_j} \mathbf{e}'_j) e^{-i \eta' \cdot (x' + \sum_{j=1}^{n-1} l_j \frac{2\pi}{\alpha_j} \mathbf{e}'_j)}.
\end{aligned}$$

We also define the operator U as follows. For a function $\varphi(x', \eta') \in C^\infty(\mathbb{R}^{n-1} \times \mathbb{R}^{n-1})$ such that $\varphi(x', \eta')$ is Q -periodic in x' and $\varphi(x', \eta') e^{i \eta' \cdot x'}$ is Q' -periodic in η' , we define $(U\varphi)(x')$ ($x' \in \mathbb{R}^{n-1}$) by

$$(U\varphi)(x') = \frac{1}{|Q'|^{\frac{1}{2}}} \int_{Q'} \varphi(x', \eta') e^{i x' \cdot \eta'} d\eta'.$$

Note that $\varphi(x', \eta' + \alpha_j \mathbf{e}'_j) = \varphi(x', \eta') e^{-i \alpha_j \mathbf{e}'_j \cdot x'}$ ($j = 1, \dots, n-1$).

The operators T and U have the following properties. See, e.g., [23] for the details.

Proposition 2.2. (i) $(T\varphi)(x', \eta')$ is Q -periodic in x' and $(T\varphi)(x', \eta')e^{i\eta' \cdot x'}$ is Q' -periodic in η' .

(ii) T is uniquely extended to an isometric operator from $L^2(\mathbb{R}^{n-1})$ to $L^2(Q'; L^2(Q))$.

(iii) U is the inverse operator of T .

(iv) Let ψ be Q -periodic in x' . Then it holds that $T(\psi\varphi) = \psi T\varphi$.

(v) $T(\partial_{x_j}\varphi) = (\partial_{x_j} + i\eta_j)T\varphi$ ($j = 1, \dots, n-1$) and T defines an isomorphism from $H^l(\mathbb{R}^{n-1})$ to $L^2(Q'; H_{per}^l(Q))$. Here $H_{per}^l(Q)$ denotes the space of Q -periodic functions in $H^l(Q)$, as in the case of $H_{per}^l(\Omega_{per})$.

We introduce the following weighted inner product:

$$\langle u_1, u_2 \rangle = \int_{\Omega_{per}} \phi_1 \phi_2 \frac{P'(\rho_s)}{\gamma^4 \rho_s} dx + \int_{\Omega_{per}} w_1 w_2 \rho_s dx$$

for $u_j = {}^\top(\phi_j, w_j)$ ($j = 1, 2$), where $\rho_s = \rho_s(x)$ is the density of the stationary solution. See Proposition 3.1 below. The associated norm $||| \cdot |||_2$ is defined by

$$|||u|||_2^2 \equiv \left(\left\| \sqrt{\frac{P'(\rho_s)}{\gamma^4 \rho_s}} \phi \right\|_2^2 + \|\sqrt{\rho_s} w\|_2^2 \right).$$

As we will see in Proposition 3.1, $\rho_s(x)$ and $\frac{P'(\rho_s(x))}{\rho_s(x)}$ are strictly positive in Ω_{per} if ν and $\frac{\gamma^2}{\nu}$ are suitably large, and $||| \cdot |||_2$ is equivalent to $\| \cdot \|_2$.

We introduce the Bogovskii lemma [4]. (See also [8, Chapter 3].)

Lemma 2.3. *There exists a bounded linear operator $\mathcal{B} : L_{per,*}^2 \rightarrow H_{per,0}^1$ such that*

$$\operatorname{div} \mathcal{B}g = g, \quad g \in L_{per,*}^2$$

$$\|\nabla \mathcal{B}g\|_2 \leq C\|g\|_2,$$

where C is a positive constant depending only on Ω_{per} . Furthermore, if $g = \operatorname{div} \mathbf{g}$ with $\mathbf{g} = {}^\top(g^1, \dots, g^n)$ satisfying $g^j|_{\Sigma_{j,-}} = g^j|_{\Sigma_{j,+}}$ ($j = 1, \dots, n-1$), $g^n|_{\Sigma_n} = 0$, then

$$\operatorname{div} \mathcal{B}(\operatorname{div} \mathbf{g}) = \operatorname{div} \mathbf{g},$$

$$\|\mathcal{B}(\operatorname{div} \mathbf{g})\|_2 \leq C\|\mathbf{g}\|_2.$$

For a given $\delta > 0$, we introduce the following notation $((\cdot, \cdot))$ and $\langle\langle \cdot, \cdot \rangle\rangle$:

$$((u_1, u_2)) := \frac{1}{\gamma^2}(\phi_1, \phi_2) + (w_1, w_2) - \delta\{(w_1, \mathcal{B}\phi_2) + (\mathcal{B}\phi_1, w_2)\},$$

$$\langle\langle u_1, u_2 \rangle\rangle := \langle u_1, u_2 \rangle - \delta\{(w_1, \mathcal{B}\phi_2) + (\mathcal{B}\phi_1, w_2)\}$$

for $u_j = {}^\top(\phi_j, w_j)$ ($j = 1, 2$). There exists a constant $c_0 > 0$ such that if $0 \leq \delta \leq \frac{c_0}{\gamma}$, then $((\cdot, \cdot))$ and $\langle\langle \cdot, \cdot \rangle\rangle$ define inner products on $L_{per,*}^2 \times L_{per}^2$ satisfying

$$\frac{1}{2}\|u\|_2^2 \leq ((u, u)) \leq \frac{3}{2}\|u\|_2^2$$

and

$$\frac{1}{2}|||u|||_2^2 \leq \langle\langle u, u \rangle\rangle \leq \frac{3}{2}|||u|||_2^2.$$

3 Main theorem

In this section we state the main theorem of this paper. We first mention the existence of stationary solution to the compressible Navier-Stokes equation. Let us consider the following stationary problem:

$$\operatorname{div}(\rho v) = 0, \quad (3.1)$$

$$\rho(v \cdot \nabla)v - \nu \Delta v - \tilde{\nu} \nabla \operatorname{div} v + \nabla P(\rho) = \rho g, \quad (3.2)$$

$$v|_{\Sigma_n} = 0, \quad \rho|_{\Sigma_{j,-}} = \rho|_{\Sigma_{j,+}}, \quad v|_{\Sigma_{j,-}} = v|_{\Sigma_{j,+}} \quad (j = 1, \dots, n-1), \quad (3.3)$$

$$\langle \rho \rangle = 1. \quad (3.4)$$

Then it holds the following assertion.

Proposition 3.1. *Let ν and $\tilde{\nu}$ satisfy (2.4). Then there exist constants $\nu_0 > 0$ and $\tilde{\gamma}_0 > 0$ such that if $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$, there exists a solution $u_s = {}^\top(\rho_s, v_s) = {}^\top(1 + \phi_s, v_s) \in H_{per}^3 \times (H_{per}^4 \cap H_{0,per}^1)$ to problem (3.1)-(3.4), and u_s satisfies*

$$\frac{\gamma^4}{\nu} \|\phi_s\|_{H^3}^2 + \nu \|v_s\|_{H^4}^2 \leq \frac{C_0}{\nu} \quad (3.5)$$

for some constant $C_0 > 0$ depending only on ν_* and Ω_{per} . Furthermore, solution of (3.1)-(3.4) satisfying (3.5) is unique.

The unique existence of stationary solution u_s in Proposition 3.1 was proved by Valli [25]. Estimate (3.5) can be obtained by looking carefully at the computations given in [25]. One can also prove Proposition 3.1 by applying Theorem 5.1 together with estimate (5.6)_k with $k = 3$.

In this paper we consider the linearized stability of the stationary solution $u_s = {}^\top(\rho_s, v_s)$ obtained in Proposition 3.1.

The linearized equation around the stationary solution u_s is written as

$$\partial_t u + Lu = 0 \quad (3.6)$$

for $u = {}^\top(\phi, w) \in D(L)$, where $L : L^2(\Omega) \rightarrow L^2(\Omega)$ is the operator defined by

$$L = \begin{pmatrix} \operatorname{div}(\cdot v_s) & \gamma^2 \operatorname{div}(\rho_s \cdot) \\ \nabla \left(\frac{P'(\rho)}{\gamma^2 \rho_s} \cdot \right) & -\frac{\nu}{\rho_s} \Delta - \frac{\tilde{\nu}}{\rho_s} \nabla \operatorname{div} \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ \frac{1}{\gamma^2 \rho_s^2} (\nu \Delta v_s + \tilde{\nu} \nabla \operatorname{div} v_s) & v_s \cdot \nabla + {}^\top(\nabla v_s) \end{pmatrix}$$

with domain

$$D(L) = \{u = {}^\top(\phi, w) \in L^2(\Omega); w \in H_0^1(\Omega), Lu \in L^2(\Omega)\}.$$

Here ∇v_s is $n \times n$ -matrix whose (j, k) component is $\partial_{x_k} v_s^j$.

As in [11], we see that $-L$ generates a C_0 -semigroup e^{-tL} . The semigroup e^{-tL} has the following properties.

Theorem 3.2. *There exist constants $\nu_0 > 0$ and $\tilde{\gamma}_0 > 0$ such that if $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$, there exists a bounded projection $\Pi : L^2(\Omega) \rightarrow L^2(\Omega)$ satisfying $\Pi L \subset L \Pi$ and $\Pi e^{-tL} = e^{-tL} \Pi$, where $e^{-tL} \Pi$ satisfies*

$$\|e^{-tL} \Pi u_0\|_{L^2(\Omega)} \leq C(1+t)^{-\frac{n-1}{4}} \|u_0\|_{L^1(\Omega)}, \quad (3.7)$$

$$\|e^{-tL} \Pi u_0 - [e^{-tH} \sigma_0] u^{(0)}\|_{L^2(\Omega)} \leq C t^{-\frac{n-1}{4} - \frac{1}{2}} \|u_0\|_{L^1(\Omega)} \quad (3.8)$$

for $u_0 \in L^1(\Omega) \cap L^2(\Omega)$, where $H : L^2(\mathbb{R}^{n-1}) \rightarrow L^2(\mathbb{R}^{n-1})$ is the linear operator defined by

$$D(H) = H^2(\mathbb{R}^{n-1}),$$

$$H\sigma = - \sum_{j,k=1}^{n-1} a_{jk} \partial_{x_j} \partial_{x_k} \sigma + \sum_{j=1}^{n-1} a_j \partial_{x_j} \sigma \quad (\sigma \in D(H)),$$

and $\sigma_0(x')$ is given by $\sigma_0(x') = \frac{|Q|}{|\Omega_{per}|} \int_{\omega_1(x')}^{\omega_2(x')} \phi_0(x', x_n) dx_n$. Here a_j, a_{jk} ($j, k = 1, \dots, n-1$) are real constants and the matrix $(a_{jk})_{j,k=1}^{n-1}$ satisfies $\sum_{j,k=1}^{n-1} a_{jk} \eta_j \eta_k \geq \frac{\kappa_0 \gamma^2}{\nu} |\eta'|^2$ for all $\eta' \in \mathbb{R}^{n-1}$, where κ_0 is a constant independent of $\nu, \tilde{\nu}$ and γ . Moreover, $u^{(0)} = {}^\top(\phi^{(0)}(x', x_n), w^{(0)}(x', x_n))$ is Q -periodic function in x' that satisfies

$$\begin{cases} \operatorname{div}(\phi^{(0)} v_s) + \gamma^2 \operatorname{div}(\rho_s w^{(0)}) = 0, & \text{in } \Omega_{per} \\ \nabla \left(\frac{P'(\rho_s)}{\gamma^2 \rho_s} \phi^{(0)} \right) + \frac{1}{\gamma^2 \rho_s^2} (\nu \Delta v_s + \tilde{\nu} \nabla \operatorname{div} v_s) \phi^{(0)} \\ \quad - \frac{\nu}{\rho_s} \Delta w^{(0)} - \frac{\tilde{\nu}}{\rho_s} \nabla \operatorname{div} w^{(0)} + v_s \cdot \nabla w^{(0)} + w^{(0)} \cdot \nabla v_s = 0, & \text{in } \Omega_{per} \\ w^{(0)}|_{\Sigma_n} = 0, \\ \int_{\Omega_{per}} \phi^{(0)} = 1. \end{cases}$$

As for the $(I - \Pi)$ -part, we can prove the following exponential decay estimates in a similar manner to [1, 2, 15].

We denote by $\tilde{H}^1(\Omega)$ the set of all locally H^1 functions in Ω whose tangential derivatives near $\partial\Omega$ belong to $L^2(\Omega)$.

Theorem 3.3. (i) Let $u_0 = {}^\top(\phi_0, w_0) \in H^1(\Omega) \times \tilde{H}^1(\Omega)$. Suppose that ν , $\tilde{\nu}$ and γ satisfy the assumption of Theorem 3.2. Then

$$e^{-tL}(I - \Pi)u_0 \in C([0, \infty); H^1(\Omega) \times \tilde{H}^1(\Omega)) \cap C((0, \infty); H^1(\Omega) \times H_0^1(\Omega))$$

and

$$\|e^{-tL}(I - \Pi)u_0\|_{H^1(\Omega)}^2 \leq Ce^{-at} \{ \|u_0\|_{H^1(\Omega) \times \tilde{H}^1(\Omega)}^2 + t^{-\frac{1}{2}} \|w_0\|_{L^2(\Omega)} \} \quad (3.9)$$

for all $t \geq 0$.

(ii) Let $T > 0$. Suppose that $u_0 \in H^1(\Omega) \times H_0^1(\Omega)$ and $f \in L^2(0, T; H^1(\Omega) \times L^2(\Omega))$. Then there exists a unique solution $u(t)$ of

$$\begin{aligned} \partial_t u + Lu &= (I - \Pi)f, & u|_{t=0} &= (I - \Pi)u_0, \\ u &\in C([0, T]; H^1(\Omega) \times H_0^1(\Omega)) \cap L^2(0, T; H^1(\Omega) \times H^2(\Omega)), \\ u(t) &\in (I - \Pi)L^2(\Omega) & \text{for } 0 \leq t \leq T, \end{aligned}$$

and $u(t)$ satisfies

$$\begin{aligned} \|u(t)\|_{H^1(\Omega)}^2 &+ \int_0^t e^{-a(t-\tau)} \|u(\tau)\|_{H^1(\Omega) \times H^2(\Omega)}^2 d\tau \\ &\leq C \left\{ e^{-at} \|u_0\|_{H^1(\Omega)}^2 + \int_0^t e^{-a(t-\tau)} \|f\|_{H^1(\Omega) \times L^2(\Omega)}^2 d\tau \right\} \end{aligned} \quad (3.10)$$

for $t \in [0, T]$, where a and C are positive constants independent of T .

Theorem 3.3 can be proved in a similar way as in [1, 2, 15], so we omit the proof. In the remaining of this paper we give a proof of Theorem 3.2.

4 Proof of Theorem 3.2

In this section we give a proof of Theorem 3.2. We first reduce the problem to the one on Ω_{per} by the Bloch transformation. We then investigate the reduced problems depending on the values of Bloch parameter η' . In subsection 4.2 we consider the spectrum for the case $\eta' = 0$; we then develop a perturbation argument to investigate the spectrum for $|\eta'| \ll 1$ in subsection 4.3. In subsection 4.4 we establish estimates for the eigenprojections for the eigenvalues near the imaginary axis. We finally give a proof of Theorem 3.2.

Throughout this section we assume that ν and $\tilde{\nu}$ satisfy (2.4), and thus, we use relation (2.5).

4.1 Formulation

Let us consider the resolvent problem for (3.6)

$$(\lambda + L)u = f, \quad u = {}^\top(\phi, w) \in D(L). \quad (4.1)$$

Here $f = {}^\top(f_0, \tilde{f})$, $\tilde{f} = {}^\top(f_1, \dots, f_n)$ and $\lambda \in \mathbb{C}$ is a resolvent parameter. As in [15], applying the Bloch transform T to (4.1), we have

$$(\lambda + L_{\eta'})Tu = Tf \quad \text{on } \Omega_{per} \quad (4.2)$$

with the dual parameter $\eta' \in Q'$, where $L_{\eta'}$ is the operator on L_{per}^2 of the form

$$\begin{aligned} L_{\eta'} = & \begin{pmatrix} {}^\top \nabla_{\eta'}(\cdot v_s) & \gamma^2 {}^\top \nabla_{\eta'}(\rho_s \cdot) \\ \nabla_{\eta'}\left(\frac{P'(\rho_s)}{\gamma^2 \rho_s} \cdot\right) & -\frac{\nu}{\rho_s} \Delta_{\eta'} - \frac{\tilde{\nu}}{\rho_s} \nabla_{\eta'} {}^\top \nabla_{\eta'} \end{pmatrix} \\ & + \begin{pmatrix} 0 & 0 \\ \frac{1}{\gamma^2 \rho_s^2}(\nu \Delta v_s + \tilde{\nu} \nabla \operatorname{div} v_s) & v_s \cdot \nabla_{\eta'} + {}^\top \nabla v_s \end{pmatrix}. \end{aligned}$$

with domain $D(L_{\eta'})$

$$D(L_{\eta'}) = \{u = {}^\top(\phi, w) \in L_{per}^2; w \in H_{0,per}^1, L_{\eta'}u \in L_{per}^2\}.$$

Here we define $\nabla_{\eta'}$ and $\Delta_{\eta'}$ by

$$\nabla_{\eta'} = \nabla + i\tilde{\eta}', \quad \Delta_{\eta'} = \nabla_{\eta'} \cdot \nabla_{\eta'},$$

where $\tilde{\eta}' = (\eta', 0)$. It is not difficult to see that $D(L_{\eta'}) = D(L_0)$ for all $\eta' \in Q'$ and that $L_{\eta'}$ is a closed operator on L_{per}^2 .

We write $L_{\eta'}$ in the form

$$L_{\eta'} = L_0 + \sum_{j=1}^{n-1} \eta_j L_j^{(1)} + \sum_{j,k=1}^{n-1} \eta_j \eta_k L_{jk}^{(2)},$$

where

$$\begin{aligned} L_0 := & \begin{pmatrix} \operatorname{div}(\cdot v_s) & \gamma^2 \operatorname{div}(\rho_s \cdot) \\ \nabla\left(\frac{P'(\rho_s)}{\gamma^2 \rho_s} \cdot\right) & -\frac{\nu}{\rho_s} \Delta - \frac{\tilde{\nu}}{\rho_s} \nabla \operatorname{div} \end{pmatrix} \\ & + \begin{pmatrix} 0 & 0 \\ \frac{1}{\gamma^2 \rho_s^2}(\nu \Delta v_s + \tilde{\nu} \nabla \operatorname{div} v_s) & v_s \cdot \nabla + {}^\top \nabla v_s \end{pmatrix}, \\ L_j^{(1)} := & i \begin{pmatrix} v_s^j & \gamma^2 \rho_s {}^\top \mathbf{e}_j \\ \left(\frac{P'(\rho_s)}{\gamma^2 \rho_s} \cdot\right) \mathbf{e}_j & -\frac{1}{\rho_s} \left(2\nu \mathbf{e}_j \otimes \mathbf{e}_j \partial_{x_j} + \tilde{\nu} \mathbf{e}_j \operatorname{div} - \tilde{\nu} \nabla({}^\top \mathbf{e}_j)\right) + v_s^j \end{pmatrix}, \end{aligned}$$

$$L_{jk}^{(2)} := \begin{pmatrix} 0 & 0 \\ 0 & \frac{1}{\rho_s} \left(\nu \delta_{jk} I_n + \tilde{\nu} \mathbf{e}_j^\top \mathbf{e}_k \right) \end{pmatrix}.$$

We also set

$$M_{\eta'} = \sum_{j=1}^n \eta_j L_j^{(1)} + \sum_{j,k=1}^n \eta_j \eta_k L_{jk}^{(2)},$$

i.e.,

$$M_{\eta'} = \begin{pmatrix} i\tilde{\eta}' \cdot v_s & i\gamma^2 \rho_s^\top \tilde{\eta}' \\ i\frac{P'(\rho_s)}{\gamma^2 \rho_s} \tilde{\eta}' & \frac{\nu}{\rho_s} \left(|\eta'|^2 - 2i\tilde{\eta}' \cdot \nabla \right) I_n - i\frac{\tilde{\nu}}{\rho_s} \tilde{\eta}'^\top \nabla_{\eta'} - i\frac{\tilde{\nu}}{\rho_s} \nabla^\top \tilde{\eta}' + (v_s \cdot i\tilde{\eta}') I_n \end{pmatrix}.$$

By (4.2), if $\lambda \in \rho(-L_{\eta'})$ for all $\eta' \in Q^*$, then we have

$$u = U(\lambda + L_{\eta'})^{-1} T f \quad \text{in } \Omega.$$

Therefore, to investigate the resolvent of $-L$, we will consider the problem for $-L_{\eta'}$:

$$\lambda u + L_{\eta'} u = f, \quad u \in D(L_{\eta'}). \quad (4.3)$$

4.2 The null space of L_0

In this subsection we investigate the null space of L_0 . We first show that 0 is an eigenvalue and $\dim \text{Ker}(L_0) = 1$.

Proposition 4.1. *There exist constants $\nu_0 > 0$ and $\tilde{\gamma}_0 > 0$ such that if $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$, then there exists a unique $u^{(0)} = {}^\top(\phi^{(0)}, w^{(0)}) \in D(L_0)$ satisfying $L_0 u^{(0)} = 0$ and $\langle \phi^{(0)} \rangle = 1$. Furthermore, $u^{(0)} = {}^\top(\phi^{(0)}, w^{(0)})$ satisfies*

$$\|\phi^{(0)} - 1\|_{H^2}^2 + \nu^2 \|w^{(0)}\|_{H^3}^2 \leq \frac{C}{\gamma^4} \quad (4.4)$$

uniformly for ν , $\tilde{\nu}$ and γ with $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$.

Proof. We consider the problem $L_0 u = 0$ with $u = {}^\top(\phi, w) \in D(L_0)$. Suppose that $\phi = 0$. Then taking the inner product of $L_0 u = 0$ with u we have

$$\begin{aligned} \nu \|\nabla w\|_2^2 + \tilde{\nu} \|\text{div} w\|_2^2 &= -\text{Re} \{ (v_s \cdot \nabla w + w \cdot \nabla v_s, \rho_s w) \} \\ &\leq C \|v_s\|_{C^1} \|\rho_s\|_\infty \|\nabla w\|_2^2. \end{aligned}$$

Therefore, if $\nu > C \|v_s\|_{C^1} \|\rho_s\|_\infty$, then $w = 0$, and hence, $u = 0$.

In what follows we assume that $\nu > C\|v_s\|_{C^1}\|\rho_s\|_\infty$. We will show that there exists a function $u = {}^\top(\phi, w)$ with $\phi \neq 0$ satisfying $L_0 u = 0$ if ν and $\frac{\gamma^2}{\nu}$ are sufficiently large. We may assume that $\langle \phi \rangle = 1$, and hence, we consider the problem

$$\begin{cases} L_0 u = 0, & u = {}^\top(\phi, w) \in D(L_0), \\ \langle \phi \rangle = 1. \end{cases} \quad (4.5)$$

To solve (4.5), we introduce the operator $\mathcal{A}[v] : H_{per,*}^2 \times H_{per}^1 \longrightarrow H_{per,*}^2 \times H_{per}^1$ defined by

$$\mathcal{A}[v] = \begin{pmatrix} \operatorname{div}(\cdot v) & \gamma^2 \operatorname{div} \\ \nabla & -\nu \Delta - \tilde{\nu} \nabla \operatorname{div} \end{pmatrix}$$

with domain

$$D(\mathcal{A}[v]) = \{u \in H_{per,*}^2 \times (H_{per}^3 \cap H_{0,per}^1); \mathcal{A}[v]u \in H_{per,*}^2 \times H_{per}^1\}.$$

We set $\phi = 1 + \tilde{\phi}$. Then (4.5) is written as

$$\mathcal{A}[v_s]\tilde{u} + B\tilde{u} = F. \quad (4.6)$$

Here $\tilde{u} = {}^\top(\tilde{\phi}, w) \in D(\mathcal{A}[v_s])$, $Bu = {}^\top((B\tilde{u})_1, (B\tilde{u})_2)$ and $F = {}^\top(F_0, \tilde{F})$, where

$$\begin{aligned} (B\tilde{u})_1 &= -\gamma^2 \operatorname{div}(\phi_s w), \\ (B\tilde{u})_2 &= -\frac{\phi_s}{1 + \phi_s}(\nu \Delta w + \tilde{\nu} \nabla \operatorname{div} w) - \nabla(\tilde{P}^{(1)}(\phi_s)\phi_s\tilde{\phi}) \\ &\quad - \frac{1}{\gamma^2} \left(\frac{1}{1 + \phi_s} \right)^2 (\nu \Delta v_s + \tilde{\nu} \nabla \operatorname{div} v_s)\tilde{\phi} - v_s \cdot \nabla w - w \cdot \nabla v_s, \\ F_0 &= -\operatorname{div} v_s, \\ \tilde{F} &= -\nabla(\tilde{P}^{(1)}(\phi_s)\phi_s) - \frac{1}{\gamma^2} \left(\frac{1}{1 + \phi_s} \right)^2 (\nu \Delta v_s + \tilde{\nu} \nabla \operatorname{div} v_s), \\ \tilde{P}^{(1)}(\phi_s) &= \int_0^1 \tilde{P}'(1 + \theta\phi_s) d\theta, \quad \tilde{P}(\rho) = \frac{P'(\rho)}{\gamma^2 \rho}. \end{aligned}$$

We set

$$\|\tilde{u}\|_{H^2 \times H^3}^* = \|\tilde{\phi}\|_{H^2} + \nu \|w\|_{H^3}$$

for $\tilde{u} = {}^\top(\tilde{\phi}, w) \in H_{per,*}^2 \times H_{per}^3$. It follows from Proposition 3.1 and (5.6)_k with $k = 2$ that if ν and $\frac{\gamma^2}{\nu}$ are sufficiently large, then

$$\|\mathcal{A}[v_s]^{-1}F\|_{H^2 \times H^3}^* \leq C \left\{ \frac{\nu}{\gamma^2} \|f^0\|_{H^2} + \|\tilde{f}\|_{H^1} \right\} \quad (4.7)$$

for $F = {}^\top(f^0, \tilde{f}) \in H_{per,*}^2 \times H_{per}^1$. By Proposition 3.1, we have

$$\begin{aligned} \|(B\tilde{u})_1\|_{H^2} &\leq C\|w\|_{H^3}, \\ \|(B\tilde{u})_2\|_{H^1} &\leq C\left\{\frac{1}{\gamma^2}\|\tilde{\phi}\|_{H^2} + \left(\frac{\nu}{\gamma^2} + \frac{1}{\nu}\right)\|w\|_{H^3}\right\} \end{aligned}$$

for $\tilde{u} = {}^\top(\tilde{\phi}, w) \in D(\mathcal{A}[v_s])$. Combining these estimates with (4.7), we see that there are positive numbers ν_0 and $\tilde{\gamma}_0$ such that if $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$, then $\|\mathcal{A}[v_s]^{-1}B\tilde{u}\|_{H^2 \times H^3}^* \leq \frac{1}{2}\|\tilde{u}\|_{H^2 \times H^3}^*$. Therefore, there exists a unique $\tilde{u} \in D(\mathcal{A}[v_s])$ satisfying $\tilde{u} = -\mathcal{A}[v_s]^{-1}B\tilde{u} + \mathcal{A}[v_s]^{-1}F$, that is, a unique solution $\tilde{u} \in D(\mathcal{A}[v_s])$ of (4.6), and $\tilde{u}$ satisfies the estimate

$$\|\tilde{u}\|_{H^2 \times H^3}^* \leq C\left\{\frac{\nu}{\gamma^2}\|v_s\|_{H^4} + \|\phi_s\|_{H^3}\right\} \leq \frac{C}{\gamma^2}.$$

This completes the proof. $\square$

We next consider the eigenspace for the eigenvalue 0. We introduce the adjoint operator $L_{\eta'}^* : L_{per}^2 \rightarrow L_{per}^2$ defined by

$$\begin{aligned} L_{\eta'}^* = & \begin{pmatrix} -v_s \cdot \nabla_{\eta'} \left(\frac{P'(\rho_s)}{\rho_s} \cdot \right) \frac{\rho_s}{P'(\rho_s)} & -\gamma^2 \nabla_{\eta'} \cdot (\rho_s \cdot) \\ -\nabla_{\eta'} \left(\frac{\rho_s}{\gamma^2 \rho_s} \cdot \right) & -\frac{\nu}{\rho_s} \Delta_{\eta'} - \frac{\tilde{\nu}}{\rho_s} \nabla_{\eta'} ({}^\top \nabla_{\eta'}) \end{pmatrix} \\ & + \begin{pmatrix} 0 & \frac{\gamma^2}{P'(\rho_s)} (\nu \Delta^\top v_s + \tilde{\nu}^\top \nabla \operatorname{div} v_s) \\ 0 & -\operatorname{div} v_s - \frac{1}{\rho_s} v_s \cdot \nabla_{\eta'} (\rho_s \cdot) + \nabla_{\eta'} v_s \end{pmatrix} \end{aligned}$$

with domain

$$D(L_{\eta'}^*) = \{u = {}^\top(\phi, w) \in L_{per}^2; w \in H_{0,per}^1, L_{\eta'}^* u \in L_{per}^2\}.$$

Then it holds that $D(L_{\eta'}^*) = D(L_0^*)$ and $\langle L_{\eta'} u_1, u_2 \rangle = \langle u_1, L_{\eta'}^* u_2 \rangle$.

Concerning the eigenspace for eigenvalue 0, we have the following

Lemma 4.2. *Let $\Pi^{(0)}$ be defined by*

$$\Pi^{(0)} u = \langle u, u^{(0)*} \rangle u^{(0)} = \langle \phi \rangle u^{(0)}$$

for $u = {}^\top(\phi, w)$, where

$$u^{(0)*} = \gamma^2 \begin{pmatrix} \frac{\gamma^2 \rho_s}{P'(\rho_s)} \\ 0 \end{pmatrix}.$$

Then the following assertions hold.

(i) $u^{(0)*}$ satisfies $u^{(0)*} \in D(L_0^*)$, $L_0^* u^{(0)*} = 0$ and $\langle u^{(0)}, u^{(0)*} \rangle = 1$.

(ii) $\Pi^{(0)}$ is a bounded projection on L_{per}^2 satisfying

$$\Pi^{(0)} L_0 \subset L_0 \Pi^{(0)} = 0, \quad \Pi^{(0)} u^{(0)} = u^{(0)} \quad \text{and} \quad \Pi^{(0)} L_{per}^2 = \text{span}\{u^{(0)}\}.$$

The assertions of Lemma 4.2 are easily verified, and so we omit the proof. We introduce some notation. In what follows we set

$$X_0 = \Pi^{(0)} L_{per}^2, \quad X_1 = (I - \Pi^{(0)}) L_{per}^2.$$

Observe that $u = {}^\top(\phi, w) \in X_1$ if and only if $\langle \phi \rangle = 0$.

Lemma 4.3. *If $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$ (by taking ν_0 and $\tilde{\gamma}_0$ suitably larger if necessary), then $\rho(-L_0|_{X_1}) \supset \{\lambda; \text{Re}\lambda \geq -\beta_0\}$ for some positive constant β_0 . Furthermore, if $\text{Re}\lambda \geq -\beta_0$, then*

$$\|((\lambda + L_0)|_{X_1})^{-1} f\|_2 \leq \frac{C}{\text{Re}\lambda + 2\beta_0} \|f\|_2, \quad (4.8)$$

$$\|\nabla \tilde{Q}((\lambda + L_0)|_{X_1})^{-1} f\|_2 \leq C \frac{1}{(\text{Re}\lambda + 2\beta_0)^{\frac{1}{2}}} \|f\|_2. \quad (4.9)$$

Proof. We first establish estimates (4.8) and (4.9). For $f = {}^\top(f^0, \tilde{f}) \in X_1$, we consider

$$(\lambda + L_0)u = f, \quad u = {}^\top(\phi, w) \in D(L_0) \cap X_1. \quad (4.10)$$

Note that $\langle \phi \rangle = \langle f^0 \rangle = 0$. We compute

$$\begin{aligned} \text{Re}\langle (\lambda + L_0)u, u \rangle &= \text{Re}\lambda \langle u, u \rangle + \text{Re}\langle L_0 u, u \rangle \\ &\quad - \delta \text{Re}\{((L_0 u)_2, \mathcal{B}\phi) + (\mathcal{B}(L_0 u)_1, w)\}, \end{aligned}$$

where $(u)_1 = \phi$ and $(u)_2 = w$ for $u = {}^\top(\phi, w)$. Since

$$\begin{aligned} -\text{Re}((L_0 u)_2, \mathcal{B}\phi) &\geq \left\| \sqrt{\frac{P'(\rho_s)}{\gamma^2 \rho_s}} \phi \right\|_2^2 - C\{\nu \|\nabla w\|_2 + \tilde{\nu} \|\text{div} w\|_2\} \|\phi\|_2 \\ &\quad - C \frac{\nu}{\gamma^2} \|v_s\|_{H^4} \|\phi\|_2^2 - C \|v_s\|_{H^4} \|\phi\|_2 \|w\|_2, \\ -\text{Re}(\mathcal{B}(L_0 u)_1, w) &\geq -\|\mathcal{B}(\text{div}(\phi v_s + \gamma^2 \rho_s w))\|_2 \|w\|_2 \\ &\geq -C\{\|v_s\|_{H^4} \|\phi\|_2 \|w\|_2 + \gamma^2 \|w\|_2^2\}, \end{aligned}$$

we have

$$\begin{aligned}
\operatorname{Re} \langle (\lambda + L_0)u, u \rangle &\geq \operatorname{Re} \lambda \langle u, u \rangle + \nu \|\nabla w\|_2^2 + \tilde{\nu} \|\operatorname{div} w\|_2^2 + \delta \left\| \sqrt{\frac{P'(\rho_s)}{\gamma^2 \rho_s}} \phi \right\|_2^2 \\
&\quad - C \|v_s\|_{H^4} \left\| \sqrt{\frac{P'(\rho_s)}{\gamma^4 \rho_s}} \phi \right\|_2^2 - \frac{C}{\gamma^2} \|v_s\|_{H^4} \|\phi\|_2^2 \\
&\quad - C \frac{\nu}{\gamma^2} \|v_s\|_{H^4} \|\phi\|_2 \|w\|_2 - C \|v_s\|_{H^4} \|w\|_2^2 \\
&\quad - C \delta \{ (\nu \|\nabla w\|_2 + \tilde{\nu} \|\operatorname{div} w\|_2) \|\phi\|_2 + \frac{\nu}{\gamma^2} \|v_s\|_{H^4} \|\phi\|_2^2 \\
&\quad + \|v_s\|_{H^4} \|\phi\|_2 \|w\|_2 + \gamma^2 \|w\|_2^2 \}.
\end{aligned}$$

Therefore, by using Proposition 3.1, we see that there exist constants $\nu_0 > 0$, $\tilde{\gamma}_0 > 0$ and $C_0 > 0$ such that if $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$, then it holds with $\delta = C_0 \min\{\frac{1}{\nu}, \frac{\nu}{\gamma^2}, \frac{1}{\gamma}\}$ that

$$\operatorname{Re} \langle (\lambda + L_0)u, u \rangle \geq \operatorname{Re} \lambda \langle u, u \rangle + \frac{\nu}{2} \|\nabla w\|_2^2 + \frac{\tilde{\nu}}{2} \|\operatorname{div} w\|_2^2 + \frac{\delta}{2} \|\phi\|_2^2.$$

Since $\operatorname{Re} \langle f, u \rangle \leq \|f\|_2 \|u\|_2 + \delta \|f\|_2 \|u\|_2 \leq C(1 + \delta) \|f\|_2 \|u\|_2$, we have

$$\operatorname{Re} \lambda \langle u, u \rangle + \frac{\nu}{2} \|\nabla w\|_2^2 + \frac{\tilde{\nu}}{2} \|\operatorname{div} w\|_2^2 + \frac{\delta}{2} \|\phi\|_2^2 \leq C(1 + \delta) \|f\|_2 \|u\|_2.$$

By the Poincaré inequality, we have

$$\operatorname{Re} \lambda \|u\|_2^2 + c_1(\nu \|w\|_2^2 + \delta \|\phi\|_2^2) + \frac{\nu}{4} \|\nabla w\|_2^2 \leq C(1 + \delta) \|f\|_2 \|u\|_2. \quad (4.11)$$

It follows that there exists $\beta_0 > 0$ with $\beta_0 = O(\min\{\nu, \frac{\nu}{\gamma^2}, \frac{1}{\nu}, \frac{1}{\gamma}\})$ such that

$$(\operatorname{Re} \lambda + 2\beta_0) \|u\|_2^2 + \nu \|\nabla w\|_2^2 \leq C \|f\|_2 \|u\|_2.$$

We thus have

$$\|u\|_2 \leq \frac{C}{\operatorname{Re} \lambda + 2\beta_0} \|f\|_2, \quad (4.12)$$

and so if $\operatorname{Re} \lambda \geq -\beta_0$, then

$$\|\nabla w\|_2 \leq \frac{C}{\nu^{\frac{1}{2}} (\operatorname{Re} \lambda + 2\beta_0)^{\frac{1}{2}}} \|f\|_2. \quad (4.13)$$

If we prove the existence of solution $u \in D(L_0) \cap X_1$ of (4.10) for all $f \in X_1$ with $\lambda \gg 1$, then we see from (4.12) and (4.13) that $\rho(-L_0|_{X_1}) \supset$

$\{\lambda; \operatorname{Re}\lambda \geq -\beta_0\}$ by a standard continuation argument. To prove the existence of solution of (4.10), we first consider the problem

$$\lambda\phi + \operatorname{div}(\phi v_s) = F^0, \quad F^0 = f^0 - \gamma^2 \operatorname{div}(\rho_s \tilde{w}) \quad (4.14)$$

for a given $\tilde{w} \in H_{0,per}^1$ and $F^0 \in L_{per,*}^2$. Applying [10, Theorem 1], we see that there exists a constant λ_0 satisfying $\lambda_0 \geq C\|v\|_{H^3}$ such that if $\lambda \geq \lambda_0$, there exists a solution ϕ of (4.14). Furthermore, ϕ satisfies

$$\|\phi\|_2 \leq \frac{C}{\lambda} \|F^0\|_2 \leq \frac{C}{\lambda} \{\|f^0\|_2 + \|\tilde{w}\|_{H^1}\}. \quad (4.15)$$

We denote this solution ϕ by $\phi(\tilde{w})$. We next consider the problem

$$\begin{cases} \lambda w - \frac{\nu}{\rho_s} \Delta w - \frac{\tilde{\nu}}{\rho_s} \nabla \operatorname{div} w = \tilde{F}, \\ w|_{\Sigma_{j,-}} = w|_{\Sigma_{j,+}}, \quad w|_{\Sigma_n} = 0. \end{cases} \quad (4.16)$$

where

$$\tilde{F} = \tilde{f} - \nabla \left(\frac{P'(\rho_s)}{\gamma^2 \rho_s} \phi \right) - \frac{1}{\gamma^2 \rho_s^2} (\nu \Delta v_s + \tilde{\nu} \operatorname{div} v_s) - v_s \cdot \nabla \tilde{w} - \tilde{w} \cdot \nabla v_s,$$

with $\phi = \phi(\tilde{w})$. Since $\operatorname{Re}(-\frac{\nu}{\rho_s} \Delta w - \frac{\tilde{\nu}}{\rho_s} \nabla \operatorname{div} w, \rho_s w) = \nu \|\nabla w\|_2^2 + \tilde{\nu} \|\operatorname{div} w\|_2^2$, there exists a unique solution $w \in H_{0,per}^1$ of (4.16) such that

$$\lambda \|w\|_2^2 + \|\nabla w\|_2^2 \leq C \|\tilde{F}\|_{H^{-1}}^2 \leq C \{\|\tilde{f}\|_2^2 + \frac{1}{\lambda} \|f^0\|_2^2 + \frac{1}{\lambda} \|\nabla \tilde{w}\|_2^2 + \|\tilde{w}\|_2^2\} \quad (4.17)$$

for $\lambda \geq \lambda_0$. We define the operator $\Gamma : H_{0,per}^1 \rightarrow H_{0,per}^1$ by the solution operator for (4.16), i.e., $\Gamma(\tilde{w}) = w$ for $\tilde{w} \in H_{0,per}^1$, with w being the solution of (4.16).

We introduce the norm $\|\cdot\|_{(\lambda)} = (\lambda \|\cdot\|_2^2 + \|\nabla \cdot\|_2^2)^{\frac{1}{2}}$ of $H_{0,per}^1$. By (4.2), Γ satisfies $\|\Gamma(\tilde{w}_1) - \Gamma(\tilde{w}_2)\|_{(\lambda)} \leq \frac{C}{\lambda} \|\tilde{w}_1 - \tilde{w}_2\|_{(\lambda)}$ for $\tilde{w}_1, \tilde{w}_2 \in H_{0,per}^1$. Let $\lambda \geq \lambda_1 = \max\{2C, \lambda_0\}$. It then follows that $\|\Gamma(\tilde{w}_1) - \Gamma(\tilde{w}_2)\|_{(\lambda)} \leq \frac{1}{2} \|\tilde{w}_1 - \tilde{w}_2\|_{(\lambda)}$, and Γ is a contraction mapping on $H_{0,per}^1$. Therefore, there exists a unique $w \in H_{0,per}^1$ satisfying $\Gamma(w) = w$. For this fixed point w , we set $\phi = \phi(w)$. It then follows that $u = {}^\top(\phi, w) \in D(L_0|_{X_1})$ is a solution of (4.10) with $\lambda \geq \lambda_1$. We thus conclude that $\rho(-L_0|_{X_1}) \supset \{\lambda; \operatorname{Re}\lambda > -\beta_0\}$. This completes the proof. $\square$

Theorem 4.4. *If $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$ (by taking ν_0 and $\tilde{\gamma}_0$ suitably larger if necessary), then*

- (i) $X_1 = \operatorname{R}(L_0)$ and $X_0 = \operatorname{Ker}(L_0)$,

$$(ii) \quad L_{per}^2 = \text{Ker}(L_0) \oplus \text{R}(L_0),$$

(iii) 0 is a simple eigenvalue of $-L_0$.

Proof. Let $f = {}^\top(f^0, \tilde{f}) \in \text{R}(L_0)$. There exists a function $u \in D(L_0)$ such that $L_0 u = f$. Applying $\Pi^{(0)}$, we have $0 = \Pi^{(0)} L_0 u = \Pi^{(0)} f = \langle f^0 \rangle u^{(0)}$. This implies $\langle f^0 \rangle = 0$, and hence, $f \in X_1$. This shows $\text{R}(L_0) \subset X_1$. Let $f = {}^\top(f^0, \tilde{f}) \in X_1$. By Lemma 4.3, there exists a unique function $u = {}^\top(\phi, w) \in D(L_0)$ satisfying $L_0 u = f$ and $\langle \phi \rangle = 0$. This shows $X_1 \subset \text{R}(L_0)$, and (i) is proved. (ii) follows from (i), and (iii) follows from and Lemma 4.2. This completes the proof. $\square$

Theorem 4.5. *If $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$ (by taking ν_0 and $\tilde{\gamma}_0$ suitably larger if necessary), then $\rho(-L_0) \supset \{\lambda; \text{Re} \lambda \geq -\beta_0\} \setminus \{0\}$. Furthermore, if $\text{Re} \lambda \geq -\beta_0$ and $\lambda \neq 0$, then*

$$\begin{aligned} (\lambda + L_0)^{-1} f &= \frac{1}{\lambda} \Pi^{(0)} f + ((\lambda + L_0)|_{X_1})^{-1} \Pi_1 f, \\ \|(\lambda + L_0)^{-1} f\|_2 &\leq C \left(\frac{1}{|\lambda|} + \frac{1}{\text{Re} \lambda + 2\beta_0} \right) \|f\|_2, \\ \|\nabla \tilde{Q}(\lambda + L_0)^{-1} f\|_2 &\leq C \left(\frac{1}{|\lambda|} + \frac{1}{(\text{Re} \lambda + 2\beta_0)^{\frac{1}{2}}} \right) \|f\|_2. \end{aligned}$$

Proof. By Theorem 4.4, problem $(\lambda + L_0)u = f$ is equivalent to

$$\begin{cases} \lambda \Pi^{(0)} u = \Pi^{(0)} f, \\ (\lambda + L_0)u_1 = f_1, \end{cases}$$

where $u_1 = (I - \Pi^{(0)})u \in D(L_0) \cap X_1$ and $f_1 = (I - \Pi^{(0)})f \in X_1$. Applying Lemma 4.3 we see that if $\text{Re} \lambda \geq -\beta_0$ and $\lambda \neq 0$, then

$$(\lambda + L_0)^{-1} f = \frac{1}{\lambda} \Pi^{(0)} f + ((\lambda + L_0)|_{X_1})^{-1} f_1,$$

and

$$\begin{aligned} \|(\lambda + L_0)^{-1} f\|_2 &\leq C \left(\frac{1}{|\lambda|} + \frac{1}{\text{Re} \lambda + 2\beta_0} \right) \|f\|_2, \\ \|\nabla \tilde{Q}(\lambda + L_0)^{-1} f\|_2 &\leq C \left(\frac{1}{|\lambda|} + \frac{1}{(\text{Re} \lambda + 2\beta_0)^{\frac{1}{2}}} \right) \|f\|_2. \end{aligned}$$

This completes the proof. $\square$

As for the spectrum of $-L_0^*$, we have the following

Proposition 4.6. Define $\Pi^{(0)*}$ by

$$\Pi^{(0)*}u = \langle u, u^{(0)} \rangle u^{(0)*}.$$

If $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$ (by taking ν_0 and $\tilde{\gamma}_0$ suitably larger if necessary), then the following assertions hold true.

- (i) $\rho(-L_0^*) \supset \{\lambda; \operatorname{Re}\lambda \geq -\beta_0\} \setminus \{0\}$, 0 is a simple eigenvalue of $-L_0^*$,
- (ii) $\Pi^{(0)*}L_0^* \subset L_0^*\Pi^{(0)*} = 0$, $\operatorname{Ker}(L_0^*) = \Pi^{(0)*}L_{per}^2 = \operatorname{span}\{u^{(0)*}\}$,
- (iii) $\operatorname{R}(L_0^*) = (I - \Pi^{(0)*})L_{per}^2$,
- (iv) $L_{per}^2 = \operatorname{Ker}(L_0^*) \oplus \operatorname{R}(L_0^*)$,

Moreover, if λ satisfying $\operatorname{Re}\lambda \geq -\beta_0$ and $\lambda \neq 0$, then

$$\begin{aligned} (\lambda + L_0^*)^{-1}f &= \frac{1}{\lambda}\Pi^{(0)*}f + ((\lambda + L_0^*)|_{(I-\Pi^{(0)*})L_{per}^2})^{-1}(I - \Pi^{(0)*})f, \\ \|(\lambda + L_0^*)^{-1}f\|_2 &\leq C\left(\frac{1}{|\lambda|} + \frac{1}{\operatorname{Re}\lambda + 2\beta_0}\right)\|f\|_2, \end{aligned} \quad (4.18)$$

$$\|\nabla \tilde{Q}(\lambda + L_0^*)^{-1}f\|_2 \leq C\left(\frac{1}{|\lambda|^{\frac{1}{2}}} + \frac{1}{|\lambda|} + \frac{1}{(\operatorname{Re}\lambda + 2\beta_0)^{\frac{1}{2}}} + \frac{1}{\operatorname{Re}\lambda + 2\beta_0}\right)\|f\|_2. \quad (4.19)$$

Proof. By the closed range theorem, we have (i)-(iv). Furthermore, if $\operatorname{Re}\lambda \geq -\beta_0$ and $\lambda \neq 0$, we see from Theorem 4.5 that

$$\begin{aligned} \langle (\lambda + L_0^*)^{-1}f, g \rangle &= \langle f, (\lambda + L_0)^{-1}g \rangle \\ &\leq \|f\|_2 \|(\lambda + L_0)^{-1}g\|_2 \\ &\leq C\left(\frac{1}{|\lambda|} + \frac{1}{\operatorname{Re}\lambda + 2\beta_0}\right)\|f\|_2 \|g\|_2, \end{aligned}$$

and hence,

$$\|(\lambda + L_0^*)^{-1}f\|_2 \leq C\left(\frac{1}{|\lambda|} + \frac{1}{\operatorname{Re}\lambda + 2\beta_0}\right)\|f\|_2. \quad (4.20)$$

Computing $\langle (\lambda + L_0^*)u, u \rangle$, we have

$$\operatorname{Re}\lambda \|u\|_2^2 - c\|u\|_2^2 + \nu\|\nabla w\|_2^2 \leq \|f\|_2 \|u\|_2,$$

where c is a positive constant, and hence, if $\operatorname{Re}\lambda \geq -\beta_0$ and $\lambda \neq 0$, then

$$\nu\|\nabla w\|_2^2 \leq C\left\{\left(\frac{1}{|\lambda|} + \frac{1}{\operatorname{Re}\lambda + 2\beta_0}\right) + \left(\frac{1}{|\lambda|} + \frac{1}{\operatorname{Re}\lambda + 2\beta_0}\right)^2\right\}\|f\|_2^2.$$

This completes the proof. $\square$

4.3 Perturbation argument

In this subsection we develop a perturbation argument to investigate the spectrum of $-L_{\eta'}$ for $|\eta'| \ll 1$.

Proposition 4.7. *If $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$ (by taking ν_0 and $\tilde{\gamma}_0$ suitably larger if necessary), then there exists a constant $r_0 > 0$ such that if $|\eta'| \leq r_0$, then*

$$\Sigma_1 := \left\{ \lambda; \operatorname{Re} \lambda > -\frac{\beta_0}{2} \right\} \setminus \left\{ \lambda; |\lambda| \leq \frac{\beta_0}{4} \right\} \subset \rho(-L_{\eta'}).$$

Furthermore,

$$\|(\lambda + L_{\eta'})^{-1}f\|_2 \leq \frac{C}{\operatorname{Re} \lambda + \beta_0} \|f\|_2, \quad (4.21)$$

$$\|\nabla \tilde{Q}(\lambda + L_{\eta'})^{-1}f\|_2 \leq \frac{C}{(\operatorname{Re} \lambda + \beta_0)^{\frac{1}{2}}} \|f\|_2 \quad (4.22)$$

for $\lambda \in \Sigma_1$.

Proof. Recall that $L_{\eta'} = L_0 + M_{\eta'}$. By Theorem 4.5, we have

$$\|(\lambda + L_0)^{-1}f\|_2 + \|\nabla \tilde{Q}(\lambda + L_0)^{-1}f\|_2 \leq C\|f\|_2$$

for $\lambda \in \Sigma_1$, and hence, since $\|M_{\eta'}u\|_2 \leq C|\eta'|(\|u\|_2 + \|\nabla w\|_2)$, it holds that $\|M_{\eta'}(\lambda + L_0)^{-1}f\|_2 \leq 2C|\eta'|\|f\|_2$. Therefore, if η' satisfies $|\eta'| < \frac{1}{2C}$, then $\lambda \in \rho(-L_{\eta'})$ and

$$\begin{aligned} (\lambda + L_{\eta'})^{-1} &= (\lambda + L_0)^{-1} \sum_{N=0}^{\infty} (-1)^N (M_{\eta'}(\lambda + L_0)^{-1})^N, \\ \|(\lambda + L_{\eta'})^{-1}f\|_2 &\leq C' \left(\frac{1}{|\lambda|} + \frac{1}{\operatorname{Re} \lambda + \beta_0} \right) \sum_{N=0}^{\infty} \|M_{\eta'}(\lambda + L_0)^{-1}\|_2^N \|f\|_2 \\ &\leq \frac{C}{\operatorname{Re} \lambda + \beta_0} \|f\|_2, \\ \|\nabla \tilde{Q}(\lambda + L_{\eta'})^{-1}f\|_2 &\leq C \left(\frac{1}{|\lambda|} + \frac{1}{(\operatorname{Re} \lambda + \beta_0)^{\frac{1}{2}}} \right) \sum_{N=0}^{\infty} \|M_{\eta'}(\lambda + L_0)^{-1}\|_2^N \|f\|_2 \\ &\leq \frac{C}{(\operatorname{Re} \lambda + \beta_0)^{\frac{1}{2}}} \|f\|_2. \end{aligned}$$

This completes the proof. $\square$

We next establish an asymptotic formula for eigenvalues of $-L_{\eta'}$ near the imaginary axis.

Proposition 4.8. *If $\nu \geq \nu_0$, $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$ and $|\eta'| \leq r_0$ (by taking ν_0 , $\tilde{\gamma}_0$ and r_0^{-1} suitably larger if necessary), then*

$$\sigma(-L_{\eta'}) \cap \{|\lambda| < \frac{\beta_0}{2}\} = \{\lambda_{\eta'}\}.$$

Here $\lambda_{\eta'}$ is a simple eigenvalue that satisfies

$$\lambda_{\eta'} = -\kappa(\eta') + O(|\eta'|^3) \quad (4.23)$$

as $\eta' \rightarrow 0$. Here

$$\kappa(\eta') = i \sum_{j=1}^{n-1} a_j \eta_j + \sum_{j,k=1}^{n-1} a_{jk} \eta_j \eta_k, \quad (4.24)$$

where $a_j \in \mathbb{R}$ and $a_{jk} \in \mathbb{R}$; a_{jk} , $j, k = 1, \dots, n-1$, satisfy

$$\sum_{j,k=1}^{n-1} a_{jk} \eta_j \eta_k \geq \frac{\kappa_0 \gamma^2}{\nu} |\eta'|^2 \quad (\eta' \in \mathbb{R}^{n-1}) \quad (4.25)$$

with some constant $\kappa_0 > 0$ independent of ν , $\tilde{\nu}$ and γ . As a consequence,

$$\operatorname{Re} \lambda_{\eta'} \leq -\frac{\kappa_0 \gamma^2}{2\nu} |\eta'|^2. \quad (4.26)$$

Proof. Since

$$\begin{aligned} \|L_j^{(1)} u\|_2 &\leq C\{\|u\|_2 + \|\nabla w\|_2\} \leq C\{\|L_0 u\|_2 + \|u\|_2\}, \\ \|L_{jk}^{(2)} u\|_2 &\leq C\|w\|_2 \leq C\{\|L_0 u\|_2 + \|u\|_2\}, \end{aligned}$$

in view of Theorem 4.5, we can apply the analytic perturbation theory to see that $\sigma(-L_{\eta'}) \cap \{|\lambda| < \frac{\beta_0}{2}\}$ consist of a simple eigenvalue, say $\lambda_{\eta'}$, for sufficiently small η' , and that $\lambda_{\eta'}$ is expanded as

$$\lambda_{\eta'} = \sum_{j=1}^{n-1} \lambda_j^{(1)} \eta_j + \sum_{j,k=1}^{n-1} \lambda_{jk}^{(2)} \eta_j \eta_k + O(|\eta'|^3)$$

as $\eta' \rightarrow 0$, where

$$\begin{aligned} \lambda_j^{(1)} &= -\langle L_j^{(1)} u^{(0)}, u^{(0)*} \rangle, \\ \lambda_{jk}^{(2)} &= -\frac{1}{2} \langle (L_{jk}^{(2)} + L_{kj}^{(2)}) u^{(0)}, u^{(0)*} \rangle + \frac{1}{2} \langle (L_j^{(1)} S L_k^{(1)} + L_k^{(1)} S L_j^{(1)}) u^{(0)}, u^{(0)*} \rangle. \end{aligned}$$

Here $S = [(I - \Pi^{(0)})L_0(I - \Pi^{(0)})]^{-1}$. See, e.g., [16, Chap. VII], [23, Chap. XII].

By the definition of $L_j^{(1)}$, $u^{(0)}$ and $u^{(0)*}$, we have

$$\lambda_j^{(1)} = -i\langle v_s^j \phi^{(0)} + \gamma^2 \rho_s w^{(0),j} \rangle.$$

As for $\lambda_{jk}^{(2)}$, since $\frac{1}{2}\langle (L_{jk}^{(2)} + L_{kj}^{(2)})u^{(0)}, u^{(0)*} \rangle = 0$, it holds that

$$\lambda_{jk}^{(2)} = \frac{1}{2}\langle (L_j^{(1)}SL_k^{(1)} + L_k^{(1)}SL_j^{(1)})u^{(0)}, u^{(0)*} \rangle$$

We set $u_1^{(k)} = {}^\top(\phi_1^{(k)}, w_1^{(k)}) = SL_k^{(1)}u^{(0)}$. Then $u_1^{(k)}$ is a solution of

$$\begin{cases} L_0 u_1^{(k)} = (I - \Pi^{(0)})L_k^{(1)}u^{(0)}, \\ \langle \phi_1^{(k)} \rangle = 0. \end{cases} \quad (4.27)$$

To prove (4.25), we prepare some estimates of $u_1^{(k)}$.

Lemma 4.9. *If $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$ (by taking ν_0 and $\tilde{\gamma}_0$ suitably larger if necessary), then*

$$\|\phi_1^{(k)}\|_{H^1} \leq C, \quad \|w_1^{(k)}\|_{H^2} \leq \frac{C}{\nu}. \quad (4.28)$$

Proof. Since $\|(I - \Pi^{(0)})L_k^{(1)}u^{(0)}\|_{L^2 \times H^1} \leq C$, it follows from Proposition 3.1 and (5.6)_k with $k = 1$ that

$$\frac{1}{\nu}\|\phi_1^{(k)}\|_{H^1}^2 + \nu\|w_1^{(k)}\|_{H^2}^2 \leq \frac{C}{\nu}\left\{1 + \left(\frac{1}{\nu^2} + \frac{\nu^2}{\gamma^4}\right)\|w_1^{(k)}\|_{H^2}^2 + \frac{1}{\gamma^4}\|\phi_1^{(k)}\|_{H^1}^2\right\}.$$

Therefore, if $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$, we have $\|\phi_1^{(k)}\|_{H^1} \leq C$ and $\|w_1^{(k)}\|_{H^2} \leq \frac{C}{\nu}$. This completes the proof. $\square$

To proceed further, we introduce the functions $U_1^{(k)} = {}^\top(\Phi_1^{(k)}, W_1^{(k)}) \in H_{per,*}^2 \times (H_{per}^3 \cap H_{0,per}^1)$ ($k = 1, \dots, n-1$), which are the unique solution of the following Stokes system

$$A_S U_1^{(k)} = F^{(k)}, \quad \langle \Phi_1^{(k)} \rangle = 0. \quad (4.29)$$

Here

$$A_S = \begin{pmatrix} 0 & \text{div} \\ \nabla & -\Delta \end{pmatrix}, \quad F^{(k)} = \begin{pmatrix} 0 \\ \mathbf{e}_k \end{pmatrix}.$$

Set $\check{\phi}_1^{(k)} = i\Phi_1^{(k)}$ and $\check{w}_1^{(k)} = \frac{i}{\nu}W_1^{(k)}$. Then $\check{u}_1^{(k)} = {}^\top(\check{\phi}_1^{(k)}, \check{w}_1^{(k)})$ is the solution of

$$\mathcal{A}\check{u}_1^{(k)} = iF^{(k)}, \quad \langle \check{\phi}_1^{(k)} \rangle = 0. \quad (4.30)$$

Here

$$\mathcal{A} = \begin{pmatrix} 0 & \gamma^2 \operatorname{div} \\ \nabla & -\nu \Delta - \tilde{\nu} \nabla \operatorname{div} \end{pmatrix}.$$

Furthermore, we have the following lemma.

Lemma 4.10 ([15]). *Let $\check{\kappa}(\eta')$ be defined by*

$$\check{\kappa}(\eta') = \sum_{j,k=1}^{n-1} \check{a}_{jk} \eta_j \eta_k \quad \text{for } \eta' \in \mathbb{R}^{n-1},$$

where

$$\check{a}_{jk} = \frac{\gamma^2}{\nu} \frac{1}{|\Omega_{per}|} (\nabla W_1^{(j)}, \nabla W_1^{(k)}) = \frac{\gamma^2}{\nu} \langle \mathbf{e}_j, W_1^{(k)} \rangle.$$

Then there exists a constant $\check{\kappa}_0 > 0$ independent of ν , $\tilde{\nu}$ and γ such that

$$\check{\kappa}(\eta') \geq \frac{\check{\kappa}_0 \gamma^2}{\nu} |\eta'|^2 \quad \text{for all } \eta' \in \mathbb{R}^{n-1}. \quad (4.31)$$

To prove (4.25) we will show that $\check{\kappa}(\eta')$ is the principal part of $\operatorname{Re} \kappa(\eta')$ when ν and $\frac{\gamma^2}{\nu}$ are large enough. We estimate the difference of $u_1^{(k)}$ and $\check{u}_1^{(k)}$.

Lemma 4.11. *If $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$ (by taking ν_0 and $\tilde{\gamma}_0$ suitably larger if necessary), then*

$$\|\phi_1^{(k)} - \check{\phi}_1^{(k)}\|_2 \leq C \left(\frac{1}{\gamma^2} + \frac{1}{\nu^2} \right), \quad \|w_1^{(k)} - \check{w}_1^{(k)}\|_2 \leq C \left(\frac{1}{\nu \gamma^2} + \frac{1}{\nu^3} \right).$$

Proof. We write $L_0 u_1^{(k)} = (I - \Pi^{(0)}) L_k^{(1)} u^{(0)}$ as

$$\mathcal{A} u_1^{(k)} = -\mathcal{C} u_1^{(k)} + (I - \Pi^{(0)}) L_k^{(1)} u^{(0)}.$$

Here $\mathcal{C}$ is defined by

$$\begin{aligned} \mathcal{C} = L_0 - \mathcal{A} = & \begin{pmatrix} \operatorname{div}(\cdot v_s) & \gamma^2 \operatorname{div}(\phi_s \cdot) \\ \nabla(\tilde{P}^{(1)}(\phi_s) \phi_s \cdot) & \nu \frac{\phi_s}{1+\phi_s} \Delta + \tilde{\nu} \frac{\phi_s}{1+\phi_s} \nabla \operatorname{div} \end{pmatrix} \\ & + \begin{pmatrix} 0 & 0 \\ \frac{1}{\gamma^2 \rho_s^2} (\nu \Delta v_s + \tilde{\nu} \nabla \operatorname{div} v_s) & v_s \cdot \nabla + {}^\top \nabla v_s \end{pmatrix}. \end{aligned}$$

It then follows that

$$\mathcal{A}(u_1^{(k)} - \check{u}_1^{(k)}) = -\mathcal{C}u_1^{(k)} + (I - \Pi^{(0)})L_k^{(1)}u^{(0)} - iF^{(k)},$$

and hence,

$$((\mathcal{A}(u_1^{(k)} - \check{u}_1^{(k)}), u_1^{(k)} - \check{u}_1^{(k)})) = I_1 + I_2,$$

where

$$\begin{aligned} I_1 &= -((\mathcal{C}u_1^{(k)}, u_1^{(k)} - \check{u}_1^{(k)})), \\ I_2 &= (((I - \Pi^{(0)})L_k^{(1)}u^{(0)} - iF^{(k)}, u_1^{(k)} - \check{u}_1^{(k)})). \end{aligned}$$

The left-hand side is estimated as

$$\begin{aligned} &((\mathcal{A}(u_1^{(k)} - \check{u}_1^{(k)}), u_1^{(k)} - \check{u}_1^{(k)})) \\ &\geq \nu \|\nabla(w_1^{(k)} - \check{w}_1^{(k)})\|_2^2 + \tilde{\nu} \|\operatorname{div}(w_1^{(k)} - \check{w}_1^{(k)})\|_2^2 + \frac{\delta}{2} \|\phi_1^{(k)} - \check{\phi}_1^{(k)}\|_2^2 \\ &\quad - C\delta \{ \nu^2 \|\nabla(w_1^{(k)} - \check{w}_1^{(k)})\|_2^2 + \tilde{\nu}^2 \|\operatorname{div}(w_1^{(k)} - \check{w}_1^{(k)})\|_2^2 + \gamma^2 \|w_1^{(k)} - \check{w}_1^{(k)}\|_2^2 \}. \end{aligned}$$

Taking $\delta = \frac{1}{2C} \min\{\frac{1}{\nu}, \frac{\nu}{\gamma^2}\}$, we have

$$\begin{aligned} &((\mathcal{A}(u_1^{(k)} - \check{u}_1^{(k)}), u_1^{(k)} - \check{u}_1^{(k)})) \\ &\geq \frac{\nu}{2} \|\nabla(w_1^{(k)} - \check{w}_1^{(k)})\|_2^2 + \frac{\tilde{\nu}}{2} \|\operatorname{div}(w_1^{(k)} - \check{w}_1^{(k)})\|_2^2 + \frac{\delta}{2} \|\phi_1^{(k)} - \check{\phi}_1^{(k)}\|_2^2. \end{aligned} \quad (4.32)$$

By Proposition 3.1 and Lemma 4.9, we see that I_1 is estimated as

$$|I_1| \leq C \left\{ \left(\frac{1}{\nu\gamma^2} + \frac{\delta}{\nu^2} \right) \|\phi_1^{(k)} - \check{\phi}_1^{(k)}\|_2 + \left(\frac{1}{\gamma^2} + \frac{1}{\nu^2} \right) \|w_1^{(k)} - \check{w}_1^{(k)}\|_2 \right\}. \quad (4.33)$$

As for I_2 , we observe that

$$(I - \Pi^{(0)})L_k^{(1)}u^{(0)} - iF^{(k)} = \tilde{L}_k^{(1)}u^{(0)} - \Pi^{(0)}L_k^{(1)}u^{(0)},$$

where

$$\begin{aligned} \tilde{L}_k^{(1)}u^{(0)} &= L_k^{(1)}u^{(0)} - iF^{(k)} \\ &= i \left(\begin{pmatrix} v_s^k \\ \left(\frac{P'(\rho_s)}{\gamma^2 \rho_s} - 1 \right) \mathbf{e}_k \end{pmatrix} - \frac{1}{\rho_s} \left(2\nu \mathbf{e}_k \otimes \mathbf{e}_k \partial_{x_k} + \tilde{\nu} \mathbf{e}_k \operatorname{div} - \tilde{\nu} \nabla(\top \mathbf{e}_k) \right) + v_s^k \right) u^{(0)} \end{aligned}$$

Therefore, by Proposition 3.1, Proposition 4.1 and Lemma 4.9, we have

$$|I_2| \leq C \left\{ \frac{1}{\nu\gamma^2} \|\phi_1^{(k)} - \check{\phi}_1^{(k)}\|_2 + \frac{1}{\gamma^2} \|w_1^{(k)} - \check{w}_1^{(k)}\|_2 \right\}. \quad (4.34)$$

From (4.32), (4.33) and (4.34), we conclude

$$\|\phi_1^{(k)} - \check{\phi}_1^{(k)}\|_2 \leq C\left(\frac{1}{\gamma^2} + \frac{1}{\nu^2}\right), \quad \|w_1^{(k)} - \check{w}_1^{(k)}\|_2 \leq C\left(\frac{1}{\nu\gamma^2} + \frac{1}{\nu^3}\right).$$

This completes the proof. $\square$

Proof of (4.25). We now turn to the proof of (4.25). Since $\rho_s = 1 + \phi_s$ and $\check{w}_1^{(k)} = \frac{i}{\nu} W_1^{(k)}$, by Lemma 4.10 and 4.11, we have

$$\begin{aligned} \sum_{j,k=1}^{n-1} \eta_j \eta_k \langle Q_0 L_j^{(1)} S L_k^{(1)} u^{(0)} \rangle &= i \sum_{j,k=1}^{n-1} \eta_j \eta_k \langle v_s^j \phi_1^{(k)} + \gamma^2 \rho_s \mathbf{e}_j \cdot w_1^{(k)} \rangle \\ &= - \sum_{j,k=1}^{n-1} \eta_j \eta_k \left(\frac{\gamma^2}{\nu} \langle \mathbf{e}_j \cdot W_1^{(k)} \rangle - i \gamma^2 \langle \mathbf{e}_j \cdot (w_1^{(k)} - \check{w}_1^{(k)}) \rangle \right) \\ &\quad + i \sum_{j,k=1}^{n-1} \eta_j \eta_k \langle v_s^j \phi_1^{(k)} + \gamma^2 \phi_s \mathbf{e}_j \cdot w_1^{(k)} \rangle \\ &\leq - \frac{\gamma^2}{\nu} \left\{ \frac{\check{\kappa}_0}{2} - C \left(\frac{1}{\nu} + \frac{1}{\gamma^2} \right) \right\} |\eta'|^2, \end{aligned}$$

and we have

$$\sum_{j,k=1}^{n-1} \lambda_{jk}^{(2)} \eta_j \eta_k \leq - \frac{\gamma^2}{\nu} \left\{ \frac{\check{\kappa}_0}{2} - C \left(\frac{1}{\nu} + \frac{1}{\gamma^2} \right) \right\} |\eta'|^2.$$

If $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$, it holds that

$$\sum_{j,k=1}^{n-1} \lambda_{jk}^{(2)} \eta_j \eta_k \leq - \frac{\check{\kappa}_0 \gamma^2}{4\nu} |\eta'|^2. \quad (4.35)$$

This completes the proof. $\square$

4.4 Boundedness of eigenprojections

In this subsection we estimate the eigenprojections $\Pi_{\eta'}$ for eigenvalues $\lambda_{\eta'}$ of $-L_{\eta'}$. By Proposition 4.6, we have the following resolvent estimates for $-L_{\eta'}^*$ with $|\eta'| \leq r_0$.

Proposition 4.12. *If $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$ (by taking ν_0 and $\tilde{\gamma}_0$ suitably larger if necessary), then there exists a constant $r_0 > 0$ such that if $|\eta'| \leq r_0$, then $\Sigma_1 \subset \rho(-L_{\eta'}^*)$ and*

$$\|(\lambda + L_{\eta'}^*)^{-1} f\|_2 \leq \frac{C}{\operatorname{Re} \lambda + \beta_0} \|f\|_2, \quad (4.36)$$

$$\|\nabla \tilde{Q}(\lambda + L_{\eta'}^*)^{-1} f\|_2 \leq \frac{C}{(\operatorname{Re} \lambda + \beta_0)^{\frac{1}{2}}} \|f\|_2 \quad (4.37)$$

uniformly for $\lambda \in \Sigma_1$.

Proof. We define $M_{\eta'}^*$ by $M_{\eta'}^* = L_{\eta'}^* - L_0^*$:

$$\begin{aligned} M_{\eta'}^* = & \begin{pmatrix} -iv_s \cdot \tilde{\eta}' & -i\gamma^2 \tilde{\eta}' \cdot (\rho_s \cdot) \\ -i\tilde{\eta}' \left(\frac{P'(\rho_s)}{\gamma^2 \rho_s} \cdot \right) & -\frac{\nu}{\rho_s} (|\eta'|^2 - 2i\tilde{\eta}' \cdot \nabla) I_n - i\frac{\tilde{\nu}}{\rho_s} \tilde{\eta}'^\top \nabla_{\eta'} \end{pmatrix} \\ & + \begin{pmatrix} 0 & 0 \\ 0 & -i\frac{1}{\rho_s} v_s \cdot \tilde{\eta}'(\rho_s \cdot) + i\tilde{\eta}'^\top v_s \end{pmatrix}. \end{aligned} \quad (4.38)$$

Since $\|M_{\eta'}^* u\|_2 \leq C|\eta'|(\|u\|_2 + \|\nabla w\|_2)$, Proposition 4.12 follows from Proposition 4.6 as in the proof of Proposition 4.7. $\square$

We next investigate the estimate for $(\lambda + L_0^*)^{-1} f$ in a higher order Sobolev space.

Proposition 4.13. *There exists a constant $\tilde{\beta}_0 > 0$ such that if $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$, then, for any $\beta_1 \in (0, \tilde{\beta}_0]$, the estimate*

$$\|(\lambda + L_0^*)^{-1} f\|_{H^2 \times H^3} \leq C \|f\|_{H^2 \times H^1}$$

holds uniformly for λ satisfying $\frac{\beta_1}{2} \leq |\lambda| \leq \beta_1$.

Proof. We write $(\lambda + L_0^*)u = f$ as

$$(\lambda + \tilde{L}_0^*)u + \tilde{M}^* u = f, \quad (4.39)$$

where $\tilde{L}_0^* : H_{per}^2 \times H_{per}^1 \longrightarrow H_{per}^2 \times H_{per}^1$ is defined by

$$\tilde{L}_0^* = \begin{pmatrix} -\operatorname{div}(v_s \cdot) & -\gamma^2 \operatorname{div} \\ -\nabla & -\nu \Delta - \tilde{\nu} \operatorname{div} \end{pmatrix}$$

with domain

$$D(\tilde{L}_0^*) = \{u \in H_{per}^2 \times (H_{per}^3 \cap H_{0,per}^1); \tilde{L}_0^* u \in H_{per}^2 \times H_{per}^1\}.$$

Here

$$\begin{aligned} \tilde{M}^* = & \begin{pmatrix} -v_s \cdot \nabla \left(\frac{P'(\rho_s)}{\rho_s} \right) \frac{\rho_s}{P'(\rho_s)} & -\gamma^2 \operatorname{div}(\phi_s \cdot) \\ -\nabla(\phi_s \tilde{P}^{(1)}(\phi_s) \cdot) & \frac{\phi_s}{\rho_s} (\nu \Delta + \tilde{\nu} \nabla \operatorname{div}) \end{pmatrix} \\ & + \begin{pmatrix} \operatorname{div} v_s & \frac{\gamma^2}{P'(\rho_s)} (\nu \Delta^\top v_s + \tilde{\nu}^\top \nabla \operatorname{div} v_s) \\ 0 & -\operatorname{div} v_s - \frac{1}{\rho_s} v_s \cdot \nabla(\rho_s \cdot) + \nabla^\top v_s \end{pmatrix}. \end{aligned}$$

To prove the proposition, we first show the invertibility of $\lambda + \tilde{L}_0^*$ for $|\lambda| \ll 1$ and then regard $\tilde{M}^*$ as a small perturbation of $\lambda + \tilde{L}_0^*$. To this end, we introduce the operator $\mathcal{A}^*[v_s] : H_{per,*}^2 \times H_{per}^1 \rightarrow H_{per,*}^2 \times H_{per}^1$ with domain $D(\mathcal{A}^*[v_s]) = D(\tilde{L}_0^*) \cap (H_{per,*}^2 \times H_{per}^1)$ defined by $\mathcal{A}^*[v_s]u = \tilde{L}_0^*u$ for $u \in D(\mathcal{A}^*[v_s])$. We observe that there exists a unique solution $u \in D(\mathcal{A}^*[v_s])$ of $(\lambda + \mathcal{A}^*[v_s])u = F$ for any $F \in H_{per,*}^2 \times H_{per}^1$ if $\operatorname{Re}\lambda \geq -\Lambda$ for some constant $\Lambda > 0$ and u satisfies estimate (5.5)₂ and, in particular, (5.6)_k with $k = 2$ when $\lambda = 0$. This can be seen with obvious modification of the proof of Theorem 5.1.

We also introduce the norm $\|u\|_{H^2 \times H^1}^* = \|\phi\|_{H^2} + \nu\|w\|_{H^1}$ for $u = {}^\top(\phi, w)$. Recall that $\|u\|_{H^2 \times H^3}^* = \|\phi\|_{H^2} + \nu\|w\|_{H^3}$.

As in the proof of Proposition 4.1, we see from (5.6)_k with $k = 2$, that there exist $\hat{\nu}_0 > 0$ and $\hat{\gamma}_0 > 0$ such that if $\nu \geq \hat{\nu}_0$ and $\frac{\gamma^2}{\nu} \geq \hat{\gamma}_0$, then

$$\|(\mathcal{A}^*[v_s])^{-1}F\|_{H^2 \times H^3}^* \leq C_1 \left\{ \frac{\nu}{\gamma^2} \|F^0\|_{H^2} + \|\tilde{F}\|_{H^1} \right\} \quad (4.40)$$

for $F = {}^\top(F^0, \tilde{F}) \in H_{*,per}^2 \times H_{per}^1$. By (4.40), we have

$$\|\lambda(\mathcal{A}^*[v_s])^{-1}u\|_{H^2 \times H^3}^* \leq C_2|\lambda|\|u\|_{H^2 \times H^1}^* \leq C_2|\lambda|\|u\|_{H^2 \times H^3}^*,$$

where $C_2 = C_2(\hat{\nu}_0, \hat{\gamma}_0) > 0$. Therefore, if $|\lambda| \leq \frac{1}{2C_2}$, then there exists a unique $u \in D(\mathcal{A}^*[v_s])$ satisfying $u = -\lambda(\mathcal{A}^*[v_s])^{-1}u + (\mathcal{A}^*[v_s])^{-1}F$, i.e., $(\lambda + \mathcal{A}^*[v_s])u = F$, and $u = (\lambda + \mathcal{A}^*[v_s])^{-1}F$ satisfies

$$\|u\|_{H^2 \times H^3}^* \leq 2C_2\|F\|_{H^2 \times H^1}^*. \quad (4.41)$$

We next show the invertibility of $\lambda + \tilde{L}_0^*$. We set $\tilde{u}^{(0)} = {}^\top(1, 0)$ and define $\tilde{\Pi}^{(0)}$ by $\tilde{\Pi}^{(0)}u = \langle \phi \rangle \tilde{u}^{(0)}$ for $u = {}^\top(\phi, w)$. Then $\tilde{\Pi}^{(0)}$ is a projection. We also set $\tilde{\Pi}_1 = I - \tilde{\Pi}^{(0)}$.

Consider the problem

$$(\lambda + \tilde{L}_0^*)u = F, \quad u \in D(\tilde{L}_0^*). \quad (4.42)$$

Since $\tilde{\Pi}^{(0)}\tilde{L}_0^*u = 0$, (4.42) is decomposed into the following system:

$$\begin{cases} \lambda\tilde{\Pi}^{(0)}u = \tilde{\Pi}^{(0)}F, \\ \lambda\tilde{\Pi}_1u + \mathcal{A}^*[v_s]\tilde{\Pi}_1u + \tilde{\Pi}_1\mathcal{A}^*[v_s]\tilde{\Pi}^{(0)}u = \tilde{\Pi}_1F, \\ u \in D(\tilde{L}_0^*). \end{cases} \quad (4.43)$$

By the first equation of (4.43), if $\lambda \neq 0$, then

$$\tilde{\Pi}^{(0)}u = \frac{1}{\lambda}\tilde{\Pi}^{(0)}F. \quad (4.44)$$

We set $\tilde{\beta}_0 = \frac{1}{2C_2}$. Let λ satisfy $\frac{\beta_1}{2} \leq |\lambda| \leq \beta_1$ for $\beta_1 \in (0, \tilde{\beta}_0]$. Substituting (4.44) into the second equation of (4.43), we have

$$\lambda \tilde{\Pi}_1 u + \mathcal{A}^*[v_s] \tilde{\Pi}_1 u = \tilde{\Pi}_1 F - \frac{1}{\lambda} \tilde{\Pi}_1 \mathcal{A}^*[v_s] \tilde{\Pi}^{(0)} F. \quad (4.45)$$

We set $G = \tilde{\Pi}_1 F - \frac{1}{\lambda} \tilde{\Pi}_1 \mathcal{A}^*[v_s] \tilde{\Pi}^{(0)} F$. Then $G \in \tilde{\Pi}_1[H_{per}^2 \times H_{per}^1] \subset H_{per,*}^2 \times H_{per}^1$. If λ satisfies $\frac{\beta_1}{2} \leq |\lambda| \leq \beta_1$, there exists a function $u_1 = {}^\top(\phi_1, w_1) \in D(\mathcal{A}^*[v_s])$ satisfying $(\lambda + \mathcal{A}^*[v_s])u_1 = G$. Furthermore, by (4.41) and Proposition 3.1, u_1 satisfies

$$\|u_1\|_{H^2 \times H^3}^* \leq 2C_2 \|G\|_{H^2 \times H^1}^* \leq C_3 \left(1 + \frac{1}{\nu\beta_1}\right) \|F\|_{H^2 \times H^1}^* \quad (4.46)$$

for $\frac{\beta_1}{2} \leq |\lambda| \leq \beta_1$ with some $C_3 = C_3(\hat{\nu}_0, \hat{\gamma}_0) > 0$.

We set $u = \frac{1}{\lambda} \tilde{\Pi}^{(0)} F + u_1$. Since $\tilde{\Pi}^{(0)} u_1 = \langle \phi_1 \rangle \tilde{u}^{(0)} = 0$, it holds that $\tilde{\Pi}^{(0)} u = \frac{1}{\lambda} \tilde{\Pi}^{(0)} F$, and we have $\tilde{\Pi}_1 u = u_1$. Consequently, u is a solution of (4.42). By (4.44) and (4.46), $u = (\lambda + \tilde{L}_0^*)^{-1} F$ satisfies

$$\|u\|_{H^2 \times H^3}^* \leq \frac{1}{|\lambda|} \|\tilde{\Pi}^{(0)} F\|_{H^2 \times H^3}^* + \|u_1\|_{H^2 \times H^3}^* \leq C_4 \|F\|_{H^2 \times H^1}^* \quad (4.47)$$

for $\frac{\beta_1}{2} \leq |\lambda| \leq \beta_1$. Here $C_4 = C_4(\hat{\nu}_0, \hat{\gamma}_0, \beta_1) > 0$. This implies that $(\lambda + \tilde{L}_0^*)^{-1} : H_{per}^2 \times H_{per}^1 \rightarrow H_{per}^2 \times (H_{per}^3 \cap H_{0,per}^1)$ is bounded and $u = (\lambda + \tilde{L}_0^*)^{-1} F$ satisfies (4.47).

We finally prove the invertibility of $\lambda + L_0^* = \lambda + \tilde{L}_0^* + \tilde{M}^*$. By Proposition 3.1, we have

$$\|\tilde{M}^* u\|_{H^2 \times H^1}^* \leq C \left(\frac{1}{\nu} + \frac{1}{\gamma^2 \nu} + \frac{\nu + \tilde{\nu}}{\gamma^2} \right) \|u\|_{H^2 \times H^3}^*. \quad (4.48)$$

It then follows from (4.47) and (4.48) that

$$\|(\lambda + \tilde{L}_0^*)^{-1} \tilde{M}^* u\|_{H^2 \times H^3}^* \leq CC_4 \left(\frac{1}{\nu} + \frac{1}{\gamma^2 \nu} + \frac{\nu + \tilde{\nu}}{\gamma^2} \right) \|u\|_{H^2 \times H^3}^*.$$

Therefore, there exist $\nu_0 > 0$ and $\tilde{\gamma}_0 > 0$ such that if $\nu \geq \nu_0$ and $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$, there exists a unique $u \in D(\tilde{L}_0^*)$ satisfying $u = -(\lambda + \tilde{L}_0^*)^{-1} \tilde{M}^* u + (\lambda + \tilde{L}_0^*)^{-1} f$, that is, a unique solution $u \in D(\tilde{L}_0^*)$ of (4.39), and $u = (\lambda + \tilde{L}_0^* + \tilde{M}^*)^{-1} f$ satisfies the estimate $\|u\|_{H^2 \times H^3}^* \leq 2C_4 \|f\|_{H^2 \times H^1}^*$. Since $H_{per}^2 \times (H_{per}^3 \cap H_{0,per}^1) \subset D(L_0^*)$, we have $u = (\lambda + L_0^*)^{-1} f$. This completes the proof. $\square$

We are now in a position to estimate the eigenprojections.

Proposition 4.14. *If $\nu \geq \nu_0$, $\frac{\gamma^2}{\nu} \geq \tilde{\gamma}_0$ and $|\eta'| \leq r_0$ (by taking ν_0 and $\tilde{\gamma}_0$ suitably larger if necessary), then*

$$\|\Pi_{\eta'} u\|_2 \leq C \|u\|_1, \quad (4.49)$$

$$\|(\Pi_{\eta'} - \Pi^{(0)})u\|_2 \leq C |\eta'| \|u\|_1. \quad (4.50)$$

Proof. By Proposition 4.7, we see that $\Pi_{\eta'}$ and $\Pi_{\eta'}^*$ are given by

$$\Pi_{\eta'} = \frac{1}{2\pi i} \int_{|\lambda|=\frac{\beta_0}{4}} (\lambda + L_{\eta'})^{-1} d\lambda, \quad \Pi_{\eta'}^* = \frac{1}{2\pi i} \int_{|\lambda|=\frac{\beta_0}{4}} (\lambda + L_{\eta'}^*)^{-1} d\lambda.$$

Then $u_{\eta'} = \Pi_{\eta'} u^{(0)}$ is an eigenfunction of $-L_{\eta'}$ for the eigenvalue $\lambda_{\eta'}$ and $u_{\eta'}^* = \Pi_{\eta'}^* u^{(0)*}$ is an eigenfunction of $-L_{\eta'}^*$ for $\lambda_{\eta'}^* = \bar{\lambda}_{\eta'}$. Since $\lambda_{\eta'}$ is a simple eigenvalue, the eigenprojection $\Pi_{\eta'}$ is represented as

$$\Pi_{\eta'} u = \frac{\langle u, u_{\eta'}^* \rangle}{\langle u_{\eta'}, u_{\eta'}^* \rangle} u_{\eta'}.$$

From the proof of Proposition 4.7, we see that $(\lambda + L_{\eta'})^{-1}$ is expanded as

$$(\lambda + L_{\eta'})^{-1} = (\lambda + L_0)^{-1} + R_{\eta'}(\lambda)$$

and $R_{\eta'}(\lambda)$ is estimated as

$$\|R_{\eta'}(\lambda)f\|_2 \leq C |\eta'| \|f\|_2, \quad \|\nabla \tilde{Q} R_{\eta'}(\lambda)f\|_2 \leq C |\eta'| \|f\|_2$$

uniformly for $|\eta'| \leq r_0$ and $|\lambda| = \frac{\beta_0}{4}$. We write $u_{\eta'}$ as

$$\begin{aligned} u_{\eta'} &= \frac{1}{2\pi i} \int_{|\lambda|=\frac{\beta_0}{4}} (\lambda + L_0)^{-1} u^{(0)} d\lambda + \frac{1}{2\pi i} \int_{|\lambda|=\frac{\beta_0}{4}} R_{\eta'}(\lambda) u^{(0)} d\lambda \\ &= u^{(0)} + u_{\eta'}^{(1)}, \end{aligned}$$

where $u_{\eta'}^{(1)} = \frac{1}{2\pi i} \int_{|\lambda|=\frac{\beta_0}{4}} R_{\eta'}(\lambda) u^{(0)} d\lambda$. Then $u_{\eta'}^{(1)}$ satisfies

$$\|u_{\eta'}^{(1)}\|_2 \leq C |\eta'|, \quad \|\nabla \tilde{Q} u_{\eta'}^{(1)}\|_2 \leq C |\eta'|.$$

Similarly, $u_{\eta'}^*$ is written as $u_{\eta'}^* = u^{(0)*} + u_{\eta'}^{(1)*}$, where $u_{\eta'}^{(1)*} = \frac{1}{2\pi i} \int_{|\lambda|=\frac{\beta_0}{4}} R_{\eta'}^*(\lambda) u^{(0)} d\lambda$.

By Proposition 4.13, $u_{\eta'}^{(1)*}$ is estimated as $\|u_{\eta'}^{(1)*}\|_{H^2} \leq C |\eta'|$. Thus, by a similar argument to that in [15, Theorem 4.11], we obtain the desired estimates (4.49) and (4.50). This completes the proof. $\square$

4.5 Estimates of the Π -part of the semigroup

We finally prove Theorem 3.2. We set

$$\Pi = U\chi_0\Pi_{\eta'}T, \quad \chi_0 = \begin{cases} 1 & |\eta'| \leq r_0, \\ 0 & |\eta'| \geq r_0. \end{cases}$$

We see from Proposition 2.2 that $\Pi^2 = \Pi$. Furthermore, by Proposition 4.8, we have

$$e^{-tL}\Pi u_0 = U\chi_0 e^{-tL_{\eta'}}\Pi_{\eta'}T u_0 = U\chi_0 e^{\lambda_{\eta'}t}\Pi_{\eta'}T u_0.$$

Then, in a similar manner to the proof of [15], by using Proposition 4.8 and 4.14, we obtain the desired result. This completes the proof. $\square$

5 Resolvent estimates for the principal part of L_0

In this section we give a unique existence of solution of the following problem which provides the principal part of L_0 . We consider the problem:

$$\lambda\phi + \gamma^2 \operatorname{div} w + \operatorname{div}(\phi v) = f^0, \quad (5.1)$$

$$\lambda w - \nu \Delta w - \tilde{\nu} \nabla \operatorname{div} w + \nabla \phi = \tilde{f}, \quad (5.2)$$

under the condition

$$w|_{\Sigma_n} = 0, \quad \phi|_{\Sigma_{j,-}} = \phi|_{\Sigma_{j,+}}, \quad w|_{\Sigma_{j,-}} = w|_{\Sigma_{j,+}} \quad (j = 1, \dots, n-1), \quad (5.3)$$

$$\langle \phi \rangle = 0. \quad (5.4)$$

Here $\lambda \in \mathbb{C}$ is a parameter and $v = {}^\top(v^1, \dots, v^n) \in H_{per}^4$ is a given function.

Throughout this section we assume that ν and $\tilde{\nu}$ satisfy (2.4), and thus, we use relation (2.5). The letter C denotes positive constants depending only on ν_* and Ω_{per} .

We have the following a priori estimates.

Theorem 5.1. *Let $k = 0, 1, 2, 3$. There exists a constant $\Lambda > 0$ such that if $\operatorname{Re} \lambda \geq -\Lambda$ and $\|v\|_{H^{3+(k-2)+}} \leq C \min \left\{ \frac{\gamma^2}{\nu}, \nu, \gamma \right\}$, then problem (5.1)-(5.4) as a unique solution $u = {}^\top(\phi, w) \in H_{per,*}^k \times (H_{per}^{k+1} \cap H_{0,per}^1)$ and u satisfies the following estimates (5.5) $_j$, $0 \leq j \leq k$:*

$$\begin{aligned} & \operatorname{Re}(\lambda + \Lambda)(\gamma^{-2}\|\phi\|_2^2 + \|w\|_2^2) + \frac{\nu}{2}\|\nabla w\|_2^2 + \frac{\tilde{\nu}}{2}\|\operatorname{div} w\|_2^2 + \frac{\delta_1}{2}\|\phi\|_2^2 \\ & \leq C \left\{ \left(\frac{1}{\delta_1 \gamma^4} + \frac{\delta_1^2}{\nu} \right) \|f^0\|_2^2 + \frac{1}{\nu} \|\tilde{f}\|_{H^{-1}}^2 \right\}, \end{aligned} \quad (5.5)_0$$

$$\begin{aligned} & \operatorname{Re}(\lambda + \Lambda)(\gamma^{-2}\|\phi\|_{H^1}^2 + \|w\|_{H^1}^2) + \frac{1}{\nu}\|\phi\|_{H^1}^2 + \nu\|w\|_{H^2}^2 + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\nabla w\|_2^2 \\ & \leq C\left\{\left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3}\right)\|f^0\|_{H^1}^2 + \left(\frac{1}{\nu} + \frac{\gamma^2}{\nu^3}\right)\|\tilde{f}\|_2^2\right\}, \end{aligned} \quad (5.5)_1$$

$$\begin{aligned} & \operatorname{Re}(\lambda + \Lambda)(\gamma^{-2}\|\phi\|_{H^2}^2 + \|w\|_{H^1}^2) + \frac{1}{\nu}\|\phi\|_{H^2}^2 + \nu\|w\|_{H^3}^2 + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\nabla w\|_2^2 \\ & \leq C\left\{\frac{|\lambda|^2}{(\nu + \tilde{\nu})^2}\left(\left(\frac{1}{\delta_1\gamma^4} + \frac{\delta_1^2}{\nu}\right)\|f^0\|_2^2 + \frac{1}{\nu}\|\tilde{f}\|_{H^{-1}}^2\right) \right. \\ & \quad \left. + \left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3}\right)\|f^0\|_{H^2}^2 + \frac{1}{\nu}\|\tilde{f}\|_{H^1}^2 + \frac{\gamma^2}{\nu^3}\|\tilde{f}\|_2^2\right\}, \end{aligned} \quad (5.5)_2$$

$$\begin{aligned} & \operatorname{Re}(\lambda + \Lambda)(\gamma^{-2}\|\phi\|_{H^3}^2 + \|w\|_{H^2}^2) + \frac{1}{\nu}\|\phi\|_{H^3}^2 + \nu\|w\|_{H^4}^2 + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\nabla w\|_2^2 \\ & \leq C\left\{\frac{|\lambda|^2}{(\nu + \tilde{\nu})^2}\left(\left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3} + \frac{1}{\delta_1\gamma^4} + \frac{\delta_1^2}{\nu}\right)\|f^0\|_2^2 + \left(\frac{1}{\nu} + \frac{\gamma^2}{\nu^3}\right)\|\tilde{f}\|_{H^{-1}}^2\right) \right. \\ & \quad \left. + \left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3}\right)\|f^0\|_{H^3}^2 + \frac{1}{\nu}\|\tilde{f}\|_{H^2}^2 + \frac{\gamma^2}{\nu^3}\|\tilde{f}\|_2^2\right\}. \end{aligned} \quad (5.5)_3$$

Here δ_1 is any positive number satisfying $0 < \delta_1 \leq C \min\left\{\frac{1}{\gamma}, \frac{1}{\nu}, \frac{\nu}{\gamma^2}\right\}$.

In particular, when $\lambda = 0$, $u = {}^\top(\phi, w)$ satisfies

$$\frac{1}{\nu}\|\phi\|_{H^k}^2 + \nu\|w\|_{H^{k+1}}^2 \leq C\left\{\frac{\nu + \tilde{\nu}}{\gamma^4}\|f^0\|_{H^k}^2 + \frac{1}{\nu}\|\tilde{f}\|_{H^{k-1}}^2\right\} \quad (5.6)_k$$

for $k = 1, 2, 3$.

To prove Theorem 5.1, we first establish a priori estimates (5.5)₀-(5.5)₃, and then using the a priori estimates and the result by Heywood-Padula [10], we will prove the existence of solutions.

a. A priori estimate

In this subsection we will prove the a priori estimate (5.5)₃ by the Matsumura-Nishida energy method [22].

Lemma 5.2. *Let (ϕ, w) be a solution of (5.1)-(5.4). Then*

$$\begin{aligned} & \operatorname{Re}\lambda(\gamma^{-2}\|\phi\|_2^2 + \|w\|_2^2) + \nu\|\nabla w\|_2^2 + (\nu + \tilde{\nu})\|\operatorname{div} w\|_2^2 \\ & \leq C\left\{\frac{1}{\varepsilon\gamma^4}\|f^0\|_2^2 + \frac{1}{\nu}\|\tilde{f}\|_{H^{-1}}^2 + \frac{1}{\gamma^2}\|v\|_{H^3}\|\phi\|_2^2 + \varepsilon\|\phi\|_2^2\right\}, \end{aligned} \quad (5.7)$$

$$\begin{aligned}
& \operatorname{Re} \lambda (\gamma^{-2} \|\phi\|_2^2 + \|w\|_2^2) + \frac{\nu}{2} \|\nabla w\|_2^2 + \frac{\tilde{\nu}}{2} \|\operatorname{div} w\|_2^2 + \frac{\delta_1}{2} \|\phi\|_2^2 \\
& \leq C \left\{ \left(\frac{1}{\delta_1 \gamma^4} + \frac{\delta_1^2}{\nu} \right) \|f^0\|_2^2 + \frac{1}{\nu} \|\tilde{f}\|_{H^{-1}}^2 \right\},
\end{aligned} \tag{5.8}$$

for $\varepsilon > 0$, where $\delta_1 = \frac{1}{2C} \min \left\{ \frac{1}{\gamma}, \frac{1}{\nu}, \frac{\nu}{\gamma^2} \right\}$.

Proof. Taking the inner products of (5.1) and (5.2) with $\frac{1}{\gamma^2} \phi$ and w , respectively, we have

$$\begin{aligned}
& \operatorname{Re} \lambda (\gamma^{-2} \|\phi\|_2^2 + \|w\|_2^2) + \nu \|\nabla w\|_2^2 + \tilde{\nu} \|\operatorname{div} w\|_2^2 \\
& = \operatorname{Re} \left\{ -\frac{1}{\gamma^2} (\operatorname{div}(\phi v), \phi) + \frac{1}{\gamma^2} (f^0, \phi) + (\tilde{f}, w) \right\}.
\end{aligned}$$

Since $\operatorname{Re}(\operatorname{div}(\phi v), \phi) = \frac{1}{2} \operatorname{Re}(\operatorname{div} v, |\phi|^2) \leq \|\operatorname{div} v\|_\infty \|\phi\|_2^2 \leq \|v\|_{H^3} \|\phi\|_2^2$, we have

$$\begin{aligned}
& \operatorname{Re} \lambda (\gamma^{-2} \|\phi\|_2^2 + \|w\|_2^2) + \nu \|\nabla w\|_2^2 + \tilde{\nu} \|\operatorname{div} w\|_2^2 \\
& \leq C \left\{ \frac{1}{\nu} \|\tilde{f}\|_{H^{-1}}^2 + \frac{1}{\varepsilon \gamma^4} \|f^0\|_2^2 + \frac{1}{\gamma^2} \|v\|_{H^3} \|\phi\|_2^2 + \varepsilon \|\phi\|_2^2 \right\}
\end{aligned}$$

for arbitrary $\varepsilon > 0$. We thus obtain (5.7).

We next prove (5.8). We have

$$\begin{aligned}
& \operatorname{Re} \lambda ((u, u)) + \frac{1}{\gamma^2} \operatorname{Re}(\operatorname{div}(\phi v) + \gamma^2 \operatorname{div} w, \phi) + \operatorname{Re}(\nabla \phi - \nu \Delta w - \tilde{\nu} \nabla \operatorname{div} w, w) \\
& - \delta_1 \operatorname{Re}(\nabla \phi - \nu \Delta w - \tilde{\nu} \nabla \operatorname{div} w, \mathcal{B} \phi) - \delta_1 \operatorname{Re}(\mathcal{B}(\gamma^2 \operatorname{div} w + \operatorname{div}(\phi v)), w) \\
& \geq \operatorname{Re} \lambda ((u, u)) + \nu \|\nabla w\|_2^2 + \tilde{\nu} \|\operatorname{div} w\|_2^2 - \frac{1}{\gamma^2} \|v\|_{H^3} \|\phi\|_2^2 \\
& - \delta_1 \operatorname{Re}(\nabla \phi - \nu \Delta w - \tilde{\nu} \nabla \operatorname{div} w, \mathcal{B} \phi) - \delta_1 \operatorname{Re}(\mathcal{B}(\gamma^2 \operatorname{div} w + \operatorname{div}(\phi v)), w).
\end{aligned}$$

By Lemma 2.3, we have

$$\begin{aligned}
& \operatorname{Re}(\nabla \phi - \nu \Delta w - \tilde{\nu} \nabla \operatorname{div} w, \mathcal{B} \phi) \leq -\frac{1}{2} \|\phi\|_2^2 + C \{ \nu^2 \|\nabla w\|_2^2 + \tilde{\nu}^2 \|\operatorname{div} w\|_2^2 \}, \\
& \operatorname{Re}(\mathcal{B}(\gamma^2 \operatorname{div} w + \operatorname{div}(\phi v)), w) \leq \|\mathcal{B}(\gamma^2 \operatorname{div} w + \operatorname{div}(\phi v))\|_2 \|w\|_2 \\
& \leq C \{ \gamma^2 \|w\|_2 + \|\phi v\|_2 \} \|w\|_2 \\
& \leq C \{ \gamma^2 \|w\|_2 + \|v\|_{H^4} \|\phi\|_2 \} \|w\|_2.
\end{aligned}$$

We set $\delta_1 = \frac{1}{2C} \min \left\{ \frac{1}{\gamma}, \frac{1}{\nu}, \frac{\nu}{\gamma^2} \right\}$ and $M = \min \left\{ \frac{\gamma^2}{\nu}, \nu, \gamma \right\}$. Then if $\|v\|_{H^4} \leq CM$, we have

$$\begin{aligned}
& \operatorname{Re}(\lambda + \Lambda)(\gamma^{-2} \|\phi\|_2^2 + \|w\|_2^2) + \frac{\nu}{2} \|\nabla w\|_2^2 + \frac{\tilde{\nu}}{2} \|\operatorname{div} w\|_2^2 + \frac{\delta_1}{2} \|\phi\|_2^2 \\
& \leq C \left\{ \left(\frac{1}{\delta_1 \gamma^4} + \frac{\delta_1^2}{\nu} \right) \|f^0\|_2^2 + \frac{1}{\nu} \|\tilde{f}\|_2^2 \right\}.
\end{aligned}$$

□

Lemma 5.3. *Let $\chi_0 \in C_{0,per}^\infty(\Omega_{per})$. Then*

$$\begin{aligned} & \operatorname{Re} \lambda (\gamma^{-2} \|\chi_0 \partial_x^k \phi\|_2^2 + \|\chi_0 \partial_x^k w\|_2^2) + \nu \|\chi_0 \nabla \partial_x^k w\|_2^2 + (\nu + \tilde{\nu}) \|\chi_0 \operatorname{div} \partial_x^k w\|_2^2 \\ & \leq C \left\{ \left(\nu + \tilde{\nu} + \frac{1}{\delta} \right) \|\partial_x^k w\|_2^2 + \frac{1}{\delta \gamma^4} \|f^0\|_{H^k}^2 + \frac{1}{\nu} \|\tilde{f}\|_{H^{k-1}}^2 \right. \\ & \quad \left. + \frac{1}{\gamma^2} \|v\|_{H^{3+(k-2)_+}} \|\phi\|_{H^k}^2 + \delta \|\phi\|_{H^k}^2 \right\} \end{aligned} \quad (5.9)$$

for $k = 1, 2, 3$ and arbitrary δ satisfying $0 < \delta < 1$.

Proof. Applying ∂_x to (5.1) and (5.2) and taking the inner product of the resulting equations with $\chi_0^2 \partial_x \phi$ and $\chi_0^2 \partial_x w$, respectively, we have

$$\begin{aligned} & \operatorname{Re} \lambda (\gamma^{-2} \|\chi_0 \partial_x \phi\|_2^2 + \|\chi_0 \partial_x w\|_{H^2}^2) + \nu \|\chi_0 \nabla \partial_x w\|_2^2 + \tilde{\nu} \|\chi_0 \operatorname{div} \partial_x w\|_2^2 \\ & = \operatorname{Re} \left\{ -\frac{1}{\gamma^2} (\partial_x \operatorname{div}(\phi v), \chi_0^2 \nabla \phi) + (-\nu \partial_x^2 w - \tilde{\nu} \partial_x \operatorname{div} w + \partial_x \phi, 2\chi_0 \nabla \chi_0 \partial_x w) \right. \\ & \quad \left. - (\tilde{f}, \chi_0^2 \partial_x^2 w + 2\chi_0 \nabla \chi_0 \partial_x w) + \frac{1}{\gamma^2} (\partial_x f^0, \chi_0^2 \nabla \phi) \right\}. \end{aligned}$$

Since $\operatorname{Re}(\partial_x \operatorname{div}(\phi v), \chi_0^2 \partial_x \phi) \leq C \|v\|_{H^3} \|\phi\|_{H^1}^2$, we have

$$\begin{aligned} & \operatorname{Re} \lambda (\gamma^{-2} \|\chi \partial_x \phi\|_2^2 + \|\chi \partial_x w\|_{H^2}^2) + \nu \|\chi_0 \nabla \partial_x w\|_2^2 + \tilde{\nu} \|\chi_0 \operatorname{div} \partial_x w\|_2^2 \\ & \leq C \left\{ \nu \|\nabla w\|_2^2 + \frac{1}{\delta} \|\nabla w\|_2^2 + \frac{1}{\nu} \|\tilde{f}\|_2^2 + \frac{1}{\gamma^2} \|v\|_{H^3} \|\phi\|_{H^1}^2 + \frac{1}{\gamma^4 \delta} \|f^0\|_{H^1}^2 + \delta \|\phi\|_{H^1}^2 \right\} \end{aligned}$$

for arbitrary δ satisfying $0 < \delta < 1$. We thus obtain (5.9). As for higher order derivatives, since $\operatorname{Re}(\partial_x^k \operatorname{div}(\phi v), \chi_0^2 \partial_x^k \phi) \leq C \|v\|_{H^{3+(k-2)_+}} \|\phi\|_{H^k}^2$ ($k = 2, 3$), we can prove (5.9) for $k = 2, 3$ in a similar manner. □

We next derive the estimates near the boundary. We introduce local coordinates $\Psi(y_1, \dots, y_{n-1}, y_n)$ near the boundary Σ_n . For each $\bar{x} \in \Sigma_n$, there exists an open neighborhood $\mathcal{O}$ of $\bar{x}$ such that each $x \in \mathcal{O}$ is written as

$$x = \Psi(y', y_n) \equiv \lambda(y') + y_n \mathbf{n}(\lambda(y')),$$

where $y' = (y_1, \dots, y_{n-1})$ and $\lambda(y')$ is local coordinates on $\Sigma_n \cap \mathcal{O}$ and $\mathbf{n}$ is the outward unit normal vector to Σ_n . We set $y = (y', y_n)$ and denote the tangential and normal derivatives by $\partial = (\partial_{y_1}, \dots, \partial_{y_{n-1}})$ and $\partial_N = \partial_{y_n}$, respectively.

As for the tangential derivatives, we have the following estimates.

Lemma 5.4. *Let $\chi \in C_0^\infty(\mathcal{O})$. Then*

$$\begin{aligned} & \operatorname{Re} \lambda (\gamma^{-2} \|\chi \partial^k \phi\|_2^2 + \|\chi \partial^k w\|_2^2) + \nu \|\chi \partial^k \nabla w\|_2^2 + (\nu + \tilde{\nu}) \|\chi \partial^k \operatorname{div} w\|_2^2 \\ & \leq C \left\{ \left(\nu + \tilde{\nu} + \frac{1}{\delta} \right) \|\nabla w\|_{H^{k-1}}^2 + \frac{1}{\delta \gamma^4} \|f^0\|_{H^k}^2 + \frac{1}{\nu} \|\tilde{f}\|_{H^{k-1}}^2 \right. \\ & \quad \left. + \frac{1}{\gamma^2} \|v\|_{H^{3+(k-2)_+}} \|\phi\|_{H^k}^2 + \delta \|\phi\|_{H^k}^2 \right\} \end{aligned} \quad (5.10)$$

for $k = 1, 2, 3$ and arbitrary δ satisfying $0 < \delta < 1$.

Proof. Using the local coordinates $\Psi(y)$, we rewrite (5.1)-(5.2) on $\mathcal{O} \cap \Omega_{per}$ as

$$\begin{cases} \lambda \tilde{\phi} + \gamma^2 a_{kj} \partial_{y_k} \tilde{w}^j + a_{kj} \partial_{y_k} (\tilde{\phi} \tilde{v}_s^j) = f^0 \circ \Psi, \\ \lambda \tilde{w}^j - \nu a_{ki} \partial_{y_k} (a_{si} \partial_{y_s} \tilde{w}^j) - \tilde{\nu} a_{kj} \partial_{y_k} (a_{si} \partial_{y_s} \tilde{w}^i) + a_{kj} \partial_{y_k} \tilde{\phi} = \tilde{f} \circ \Psi \end{cases} \quad (5.11)$$

on $\Psi^{-1}(\mathcal{O} \cap \Omega_{per})$, where $j = 1, \dots, n$, $\tilde{w}(y) = w(\Psi(y))$, $\tilde{\phi}(y) = \phi(\Psi(y))$, $\tilde{v}_s(y) = v_s(\Psi(y))$, $\tilde{\rho}_s(y) = \rho_s(\Psi(y))$, $\eta_n = 0$, $a_{kj} = a_{kj}(y)$ is the entry (k, j) of $(D\Psi)^{-1} = (D\Psi^{-1}) \circ \Psi$. $D\Psi = (\partial_{y_j} \Psi^k)$ is the Jacobian matrix of Ψ . Applying $\partial_{y'}^k$ to (5.11) and taking the inner product of the resulting equation with $\chi^2 \partial_{y'} \tilde{u}$, $\tilde{u} = {}^\top(\tilde{\phi}, \tilde{w})$, we have the desired estimate in a similar manner to the proof of Lemma 5.3. $\square$

Concerning the normal derivative of ϕ , we have the following estimates.

Lemma 5.5. *Let $\chi \in C_0^\infty(\mathcal{O})$. Then*

$$\begin{aligned} & \operatorname{Re} \lambda \frac{\nu + \tilde{\nu}}{\gamma^2} \|\chi \partial^k \partial_N^{l+1} \phi\|_2^2 + \|\chi \partial^k \partial_N^{l+1} \phi\|_2^2 \\ & \leq C \left\{ |\lambda|^2 \|w\|_{H^{k+l}}^2 + \nu^2 \|\chi \partial^{k+1} \partial_N^l \nabla w\|_2^2 + \frac{(\nu + \tilde{\nu})^2}{\gamma^4} \|f^0\|_{H^{k+l+1}}^2 \right. \\ & \quad \left. + \|\tilde{f}\|_{H^{k+l}}^2 + \frac{\nu + \tilde{\nu}}{\gamma^2} \|v\|_{H^{3+(k+l-2)_+}} \|\phi\|_{H^{k+l+1}}^2 \right\} \end{aligned} \quad (5.12)$$

for $k + l = 0, 1, 2$.

Proof. We apply $\frac{\nu + \tilde{\nu}}{\gamma^2} \partial_N$ to (5.1) and take the inner product of (5.2) with $\mathbf{n}$, we have

$$\begin{aligned} & \lambda \frac{\nu + \tilde{\nu}}{\gamma^2} \partial_N \phi + (\nu + \tilde{\nu}) \partial_N \operatorname{div} w + \frac{\nu + \tilde{\nu}}{\gamma^2} \partial_N \operatorname{div}(\phi v) = \frac{\nu + \tilde{\nu}}{\gamma^2} \partial_N f^0, \\ & \lambda w \cdot \mathbf{n} - \nu \Delta w \cdot \mathbf{n} - \tilde{\nu} \partial_N \operatorname{div} w + \partial_N \phi = \tilde{f} \cdot \mathbf{n}. \end{aligned} \quad (5.13)$$

Adding these two equations, we obtain

$$\begin{aligned}
& \lambda \frac{\nu + \tilde{\nu}}{\gamma^2} \partial_N \phi + \partial_N \phi \\
& = -\lambda w \cdot \mathbf{n} + \nu(\Delta w \cdot \mathbf{n} - \partial_N \operatorname{div} w) + \tilde{f} \cdot \mathbf{n} + \frac{\nu + \tilde{\nu}}{\gamma^2} \partial_N f^0 - \frac{\nu + \tilde{\nu}}{\gamma^2} \partial_N \operatorname{div}(\phi v).
\end{aligned} \tag{5.14}$$

Applying $\partial^k \partial_N^l$ ($k + l = 0, 1, 2$) to (5.14), we take the inner product of the resulting equation with $\chi^2 \partial^k \partial_N^{l+1} \phi$ ($k + l = 0, 1, 2$). By integration by parts, we have the desired estimate. $\square$

Lemma 5.6. *Let $\chi \in C_0^\infty(\mathcal{O})$. Then*

$$\begin{aligned}
& (\nu + \tilde{\nu}) \|\chi \partial^k \partial_N^{l+1} \operatorname{div} w\|_2^2 \\
& \leq C \left\{ \frac{|\lambda|^2}{\nu + \tilde{\nu}} \|w\|_{H^{k+l}}^2 + \nu \|\chi \partial^{k+1} \partial_N^l \nabla w\|_2^2 + \frac{1}{\nu + \tilde{\nu}} \|\chi \partial^k \partial_N^{l+1} \phi\|_2^2 \right. \\
& \quad \left. + \frac{\nu + \tilde{\nu}}{\gamma^4} \|f^0\|_{H^{k+l+1}}^2 + \frac{1}{\nu + \tilde{\nu}} \|\tilde{f}\|_{H^{k+l}}^2 + \frac{1}{\gamma^2} \|v\|_{H^{3+(k+l-2)_+}} \|\phi\|_{H^{k+l+1}}^2 \right\}
\end{aligned} \tag{5.15}$$

for $k + l = 0, 1, 2$.

Proof. Adding $(\nu + \tilde{\nu}) \partial_N \operatorname{div} w$ to (5.13), we have

$$(\nu + \tilde{\nu}) \partial_N \operatorname{div} w = \lambda w \cdot \mathbf{n} + \nabla \phi \cdot \mathbf{n} - \nu(\Delta w \cdot \mathbf{n} - \partial_N \operatorname{div} w) - \tilde{f} \cdot \mathbf{n}. \tag{5.16}$$

Multiplying (5.16) by $\chi^2 \partial_N \operatorname{div} w$ and integrating the resulting equation, we obtain

$$(\nu + \tilde{\nu}) \|\chi \partial_N \operatorname{div} w\|_2^2 \leq C \left\{ \frac{|\lambda|^2}{\nu + \tilde{\nu}} \|w\|_2^2 + \frac{1}{\nu + \tilde{\nu}} \|\chi \partial_N \phi\|_2^2 + \nu \|\chi \partial \nabla w\|_2^2 + \frac{1}{\nu + \tilde{\nu}} \|\tilde{f}\|_2^2 \right\}.$$

As for higher order derivatives we apply $\partial^k \partial_N^l$ to (5.16) and take the inner product of the resulting equation with $\chi^2 \partial^k \partial_N^{l+1} \operatorname{div} w$. Substituting (5.12) for the resulting equation, we have the desired estimate. $\square$

We next estimate $\|\chi \partial^k \partial_x^{l+2} w\|_2^2$ for $k + l = 0, 1$.

Lemma 5.7. *Let $\chi \in C_0^\infty(\mathcal{O})$. Then, for arbitrary δ satisfying $0 < \delta < 1$,*

$$\begin{aligned}
& \nu \|\chi \partial^k \partial_x^{l+2} w\|_2^2 + \frac{1}{\nu} \|\chi \partial^k \partial_x^{l+1} \phi\|_2^2 \\
& \leq C \left\{ \frac{\lambda^2}{\nu} \|w\|_{H^{k+l}}^2 + (\nu + \tilde{\nu}) \|\nabla w\|_{H^{k+l}}^2 + \frac{1}{\nu} \|\phi\|_{H^{k+l}}^2 + \frac{1}{\nu} \|\tilde{f}\|_{H^{k+l}}^2 \right. \\
& \quad \left. + (\nu + \tilde{\nu}) \|\chi \partial^k \partial_x^{l+1} \operatorname{div} w\|_2^2 \right\}
\end{aligned} \tag{5.17}$$

for $k + l = 1, 2$, $k \geq 1$.

Proof. Applying $\chi\partial^k(k = 1, 2)$ to (5.1) and (5.2), it holds that

$$\begin{cases} \operatorname{div}(\chi\partial^k w) = F^0 & \text{in } \mathcal{O} \cap \Omega_{per}, \\ -\Delta(\chi\partial^k w) + \frac{1}{\nu}\nabla(\chi\partial^k \phi) = \tilde{F} & \text{in } \mathcal{O} \cap \Omega_{per}, \\ (\chi\partial^k w)|_{\Sigma_{j,-}} = (\chi\partial^k w)|_{\Sigma_{j,+}}, & (j = 1, \dots, n-1), \\ (\chi\partial^k w)|_{\Sigma_n} = 0, \end{cases} \quad (5.18)$$

where

$$\begin{aligned} F^0 &= [\nabla, \chi\partial^k]w - \lambda\chi\partial^k \phi - \frac{1}{\gamma^2}\chi\partial^k \operatorname{div}(\phi v) + \frac{1}{\gamma^2}\chi\partial^k f^0, \\ \tilde{F} &= [\Delta, \chi\partial^k]w + \frac{1}{\nu}[\nabla, \chi\partial^k]\phi - \frac{\lambda}{\nu}\chi\partial^k w - \frac{\tilde{\nu}}{\nu}\chi\partial^k \nabla \operatorname{div} w + \frac{1}{\nu}\chi\partial^k \tilde{f}. \end{aligned}$$

By using the estimates for the Stokes problem (see, e.g., [8]), we have the desired estimate. $\square$

Lemma 5.8. *Let $u = {}^\top(\phi, w)$ be a solution of (5.1)-(5.4). Then*

$$\begin{aligned} &\frac{|\lambda|^2}{2}\|w\|_2^2 + \operatorname{Re}\lambda\nu\|\nabla w\|_2^2 + \operatorname{Re}\lambda\tilde{\nu}\|\operatorname{div} w\|_2^2 \\ &\leq C\left\{\frac{1}{\gamma^2}\|f^0\|_2^2 + \|\tilde{f}\|_2^2 + \frac{1}{\gamma^2}\|v\|_{H^3}^2\|\phi\|_{H^1}^2 + \gamma^2\|\operatorname{div} w\|_2^2\right\}. \end{aligned} \quad (5.19)$$

Proof. We take the inner product of (5.2) with λw , we have

$$|\lambda|^2\|w\|_2^2 + \lambda\nu\|\nabla w\|_2^2 + \lambda\tilde{\nu}\|\operatorname{div} w\|_2^2 - (\bar{\lambda}\phi, \operatorname{div} w) = (\tilde{f}, \lambda w). \quad (5.20)$$

Substituting (5.1) for $\lambda\phi$ in (5.20), the forth term of left-hand side is written as $-(\bar{\lambda}\phi, \operatorname{div} w) = \frac{\bar{\lambda}}{\lambda}(\gamma^2\operatorname{div} w + \operatorname{div}(\phi v) - f^0, \operatorname{div} w)$. It then follows that

$$\begin{aligned} &\frac{|\lambda|^2}{2}\|w\|_2^2 + \operatorname{Re}\lambda\nu\|\nabla w\|_2^2 + \operatorname{Re}\lambda\tilde{\nu}\|\operatorname{div} w\|_2^2 \\ &\leq C\left\{\|\tilde{f}\|_2^2 + \frac{1}{\gamma^2}\|f^0\|_2^2 + \frac{1}{\gamma^2}\|v\|_{H^3}^2\|\phi\|_{H^1}^2 + \gamma^2\|\operatorname{div} w\|_2^2\right\}. \end{aligned}$$

$\square$

Lemma 5.9. *Let $u = {}^\top(\phi, w)$ be a solution of (5.1)-(5.4). Then*

$$\nu\|w\|_{H^{k+2}}^2 + \frac{1}{\nu}\|\phi\|_{H^{k+1}}^2 \leq C\left\{\frac{|\lambda|^2}{\nu}\|w\|_{H^k}^2 + (\nu + \tilde{\nu})\|\operatorname{div} w\|_{H^{k+1}}^2 + \frac{1}{\nu}\|\tilde{f}\|_{H^k}^2\right\} \quad (5.21)$$

for $k = 0, 1, 2$.

Proof. We write (5.1)-(5.2) as

$$\begin{cases} \operatorname{div} w = -\lambda\phi - \frac{1}{\gamma^2}\operatorname{div}(\phi w) + \frac{1}{\gamma^2}f^0, & \text{in } \Omega_{per} \\ -\nu\Delta w + \nabla\phi = -\lambda w + \tilde{\nu}\nabla\operatorname{div} w + \tilde{f}, & \text{in } \Omega_{per} \\ w|_{\Sigma_n} = 0, \quad \phi|_{\Sigma_{j,-}} = \phi|_{\Sigma_{j,+}}, \quad w|_{\Sigma_{j,-}} = w|_{\Sigma_{j,+}} \quad (j = 1, \dots, n-1), & \text{on } \partial\Omega_{per}. \end{cases} \quad (5.22)$$

We set $\tilde{F} = \tilde{\nu}\nabla\operatorname{div} w + \tilde{f}$. Applying the estimate for Stokes problem, we have

$$\begin{aligned} \|w\|_{H^{k+2}}^2 + \frac{1}{\nu^2}\|\phi\|_{H^{k+1}}^2 &\leq C\left\{\frac{|\lambda|^2}{\nu^2}\|w\|_{H^k}^2 + \|\operatorname{div} w\|_{H^{k+1}}^2 + \frac{1}{\nu^2}\|\tilde{F}\|_{H^k}^2\right\} \\ &\leq C\left\{\frac{|\lambda|^2}{\nu^2}\|w\|_{H^k}^2 + \left(1 + \frac{\tilde{\nu}^2}{\nu^2}\right)\|\nabla\operatorname{div} w\|_{H^k}^2 + \frac{1}{\nu^2}\|\tilde{f}\|_{H^k}^2\right\} \\ &\leq C\left\{\frac{|\lambda|^2}{\nu^2}\|w\|_{H^k}^2 + \left(1 + \frac{\tilde{\nu}}{\nu}\right)\|\nabla\operatorname{div} w\|_{H^k}^2 + \frac{1}{\nu^2}\|\tilde{f}\|_{H^k}^2\right\}. \end{aligned} \quad (5.23)$$

Multiplying (5.23) by ν , we obtain (5.21). $\square$

Proof of Theorem 5.1. Estimate (5.5)₀ follows from (5.8).

Let $b_j^{(1)}$ ($j = 1, \dots, 4$) be positive numbers independent of ν , $\tilde{\nu}$ and γ and consider

$$\begin{aligned} &(5.7) + \frac{1}{\nu + \tilde{\nu}} \times (5.19) + b_1^{(1)} \times \{(5.9)_{k=1} + (5.10)_{k=1}\} \\ &+ \frac{b_2^{(1)}}{\nu} \times (5.12)_{k=l=0} + b_3^{(1)} \times (5.15)_{k=l=0} + b_4^{(1)} \times (5.21)_{k=0}. \end{aligned}$$

Taking $b_j^{(1)}$ ($j = 1, \dots, 4$) suitably small and $\varepsilon = \delta = \frac{1}{2C\nu}$, we have

$$\begin{aligned} &\operatorname{Re}\lambda(\gamma^{-2}\|\phi\|_{H^1}^2 + \|w\|_{H^1}^2 + \|\operatorname{div} w\|_2^2) + \frac{1}{2\nu}\|\phi\|_{H^1}^2 + \frac{1}{2}\nu\|w\|_{H^2}^2 \\ &+ \frac{1}{2}(\nu + \tilde{\nu})\|\operatorname{div} w\|_{H^1}^2 + \frac{1}{2}\frac{|\lambda|^2}{\nu + \tilde{\nu}}\|w\|_2^2 \\ &\leq C\left\{\nu\|\nabla w\|_2^2 + \frac{1}{\gamma^2}\|v\|_{H^3}\|\phi\|_{H^1}^2 + \frac{1}{\nu\gamma^2}\|v\|_{H^3}^2\|\phi\|_{H^1}^2 + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\operatorname{div} w\|_2^2 \right. \\ &\quad \left. + \left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu\gamma^2}\right)\|f^0\|_{H^1}^2 + \frac{1}{\nu}\|\tilde{f}\|_2^2\right\}. \end{aligned} \quad (5.24)$$

Computing $\frac{1}{2C} \times (5.24) + (5.7)$ with $\varepsilon = \frac{1}{8C\nu} + (5.7)$ with $\varepsilon = \frac{\nu}{16C\gamma^2}$, we have

$$\begin{aligned}
& \operatorname{Re}\lambda(\gamma^{-2}\|\phi\|_{H^1}^2 + \|w\|_{H^1}^2 + \|\operatorname{div}w\|_2^2) + \frac{1}{8\nu}\|\phi\|_{H^1}^2 + \frac{1}{2}\nu\|w\|_{H^2}^2 \\
& + \frac{1}{2}(\nu + \tilde{\nu})\|\operatorname{div}w\|_{H^1}^2 + \frac{1}{2}\frac{|\lambda|^2}{\nu + \tilde{\nu}}\|w\|_2^2 + \frac{\operatorname{Re}\lambda}{(\nu + \tilde{\nu})^2}(\|\phi\|_2^2 + \gamma^2\|w\|_2^2) \\
& + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\nabla w\|_2^2 + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\operatorname{div}w\|_2^2 \\
& \leq C\left\{\left(\frac{1}{\nu^2} + \frac{1}{\gamma^2}\right)\|v\|_{H^3}\|\phi\|_{H^1}^2 + \frac{1}{\nu\gamma^2}\|v\|_{H^3}^2\|\phi\|_{H^1}^2\right. \\
& \quad \left.+ \left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3}\right)\|f^0\|_{H^1}^2 + \left(\frac{1}{\nu} + \frac{\gamma^2}{\nu^3}\right)\|\tilde{f}\|_2^2\right\}.
\end{aligned}$$

If v satisfies $\|v\|_{H^3} \leq \frac{1}{32C} \min\{\frac{\gamma^2}{\nu}, \nu, \gamma\}$, then

$$\begin{aligned}
& \operatorname{Re}\lambda(\gamma^{-2}\|\phi\|_{H^1}^2 + \|w\|_{H^1}^2 + \|\operatorname{div}w\|_2^2) + \frac{1}{16\nu}\|\phi\|_{H^1}^2 + \frac{1}{2}\nu\|w\|_{H^2}^2 \\
& + \frac{1}{2}(\nu + \tilde{\nu})\|\operatorname{div}w\|_{H^1}^2 + \frac{1}{2}\frac{|\lambda|^2}{\nu + \tilde{\nu}}\|w\|_2^2 + \frac{\operatorname{Re}\lambda}{(\nu + \tilde{\nu})^2}(\|\phi\|_2^2 + \gamma^2\|w\|_2^2) \\
& + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\nabla w\|_2^2 + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\operatorname{div}w\|_2^2 \\
& \leq C\left\{\left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3}\right)\|f^0\|_{H^1}^2 + \left(\frac{1}{\nu} + \frac{\gamma^2}{\nu^3}\right)\|\tilde{f}\|_2^2\right\}.
\end{aligned} \tag{5.25}$$

Furthermore, if $\lambda \geq -\Lambda_1 := -\frac{1}{32} \min\{\frac{\gamma^2}{\nu}, \nu + \tilde{\nu}, \nu\}$, we see from (5.25) that

$$\begin{aligned}
& \operatorname{Re}(\lambda + \Lambda_1)(\gamma^{-2}\|\phi\|_{H^1}^2 + \|w\|_{H^1}^2) + \frac{1}{\nu}\|\phi\|_{H^1}^2 + \nu\|w\|_{H^2}^2 + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\nabla w\|_2^2 \\
& \leq C\left\{\left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3}\right)\|f^0\|_{H^1}^2 + \left(\frac{1}{\nu} + \frac{\gamma^2}{\nu^3}\right)\|\tilde{f}\|_2^2\right\},
\end{aligned} \tag{5.26}$$

and hence we have (5.5)₁ and

$$\nu\|w\|_{H^2}^2 \leq C\left\{\left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3}\right)\|f^0\|_{H^1}^2 + \left(\frac{1}{\nu} + \frac{\gamma^2}{\nu^3}\right)\|\tilde{f}\|_2^2\right\}. \tag{5.27}$$

Let $b_j^{(2)}$ ($j = 1, 4, 5$) and $b_{j,kl}^{(2)}$ ($j = 2, 3$) be positive numbers independent of $\nu, \tilde{\nu}$ and γ and consider

$$\begin{aligned}
& (5.25) + \frac{1}{\nu} \frac{|\lambda|^2}{\nu + \tilde{\nu}} \times (5.8) + b_1^{(2)} \times \{(5.9)_{k=2} + (5.10)_{k=2}\} \\
& + \sum_{k+l=1} b_{2,kl}^{(2)} \times (5.12) + b_{3,kl}^{(2)} \times (5.15) + b_4^{(2)} \times (5.17)_{k=1,l=0} + b_5^{(2)} \times (5.21)_{k=1}.
\end{aligned}$$

Taking $b_j^{(2)}$ ($j = 1, 4, 5$) and $b_{j,kl}^{(2)}$ ($j = 2, 3$) suitably small and $\varepsilon = \frac{1}{2C\nu}$, we have

$$\begin{aligned}
& \operatorname{Re}\lambda(\gamma^{-2}\|\phi\|_{H^2}^2 + \|w\|_{H^1}^2 + \|\chi_0\partial^2 w\|_2^2 + \|\chi\partial^2 w\|_2^2) + \frac{1}{2\nu}\|\phi\|_{H^2}^2 + \frac{1}{2}\nu\|w\|_{H^3}^2 \\
& + \frac{1}{2}(\nu + \tilde{\nu})\|\operatorname{div} w\|_{H^2}^2 + \frac{1}{2}\frac{|\lambda|^2}{\nu + \tilde{\nu}}\|w\|_{H^1}^2 + \frac{|\lambda|^2}{\nu + \tilde{\nu}}\|\operatorname{div} w\|_2^2 \\
& + \operatorname{Re}\lambda\|\operatorname{div} w\|_2^2 + \frac{\operatorname{Re}\lambda}{(\nu + \tilde{\nu})^2}\left(\|\phi\|_2^2 + \gamma^2\|w\|_2^2\right) + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\nabla w\|_2^2 \\
& + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\operatorname{div} w\|_2^2 + \frac{|\lambda|^2\operatorname{Re}\lambda}{(\nu + \tilde{\nu})^2}(\gamma^{-2}\|\phi\|_2^2 + \|w\|_2^2) + \frac{|\lambda|^2\delta_1}{(\nu + \tilde{\nu})^2}\|\phi\|_2^2 \\
& \leq C\left\{\nu\|\partial_x^2 w\|_2^2 + \left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3}\right)\|f^0\|_{H^2}^2 + \frac{1}{\nu}\|\tilde{f}\|_{H^1}^2 + \frac{\gamma^2}{\nu^3}\|\tilde{f}\|_2^2\right. \\
& \quad \left.+ \frac{1}{\gamma^2}\|v\|_{H^3}\|\phi\|_{H^2}^2 + \frac{|\lambda|^2}{(\nu + \tilde{\nu})^2}\left(\left(\frac{1}{\delta_1\gamma^4} + \frac{\delta_1^2}{\nu}\right)\|f^0\|_2^2 + \frac{1}{\nu}\|\tilde{f}\|_{H^{-1}}^2\right)\right\}.
\end{aligned} \tag{5.28}$$

Let v satisfy $\|v\|_{H^3} \leq \frac{1}{8C} \min\{\frac{\gamma^2}{\nu}, \gamma, \nu\}$. Computing $\frac{1}{2C} \times (5.28) + (5.25)$ we have

$$\begin{aligned}
& \operatorname{Re}\lambda(\gamma^{-2}\|\phi\|_{H^2}^2 + \|w\|_{H^1}^2 + \|\chi_0\partial^2 w\|_2^2 + \|\chi\partial^2 w\|_2^2) + \frac{1}{2\nu}\|\phi\|_{H^2}^2 + \frac{1}{2}\nu\|w\|_{H^3}^2 \\
& + \frac{1}{2}(\nu + \tilde{\nu})\|\operatorname{div} w\|_{H^2}^2 + \frac{1}{2}\frac{|\lambda|^2}{\nu + \tilde{\nu}}\|w\|_{H^1}^2 + \frac{|\lambda|^2}{\nu + \tilde{\nu}}\|\operatorname{div} w\|_2^2 \\
& + \operatorname{Re}\lambda\|\operatorname{div} w\|_2^2 + \frac{\operatorname{Re}\lambda}{(\nu + \tilde{\nu})^2}\left(\|\phi\|_2^2 + \gamma^2\|w\|_2^2\right) + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\nabla w\|_2^2 \\
& + \frac{\gamma^2}{\nu + \tilde{\nu}}\|\operatorname{div} w\|_2^2 + \frac{|\lambda|^2\operatorname{Re}\lambda}{(\nu + \tilde{\nu})^2}(\gamma^{-2}\|\phi\|_2^2 + \|w\|_2^2) + \frac{|\lambda|^2\delta_1}{(\nu + \tilde{\nu})^2}\|\phi\|_2^2 \\
& \leq C\left\{\frac{|\lambda|^2}{(\nu + \tilde{\nu})^2}\left(\left(\frac{1}{\delta_1\gamma^4} + \frac{\delta_1^2}{\nu}\right)\|f^0\|_2^2 + \frac{1}{\nu}\|\tilde{f}\|_{H^{-1}}^2\right)\right. \\
& \quad \left.+ \left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3}\right)\|f^0\|_{H^2}^2 + \frac{1}{\nu}\|\tilde{f}\|_{H^1}^2 + \frac{\gamma^2}{\nu^3}\|\tilde{f}\|_2^2\right\}.
\end{aligned}$$

If λ satisfies $\lambda \geq -\Lambda_2 := -\frac{1}{8} \min\{\nu, \delta_1 \gamma^2\}$, we have

$$\begin{aligned}
& \operatorname{Re} \lambda (\gamma^{-2} \|\phi\|_{H^2}^2 + \|w\|_{H^1}^2) + \frac{1}{\nu} \|\phi\|_{H^2}^2 + \nu \|w\|_{H^3}^2 + \frac{\gamma^2}{\nu + \tilde{\nu}} \|\nabla w\|_2^2 \\
& + (\nu + \tilde{\nu}) \|\operatorname{div} w\|_{H^2}^2 + \frac{|\lambda|^2}{\nu + \tilde{\nu}} \|w\|_{H^1}^2 + \frac{|\lambda|^2}{\nu + \tilde{\nu}} \|\operatorname{div} w\|_2^2 \\
& \leq C \left\{ \frac{|\lambda|^2}{(\nu + \tilde{\nu})^2} \left(\left(\frac{1}{\delta_1 \gamma^4} + \frac{\delta_1^2}{\nu} \right) \|f^0\|_2^2 + \frac{1}{\nu} \|\tilde{f}\|_{H^{-1}}^2 \right) \right. \\
& \quad \left. + \left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3} \right) \|f^0\|_{H^2}^2 + \frac{1}{\nu} \|\tilde{f}\|_{H^1}^2 + \frac{\gamma^2}{\nu^3} \|\tilde{f}\|_2^2 \right\}
\end{aligned} \tag{5.29}$$

and

$$\begin{aligned}
& \operatorname{Re}(\lambda + \Lambda_2) (\gamma^{-2} \|\phi\|_{H^2}^2 + \|w\|_{H^1}^2) + \frac{1}{\nu} \|\phi\|_{H^2}^2 + \nu \|w\|_{H^3}^2 + \frac{\gamma^2}{\nu + \tilde{\nu}} \|\nabla w\|_2^2 \\
& \leq C \left\{ \frac{|\lambda|^2}{(\nu + \tilde{\nu})^2} \left(\left(\frac{1}{\delta_1 \gamma^4} + \frac{\delta_1^2}{\nu} \right) \|f^0\|_2^2 + \frac{1}{\nu} \|\tilde{f}\|_{H^{-1}}^2 \right) \right. \\
& \quad \left. + \left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3} \right) \|f^0\|_{H^2}^2 + \frac{1}{\nu} \|\tilde{f}\|_{H^1}^2 + \frac{\gamma^2}{\nu^3} \|\tilde{f}\|_2^2 \right\}.
\end{aligned}$$

Let $b_j^{(3)}$ ($j = 1, 5$) and $b_{j,kl}^{(3)}$ ($j = 2, 3, 4$) be positive numbers independent of $\nu, \tilde{\nu}$ and γ and consider

$$\begin{aligned}
& (5.29) + \frac{1}{\nu} \frac{\lambda^2}{\nu + \tilde{\nu}} \times (5.27) + b_1^{(3)} \times \{(5.9)_{k=3} + (5.10)_{k=3}\} \\
& + \sum_{k+l=2} \left\{ \frac{b_{2,kl}^{(3)}}{\nu} \times (5.12) + b_{3,kl}^{(3)} \times (5.15) \right\} \\
& + \sum_{\substack{k+l=2 \\ k \geq 1}} b_{4,kl}^{(3)} \times (5.17) + b_5^{(3)} \times (5.21)_{k=2}.
\end{aligned}$$

Similarly, we obtain

$$\begin{aligned}
& \operatorname{Re}(\lambda + \Lambda_3) (\gamma^{-2} \|\phi\|_{H^3}^2 + \|w\|_{H^2}^2) + \frac{1}{\nu} \|\phi\|_{H^3}^2 + \nu \|w\|_{H^4}^2 + \frac{\gamma^2}{\nu + \tilde{\nu}} \|\nabla w\|_2^2 \\
& \leq C \left\{ \frac{|\lambda|^2}{(\nu + \tilde{\nu})^2} \left(\left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3} + \frac{1}{\delta_1 \gamma^4} + \frac{\delta_1^2}{\nu} \right) \|f^0\|_2^2 + \left(\frac{1}{\nu} + \frac{\gamma^2}{\nu^3} \right) \|\tilde{f}\|_{H^{-1}}^2 \right) \right. \\
& \quad \left. + \left(\frac{\nu + \tilde{\nu}}{\gamma^4} + \frac{1}{\nu^3} \right) \|f^0\|_{H^3}^2 + \frac{1}{\nu} \|\tilde{f}\|_{H^2}^2 + \frac{\gamma^2}{\nu^3} \|\tilde{f}\|_2^2 \right\}.
\end{aligned}$$

Taking $\Lambda = \max\{\Lambda_1, \Lambda_2, \Lambda_3\}$, we have (5.5)₁-(5.5)₃.

When $\lambda = 0$, we can obtain the estimate $(5.6)_k$ for $k = 1, 2, 3$ in a similar manner to above. In fact, estimate $(5.6)_k$ with $k = 1$ is obtained as follows. We do not use (5.19) in the computation for (5.24) and obtain

$$\begin{aligned} & \frac{1}{2\nu} \|\phi\|_{H^1}^2 + \frac{1}{2} \nu \|w\|_{H^2}^2 + \frac{1}{2} (\nu + \tilde{\nu}) \|\operatorname{div} w\|_{H^1}^2 \\ & \leq C \left\{ \nu \|\nabla w\|_2^2 + \frac{1}{\gamma^2} \|v\|_{H^3} \|\phi\|_{H^1}^2 + \frac{\nu + \tilde{\nu}}{\gamma^4} \|f^0\|_{H^1}^2 + \frac{1}{\nu} \|\tilde{f}\|_2^2 \right\}, \end{aligned} \quad (5.30)$$

instead of (5.24). Computing $\frac{1}{2C} \times (5.30) + (5.7)$ with $\varepsilon = \frac{1}{8C\nu}$, we have

$$\begin{aligned} & \frac{1}{2\nu} \|\phi\|_{H^1}^2 + \frac{1}{2} \nu \|w\|_{H^2}^2 + \frac{1}{2} (\nu + \tilde{\nu}) \|\operatorname{div} w\|_{H^1}^2 \\ & \leq C \left\{ \frac{1}{\gamma^2} \|v\|_{H^3} \|\phi\|_{H^1}^2 + \frac{\nu + \tilde{\nu}}{\gamma^4} \|f^0\|_{H^1}^2 + \frac{1}{\nu} \|\tilde{f}\|_2^2 \right\}. \end{aligned} \quad (5.31)$$

The desired estimate $(5.6)_k$ with $k = 1$ follows from (5.31) by taking $\|v\|_{H^3}$ suitably small. One can prove $(5.6)_k$ for $k = 2, 3$ in a similar manner. $\square$

b. Existence of solution of (5.1)-(5.4)

In this subsection we prove the existence of a solution $u = {}^\top(\phi, w)$ of (5.1)-(5.4).

Let $k = 0, 1, 2, 3$. We write (5.1)-(5.2) in the form

$$(\lambda + \mathcal{A}_k[v])u = f \quad (5.32)$$

for $u = {}^\top(\phi, w) \in D(\mathcal{A}_k[v])$ and $f = {}^\top(f^0, \tilde{f}) \in H_{per}^k \times H_{per}^{k-1}$. Here $\mathcal{A}_k[v] : H_{per,*}^k \times H_{per}^{k-1} \longrightarrow H_{per,*}^k \times H_{per}^{k-1}$ is the operator defined by

$$\mathcal{A}_k[v] = \begin{pmatrix} \operatorname{div}(\cdot v) & \gamma^2 \operatorname{div} \\ \nabla & -\nu \Delta - \tilde{\nu} \nabla \operatorname{div} \end{pmatrix}$$

with domain

$$D(\mathcal{A}_k[v]) = \{u \in H_{per,*}^k \times (H_{per}^{k+1} \cap H_{0,per}^1); \mathcal{A}_k[v]u \in H_{per,*}^k \times H_{per}^{k-1}\}.$$

If we prove the existence of solution $u \in D(\mathcal{A}_k[v])$ of (5.32) for all $f \in H_{per,*}^k \times H_{per}^{k+1}$ with $\lambda \gg 1$, then we see from estimates (5.5)₀-(5.5)₃ that $\{\lambda \in \mathbb{C}; \operatorname{Re} \lambda \geq -\Lambda\} \subset \rho(-\mathcal{A}_k[v])$ by a standard continuation argument.

The existence of solution of (5.32) for $\lambda \gg 1$ can be shown in a similar manner to the proof of Lemma 4.3. We first consider the problem

$$\lambda \phi + \operatorname{div}(\phi v) = F^0, \quad F^0 = f^0 - \gamma^2 \operatorname{div} \tilde{w} \quad (5.33)$$

for a given $\tilde{w} \in H_{per}^{k+1} \cap H_{0,per}^1$ and $f^0 \in H_{per,*}^k$. Applying the results of [10], we see that there exists a constant λ_0 satisfying $\lambda_0 \geq \|v\|_{H^3}$ such that if $\lambda \geq \lambda_0$, then the solution operator for (5.33) is an isomorphism on $H_{per,*}^k$ with operator norm bounded by $\frac{C}{\lambda}$ uniformly for $\lambda \gg 1$. We denote this solution operator by $\phi(\tilde{w})$. We next consider the problem

$$\begin{cases} \lambda w - \nu \Delta w - \tilde{\nu} \nabla \operatorname{div} w = \tilde{F}, \\ w|_{\Sigma_{j,-}} = w|_{\Sigma_{j,+}}, \quad w|_{\Sigma_n} = 0, \end{cases} \quad (5.34)$$

where $\tilde{F} = \tilde{f} - \nabla \phi$ with $\phi = \phi(\tilde{w})$. Since $-\nu \Delta w - \tilde{\nu} \nabla \operatorname{div} w$ is strongly elliptic, the solution operator for (5.34) provides an isomorphism from H_{per}^{k-1} to $H_{per}^{k+1} \cap H_{0,per}^1$. One can then define the operator $\Gamma : H_{per}^{k+1} \cap H_{0,per}^1 \rightarrow H_{per}^{k+1} \cap H_{0,per}^1$ by the solution operator for (5.34), i.e., $\Gamma(\tilde{w}) = w$ for $\tilde{w} \in H_{per}^{k+1} \cap H_{0,per}^1$, where w is the solution of (5.34). In a similar manner to the proof of Lemma 4.3, one can prove that there exists a unique $w \in H_{per}^{k+1} \cap H_{0,per}^1$ satisfying $\Gamma(w) = w$ if $\lambda \gg 1$. For this fixed point w , we set $\phi = \phi(w)$. Then by (5.33)-(5.34), we see that $u = {}^\top(\phi, w) \in D(\mathcal{A}_k[v])$ is a solution of (5.32) with $\lambda \gg 1$. This completes the proof. $\square$

Remark 5.10. *The unique existence of (5.33) for $k = 0, 1, 2$ follows from [10, Theorems 1, 2, 9] respectively. As for the case $k = 3$, it was proved in [10, Theorem 11] under the assumption $v \in W^{m,p}$ for some $p > n$, where n is the space dimension. This condition for v can be replaced by $v \in H^4$ in our case $k = 3$. In fact, one can show the unique existence of solution $\phi \in H^3$ of (5.33) when $v \in H^4$ as in the proof of [10, Theorem 11]. The point is to guarantee $\phi \nabla \operatorname{div} v \in H^2$ for $\phi \in H^2$, which follows from $v \in H^4$.*

Acknowledgements. Y. Kagei was partly supported by JSPS KAKENHI Grant Number 24340028, 16H03947, 15K13449, 24224003.

References

- [1] Aoyama, R., Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a parallel flow in a cylindrical domain, Kyushu J. Math., **69** (2015), pp. 293–343.
- [2] Aoyama, R. and Kagei, Y., Spectral properties of the semigroup for the linearized compressible Navier-Stokes equation around a parallel flow in a cylindrical domain, Adv. Differential Equations, **21** (2016), pp. 265–300.

- [3] Aoyama, R. and Kagei, Y., Large time behavior of solutions to the compressible Navier-Stokes equations around a parallel flow in a cylindrical domain, *Nonlinear Anal.*, **127** (2015), pp. 362–396.
- [4] Bogovskii, M. E., Solution of the first boundary value problem for an equation of continuity of an incompressible medium, *Soviet Math. Dokl.*, **20** (1979), pp. 1094–1098.
- [5] Březina, J., Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a time-periodic parallel flow, *SIAM J. Math. Anal.* **45** (2013), pp. 3514–3574.
- [6] Březina, J. and Kagei, Y., Spectral properties of the linearized compressible Navier-Stokes equation around time-periodic parallel flow, *J. Differential Equations* **255** (2013), pp. 1132–1195.
- [7] Deckelnick, K., Decay estimates for the compressible Navier-Stokes equations in unbounded domains, *Math. Z.* **209** (1992), pp. 115–130.
- [8] Galdi G. P., *An Introduction to the Mathematical Theory of the Navier-Stokes Equations*, Vol. 1, Springer-Verlag, New York (1994).
- [9] Hoff, D. and Zumbrun, K., Multi-dimensional diffusion waves for the Navier-Stokes equations of compressible flow, *Indiana Univ. Math. J.* **44** (1995), pp. 604–676.
- [10] Heywood, J.G. and Padula M., On the steady transport equation. in *Fundamental Directions in Mathematical Fluid Mechanics*, ed. by Galdi, G.P., Heywood, J.G., Rannacher, R. Birkhäuser, (2000), pp. 149–170.
- [11] Iooss, G. and Padula, M., Structure of the linearized problem for compressible parallel fluid flows, *Ann. Univ. Ferrara Sez.*, **VII 43** (1998), pp. 157–171.
- [12] Kagei, Y., Asymptotic behavior of solutions of the compressible Navier-Stokes equation around a parallel flow, *Arch. Rational Mech. Anal.*, **205** (2012), pp. 585–650.
- [13] Kagei Y., Nagafuchi Y. and Sudou T., Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow, *Journal of Math-for-Industry*, **2** (2010A), pp. 39–56. Correction to "Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow" in *J. Math-for-Ind.*, **2** (2010A), pp. 39–56, *J. Math-for-Ind.*, **2** (2010B), pp. 235.

- [14] Kagei, Y. and Kobayashi, T., Asymptotic Behavior of Solutions of the Compressible Navier-Stokes Equation on the Half Space, Arch. Rational Mech. Anal., **177** (2005), pp. 231–330.
- [15] Kagei, Y., Makio, N., Spectral properties of the linearized semigroup of the compressible Navier-Stokes equation on a periodic layer, Publ. Res. Inst. Math. Sci., **51**, (2015), pp. 337–372.
- [16] Kato, T., *Perturbation theory for linear operators*, Springer-Verlag, Berlin, Heidelberg, New York, (1980).
- [17] Kawashima, S., Systems of a hyperbolic-parabolic composite type, with applications to the equations of magnethydrodynamics, Ph. D. Thesis, Kyoto University (1983).
- [18] Kobayashi, T., Some estimates of solutions for the equations of motion of compressible viscous fluid in an exterior domain in $\mathbf{R}^3$, J. Differential Equations **184** (2002), pp. 587-619.
- [19] Kobayashi, T. and Shibata, Y., Decay estimates of solutions for the equations of motion of compressible viscous and heat-conductive gases in an exterior domain in $\mathbf{R}^3$, Comm. Math. Phys. **200** (1999), pp. 621-659.
- [20] Matsumura, A., An energy method for the equations of motion of compressible viscous and heat-conductive fluids, University of Wisconsin-Madison, MRC Technical Summary Report # 2194 (1981), pp. 1-16.
- [21] Matsumura, A. and Nishida, T., The initial value problem for the equations of motion of compressible viscous and heat-conductive fluids, Proc. Japan Acad. Ser. A **55** (1979), pp. 337-342.
- [22] Matsumura, A. and Nishida T., Initial boundary value problems for the equations of motion of compressible viscous and heat-conductive fluids, Commun. Math. Phys. **89** (1983), pp. 445-464.
- [23] Reed, M., Simon, B., *Methods of modern mathematical physics IV*. Academic Press, (1979).
- [24] Shibata, Y. and Tanaka, K., Rate of convergence of non-stationary flow to the steady flow of compressible viscous fluid, Comput. Math. Appl. **53** (2007), pp. 605-623.
- [25] Valli, A., On the existence of stationary solutions to compressible Navier-Stokes equations, Ann. Inst. H Poincaré, **4** (1987), pp. 99-113.

List of MI Preprint Series, Kyushu University

The Global COE Program

Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p-adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials
- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials

- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings
- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric τ -functions of the q-Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications

- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain
- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection
- MI2009-26 Manabu YOSHIDA
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance

- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA
Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI
Finite element computation for scattering problems of micro-hologram using DtN map
- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO
Abstract collision systems on groups

- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA
An algebraic approach to underdetermined experiments
- MI2010-10 Kei HIROSE & Sadanori KONISHI
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE
Lagrangian approach to weakly nonlinear stability of an elliptical flow
- MI2010-21 Hiroki MASUDA
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiro FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra

- MI2010-25 Toshimitsu TAKAESU
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI
Spin-spin correlation functions of the q -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and spike detection via Gaussian basis expansions
- MI2010-32 Nobutaka NAKAZONO
Hypergeometric τ functions of the q -Painlevé systems of type $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA
CAP representations of inner forms of $Sp(4)$ with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER
Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide FUKUMOTO & Alexander B. SAMOKHIN
Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI
Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI
Anderson-Darling test and the Malliavin calculus
- MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling via Compressed Sensing

- MI2011-5 Hiroshi INOUE
Implications in Compressed Sensing and the Restricted Isometry Property
- MI2011-6 Daeju KIM & Sadanori KONISHI
Predictive information criterion for nonlinear regression model based on basis expansion methods
- MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI
Group variable selection via relevance vector machine
- MI2011-8 Jan BREZINA & Yoshiyuki KAGEI
Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow
Group variable selection via relevance vector machine
- MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC
Recursive structures in the multispecies TASEP
- MI2011-10 Kazunori YASUTAKE
On projective space bundle with nef normalized tautological line bundle
- MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA
An explicit formula for the discrete power function associated with circle patterns of Schramm type
- MI2011-12 Yoshiyuki KAGEI
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow
- MI2011-13 Vladimír CHALUPECKÝ & Adrian MUNTEAN
Semi-discrete finite difference multiscale scheme for a concrete corrosion model: approximation estimates and convergence
- MI2011-14 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves
- MI2011-15 Hiroshi INOUE
A generalization of restricted isometry property and applications to compressed sensing
- MI2011-16 Yu KAWAKAMI
A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space
- MI2011-17 Naoyuki KAMIYAMA
Matroid intersection with priority constraints
- MI2012-1 Kazufumi KIMOTO & Masato WAKAYAMA
Spectrum of non-commutative harmonic oscillators and residual modular forms
- MI2012-2 Hiroki MASUDA
Mighty convergence of the Gaussian quasi-likelihood random fields for ergodic Levy driven SDE observed at high frequency

- MI2012-3 Hiroshi INOUE
A Weak RIP of theory of compressed sensing and LASSO
- MI2012-4 Yasuhide FUKUMOTO & Youich MIE
Hamiltonian bifurcation theory for a rotating flow subject to elliptic straining field
- MI2012-5 Yu KAWAKAMI
On the maximal number of exceptional values of Gauss maps for various classes of surfaces
- MI2012-6 Marcio GAMEIRO, Yasuaki HIRAOKA, Shunsuke IZUMI, Miroslav KRAMAR, Konstantin MISCHAIKOW & Vidit NANDA
Topological Measurement of Protein Compressibility via Persistence Diagrams
- MI2012-7 Nobutaka NAKAZONO & Seiji NISHIOKA
Solutions to a q -analog of Painlevé III equation of type $D_7^{(1)}$
- MI2012-8 Naoyuki KAMIYAMA
A new approach to the Pareto stable matching problem
- MI2012-9 Jan BREZINA & Yoshiyuki KAGEI
Spectral properties of the linearized compressible Navier-Stokes equation around time-periodic parallel flow
- MI2012-10 Jan BREZINA
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a time-periodic parallel flow
- MI2012-11 Daeju KIM, Shuichi KAWANO & Yoshiyuki NINOMIYA
Adaptive basis expansion via the extended fused lasso
- MI2012-12 Masato WAKAYAMA
On simplicity of the lowest eigenvalue of non-commutative harmonic oscillators
- MI2012-13 Masatoshi OKITA
On the convergence rates for the compressible Navier- Stokes equations with potential force
- MI2013-1 Abuduwaili PAERHATI & Yasuhide FUKUMOTO
A Counter-example to Thomson-Tait-Chetayev's Theorem
- MI2013-2 Yasuhide FUKUMOTO & Hirofumi SAKUMA
A unified view of topological invariants of barotropic and baroclinic fluids and their application to formal stability analysis of three-dimensional ideal gas flows
- MI2013-3 Hiroki MASUDA
Asymptotics for functionals of self-normalized residuals of discretely observed stochastic processes
- MI2013-4 Naoyuki KAMIYAMA
On Counting Output Patterns of Logic Circuits
- MI2013-5 Hiroshi INOUE
RIPless Theory for Compressed Sensing

- MI2013-6 Hiroshi INOUE
Improved bounds on Restricted isometry for compressed sensing
- MI2013-7 Hidetoshi MATSUI
Variable and boundary selection for functional data via multiclass logistic regression modeling
- MI2013-8 Hidetoshi MATSUI
Variable selection for varying coefficient models with the sparse regularization
- MI2013-9 Naoyuki KAMIYAMA
Packing Arborescences in Acyclic Temporal Networks
- MI2013-10 Masato WAKAYAMA
Equivalence between the eigenvalue problem of non-commutative harmonic oscillators and existence of holomorphic solutions of Heun's differential equations, eigenstates degeneration, and Rabi's model
- MI2013-11 Masatoshi OKITA
Optimal decay rate for strong solutions in critical spaces to the compressible Navier-Stokes equations
- MI2013-12 Shuichi KAWANO, Ibuki HOSHINA, Kazuki MATSUDA & Sadanori KONISHI
Predictive model selection criteria for Bayesian lasso
- MI2013-13 Hayato CHIBA
The First Painleve Equation on the Weighted Projective Space
- MI2013-14 Hidetoshi MATSUI
Variable selection for functional linear models with functional predictors and a functional response
- MI2013-15 Naoyuki KAMIYAMA
The Fault-Tolerant Facility Location Problem with Submodular Penalties
- MI2013-16 Hidetoshi MATSUI
Selection of classification boundaries using the logistic regression
- MI2014-1 Naoyuki KAMIYAMA
Popular Matchings under Matroid Constraints
- MI2014-2 Yasuhide FUKUMOTO & Youichi MIE
Lagrangian approach to weakly nonlinear interaction of Kelvin waves and a symmetry-breaking bifurcation of a rotating flow
- MI2014-3 Reika AOYAMA
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Parallel flow in a cylindrical domain
- MI2014-4 Naoyuki KAMIYAMA
The Popular Condensation Problem under Matroid Constraints

- MI2014-5 Yoshiyuki KAGEI & Kazuyuki TSUDA
Existence and stability of time periodic solution to the compressible Navier-Stokes equation for time periodic external force with symmetry
- MI2014-6 This paper was withdrawn by the authors.
- MI2014-7 Masatoshi OKITA
On decay estimate of strong solutions in critical spaces for the compressible Navier-Stokes equations
- MI2014-8 Rong ZOU & Yasuhide FUKUMOTO
Local stability analysis of azimuthal magnetorotational instability of ideal MHD flows
- MI2014-9 Yoshiyuki KAGEI & Naoki MAKIO
Spectral properties of the linearized semigroup of the compressible Navier-Stokes equation on a periodic layer
- MI2014-10 Kazuyuki TSUDA
On the existence and stability of time periodic solution to the compressible Navier-Stokes equation on the whole space
- MI2014-11 Yoshiyuki KAGEI & Takaaki NISHIDA
Instability of plane Poiseuille flow in viscous compressible gas
- MI2014-12 Chien-Chung HUANG, Naonori KAKIMURA & Naoyuki KAMIYAMA
Exact and approximation algorithms for weighted matroid intersection
- MI2014-13 Yusuke SHIMIZU
Moment convergence of regularized least-squares estimator for linear regression model
- MI2015-1 Hidetoshi MATSUI & Yuta UMEZU
Sparse regularization for multivariate linear models for functional data
- MI2015-2 Reika AOYAMA & Yoshiyuki KAGEI
Spectral properties of the semigroup for the linearized compressible Navier-Stokes equation around a parallel flow in a cylindrical domain
- MI2015-3 Naoyuki KAMIYAMA
Stable Matchings with Ties, Master Preference Lists, and Matroid Constraints
- MI2015-4 Reika AOYAMA & Yoshiyuki KAGEI
Large time behavior of solutions to the compressible Navier-Stokes equations around a parallel flow in a cylindrical domain
- MI2015-5 Kazuyuki TSUDA
Existence and stability of time periodic solution to the compressible Navier-Stokes-Korteweg system on R^3
- MI2015-6 Naoyuki KAMIYAMA
Popular Matchings with Ties and Matroid Constraints

- MI2015-7 Shoichi EGUCHI & Hiroki MASUDA
Quasi-Bayesian model comparison for LAQ models
- MI2015-8 Yoshiyuki KAGEI & Ryouta OOMACHI
Stability of time periodic solution of the Navier-Stokes equation on the half-space under oscillatory moving boundary condition
- MI2016-1 Momonari KUDO
Analysis of an algorithm to compute the cohomology groups of coherent sheaves and its applications
- MI2016-2 Yoshiyuki KAGEI & Masatoshi OKITA
Asymptotic profiles for the compressible Navier-Stokes equations on the whole space
- MI2016-3 Shota ENOMOTO & Yoshiyuki KAGEI
Asymptotic behavior of the linearized semigroup at space-periodic stationary solution of the compressible Navier-Stokes equation