On Chorin's method for stationary solutions of the Oberbeck-Boussinesq equation

Kagei, Yoshiyuki
Faculty of Mathematics, Kyushu University

Nishida, Takaaki
Department of Applied Complex System, Kyoto University

<http://hdl.handle.net/2324/1655026>

出版情報 : MI Preprint Series. 2016-5, 2016-04-22. 九州大学大学院数理学研究院
バージョン :
権利関係 :

MI Preprint Series

**Mathematics for Industry
Kyushu University**

On Chorin's method for stationary solutions of the Oberbeck-Boussinesq equation

**Yoshiyuki Kagei
& Takaaki Nishida**

MI 2016-5

(Received April 22, 2016)

Institute of Mathematics for Industry
Graduate School of Mathematics
Kyushu University
Fukuoka, JAPAN

On Chorin's method for stationary solutions of the Oberbeck-Boussinesq equation

Yoshiyuki Kagei¹ and Takaaki Nishida²

¹ Faculty of Mathematics,
Kyushu University,
Nishi-ku, Motooka 744,
Fukuoka 819-0395, Japan

² Department of Applied Complex System,
Kyoto University,
Yoshida Honmachi, Sakyo-ku,
Kyoto, 606-8317, Japan

Abstract

Stability of stationary solutions of the Oberbeck-Boussinesq system (OB) and the corresponding artificial compressible system is considered. The latter system is obtained by adding the time derivative of the pressure with small parameter $\varepsilon > 0$ to the continuity equation of (OB), which was proposed by A. Chorin to find stationary solutions of (OB) numerically. Both systems have the same sets of stationary solutions and the system (OB) is obtained from the artificial compressible one as the limit $\varepsilon \rightarrow 0$ which is a singular limit. It is proved that if a stationary solution of the artificial compressible system is stable for sufficiently small $\varepsilon > 0$, then it is also stable as a solution of (OB). The converse is proved provided that the velocity field of the stationary solution satisfies some smallness condition.

Mathematics Subject Classification (2000). 35Q30, 76N15.

Keywords. Oberbeck-Boussinesq equation, artificial compressible system, Chorin's method, singular perturbation, stability, cellular convection patterns.

1 Introduction

This paper is concerned with the Oberbeck-Boussinesq equation

$$\operatorname{div} \mathbf{v} = 0, \quad (1.1)$$

$$\operatorname{Pr}^{-1} (\partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v}) - \Delta \mathbf{v} + \nabla p - \operatorname{Ra} \theta \mathbf{e}_3 = \mathbf{0}, \quad (1.2)$$

$$\partial_t \theta + \mathbf{v} \cdot \nabla \theta - \Delta \theta - \operatorname{Ra} \mathbf{v} \cdot \mathbf{e}_3 = 0, \quad (1.3)$$

and the artificial compressible system for (1.1)–(1.3):

$$\varepsilon^2 \partial_t p + \operatorname{div} \mathbf{v} = 0, \quad (1.4)$$

$$\operatorname{Pr}^{-1} (\partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v}) - \Delta \mathbf{v} + \nabla p - \operatorname{Ra} \theta \mathbf{e}_3 = \mathbf{0}, \quad (1.5)$$

$$\partial_t \theta + \mathbf{v} \cdot \nabla \theta - \Delta \theta - \operatorname{Ra} \mathbf{v} \cdot \mathbf{e}_3 = 0. \quad (1.6)$$

Here $\mathbf{v} = {}^\top(v^1(x, t), v^2(x, t), v^3(x, t))$, $p = p(x, t)$ and $\theta = \theta(x, t)$ denote the unknown velocity field, pressure and temperature deviation from the heat conductive state, respectively, at time $t > 0$ and position $x \in \mathbb{R}^3$; $\mathbf{e}_3 = {}^\top(0, 0, 1) \in \mathbb{R}^3$; $\operatorname{Pr} > 0$ and $\operatorname{Ra} > 0$ are non-dimensional parameters, called Prandtl and Rayleigh numbers, respectively; and $\varepsilon > 0$ is a small parameter, called artificial Mach number. Here and in what follows, the superscript ${}^\top \cdot$ stands for the transposition. The systems (1.1)–(1.3) and (1.4)–(1.6) are considered in the infinite layer Ω :

$$\Omega = \{x = (x', x_3); x' = (x_1, x_2) \in \mathbb{R}^2, 0 < x_3 < 1\}.$$

The Oberbeck-Boussinesq equation (1.1)–(1.3) is a system of equations which describes convection phenomena of viscous fluid occupying Ω heated from below (heated at $x_3 = 0$) under the gravitational force. It is well known ([1, 6, 7, 10]) that under the boundary condition

$$\mathbf{v}|_{x_3=0,1} = \mathbf{0}, \quad \theta|_{x_3=0,1} = 0, \quad (1.7)$$

there exists a critical number $\operatorname{Ra}_c > 0$ such that when $\operatorname{Ra} < \operatorname{Ra}_c$, the heat conductive state $\mathbf{v} = \mathbf{0}$, $\theta = 0$ is stable, while, when $\operatorname{Ra} > \operatorname{Ra}_c$, the heat conductive state is unstable and convective cellular stationary solutions bifurcate from the heat conductive state.

A. Chorin ([2, 3, 4]) proposed the artificial compressible system such as (1.4)–(1.6) to find stationary solutions of equations for viscous incompressible fluid numerically. In the context of the Oberbeck-Boussinesq equation (1.1)–(1.3), the idea is stated as follows. Obviously, the sets of stationary solutions of (1.1)–(1.3) and (1.4)–(1.6) are the same ones. If solutions of the artificial compressible system (1.4)–(1.6) converge to a function $u_s = {}^\top(p_s, \mathbf{v}_s, \theta_s)$ as $t \rightarrow \infty$, then the limit u_s is a stationary solution of (1.4)–(1.6) which is thus a stationary solution of (1.1)–(1.3). By using this method, Chorin numerically obtained stationary cellular convection solutions of (1.1)–(1.3).

Since the limit u_s in Chorin's method described above is a large time limit of solutions of (1.4)–(1.6), u_s is stable as a solution of (1.4)–(1.6). It is of interest to consider whether u_s is stable as a solution of (1.1)–(1.3), in other words, whether u_s represents an observable stationary flow in the real world, and, conversely, what kind of stationary flows can be computed by Chorin's method. These questions are to be formulated as stability problem for stationary solutions of the systems (1.1)–(1.3) and (1.4)–(1.6). Since the system (1.1)–(1.3) is obtained from (1.4)–(1.6) as the limit $\varepsilon \rightarrow 0$, one could expect that solutions of (1.1)–(1.3) would be approximated by solutions of (1.4)–(1.6) when $\varepsilon \ll 1$. However, this limiting process is a singular limit, and hence, it is not straightforward to conclude that stability properties of u_s

as a solution of (1.1)–(1.3) are the same as those as a solution of (1.4)–(1.6) even when $0 < \varepsilon \ll 1$.

The purpose of this paper is to investigate stability relations of stationary solutions between the systems (1.1)–(1.3) and (1.4)–(1.6) when ε is sufficiently small. We investigate the spectra of the linearized operators around a stationary solution of (1.1)–(1.3) and (1.4)–(1.6) for $0 < \varepsilon \ll 1$.

We first show that if a stationary solution $u_s = {}^\top(p_s, \mathbf{v}_s, \theta_s)$ of (1.4)–(1.6) is asymptotically stable for sufficiently small ε , then so is u_s as a stationary solution of (1.1)–(1.3). More precisely, we consider (1.1)–(1.3) and (1.4)–(1.6) under the boundary condition (1.7) and the periodicity condition

$$p, \mathbf{v} \text{ and } \theta \text{ are } \mathcal{Q}\text{-periodic in } (x_1, x_2), \quad (1.8)$$

where $\mathcal{Q} = [-\pi/\alpha_1, \pi/\alpha_1] \times [-\pi/\alpha_2, \pi/\alpha_2]$. Here $\alpha_j, j = 1, 2$, are positive constants. We denote the basic period domain by $\Omega_{per} = \mathcal{Q} \times (0, 1)$. We introduce the linearized operators around u_s associated with (1.1)–(1.3) and (1.4)–(1.6). Let $L : L_{per,\sigma}^2 \times L_{per}^2 \rightarrow L_{per,\sigma}^2 \times L_{per}^2$ be the operator acting on $\mathbf{U} = {}^\top(\mathbf{w}, \theta) \in D(L)$ defined by

$$L = \begin{pmatrix} -\text{Pr}\mathbb{P}\Delta + \mathbb{P}(\mathbf{v}_s \cdot \nabla + {}^\top(\nabla \mathbf{v}_s)) & -\text{PrRa}\mathbb{P}\mathbf{e}_3 \\ {}^\top(\nabla \theta_s) - \text{Ra}^\top \mathbf{e}_3 & -\Delta + \mathbf{v}_s \cdot \nabla \end{pmatrix}$$

with domain $D(L) = [(H_{per}^2 \cap H_{0,per}^1)^3 \cap L_{per,\sigma}^2] \times [H_{per}^2 \cap H_{0,per}^1]$. Here $L_{per}^2, H_{per}^k, \dots$, denote $L^2, H^k, \dots$ spaces over Ω_{per} with periodicity condition in x' , $\mathbb{P}$ denotes the Helmholtz projection from $(L_{per}^2)^3$ to $L_{per,\sigma}^2$, where $L_{per,\sigma}^2$ denotes the set of all solenoidal vector fields $\mathbf{w} = {}^\top(w^1, w^2, w^3)$ in $(L_{per}^2)^3$ with $w^3|_{x_3=0,1} = 0$. We define the operator $L_\varepsilon : H_{per,*}^1 \times (L_{per}^2)^3 \times L_{per}^2 \rightarrow H_{per,*}^1 \times (L_{per}^2)^3 \times L_{per}^2$, acting on $u = {}^\top(p, \mathbf{w}, \theta) \in D(L_\varepsilon)$, by

$$L_\varepsilon = \begin{pmatrix} 0 & \frac{1}{\varepsilon^2} \text{div} & 0 \\ \text{Pr}\nabla & -\text{Pr}\Delta + \mathbf{v}_s \cdot \nabla + {}^\top(\nabla \mathbf{v}_s) & -\text{PrRa}\mathbf{e}_3 \\ 0 & {}^\top(\nabla \theta_s) - \text{Ra}^\top \mathbf{e}_3 & -\Delta + \mathbf{v}_s \cdot \nabla \end{pmatrix}$$

with domain $D(L_\varepsilon) = H_{per,*}^1 \times [H_{per}^2 \cap H_{0,per}^1]^3 \times [H_{per}^2 \cap H_{0,per}^1]$. Here $H_{per,*}^1 = \{p \in H_{per}^1; \int_{\Omega_{per}} p(x) dx = 0\}$.

We prove that if there exists a positive number b_0 such that $\rho(-L_{\varepsilon_n}) \supset \{\lambda \in \mathbb{C}; \text{Re } \lambda \geq -b_0\}$ for some sequence $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$, then there exists a constant $b_1 > 0$ such that $\rho(-L) \supset \{\lambda \in \mathbb{C}; \text{Re } \lambda \geq -b_1\}$. Therefore, a stationary solution obtained by Chorin's method with $0 < \varepsilon \ll 1$ is stable as a solution of the Oberbeck-Boussinesq system (1.1)–(1.3). Furthermore, we prove an instability result: if $\sigma(-L) \cap \{\lambda \in \mathbb{C}; \text{Re } \lambda > 0\} \neq \emptyset$, then $\sigma(-L_\varepsilon) \cap \{\lambda \in \mathbb{C}; \text{Re } \lambda > 0\} \neq \emptyset$ for $0 < \varepsilon \ll 1$. This shows that unstable stationary solutions of (1.1)–(1.3) cannot be obtained by Chorin's method with $0 < \varepsilon \ll 1$.

As a converse of the above result, we prove that if

$$\rho(-L) \supset \{\lambda \in \mathbb{C}; \text{Re } \lambda \geq -b_0\} \quad (1.9)$$

for some constant $b_0 > 0$, then there exist constants $\delta_0 > 0$ and $b_1 > 0$ such that

$$\rho(-L_\varepsilon) \supset \{\lambda \in \mathbb{C}; \text{Re } \lambda \geq -b_1\} \quad (1.10)$$

for $0 < \varepsilon \ll 1$, provided that

$$\inf_{\mathbf{w} \in H_{0,per}^1, \mathbf{w} \neq \mathbf{0}} \frac{\operatorname{Re}(\mathbf{w} \cdot \nabla \mathbf{v}_s, \mathbf{w})_{L^2}}{\|\nabla \mathbf{w}\|_{L^2}^2} \geq -\delta_0. \quad (1.11)$$

This gives a sufficient condition for u_s to be computed by Chorin's method with $0 < \varepsilon \ll 1$. We note that no smallness condition for the temperature θ_s of u_s is required.

This result is applicable to stable bifurcating cellular convective patterns such as roll pattern, hexagonal pattern and etc., when $\operatorname{Ra} \sim \operatorname{Ra}_c$. In fact, the velocity fields of bifurcating convective patterns are small when $\operatorname{Ra} \sim \operatorname{Ra}_c$ since they bifurcate from $\mathbf{v} = \mathbf{0}$, $\theta = 0$ when Ra crosses Ra_c , and hence, condition (1.11) is satisfied.

To prove the first result, we show that an eigenvalue λ of $-L_\varepsilon$ with $\operatorname{Im} \lambda = O(1)$ can be obtained as a perturbation of an eigenvalue of $-L$ as $\varepsilon \rightarrow 0$. In fact, it is proved that if λ_0 is an eigenvalue of $-L$, then, for any neighborhood of λ_0 , there exists an eigenvalue of $-L_\varepsilon$ for $0 < \varepsilon \ll 1$ and the total projection P_ε on the neighborhood satisfies $P_\varepsilon = P_0 + O(\varepsilon)$, where P_0 is a projection whose velocity-temperature part is the eigenprojection for the eigenvalue λ_0 of $-L$ (Theorem 5.2 (ii)). Based on this observation we show the first result by a contradiction argument. In particular, we find that if $-L$ has an eigenvalue with positive real part, then so does $-L_\varepsilon$ with $0 < \varepsilon \ll 1$, which gives the instability result.

To prove the converse, we investigate the resolvent $(\lambda + L_\varepsilon)^{-1}$ according to the cases for λ near the imaginary axis with $|\operatorname{Im} \lambda| \geq O(\varepsilon^{-1})$, $|\operatorname{Im} \lambda| \leq O(\varepsilon^{-1})$ and $|\operatorname{Im} \lambda| = O(\varepsilon^{-1})$ under the condition (1.9). By a standard energy method one can show that $-L_\varepsilon$ is a sectorial operator and $\{\lambda; \operatorname{Re} \lambda \geq -b, |\operatorname{Im} \lambda| \geq O(\varepsilon^{-1})\} \subset \rho(-L_\varepsilon)$ with some $b > 0$ for $0 < \varepsilon \ll 1$. We show that the spectrum of $-L_\varepsilon$ near the imaginary axis with $|\operatorname{Im} \lambda| \leq O(\varepsilon^{-1})$ can be treated as a kind of regular perturbation of that of $-L$, which concludes that $\{\lambda; \operatorname{Re} \lambda \geq -b, |\operatorname{Im} \lambda| \leq O(\varepsilon^{-1})\} \subset \rho(-L_\varepsilon)$. The spectrum of $-L_\varepsilon$ in the region $|\operatorname{Im} \lambda| = O(\varepsilon^{-1})$ stems from the ‘‘compressible’’ aspect of $-L_\varepsilon$ and does not appear in the spectrum of $-L$. We prove $\{\lambda; \operatorname{Re} \lambda \geq -b, |\operatorname{Im} \lambda| = O(\varepsilon^{-1})\} \subset \rho(-L_\varepsilon)$ by an energy method provided that the condition (1.11) holds.

One more remark is added in order. Due to the translation invariance in x_1 and x_2 variables, 0 is an eigenvalue of $-L_\varepsilon$ whenever $\partial_{x_1} u_s \neq 0$ or $\partial_{x_2} u_s \neq 0$. In this case nonzero $\partial_{x_j} u_s$ are eigenfunctions for the eigenvalue 0. Correspondingly, 0 is an eigenvalue of $-L$ whenever $\partial_{x_1} \mathbf{U}_s \neq 0$ or $\partial_{x_2} \mathbf{U}_s \neq 0$. We formulate the main results of this paper by taking this translation invariance into account. We note that under the translation invariance one can show the orbital stability of $\mathbf{U}_s$ and u_s for the corresponding nonlinear problems.

This paper is organized as follows. In section 2 we introduce notations used in this paper. In section 3 we state the main theorems of this paper. Section 4 is devoted to the study of the null spaces of L and L_ε for $0 < \varepsilon \ll 1$. We then prove the first results in section 5. The converse is proved in section 6; and we give the estimates for the derivatives of the pressure component of $(\lambda + L_\varepsilon)^{-1}$ in section 7.

2 Preliminaries

In this section we introduce notation used in this paper.

For given $\alpha_1, \alpha_2 > 0$, we denote the basic period cell by

$$\mathcal{Q} = \left[-\frac{\pi}{\alpha_1}, \frac{\pi}{\alpha_1}\right) \times \left[-\frac{\pi}{\alpha_2}, \frac{\pi}{\alpha_2}\right).$$

We denote the basic period domain by Ω_{per} :

$$\Omega_{per} = \mathcal{Q} \times (0, 1).$$

We denote by C_{per}^∞ the space of restrictions of functions in $C^\infty(\overline{\Omega})$ which are $\mathcal{Q}$ -periodic in $x' = (x_1, x_2)$. We also denote by $C_{0,per}^\infty$ the space of restrictions of functions in C^∞ which are $\mathcal{Q}$ -periodic in $x' = (x_1, x_2)$ and vanish near $x_3 = 0, 1$. For $1 \leq r \leq \infty$ we denote by $L^r(\Omega_{per})$ the usual Lebesgue space over Ω_{per} , and its norm is denoted by $\|\cdot\|_r$. The k th order L^2 Sobolev space over Ω_{per} is denoted by $H^k(\Omega_{per})$, and its norm is denoted by $\|\cdot\|_{H^k}$.

We set

$$\begin{aligned} L_{per}^2 &= \text{the } L^2(\Omega_{per})\text{-closure of } C_{0,per}^\infty, \\ H_{per}^k &= \text{the } H^k(\Omega_{per})\text{-closure of } C_{per}^\infty, \\ H_{0,per}^1 &= \text{the } H^1(\Omega_{per})\text{-closure of } C_{0,per}^\infty. \end{aligned}$$

We note that if $f \in H_{0,per}^1$, then $f|_{x_j=-\pi/\alpha_j} = f|_{x_j=\pi/\alpha_j}$ and $f|_{x_3=0,1} = 0$. The inner product of $f_j \in L_{per}^2$ ($j = 1, 2$) is denoted by

$$(f_1, f_2) = \int_{\Omega_{per}} f_1(x) \overline{f_2(x)} dx,$$

where $\bar{z}$ denotes the complex conjugate of z .

The mean value of a function $\phi(x)$ over Ω_{per} is denoted by $\langle \phi \rangle$:

$$\langle \phi \rangle = \frac{1}{|\Omega_{per}|} \int_{\Omega_{per}} \phi(x) dx.$$

The set of all $\phi \in L_{per}^2$ with $\langle \phi \rangle = 0$ is denoted by $L_{per,*}^2$, i.e.,

$$L_{per,*}^2 = \{\phi \in L_{per}^2 : \langle \phi \rangle = 0\}.$$

Furthermore, we set

$$H_{per,*}^k = H_{per}^k \cap L_{per,*}^2.$$

We denote by $C_{0,per,\sigma}^\infty$ the set of all vector fields $\mathbf{v}$ in $(C_{0,per}^\infty)^3$ with $\text{div } \mathbf{v} = 0$. We set

$$L_{per,\sigma}^2 = \text{the } L^2(\Omega_{per})^3\text{-closure of } C_{0,per,\sigma}^\infty.$$

It is known that $(L_{per}^2)^3 = L_{per,\sigma}^2 \oplus G_{per}^2$, where $G_{per}^2 = \{\nabla p; p \in H_{per}^1\}$ is the orthogonal complement of $L_{per,\sigma}^2$. The orthogonal projection $\mathbb{P}$ on $L_{per,\sigma}^2$ is called

the Helmholtz projection. We define the projection $\mathbf{P}$ from $(L_{per}^2)^3 \times L_{per}^2$ onto $L_{per,\sigma}^2 \times L_{per}^2$ by

$$\mathbf{P} = \begin{pmatrix} \mathbb{P} & 0 \\ 0 & I \end{pmatrix}.$$

For simplicity the set of all vector fields in $(L_{per}^2)^3$ (resp. $(H_{0,per}^1)^3, (H_{per}^k)^3$) are frequently denoted by L_{per}^2 (resp. $H_{0,per}^1, H_{per}^k$) if no confusion will occur.

We also use notation L_{per}^2 for the set of all $u = {}^\top(p, \mathbf{w}, \theta)$ with $p \in L_{per}^2$, $\mathbf{w} = {}^\top(w^1, w^2, w^3) \in L_{per}^2$ and $\theta \in L_{per}^2$ if no confusion will occur.

Let ε be a positive number. We introduce an inner product $\langle\langle u_1, u_2 \rangle\rangle_\varepsilon$ for $u_j = {}^\top(p_j, \mathbf{w}_j, \theta_j)$ ($j = 1, 2$) defined by

$$\langle\langle u_1, u_2 \rangle\rangle_\varepsilon = \varepsilon^2(p_1, p_2) + \text{Pr}^{-1}(\mathbf{w}_1, \mathbf{w}_2) + (\theta_1, \theta_2).$$

We also define the inner product $\langle \mathbf{U}_1, \mathbf{U}_2 \rangle$ for $\mathbf{U}_j = {}^\top(\mathbf{w}_j, \theta_j)$ ($j = 1, 2$) by

$$\langle \mathbf{U}_1, \mathbf{U}_2 \rangle = \text{Pr}^{-1}(\mathbf{w}_1, \mathbf{w}_2) + (\theta_1, \theta_2).$$

We denote the resolvent set of a closed operator A by $\rho(A)$ and the spectrum of A by $\sigma(A)$. The kernel and the range of A are denoted by $\text{Ker}(A)$ and $R(A)$, respectively.

3 Main Results

In this section we state the main results of this paper. Let $u_s = {}^\top(p_s, \mathbf{v}_s, \theta_s)$ be a stationary solution of (1.1)–(1.3) under the boundary conditions (1.7) and (1.8) satisfying $\int_{\Omega_{per}} p_s(x) dx = 0$. We consider the linearized problem for the Oberbeck-Boussinesq system (1.1)–(1.3) around $u_s = {}^\top(p_s, \mathbf{v}_s, \theta_s)$:

$$\text{div } \mathbf{w} = 0, \quad (3.1)$$

$$\text{Pr}^{-1} \partial_t \mathbf{w} - \Delta \mathbf{w} + \text{Pr}^{-1}(\mathbf{v}_s \cdot \nabla \mathbf{w} + \mathbf{w} \cdot \nabla \mathbf{v}_s) + \nabla p - \text{Ra} \theta \mathbf{e}_3 = \mathbf{0}, \quad (3.2)$$

$$\partial_t \theta - \Delta \theta + \mathbf{v}_s \cdot \nabla \theta + \mathbf{w} \cdot \nabla \theta_s - \text{Ra} \mathbf{w} \cdot \mathbf{e}_3 = 0 \quad (3.3)$$

under the boundary condition

$$\mathbf{w}|_{x_3=0,1} = \mathbf{0}, \quad \theta|_{x_3=0,1} = 0, \quad (3.4)$$

and the periodicity condition

$$p, \mathbf{w} \text{ and } \theta \text{ are } \mathcal{Q}\text{-periodic in } (x_1, x_2). \quad (3.5)$$

The linearized problem for the artificial compressible system is written as

$$\varepsilon^2 \partial_t p + \text{div } \mathbf{w} = 0, \quad (3.6)$$

$$\text{Pr}^{-1} \partial_t \mathbf{w} - \Delta \mathbf{w} + \text{Pr}^{-1}(\mathbf{v}_s \cdot \nabla \mathbf{w} + \mathbf{w} \cdot \nabla \mathbf{v}_s) + \nabla p - \text{Ra} \theta \mathbf{e}_3 = \mathbf{0}, \quad (3.7)$$

$$\partial_t \theta - \Delta \theta + \mathbf{v}_s \cdot \nabla \theta + \mathbf{w} \cdot \nabla \theta_s - \text{Ra} \mathbf{w} \cdot \mathbf{e}_3 = 0 \quad (3.8)$$

with the boundary conditions (3.4) and (3.5).

By applying the projection $\mathbf{P}$, problem (3.1)–(3.5) is written as

$$\Pr^{-1}\partial_t \mathbf{w} - \mathbb{P}\Delta \mathbf{w} + \Pr^{-1}\mathbb{P}(\mathbf{v}_s \cdot \nabla \mathbf{w} + \mathbf{w} \cdot \nabla \mathbf{v}_s) - \text{Ra}\mathbb{P}\theta \mathbf{e}_3 = \mathbf{0}, \quad (3.9)$$

$$\partial_t \theta - \Delta \theta + \mathbf{v}_s \cdot \nabla \theta + \mathbf{w} \cdot \nabla \theta_s - \text{Ra} \mathbf{w} \cdot \mathbf{e}_3 = 0 \quad (3.10)$$

with the boundary conditions (3.4) and (3.5). We introduce the linearized operator around $\mathbf{U}_s = {}^\top(\mathbf{v}_s, \theta_s)$ associated with problem (3.9)–(3.10) under (3.4) and (3.5). We define the operator $L : L_{per,\sigma}^2 \times L_{per}^2 \rightarrow L_{per,\sigma}^2 \times L_{per}^2$ by

$$L = \begin{pmatrix} -\Pr\mathbb{P}\Delta + \mathbb{P}(\mathbf{v}_s \cdot \nabla + {}^\top(\nabla \mathbf{v}_s)) & -\Pr\text{Ra}\mathbb{P}\mathbf{e}_3 \\ {}^\top(\nabla \theta_s) - \text{Ra}{}^\top \mathbf{e}_3 & -\Delta + \mathbf{v}_s \cdot \nabla \end{pmatrix}$$

with domain $D(L) = [(H_{per}^2 \cap H_{0,per}^1)^3 \cap L_{per,\sigma}^2] \times [H_{per}^2 \cap H_{0,per}^1]$.

We also introduce the linearized operator around $u_s = {}^\top(p_s, \mathbf{w}_s, \theta_s)$ associated with (3.6)–(3.8) under (3.4) and (3.5). We define the operator $L_\varepsilon : H_{per,*}^1 \times (L_{per}^2)^3 \times L_{per}^2 \rightarrow H_{per,*}^1 \times (L_{per}^2)^3 \times L_{per}^2$ by

$$L_\varepsilon = \begin{pmatrix} 0 & \frac{1}{\varepsilon^2} \text{div} & 0 \\ \Pr \nabla & -\Pr \Delta + \mathbf{v}_s \cdot \nabla + {}^\top(\nabla \mathbf{v}_s) & -\Pr \text{Ra} \mathbf{e}_3 \\ 0 & {}^\top(\nabla \theta_s) - \text{Ra}{}^\top \mathbf{e}_3 & -\Delta + \mathbf{v}_s \cdot \nabla \end{pmatrix}$$

with domain $D(L_\varepsilon) = H_{per,*}^1 \times [H_{per}^2 \cap H_{0,per}^1]^3 \times [H_{per}^2 \cap H_{0,per}^1]$.

Before going further we make one observation. Due to the translation invariance in x_1 and x_2 variables, 0 is an eigenvalue of $-L_\varepsilon$ whenever $\partial_{x_1} u_s \neq 0$ or $\partial_{x_2} u_s \neq 0$. In this case nonzero $\partial_{x_j} u_s$ are eigenfunctions for the eigenvalue 0. Correspondingly, 0 is an eigenvalue of $-L$ whenever $\partial_{x_1} \mathbf{U}_s \neq 0$ or $\partial_{x_2} \mathbf{U}_s \neq 0$. For simplicity we consider the case $\partial_{x_j} u_s \neq 0$ (and hence $\partial_{x_j} \mathbf{U}_s \neq 0$) for $j = 1, 2$.

Theorem 3.1. *Let $\partial_{x_j} u_s \neq 0$ for $j = 1, 2$. If there exists a positive number b_0 such that $\rho(-L_{\varepsilon_n}) \supset \{\lambda \in \mathbb{C}; \text{Re } \lambda \geq -b_0\} \setminus \{0\}$ for some sequence $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$ and 0 is a semisimple eigenvalue of $-L_{\varepsilon_n}$ with $\text{Ker}(-L_{\varepsilon_n}) = \text{span}\{\partial_{x_1} u_s, \partial_{x_2} u_s\}$, then there exists a constant $b_1 > 0$ such that $\rho(-L) \supset \{\lambda \in \mathbb{C}; \text{Re } \lambda \geq -b_1\} \setminus \{0\}$ and 0 is a semisimple eigenvalue of $-L$ with $\text{Ker}(-L) = \text{span}\{\partial_{x_1} \mathbf{U}_s, \partial_{x_2} \mathbf{U}_s\}$.*

Theorem 3.1 shows that if u_s is obtained by Chorin's method with $0 < \varepsilon \ll 1$, then it is stable as a solution of the Oberbeck-Boussinesq system. In particular, we have the following instability result.

Theorem 3.2. *Let $\partial_{x_j} u_s \neq 0$ for $j = 1, 2$. If $\sigma(-L) \cap \{\lambda \in \mathbb{C}; \text{Re } \lambda > 0\} \neq \emptyset$, then $\sigma(-L_\varepsilon) \cap \{\lambda \in \mathbb{C}; \text{Re } \lambda > 0\} \neq \emptyset$ for sufficiently small ε .*

By Theorem 3.2, we see that unstable stationary solutions of the Oberbeck-Boussinesq system cannot be obtained by Chorin's method with $0 < \varepsilon \ll 1$.

We next give a sufficient condition for u_s to be computed by Chorin's method with $0 < \varepsilon \ll 1$.

Theorem 3.3. *Let $\partial_{x_j} \mathbf{U}_s \neq \mathbf{0}$ for $j = 1, 2$. Suppose that $\rho(-L) \supset \{\lambda \in \mathbb{C}; \operatorname{Re} \lambda \geq -b_0\} \setminus \{0\}$ for some constant $b_0 > 0$ and 0 is a semisimple eigenvalue of $-L$ with $\operatorname{Ker}(-L) = \operatorname{span}\{\partial_{x_1} \mathbf{U}_s, \partial_{x_2} \mathbf{U}_s\}$. Then there exist constants $\varepsilon_0 > 0$, $\delta_0 > 0$ and $b_1 > 0$ such that if*

$$\inf_{\mathbf{w} \in H_{per,0}^1, \mathbf{w} \neq \mathbf{0}} \frac{\operatorname{Re}(\mathbf{w} \cdot \nabla \mathbf{v}_s, \mathbf{w})}{\|\nabla \mathbf{w}\|_2^2} \geq -\delta_0, \quad (3.11)$$

then $\rho(-L_\varepsilon) \supset \{\lambda \in \mathbb{C}; \operatorname{Re} \lambda \geq -b_1\} \setminus \{0\}$ for all $0 < \varepsilon \leq \varepsilon_0$ and 0 is a semisimple eigenvalue of $-L_\varepsilon$ with $\operatorname{Ker}(-L_\varepsilon) = \operatorname{span}\{\partial_{x_1} u_s, \partial_{x_2} u_s\}$.

In Theorem 3.3 we require smallness condition for the velocity field $\mathbf{v}_s$ only but not for the temperature θ_s .

Remark 3.4. *It is known that $-L$ is sectorial. Furthermore, $-L_\varepsilon$ is also sectorial for each $\varepsilon > 0$. (See Proposition 6.1 below.) Consequently, under the assumptions of Theorems 3.1 and 3.3, $u_s = {}^\top(p_s, \mathbf{v}_s, \theta_s)$ is asymptotically (orbitally) stable as a solution of both problems. More precisely, a solution of the nonlinear problem near u_s converges to a stationary solution $u_s(\cdot + a', \cdot)$ for some $a' \in \mathbb{R}^2$ as $t \rightarrow \infty$.*

Remark 3.5. *Theorems 3.1 and 3.3 also hold with slight modifications when $\partial_{x_j} u_s = 0$ (and hence $\partial_{x_j} \mathbf{U}_s = 0$) for both $j = 1, 2$ or for one of $j = 1, 2$. For example, if $\partial_{x_1} \mathbf{U}_s \neq \mathbf{0}$ and $\partial_{x_2} \mathbf{U}_s = \mathbf{0}$, then Theorem 3.3 holds with $\operatorname{Ker}(-L) = \operatorname{span}\{\partial_{x_1} \mathbf{U}_s, \partial_{x_2} \mathbf{U}_s\}$ and $\operatorname{Ker}(-L_\varepsilon) = \operatorname{span}\{\partial_{x_1} u_s, \partial_{x_2} u_s\}$ replaced by $\operatorname{Ker}(-L) = \operatorname{span}\{\partial_{x_1} \mathbf{U}_s\}$ and $\operatorname{Ker}(-L_\varepsilon) = \operatorname{span}\{\partial_{x_1} u_s\}$, respectively. Likewise, the theorems hold for other cases with similar modifications.*

Since the velocity fields of cellular stationary convective patterns bifurcating from the heat conductive state are small when $\operatorname{Ra} \sim \operatorname{Ra}_c$, we apply Theorem 3.3 and Remark 3.5 to obtain the following corollary.

Corollary 3.6. *If u_s is a stable convective pattern of (1.1)–(1.3) bifurcating from the heat conductive state, then it is also stable as a solution of (1.4)–(1.6) for $0 < \varepsilon \ll 1$ when $\operatorname{Ra} \sim \operatorname{Ra}_c$.*

To prove Theorems 3.1–3.3, we will first investigate the null spaces of L and L_ε in section 4. Theorems 3.1 and 3.2 will be proved in section 5. The proof of Theorem 3.3 will be given in sections 6 and 7.

4 The null spaces of L and L_ε

In this section we investigate the relation of the null spaces of L and L_ε . We introduce the adjoint operator $L^* : L_{per,\sigma}^2 \times L_{per}^2 \rightarrow L_{per,\sigma}^2 \times L_{per}^2$ of L :

$$L^* = \begin{pmatrix} -\operatorname{Pr}\mathbb{P}\Delta + \mathbb{P}(-\mathbf{v}_s \cdot \nabla + (\nabla \mathbf{v}_s)) & \operatorname{Pr}\mathbb{P}((\nabla \theta_s) - \operatorname{Ra} \mathbf{e}_3) \\ -\operatorname{Ra} {}^\top \mathbf{e}_3 & -\Delta - \mathbf{v}_s \cdot \nabla \end{pmatrix}$$

with domain $D(L^*) = [(H_{per}^2 \cap H_{0,per}^1)^3 \cap L_{per,\sigma}^2] \times [H_{per}^2 \cap H_{0,per}^1]$.

In what follows we assume that 0 is a semisimple eigenvalue of $-L$ with $\text{Ker}(-L) = \text{span}\{\partial_{x_1}\mathbf{U}_s, \partial_{x_2}\mathbf{U}_s\}$.

Since $-L$ is a sectorial operator with compact resolvent and 0 is a semisimple eigenvalue of $-L$, we have the following resolvent estimate for $-L$.

Proposition 4.1. *Set $\mathbf{U}_j^{(0)} = \partial_{x_j}\mathbf{U}_s$ for $j = 1, 2$. Then the following assertions hold.*

(i) *There exist $\mathbf{U}_j^* = {}^\top(\mathbf{w}_j^*, \theta_j^*) \in D(L^*)$ such that $L^*\mathbf{U}_j^* = \mathbf{0}$ and $\langle \mathbf{U}_j^{(0)}, \mathbf{U}_j^* \rangle = \delta_{jk}$ for $j, k = 1, 2$. Furthermore,*

$$L_{per,\sigma}^2 \times L_{per}^2 = X_0 \oplus X_1,$$

where $X_0 = \text{Ker}(-L)$ and

$$X_1 = R(-L) = \{\mathbf{U} \in L_{per,\sigma}^2 \times L_{per}^2; \langle \mathbf{U}, \mathbf{U}_j^* \rangle = 0, j = 1, 2\}.$$

The eigenprojection $\mathbf{\Pi}_0$ for the eigenvalue 0 of $-L$ is given by

$$\mathbf{\Pi}_0\mathbf{U} = \langle \mathbf{U}, \mathbf{U}_1^* \rangle \mathbf{U}_1^{(0)} + \langle \mathbf{U}, \mathbf{U}_2^* \rangle \mathbf{U}_2^{(0)}.$$

(ii) Set $\mathbf{\Pi}_0^c = I - \mathbf{\Pi}_0$. There exist constants $a_0 > 0$ and $c_0 \in \mathbb{R}$ such that

$$\Sigma \setminus \{0\} \subset \rho(-L),$$

where

$$\Sigma := \{\lambda \in \mathbb{C}; \text{Re } \lambda \geq -a_0|\text{Im } \lambda|^2 + c_0\},$$

and the estimates

$$\|(\lambda + L)^{-1}\mathbf{F}\|_2 \leq C \left\{ \frac{1}{|\lambda|} \|\mathbf{\Pi}_0\mathbf{F}\|_2 + \frac{1}{|\lambda| + 1} \|\mathbf{\Pi}_0^c\mathbf{F}\|_2 \right\},$$

$$\|\partial_x^2(\lambda + L)^{-1}\mathbf{F}\|_2 \leq C \left\{ \frac{1}{|\lambda|} \|\mathbf{\Pi}_0\mathbf{F}\|_2 + \|\mathbf{\Pi}_0^c\mathbf{F}\|_2 \right\}$$

hold uniformly for $\lambda \in \Sigma \setminus \{0\}$. Furthermore, if $\mathbf{\Pi}_0\mathbf{F} = 0$, then $\mathbf{\Pi}_0(\lambda + L)^{-1}\mathbf{F} = 0$ and the above estimates hold for any $\lambda \in \Sigma$.

We next introduce an operator $\mathcal{L}_{\varepsilon,\lambda} : H_{per,*}^1 \times (L_{per}^2)^3 \times L_{per}^2 \rightarrow H_{per,*}^1 \times (L_{per}^2)^3 \times L_{per}^2$ defined by

$$D(\mathcal{L}_{\varepsilon,\lambda}) = H_{per,*}^1 \times [H_{per}^2 \cap H_{0,per}^1]^3 \times [H_{per}^2 \cap H_{0,per}^1],$$

$$\mathcal{L}_{\varepsilon,\lambda} = \begin{pmatrix} 0 & \frac{1}{\varepsilon^2} \text{div} & 0 \\ \text{Pr} \nabla & \lambda - \text{Pr} \Delta + \mathbf{v}_s \cdot \nabla + {}^\top(\nabla \mathbf{v}_s) & -\text{Pr} \text{Ra} \mathbf{e}_3 \\ 0 & {}^\top(\nabla \theta_s) - \text{Ra} {}^\top \mathbf{e}_3 & \lambda - \Delta + \mathbf{v}_s \cdot \nabla \end{pmatrix},$$

and its adjoint $\mathcal{L}_{\varepsilon,\lambda}^* : H_{per,*}^1 \times (L_{per}^2)^3 \times L_{per}^2 \rightarrow H_{per,*}^1 \times (L_{per}^2)^3 \times L_{per}^2$ by

$$D(\mathcal{L}_{\varepsilon,\lambda}^*) = H_{per,*}^1 \times [H_{per}^2 \cap H_{0,per}^1]^3 \times [H_{per}^2 \cap H_{0,per}^1],$$

$$\mathcal{L}_{\varepsilon,\lambda}^* = \begin{pmatrix} 0 & -\frac{1}{\varepsilon^2} \operatorname{div} & 0 \\ -\operatorname{Pr} \nabla & \bar{\lambda} - \operatorname{Pr} \Delta - \mathbf{v}_s \cdot \nabla + (\nabla \mathbf{v}_s)^\top & \operatorname{Pr}(\nabla \theta_s) - \operatorname{Pr} \operatorname{Ra} \mathbf{e}_3 \\ 0 & -\operatorname{Ra}^\top \mathbf{e}_3 & \bar{\lambda} - \Delta - \mathbf{v}_s \cdot \nabla \end{pmatrix}.$$

Note that

$$\mathcal{L}_{\varepsilon,0} = L_\varepsilon.$$

We also introduce the operators $\mathbf{A}_\lambda$ and $\mathcal{A}_\lambda$ by

$$\mathbf{A}_\lambda \mathbf{w} = \begin{pmatrix} \lambda - \operatorname{Pr} \Delta + \mathbf{v}_s \cdot \nabla + (\nabla \mathbf{v}_s)^\top \\ \operatorname{Pr}(\nabla \theta_s) - \operatorname{Ra}^\top \mathbf{e}_3 \end{pmatrix} \mathbf{w},$$

and

$$\mathcal{A}_\lambda \mathbf{w} = \begin{pmatrix} 0 \\ \lambda - \operatorname{Pr} \Delta + \mathbf{v}_s \cdot \nabla + (\nabla \mathbf{v}_s)^\top \\ \operatorname{Pr}(\nabla \theta_s) - \operatorname{Ra}^\top \mathbf{e}_3 \end{pmatrix} \mathbf{w}.$$

We first prepare the following lemma.

Lemma 4.2. *There exists a bounded linear operator $\mathbf{V} : H_{per,*}^1 \rightarrow [H_{per}^2 \cap H_{0,per}^1]^3$ such that*

$$\operatorname{div} \mathbf{V} f = f, \quad \|\mathbf{V} f\|_{H^2} \leq C \|f\|_{H^1}$$

for $f \in H_{per,*}^1$.

This lemma follows from the solvability for the nonhomogeneous Stokes problem $\operatorname{div} \mathbf{v} = f$, $-\Delta \mathbf{v} + \nabla p = 0$ under the boundary conditions $v|_{x_3=0,1} = \mathbf{0}$, and $p, \mathbf{v}$ being $\mathcal{Q}$ -periodic in x' . See, e.g., [5].

In what follows we set

$$Y := H_{per,*}^1 \times (L_{per}^2)^3 \times L_{per}^2.$$

Proposition 4.3. *Let $\varepsilon > 0$.*

(i) *Let $u_j^{(0)} = \partial_{x_j} u_s$ ($j = 1, 2$). Then*

$$\operatorname{Ker}(\mathcal{L}_{\varepsilon,0}) = \operatorname{Ker}(L_\varepsilon) = \operatorname{span} \{u_1^{(0)}, u_2^{(0)}\}.$$

(ii) *For each $j = 1, 2$, there exists a unique $p_j^* \in H_{per,*}^1$ such that $\mathcal{L}_{\varepsilon,0}^* u_j^* = 0$, $u_j^* = {}^\top(p_j^*, \mathbf{w}_j^*, \theta_j^*)$, where ${}^\top(\mathbf{w}_j^*, \theta_j^*)$ ($j = 1, 2$) are the functions given in Proposition 4.1 (i). Furthermore,*

$$R(\mathcal{L}_{\varepsilon,0}) = R(L_\varepsilon) = \{u \in Y; \langle u, u_j^* \rangle_\varepsilon = 0, j = 1, 2\}.$$

(iii) *There exists a positive number ε_1 such that if $|\varepsilon| \leq \varepsilon_1$, then*

$$Y = \operatorname{Ker}(L_\varepsilon) \oplus R(L_\varepsilon).$$

Therefore, 0 is a semisimple eigenvalue of L_ε . Furthermore, if $F = {}^\top(f, \mathbf{g}, h) \in R(L_\varepsilon)$, then there exists a unique solution $u = {}^\top(p, \mathbf{w}, \theta) \in D(L_\varepsilon) \cap R(L_\varepsilon)$ of $L_\varepsilon u = F$ and u satisfies

$$\|u\|_{H^1 \times H^2 \times H^2} \leq C \{\varepsilon^2 \|f\|_{H^1} + \|\mathbf{F}\|_2\},$$

where $\mathbf{F} = {}^\top(\mathbf{g}, h)$.

Proof. Let $u = {}^\top(p, \mathbf{w}, \theta) \in D(\mathcal{L}_{\varepsilon,0})$ satisfy $\mathcal{L}_{\varepsilon,0}u = 0$. Then $\mathbf{U} = {}^\top(\mathbf{w}, \theta)$ satisfies $\mathbf{U} \in D(L)$ and $L\mathbf{U} = \mathbf{0}$. By assumption, there exist $a_j \in \mathbb{C}$ such that $\mathbf{U} = a_1\partial_{x_1}\mathbf{U}_s + a_2\partial_{x_2}\mathbf{U}_s$. On the other hand, $\mathcal{L}_{\varepsilon,0}(a_1\partial_{x_1}u_s + a_2\partial_{x_2}u_s) = 0$. We thus obtain

$$\nabla p = \nabla(a_1\partial_{x_1}p_s + a_2\partial_{x_2}p_s).$$

Since p and $a_1\partial_{x_1}p_s + a_2\partial_{x_2}p_s$ are in $H_{per,*}^1$, we have $p = a_1\partial_{x_1}p_s + a_2\partial_{x_2}p_s$, and hence, $u = a_1\partial_{x_1}u_s + a_2\partial_{x_2}u_s$. This shows $\text{Ker}(\mathcal{L}_{\varepsilon,0}) = \text{span}\{u_1^{(0)}, u_2^{(0)}\}$ and (i) is proved.

As for (ii), since $L^*\mathbf{U}_j^* = \mathbf{0}$, there exists a unique $p_j^* \in H_{per,*}^1$ such that $u_j^* = {}^\top(p_j^*, \mathbf{U}_j^*)$ satisfies $u_j \in D(\mathcal{L}_{\varepsilon,0}^*)$ and $\mathcal{L}_{\varepsilon,0}^*u_j^* = 0$.

Let us prove $R(\mathcal{L}_{\varepsilon,0}) = \{u \in Y; \langle \langle u, u_j^* \rangle \rangle_\varepsilon = 0, j = 1, 2\}$. Set $\tilde{\mathbf{w}} = \mathbf{w} - \varepsilon^2\mathbf{V}f$. Then

$$\mathcal{L}_{\varepsilon,0}\tilde{u} = \tilde{F} - \varepsilon^2\mathcal{A}_0\mathbf{V}f, \quad (4.1)$$

where $\tilde{u} = {}^\top(p, \tilde{\mathbf{w}}, \theta)$ and $\tilde{F} = {}^\top(0, \mathbf{g}, h)$. Therefore, we have

$$L\tilde{\mathbf{U}} = \mathbf{P}[\mathbf{F} - \varepsilon^2\mathbf{A}_0\mathbf{V}f], \quad (4.2)$$

where $\tilde{\mathbf{U}} = {}^\top(\tilde{\mathbf{w}}, \theta)$ and $\mathbf{F} = {}^\top(\mathbf{g}, h)$. Since $\langle \langle F, u_j^* \rangle \rangle_\varepsilon = 0$ for $j = 1, 2$, we have

$$\begin{aligned} & \langle \mathbf{P}[\mathbf{F} - \varepsilon^2\mathbf{A}_0\mathbf{V}f], \mathbf{U}_j^* \rangle \\ &= \langle \mathbf{F}, \mathbf{U}_j^* \rangle - \varepsilon^2\text{Pr}^{-1}(-\text{Pr}\Delta\mathbf{V}f + \mathbf{v}_s \cdot \nabla\mathbf{V}f + \mathbf{V}f \cdot \nabla\mathbf{v}_s, \mathbf{w}_j^*) \\ & \quad - \varepsilon^2(\mathbf{V}f \cdot \nabla\theta_s - \text{Ra}\mathbf{V}f \cdot \mathbf{e}_3, \theta_j^*) \\ &= \langle \mathbf{F}, \mathbf{U}_j^* \rangle \\ & \quad - \varepsilon^2\text{Pr}^{-1}(\mathbf{V}f, -\text{Pr}\Delta\mathbf{w}_j^* - \mathbf{v}_s \cdot \nabla\mathbf{w}_j^* + (\nabla\mathbf{v}_s)\mathbf{w}_j^* + \text{Pr}(\nabla\theta_s)\theta_j^* - \text{PrRa}\theta_j^*\mathbf{e}_3) \\ &= \langle \mathbf{F}, \mathbf{U}_j^* \rangle - \varepsilon^2\text{Pr}^{-1}(\mathbf{V}f, \text{Pr}\nabla p_j^*) \\ &= \langle \mathbf{F}, \mathbf{U}_j^* \rangle + \varepsilon^2(\text{div}\mathbf{V}f, p_j^*) \\ &= \langle \langle F, u_j^* \rangle \rangle_\varepsilon = 0. \end{aligned}$$

Applying Proposition 4.1, we see that there exists a unique solution $\tilde{\mathbf{U}} = {}^\top(\tilde{\mathbf{w}}, \theta) \in D(L) \cap X_1$ of (4.2) with estimate

$$\|\tilde{\mathbf{U}}\|_{H^2} \leq C\{\varepsilon^2\|f\|_{H^1} + \|\mathbf{F}\|_2\}.$$

It then follows that there exists a unique $p \in H_{per,*}^1$ such that $\tilde{u} = {}^\top(p, \tilde{\mathbf{w}}, \theta) \in D(\mathcal{L}_{\varepsilon,0})$ is a solution of (4.1). Setting $u = {}^\top(p, \tilde{\mathbf{w}} + \varepsilon^2\mathbf{V}f, \theta)$, we see that $u \in D(\mathcal{L}_{\varepsilon,0})$ and $\mathcal{L}_{\varepsilon,0}u = F$ with estimate

$$\|u\|_{H^1 \times H^2 \times H^2} \leq C\{\varepsilon^2\|f\|_{H^1} + \|\mathbf{F}\|_2\}. \quad (4.3)$$

We thus conclude that $R(\mathcal{L}_{\varepsilon,0}) = \{u \in Y; \langle \langle u, u_j^* \rangle \rangle_\varepsilon = 0, j = 1, 2\}$. Note that $R(L_\varepsilon)$ is closed.

As for (iii), we first show that there exists $u_{j,\varepsilon}^{(0)} \in D(L_\varepsilon)$ ($j = 1, 2$) such that

$$\text{Ker}(L_\varepsilon) = \text{span}\{u_{1,\varepsilon}^{(0)}, u_{2,\varepsilon}^{(0)}\}, \quad \langle\langle u_{j,\varepsilon}^{(0)}, u_k^* \rangle\rangle_\varepsilon = \delta_{jk} \quad (j, k = 1, 2).$$

We look for $u_{j,\varepsilon}^{(0)}$ in the form $u_{j,\varepsilon}^{(0)} = a_{1j}u_1^{(0)} + a_{2j}u_2^{(0)}$ ($j = 1, 2$). By the conditions $\langle\langle u_{j,\varepsilon}^{(0)}, u_k^* \rangle\rangle = \delta_{jk}$, we have

$$B_\varepsilon \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (4.4)$$

where

$$B_\varepsilon = \begin{pmatrix} \langle\langle u_1^{(0)}, u_1^* \rangle\rangle_\varepsilon & \langle\langle u_2^{(0)}, u_1^* \rangle\rangle_\varepsilon \\ \langle\langle u_1^{(0)}, u_2^* \rangle\rangle_\varepsilon & \langle\langle u_2^{(0)}, u_2^* \rangle\rangle_\varepsilon \end{pmatrix}.$$

Since $\langle\mathbf{U}_j^{(0)}, \mathbf{U}_k^*\rangle = \delta_{jk}$, we see that

$$B_\varepsilon = \begin{pmatrix} 1 + \varepsilon^2(p_1^{(0)}, p_1^*) & \varepsilon^2(p_2^{(0)}, p_1^*) \\ \varepsilon^2(p_1^{(0)}, p_2^*) & 1 + \varepsilon^2(p_2^{(0)}, p_2^*) \end{pmatrix}.$$

Therefore,

$$B_\varepsilon \rightarrow \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

as $\varepsilon \rightarrow 0$, and hence, B_ε has its inverse B_ε^{-1} for $0 < \varepsilon \ll 1$. We thus obtain the desired $u_{j,\varepsilon}^{(0)}$ ($j = 1, 2$).

Set

$$P_{0,\varepsilon}u = \sum_{j=1}^2 \langle\langle u, u_j^* \rangle\rangle_\varepsilon u_{j,\varepsilon}^{(0)}.$$

Then $P_{0,\varepsilon}$ is the projection on $\text{Ker}(L_\varepsilon)$, i.e., the eigenprojection for the eigenvalue 0 of $-L_\varepsilon$. It is easy to see that $R(I - P_{0,\varepsilon}) = \{u \in Y; \langle\langle u, u_j^* \rangle\rangle_\varepsilon = 0, j = 1, 2\}$. This, together with (ii), implies that $R(L_\varepsilon) = R(I - P_{0,\varepsilon})$, and hence, $Y = \text{Ker}(L_\varepsilon) \oplus R(L_\varepsilon)$. This decomposition and (ii), together with (4.3), yield the unique existence of solution $u \in D(L_\varepsilon) \cap R(L_\varepsilon)$ of $L_\varepsilon u = F$ for $F \in R(L_\varepsilon)$ with the desired estimate. This completes the proof $\square$

5 Proof of Theorems 3.1 and 3.2

In this section we prove Theorems 3.1 and 3.2. We introduce some operators. We define $\mathbb{L}_\lambda : (H_{per}^2)^3 \times L_{per}^2 \rightarrow (L_{per}^2)^3$ by

$$\mathbb{L}_\lambda \mathbf{U} = \lambda \mathbf{w} - \text{Pr} \Delta \mathbf{w} + \mathbf{v}_s \cdot \nabla \mathbf{w} + \mathbf{w} \cdot \nabla \mathbf{v}_s - \text{Pr} \text{Ra} \theta \mathbf{e}_3$$

for $\mathbf{U} = {}^\top(\mathbf{w}, \theta) \in (H_{per}^2)^3 \times L_{per}^2$.

Let $\lambda \in \rho(-L)$. For any $\mathbf{F} = {}^\top(\mathbf{g}, h) \in L_{per,\sigma}^2 \times L_{per}^2$, there exists a unique $p \in H_{per,*}^1$ satisfying

$$\nabla p = \mathbb{I} \mathbf{F} - \mathbb{L}_\lambda (\lambda + L)^{-1} \mathbf{F}, \quad (5.1)$$

and

$$\|p\|_{H^1} \leq C\{|\lambda|C_{0,\lambda} + C_{2,\lambda} + 1\}\|\mathbf{F}\|_2,$$

where $\mathbb{I}$ is the $n \times (n+1)$ matrix given by $\mathbb{I} = (I, \mathbf{0})$ with I being the $n \times n$ identity matrix, i.e., $\mathbb{I}\mathbf{F} = \mathbf{g}$, and

$$C_{k,\lambda} = \sup \left\{ \frac{\|(\lambda + L)^{-1}\mathbf{F}\|_{H^k \times H^k}}{\|\mathbf{F}\|_2}; \mathbf{F} \in L_{per,\sigma}^2 \times L_{per}^2, \mathbf{F} \neq \mathbf{0} \right\} \quad (k = 0, 2).$$

We denote this p by $p_\lambda[\mathbf{F}]$. The correspondence $\mathbf{F} \in L_{per,\sigma}^2 \times L_{per}^2 \mapsto p_\lambda[\mathbf{F}] \in H_{per,*}^1$ is linear, and hence, defines a bounded linear operator.

We first give an expression of $\mathcal{L}_{\varepsilon,\lambda}^{-1}$ in terms of $(\lambda + L)^{-1}$.

Proposition 5.1. *Let $\lambda \in \rho(-L)$. Then*

$$\mathcal{L}_{\varepsilon,\lambda}^{-1}F = \begin{pmatrix} p_\lambda[\mathbf{P}(\mathbf{F} - \varepsilon^2 \mathbf{A}_\lambda \mathbf{V}f)] \\ (\lambda + L)^{-1}\mathbf{P}(\mathbf{F} - \varepsilon^2 \mathbf{A}_\lambda \mathbf{V}f) + \varepsilon^2 \mathbf{V}f \end{pmatrix}$$

for $F = {}^\top(f, \mathbf{g}, h) \in H_{per,*}^1 \times L_{per}^2 \times L_{per}^2$.

Proof. For a given $F = {}^\top(f, \mathbf{g}, h) \in H_{per,*}^1 \times L_{per}^2 \times L_{per}^2$ we consider the problem $\mathcal{L}_{\varepsilon,\lambda}u = F$. As in the proof of Proposition 4.3, we set $\tilde{u} = u - {}^\top(0, \varepsilon^2 \mathbf{V}f, 0)$. Then we have $\mathcal{L}_{\varepsilon,\lambda}\tilde{u} = \tilde{F} - \varepsilon^2 \mathcal{A}_\lambda \mathbf{V}f$ with $\tilde{F} = {}^\top(0, \mathbf{g}, h)$. We thus arrive at

$$(\lambda + L)\tilde{U} = \mathbf{P}(\mathbf{F} - \varepsilon^2 \mathbf{A}_\lambda \mathbf{V}f),$$

where $\mathbf{F} = {}^\top(\mathbf{g}, h)$. Since $\lambda \in \rho(-L)$ we have

$${}^\top(\tilde{\mathbf{w}}, \theta) := \tilde{U} = (\lambda + L)^{-1}\mathbf{P}(\mathbf{F} - \varepsilon^2 \mathbf{A}_\lambda \mathbf{V}f).$$

It then follows that $\tilde{u} = {}^\top(p, \tilde{\mathbf{w}}, \theta)$ with

$$p = p_\lambda[\mathbf{P}(\mathbf{F} - \varepsilon^2 \mathbf{A}_\lambda \mathbf{V}f)]$$

satisfies $\mathcal{L}_{\varepsilon,\lambda}\tilde{u} = \tilde{F} - \varepsilon^2 \mathcal{A}_\lambda \mathbf{V}f$. Setting $u = {}^\top(p, \mathbf{w}, \theta)$ with $\mathbf{w} = \tilde{\mathbf{w}} + \varepsilon^2 \mathbf{V}f$ we have the desired expression. This completes the proof. $\square$

We next give a perturbation result of eigenvalues of $-L$.

Theorem 5.2. *Let $\lambda_0 \in \sigma(-L)$. Then λ_0 is an eigenvalue of $-L$ with finite multiplicity and the following assertions hold.*

(i) *There exist $C > 0$, $m \in \mathbb{N}$ and $r_1 > 0$ such that*

$$\{\lambda; 0 < |\lambda - \lambda_0| \leq r_1\} \subset \rho(-L)$$

and

$$\|(\lambda + L)^{-1}\mathbf{g}\|_{H^2} \leq \frac{C}{|\lambda - \lambda_0|^m} \|\mathbf{g}\|_2$$

for λ with $0 < |\lambda - \lambda_0| \leq r_1$.

(ii) For any $r_0 \in (0, r_1]$, there exists a positive number $\varepsilon_0 = \varepsilon_0(r_0, r_1, m, |\lambda_0|)$ such that $\sigma(-L_\varepsilon) \cap \{\lambda; |\lambda - \lambda_0| < r_0\} \neq \emptyset$ for $0 < \varepsilon \leq \varepsilon_0$, and if $|\lambda - \lambda_0| = r_0$, then $\lambda \in \rho(-L_\varepsilon)$ for $0 < \varepsilon \leq \varepsilon_0$ and the total projection P_ε , the sum of eigenprojections for eigenvalues of $-L_\varepsilon$ lying inside the circle $|\lambda - \lambda_0| = r_0$, satisfies

$$P_\varepsilon F = \frac{1}{2\pi i} \int_{|\lambda - \lambda_0| = r_0} (\lambda + L_\varepsilon)^{-1} F d\lambda = P_0 F + O(\varepsilon^2) F$$

in $H_{per,*}^1 \times L^2 \times L^2$ as $\varepsilon \rightarrow 0$. Here P_0 is the projection defined by

$$P_0 F = \begin{pmatrix} \Pi_{\lambda_0} \mathbf{P} \mathbf{F} \\ \Pi_{\lambda_0} \mathbf{P} \mathbf{F} \end{pmatrix}$$

for $F = {}^\top(f, \mathbf{g}, h) \in H_{per,*}^1 \times L^2 \times L^2$, where $\mathbf{F} = {}^\top(\mathbf{g}, h)$, Π_{λ_0} is the eigenprojection for the eigenvalue λ_0 of $-L$ given by

$$\Pi_{\lambda_0} \mathbf{P} \mathbf{F} = \frac{1}{2\pi i} \int_{|\lambda - \lambda_0| = r_0} (\lambda + L)^{-1} \mathbf{P} \mathbf{F} d\lambda$$

and

$$\Pi_{\lambda_0} \mathbf{P} \mathbf{F} = -\frac{1}{2\pi i} \int_{|\lambda - \lambda_0| = r_0} p_\lambda[\mathbf{P} \mathbf{F}] d\lambda,$$

i.e., $\Pi_{\lambda_0} \mathbf{F} \in H_{per,*}^1$ is the function satisfying

$$\nabla \Pi_{\lambda_0} \mathbf{P} \mathbf{F} = -\frac{1}{2\pi i} \int_{|\lambda - \lambda_0| = r_0} (\mathbb{I} - \mathbb{L}_\lambda (\lambda + L)^{-1}) \mathbf{P} \mathbf{F} d\lambda.$$

(iii) It holds that $\dim R(P_\varepsilon) = \dim R(P_0) = \dim R(\Pi_{\lambda_0})$ for $0 < \varepsilon \leq \varepsilon_0$.

Remark 5.3. Let $\tilde{m}$ be the algebraic multiplicity of λ_0 . Then one can take a basis of $R(\Pi_{\lambda_0})$ consisting of functions $\mathbf{U}_j, (\lambda_0 + L)\mathbf{U}_j, \dots, (\lambda_0 + L)^{m_j-1}\mathbf{U}_j$, $j = 1, \dots, r$, with $(\lambda_0 + L)^{m_j}\mathbf{U}_j = \mathbf{0}$ and $m_1 + \dots + m_r = \tilde{m}$. We denote by $p_{j\ell} \in H_{per,*}^1$ the pressure satisfying

$$\mathbb{L}_{\lambda_0}(\lambda_0 + L)^{m_j-\ell-1}\mathbf{U}_j = \mathbb{I}(\lambda_0 + L)^{m_j-\ell}\mathbf{U}_j - \nabla p_{j\ell}$$

for $1 \leq j \leq r$, $0 \leq \ell \leq m_j - 1$. It then follows that a set of functions ${}^\top(p_{j\ell}, (\lambda_0 + L)^{m_j-\ell}\mathbf{U}_j)$, $1 \leq j \leq r$, $0 \leq \ell \leq m_j - 1$, forms a basis of $R(P_0)$.

Proof. Since $-L$ has compact resolvent, its spectrum consists of discrete eigenvalues with finite multiplicities. Let the algebraic multiplicity of λ_0 be $\tilde{m}$. Then we see that there exist a $m \in \mathbb{N}$, $1 \leq m \leq \tilde{m}$, and a constant $r_1 > 0$ such that $\{\lambda; 0 < |\lambda - \lambda_0| \leq r_1\} \subset \rho(-L)$ and

$$\|(\lambda + L)^{-1} \mathbf{g}\|_{H^2} \leq \frac{C}{|\lambda - \lambda_0|^m} \|\mathbf{g}\|_2$$

for λ with $0 < |\lambda - \lambda_0| \leq r_1$. This proves (i).

As for (ii), we set

$$J = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

and write $\lambda + L_\varepsilon$ as

$$\lambda + L_\varepsilon = \mathcal{L}_{\varepsilon, \lambda} + \lambda J = \mathcal{L}_{\varepsilon, \lambda}(I + \lambda \mathcal{L}_{\varepsilon, \lambda}^{-1} J).$$

Since $JF = {}^\top(f, \mathbf{0}, 0)$ for $F = {}^\top(f, \mathbf{g}, h)$, we see from (i) and Proposition 5.1 that

$$\|\mathcal{L}_{\varepsilon, \lambda}^{-1} JF\|_{H^1 \times H^2 \times H^2} \leq C\varepsilon^2 \frac{(|\lambda_0| + r_0^m + 1)^2}{r_0^m} \|f\|_{H^1}$$

for $|\lambda - \lambda_0| = r_0$. Therefore, if

$$\varepsilon^2 \leq \frac{r_0^m}{4C(|\lambda_0| + r_0^m + 1)^2} \frac{1}{|\lambda_0| + r_0},$$

then

$$\|\lambda \mathcal{L}_{\varepsilon, \lambda}^{-1} JF\|_{H^1 \times H^2 \times H^2} \leq C\varepsilon^2 (|\lambda_0| + r_0) \frac{(|\lambda_0| + r_0^m + 1)^2}{r_0^m} \|f\|_{H^1} \leq \frac{1}{2} \|F\|_{H^1 \times L^2 \times L^2}$$

for $|\lambda - \lambda_0| = r_0$. It then follows that $(I + \lambda \mathcal{L}_{\varepsilon, \lambda}^{-1} J)$ is boundedly invertible both on Y and $D(L_\varepsilon)$ with estimates

$$\|(I + \lambda \mathcal{L}_{\varepsilon, \lambda}^{-1} J)^{-1} F\|_{H^1 \times L^2 \times L^2} \leq 2 \|F\|_{H^1 \times L^2 \times L^2}$$

for $F \in Y$ and

$$\|(I + \lambda \mathcal{L}_{\varepsilon, \lambda}^{-1} J)^{-1} F\|_{H^1 \times H^2 \times H^2} \leq 2 \|F\|_{H^1 \times H^2 \times H^2}$$

for $F \in D(L_\varepsilon)$. We thus find that $\lambda + L_\varepsilon = \mathcal{L}_{\varepsilon, \lambda} + \lambda J$ has a bounded inverse $(\lambda + L_\varepsilon)^{-1} = (\mathcal{L}_{\varepsilon, \lambda} + \varepsilon^2 \lambda J)^{-1}$ on Y which satisfies

$$\begin{aligned} (\lambda + L_\varepsilon)^{-1} &= (I + \lambda \mathcal{L}_{\varepsilon, \lambda}^{-1} J)^{-1} \mathcal{L}_{\varepsilon, \lambda}^{-1} \\ &= \sum_{N=0}^{\infty} (-\lambda)^N (\mathcal{L}_{\varepsilon, \lambda}^{-1} J)^N \mathcal{L}_{\varepsilon, \lambda}^{-1} \\ &= \mathcal{L}_{\varepsilon, \lambda}^{-1} - \lambda \mathcal{L}_{\varepsilon, \lambda}^{-1} J \sum_{N=0}^{\infty} (-\lambda)^N (\mathcal{L}_{\varepsilon, \lambda}^{-1} J)^N \mathcal{L}_{\varepsilon, \lambda}^{-1} \end{aligned}$$

with estimates

$$\|(\lambda + L_\varepsilon)^{-1} F\|_{H^1 \times L^2 \times L^2} \leq C \|F\|_{H^1 \times L^2 \times L^2}$$

and

$$\left\| \lambda \mathcal{L}_{\varepsilon, \lambda}^{-1} J \sum_{N=0}^{\infty} (-\lambda)^N (\mathcal{L}_{\varepsilon, \lambda}^{-1} J)^N \mathcal{L}_{\varepsilon, \lambda}^{-1} F \right\|_{H^1 \times L^2 \times L^2} \leq C\varepsilon^2 \|F\|_{H^1 \times L^2 \times L^2}.$$

Using these estimates with Proposition 5.1 and the assertion (i), we obtain the desired result (ii).

As for (iii), we first observe that P_0 is a projection, i.e., $P_0^2 = P_0$. By (ii) we have $P_\varepsilon \rightarrow P_0$ in the operator norm as $\varepsilon \rightarrow 0$. It then follows from [8, Chap. I, Lemma 4.10] that $\dim R(P_\varepsilon) = \dim R(P_0)$ for $0 < \varepsilon \ll 1$. On the other hand, $\dim R(P_0) = \dim R(\Pi_{\lambda_0})$. We thus obtain the desired result. This completes the proof. $\square$

Proof of Theorems 3.1 and 3.2. Theorem 3.2 is an immediate consequence of Theorem 5.2 (ii). To prove Theorems 3.1, we first observe that $\dim \text{Ker}(L) = \dim \text{Ker}(\mathcal{L}_{\varepsilon,0})$. Suppose that 0 is not a semisimple eigenvalue of L . Then

$$\dim R(\Pi_0) > \dim \text{Ker}(L) = \dim \text{Ker}(\mathcal{L}_{\varepsilon,0}) = \dim \text{Ker}(L_\varepsilon). \quad (5.2)$$

Since 0 is a semisimple eigenvalue of L_ε , we have $\dim \text{Ker}(L_\varepsilon) = \dim R(P_\varepsilon)$. This, together with (5.2), implies that $\dim R(\Pi_0) > \dim R(P_\varepsilon)$, which contradicts to Theorem 5.2 (iii) with $\lambda_0 = 0$. We thus conclude that 0 is a semisimple eigenvalue of $-L$.

Suppose that there exists $\lambda_0 \in \sigma(-L)$ with $\lambda_0 \neq 0$ and $\text{Re } \lambda_0 \geq 0$. Then, by Proposition 4.1 (ii), we have $\text{Im } \lambda_0 = O(1)$. Let $r_0 = (\text{Re } \lambda_0 + b_0)/2 > 0$. We see from Theorem 5.2 (ii) that there exist $\varepsilon_0 > 0$ such that $\{\lambda; |\lambda - \lambda_0| < r_0\} \cap \sigma(-L_\varepsilon) \neq \emptyset$ for all $0 < \varepsilon \leq \varepsilon_0$. But if $|\lambda - \lambda_0| < r_0$, then

$$\text{Re } \lambda = \text{Re}(\lambda - \lambda_0) + (\text{Re } \lambda_0 + b_0) - b_0 \geq r_0 - b_0 > -b_0.$$

This contradicts to the assumption $\{\lambda; \text{Re } \lambda \geq -b_0\} \setminus \{0\} \subset \rho(-L_{\varepsilon_n})$ with $\varepsilon_n \rightarrow 0$. The proof of Theorem 3.1 is complete. $\square$

6 Proof of Theorem 3.3

In this section we give a proof of Theorem 3.3. We consider the resolvent problem for $-L_\varepsilon$:

$$\lambda u + L_\varepsilon u = F, \quad (6.1)$$

where $u = {}^\top(p, \mathbf{w}, \theta) \in D(L_\varepsilon)$ and $F = {}^\top(f, \mathbf{g}, h) \in Y$. Problem (6.1) is written as

$$\varepsilon^2 \lambda p + \text{div } \mathbf{w} = \varepsilon^2 f, \quad (6.2)$$

$$\text{Pr}^{-1} \lambda \mathbf{w} - \Delta \mathbf{w} + \text{Pr}^{-1}(\mathbf{v}_s \cdot \nabla \mathbf{w} + \mathbf{w} \cdot \nabla \mathbf{v}_s) + \nabla p - \text{Ra} \theta \mathbf{e}_3 = \text{Pr}^{-1} \mathbf{g}, \quad (6.3)$$

$$\lambda \theta - \Delta \theta + \mathbf{v}_s \cdot \nabla \theta + \mathbf{w} \cdot \nabla \theta_s - \text{Ra} \mathbf{w} \cdot \mathbf{e}_3 = h, \quad (6.4)$$

and $u = {}^\top(p, \mathbf{w}, \theta)$ satisfies the boundary conditions (3.4) and (3.5).

Proposition 6.1. *There exist constants $a_\varepsilon = O(1) > 0$ and $b_\varepsilon = O(1) > 0$ as $\varepsilon \rightarrow 0$ such that $\{\lambda \in \mathbb{C}; \text{Re } \lambda \geq -a_\varepsilon \varepsilon^2 |\text{Im } \lambda|^2 + b_\varepsilon\} \subset \rho(-L_\varepsilon)$ for all $0 < \varepsilon \leq 1$.*

Proof. We set

$$|||u|||_2 = (\varepsilon^2 \|p\|_2^2 + \text{Pr}^{-1} \|\mathbf{w}\|_2^2 + \|\theta\|_2^2)^{\frac{1}{2}}$$

for $u = {}^\top(p, \mathbf{w}, \theta)$. Taking the inner product $\langle \cdot, \cdot \rangle_\varepsilon$ of (6.1) with u , we have

$$\begin{aligned} & \lambda |||u|||_2^2 + \|\nabla \mathbf{w}\|_2^2 + \|\nabla \theta\|_2^2 \\ &= 2\text{RaRe}(\theta, w^3) + 2i\text{Im}(p, \text{div } \mathbf{w}) - \text{Pr}^{-1}(\mathbf{v}_s \cdot \nabla \mathbf{w}, \mathbf{w}) - (\mathbf{v}_s \cdot \nabla \theta, \theta) \\ & \quad - \text{Pr}^{-1}(\mathbf{w} \cdot \nabla \mathbf{v}_s, \mathbf{w}) - (\mathbf{w} \cdot \nabla \theta_s, \theta) + \langle F, u \rangle_\varepsilon. \end{aligned} \quad (6.5)$$

Noting that $\text{Re}(\mathbf{v}_s \cdot \nabla \mathbf{w}, \mathbf{w}) = \text{Re}(\mathbf{v}_s \cdot \nabla \theta, \theta) = 0$, we see from the real part of (6.5) that

$$(\text{Re } \lambda - b_2) |||u|||_2^2 + \|\nabla \mathbf{w}\|_2^2 + \|\nabla \theta\|_2^2 \leq |||F|||_2 |||u|||_2, \quad (6.6)$$

where $b_2 = 3\{\|\nabla \mathbf{v}_s\|_\infty + (\text{Pr} + 1)(2\text{Ra} + \|\nabla \theta_s\|_\infty)\}$. The imaginary part of (6.5) yields

$$\begin{aligned} |\text{Im } \lambda| |||u|||_2^2 &\leq 2\|p\|_2 \|\text{div } \mathbf{w}\|_2 + \text{Pr}^{-1} \|\mathbf{v}_s\|_\infty \|\nabla \mathbf{w}\|_2 \|\mathbf{w}\|_2 \\ & \quad + \|\mathbf{v}_s\|_\infty \|\nabla \theta\|_2 \|\theta\|_2 + \text{Pr}^{-1} \|\nabla \mathbf{v}_s\|_\infty \|\mathbf{w}\|_2^2 + \|\nabla \theta_s\|_\infty \|\mathbf{w}\|_2 \|\theta\|_2 \\ & \quad + |||F|||_2 |||u|||_2 \\ &\leq \left\{ \left(2\varepsilon^{-1} + \text{Pr}^{-\frac{1}{2}} \|\mathbf{v}_s\|_\infty \right) \|\nabla \mathbf{w}\|_2 + \|\mathbf{v}_s\|_\infty \|\nabla \theta\|_2 \right. \\ & \quad \left. + \text{Pr}^{-\frac{1}{2}} \|\nabla \mathbf{v}_s\|_\infty \|\mathbf{w}\|_2 + \text{Pr}^{\frac{1}{2}} \|\nabla \theta_s\|_\infty \|\theta\|_2 + |||F|||_2 \right\} |||u|||_2, \end{aligned}$$

and hence,

$$\begin{aligned} |\text{Im } \lambda|^2 |||u|||_2^2 &\leq 36 \left\{ (4\varepsilon^{-2} + \text{Pr}^{-1} \|\mathbf{v}_s\|_\infty^2) \|\nabla \mathbf{w}\|_2^2 + \|\mathbf{v}_s\|_\infty^2 \|\nabla \theta\|_2^2 \right. \\ & \quad \left. + \text{Pr}^{-1} \|\nabla \mathbf{v}_s\|_\infty^2 \|\mathbf{w}\|_2^2 + \text{Pr} \|\nabla \theta_s\|_\infty^2 \|\theta\|_2^2 + |||F|||_2^2 \right\}. \end{aligned} \quad (6.7)$$

It then follows from (6.6) and (6.7) that

$$\begin{aligned} & (\text{Re } \lambda + a_\varepsilon \varepsilon^2 |\text{Im } \lambda|^2 - b_\varepsilon) |||u|||_2^2 + \frac{1}{2} (\|\nabla \mathbf{w}\|_2^2 + \|\nabla \theta\|_2^2) \\ & \leq (\delta^{-1} + 36a_\varepsilon \varepsilon^2) |||F|||_2^2 \end{aligned} \quad (6.8)$$

for all $\delta > 0$, where

$$a_\varepsilon = \frac{1}{72(4 + \varepsilon^2(\text{Pr}^{-1} + 1)\|\mathbf{v}_s\|_\infty^2)}$$

and

$$b_\varepsilon = \delta + b_2 + 36a_\varepsilon \varepsilon^2 (\|\nabla \mathbf{v}_s\|_\infty^2 + \text{Pr} \|\nabla \theta_s\|_\infty^2).$$

It remains to estimate $\|\partial_x p\|_2$. To this end we use the estimates in section 7. We introduce quantities $D(\mathbf{w}, \theta)$ and $M(\mathbf{w}, \theta)$ defined by

$$\begin{aligned} D(\mathbf{w}, \theta) &= \|\nabla \mathbf{w}\|_2^2 + \|\nabla \theta\|_2^2, \\ M(\mathbf{w}, \theta) &= \|\mathbf{v}_s\|_\infty^2 (\text{Pr}^{-2} \|\nabla \mathbf{w}\|_2^2 + \|\nabla \theta\|_2^2) + (\text{Pr}^{-2} \|\nabla \mathbf{w}_s\|_\infty^2 + \|\nabla \theta_s\|_\infty^2) \|\mathbf{w}\|_2^2 \\ & \quad + \text{Ra}(1 + \text{Ra})(\|\mathbf{w}\|_2^2 + \|\theta\|_2^2). \end{aligned}$$

We set

$$\Sigma_1 = \{\lambda \in ; \operatorname{Re} \lambda + a_\varepsilon \varepsilon^2 |\operatorname{Im} \lambda| - b_\varepsilon \geq 0\}$$

and

$$\Sigma_2 = \{\lambda \in ; \operatorname{dist}(\lambda, \Sigma_1) \geq 1\}.$$

In what follows we assume that $\lambda \in \Sigma_2$. It then follows that

$$M(\mathbf{w}, \theta) \leq C \|F\|_2^2.$$

Here C is a positive constant depending on Ra and Pr but not on ε .

We consider $(1 + \varepsilon^{-2}) \times (6.8) + d_0 \times (7.8)$. Then taking $d_0 > 0$ suitably small, we see that

$$\begin{aligned} & (\operatorname{Re} \lambda + a_\varepsilon \varepsilon^2 |\operatorname{Im} \lambda|^2 - b_\varepsilon) \|u\|_2^2 + (\operatorname{Re} \lambda + |\operatorname{Im} \lambda|) D(\mathbf{w}, \theta) \\ & + (\operatorname{Re} \lambda + \varepsilon^2 |\operatorname{Im} \lambda|^2) (\|\partial_{x'} u\|_2^2 + \varepsilon^2 \|\partial_{x_3} p\|_2^2) \\ & + b_0 \left(\left(1 + \frac{1}{\varepsilon^2}\right) D(\mathbf{w}, \theta) + \|\varepsilon^2 \lambda p\|_{H^1}^2 + \frac{1}{2} |\lambda|^2 \|u\|_2^2 \right) \\ & + b_0 (\|\partial_x^2 \mathbf{w}\|_2^2 + \|\partial_x^2 \theta\|_2^2 + \|\partial_x p\|_2^2) \\ & \leq C(1 + \varepsilon^2) \left\{ \varepsilon^2 \|\partial_x f\|_2^2 + \left(1 + \frac{1}{\varepsilon^2}\right) \|F\|_2^2 \right\} \end{aligned} \quad (6.9)$$

with some positive constants b_0 and C uniformly for $0 < \varepsilon \leq 1$. This completes the proof. $\square$

We next show that the spectrum of $-L_\varepsilon$ in a disc with radius $O(\varepsilon^{-1})$ can be viewed as a perturbation of the one of $-L$. From the assumption of Theorem 3.3, we see that one can take the constant c_0 in Proposition 4.1 (ii) so that $c_0 < 0$ by changing $a_0 > 0$ suitably. In what follows we fix these $a_0 > 0$ and $c_0 < 0$. Using Proposition 4.1 (ii), Lemma 4.2, (5.1) and Proposition 5.1, we have the following estimates for $\mathcal{L}_{\varepsilon, \lambda}^{-1}$.

Proposition 6.2. *Let $\varepsilon > 0$. If $\lambda \in \Sigma \setminus \{0\}$, then $\mathcal{L}_{\varepsilon, \lambda}$ has a bounded inverse $\mathcal{L}_{\varepsilon, \lambda}^{-1}$ and ${}^\top(p, \mathbf{v}, \theta) = \mathcal{L}_{\varepsilon, \lambda}^{-1} F$ for $F = {}^\top(f, \mathbf{g}, h) \in Y$ satisfies*

$$\|U\|_2 \leq \frac{C}{|\lambda|} \sum_{j=1}^2 |\langle \langle F, u_j^* \rangle \rangle_\varepsilon| + C \left\{ \varepsilon^2 \|f\|_{H^1} + \frac{1}{|\lambda| + 1} \|F\|_2 \right\},$$

$$\|\partial_x^2 U\|_2 + \|\partial_x p\|_2 \leq \frac{C}{|\lambda|} \sum_{j=1}^2 |\langle \langle F, u_j^* \rangle \rangle_\varepsilon| + C \left\{ \varepsilon^2 (|\lambda| + 1) \|f\|_{H^1} + \|F\|_2 \right\},$$

where $U = {}^\top(\mathbf{w}, \theta)$ and $F = {}^\top(\mathbf{g}, h)$. Furthermore, if $F \in Y_{1, \varepsilon}$, then ${}^\top(p, \mathbf{v}, \theta) = \mathcal{L}_{\varepsilon, \lambda}^{-1} F$ exists uniquely in $Y_{1, \varepsilon}$ and the above estimates hold with $\lambda = 0$ and $\langle \langle F, u_j^* \rangle \rangle_\varepsilon = 0$ ($j = 1, 2$).

Proof. As in the proof of Proposition 4.3 (ii), we see that

$$\Pi_0 \mathbf{P}[\mathbf{F} - \varepsilon^2 \mathbf{A}_\lambda \mathbf{V} f] = \sum_{j=1}^2 \left\{ \langle \langle F, u_j^* \rangle \rangle_\varepsilon - \frac{\varepsilon^2 \lambda}{\text{Pr}} (\mathbf{V} f, \mathbf{w}_j^*) \right\} \mathbf{U}_j^{(0)}.$$

Therefore, applying Proposition 4.1 (ii), Lemma 4.2 and Proposition 5.1 together with (5.1), we have the desired result. This completes the proof. $\square$

Proposition 6.3. *There exist positive numbers ε_1 and a_1 such that*

$$\Sigma \cap \{\lambda \in \mathbb{C}; \text{Re } \lambda \geq -b_0, |\lambda| \leq a_1 \varepsilon^{-1}\} \subset \rho(-L_\varepsilon|_{Y_{1,\varepsilon}})$$

for all $0 < \varepsilon \leq \varepsilon_1$. Here

$$Y_{1,\varepsilon} = R(L_\varepsilon) = \{u \in Y; \langle \langle u, u_j^* \rangle \rangle_\varepsilon = 0, j = 1, 2\}.$$

Proof. As in the proof of Theorem 5.2, we write the resolvent problem

$$(\lambda + L_\varepsilon)u = F$$

on $Y_{1,\varepsilon}$ as

$$\mathcal{L}_{\varepsilon,\lambda} u + \lambda J u = F, \tag{6.10}$$

where $F = {}^\top(f, \mathbf{g}, h) \in Y_{1,\varepsilon}$. If $\lambda \in \Sigma$, then it follows from Proposition 6.2 that (6.10) is written as

$$\mathcal{L}_{\varepsilon,\lambda}(I + \lambda \mathcal{L}_{\varepsilon,\lambda}^{-1} J)u = F,$$

and, furthermore, we have

$$\|\mathcal{L}_{\varepsilon,\lambda}^{-1} J F\|_{H^1 \times H^2 \times H^2} \leq \varepsilon^2 C_1 (|\lambda| + 1) \|f\|_{H^1}$$

for all $F = {}^\top(f, \mathbf{g}, h) \in Y_{1,\varepsilon}$. It then follows that there exists $\varepsilon_1 > 0$ such that if $\lambda \in \Sigma$ and $|\lambda| \leq 1/(4\sqrt{C_1}\varepsilon)$, then $\mathcal{L}_{\varepsilon,\lambda}^{-1} J F \in D(\mathcal{L}_{\varepsilon,\lambda}) = D(L_\varepsilon)$ and $\|\lambda \mathcal{L}_{\varepsilon,\lambda}^{-1} J F\|_{H^1 \times H^2 \times H^2} \leq \frac{1}{2} \|F\|_{H^1 \times L^2 \times L^2}$ for $0 < \varepsilon \leq \varepsilon_1$. Therefore, $(I + \lambda \mathcal{L}_{\varepsilon,\lambda}^{-1} J)$ is boundedly invertible both on $Y_{1,\varepsilon}$ and $Y_{1,\varepsilon} \cap D(L_\varepsilon)$ with estimates

$$\|(I + \lambda \mathcal{L}_{\varepsilon,\lambda}^{-1} J)^{-1} F\|_{H^1 \times L^2 \times L^2} \leq 2 \|F\|_{H^1 \times L^2 \times L^2}$$

for $F \in Y_{1,\varepsilon}$ and

$$\|(I + \lambda \mathcal{L}_{\varepsilon,\lambda}^{-1} J)^{-1} F\|_{H^1 \times H^2 \times H^2} \leq 2 \|F\|_{H^1 \times H^2 \times H^2}$$

for $F \in Y_{1,\varepsilon} \cap D(L_\varepsilon)$. We thus find that $\lambda + L_\varepsilon = \mathcal{L}_{\varepsilon,\lambda} + \lambda J$ has a bounded inverse $(\lambda + L_\varepsilon)^{-1} = (\mathcal{L}_{\varepsilon,\lambda} + \varepsilon^2 \lambda J)^{-1}$ on $Y_{1,\varepsilon}$ which satisfies

$$(\lambda + L_\varepsilon)^{-1} = \mathcal{L}_{\varepsilon,\lambda}^{-1} - \lambda \mathcal{L}_{\varepsilon,\lambda}^{-1} J \sum_{N=0}^{\infty} (-\lambda)^N (\mathcal{L}_{\varepsilon,\lambda}^{-1} J)^N \mathcal{L}_{\varepsilon,\lambda}^{-1}$$

and

$$\begin{aligned} \|(\lambda + L_\varepsilon)^{-1} F\|_{H^1 \times L^2 \times L^2} &\leq 2C_1 \{ \varepsilon^2 (|\lambda| + 1) \|f\|_{H^1} + \|\mathbf{F}\|_2 \} \\ &\leq 2C_1 \left\{ \varepsilon \left(\frac{1}{4\sqrt{C_1}} + \varepsilon \right) \|f\|_{H^1} + \|\mathbf{F}\|_2 \right\} \end{aligned}$$

with $\mathbf{F} = {}^\top(\mathbf{g}, h)$. This completes the proof. $\square$

Theorem 3.3 follows from Propositions 4.3, 6.1 and 6.3 if $\sqrt{b_\varepsilon/a_\varepsilon} < a_1$ for $0 < \varepsilon \ll 1$. In the case $\sqrt{b_\varepsilon/a_\varepsilon} \geq a_1$, there is some range of λ near the imaginary axis with $|\operatorname{Im} \lambda| = O(\varepsilon^{-1})$ to be proved that it belongs to $\rho(-L_\varepsilon)$.

To prove Theorem 3.3 when $\sqrt{b_\varepsilon/a_\varepsilon} \geq a_1$, we first prepare an estimate for the θ -component. We recall that the Poincaré inequality

$$\|\nabla \theta\|_2 \geq \beta \|\theta\|_2$$

holds for $\theta \in H_{0,per}^1$ with some positive constant β .

Proposition 6.4. *Let ${}^\top(p, \mathbf{w}, \theta)$ be a solution of (6.2)–(6.4) under boundary conditions (3.4) and (3.5). Then if $\operatorname{Re} \lambda \geq -\frac{\beta^2}{2}$, the following estimates hold:*

$$\begin{aligned} \|\theta\|_2 &\leq \frac{1}{|\operatorname{Im} \lambda|} \left(1 + \frac{2\|\mathbf{v}_s\|_\infty}{\beta} \right) \{ (\|\nabla \theta_s\|_\infty + \operatorname{Ra}) \|\mathbf{w}\|_2 + \|h\|_2 \}, \\ \|\nabla \theta\|_2 &\leq \frac{2}{\beta} \{ (\|\nabla \theta_s\|_\infty + \operatorname{Ra}) \|\mathbf{w}\|_2 + \|h\|_2 \}. \end{aligned}$$

Proof. We take the inner product of (6.4) with θ to obtain

$$\begin{aligned} \lambda \|\theta\|_2^2 + \|\nabla \theta\|_2^2 &= - (i \operatorname{Im} (\mathbf{v}_s \cdot \nabla \theta, \theta) + (\mathbf{w} \cdot \nabla \theta_s, \theta) - \operatorname{Ra} (w^3, \theta)) + (h, \theta). \end{aligned} \tag{6.11}$$

Using the Poincaré inequality we see from the real part of (6.11) that

$$\left(\operatorname{Re} \lambda + \frac{\beta^2}{2} \right) \|\theta\|_2^2 + \frac{1}{2} \|\nabla \theta\|_2^2 \leq \{ (\|\nabla \theta_s\|_\infty + \operatorname{Ra}) \|\mathbf{w}\|_2 + \|h\|_2 \} \|\theta\|_2.$$

This implies that if $\operatorname{Re} \lambda \geq -\frac{\beta^2}{2}$, then

$$\|\nabla \theta\|_2 \leq \frac{2}{\beta} \{ (\|\nabla \theta_s\|_\infty + \operatorname{Ra}) \|\mathbf{w}\|_2 + \|h\|_2 \}. \tag{6.12}$$

Furthermore, the imaginary part of (6.11), together with (6.12), implies that

$$\begin{aligned} |\operatorname{Im} \lambda| \|\theta\|_2^2 &= | - \operatorname{Im} ((\mathbf{v}_s \cdot \nabla \theta, \theta) + (\mathbf{w} \cdot \nabla \theta_s, \theta) - \operatorname{Ra} (w^3, \theta)) + \operatorname{Im} (h, \theta) | \\ &\leq \{ \|\mathbf{v}_s\|_\infty \|\nabla \theta\|_2 + (\|\nabla \theta_s\|_\infty + \operatorname{Ra}) \|\mathbf{w}\|_2 + \|h\|_2 \} \|\theta\|_2 \\ &\leq \left(1 + \frac{2\|\mathbf{v}_s\|_\infty}{\beta} \right) \{ (\|\nabla \theta_s\|_\infty + \operatorname{Ra}) \|\mathbf{w}\|_2 + \|h\|_2 \} \|\theta\|_2, \end{aligned}$$

from which we have the desired estimate for $\|\theta\|_2$. This completes the proof. $\square$

We are now in a position to complete the proof of Theorem 3.3.

Proposition 6.5. For given $\mu_* > 0$ and $\eta_* > 0$ there exist constants $\varepsilon_1 > 0$ and $c_2 > 0$ such that if

$$\inf_{\mathbf{w} \in H_{0,per}^1, \mathbf{w} \neq \mathbf{0}} \frac{\operatorname{Re}(\mathbf{w} \cdot \nabla \mathbf{v}_s, \mathbf{w})}{\|\nabla \mathbf{w}\|_2^2} \geq -\frac{\operatorname{Pr}}{32},$$

then

$$\left\{ \lambda = \mu + i\frac{\eta}{\varepsilon}; -c_2 \leq \mu \leq \mu_*, |\eta| \geq \eta_* \right\} \subset \rho(-L_\varepsilon)$$

for all $0 < \varepsilon \leq \varepsilon_1$. Here ε_1 and c_2 are positive constants depending only on Pr , Ra , $\|\mathbf{v}_s\|_{C^1}$, $\|\nabla \theta_s\|_\infty$, β , μ_* and η_*

Proof. We see from (6.2) that

$$p = -\frac{1}{\varepsilon^2 \lambda} \operatorname{div} \mathbf{w} + \frac{1}{\lambda} f. \quad (6.13)$$

Substituting (6.13) into (6.3), we have

$$\frac{\varepsilon^2 \lambda^2}{\operatorname{Pr}} \mathbf{w} - \varepsilon^2 \lambda \Delta \mathbf{w} - \nabla \operatorname{div} \mathbf{w} + \frac{\varepsilon^2 \lambda}{\operatorname{Pr}} (\mathbf{v}_s \cdot \nabla \mathbf{w} + \mathbf{w} \cdot \nabla \mathbf{v}_s) - \varepsilon^2 \lambda \operatorname{Ra} \theta \mathbf{e}_3 = \varepsilon^2 \mathbf{G}_\lambda, \quad (6.14)$$

where $\mathbf{G}_\lambda = \frac{\lambda}{\operatorname{Pr}} \mathbf{g} - \nabla f$.

Let $\lambda = \mu + i\frac{\eta}{\varepsilon}$ with $|\eta| \geq \eta_*(> 0)$. Without loss of generality we may assume $\eta \geq \eta_*$. Taking the inner product of (6.14) with $\mathbf{w}$, we have

$$\begin{aligned} & \frac{\varepsilon^2 \lambda^2}{\operatorname{Pr}} \|\mathbf{w}\|_2^2 + \varepsilon^2 \lambda \|\nabla \mathbf{w}\|_2^2 + \|\operatorname{div} \mathbf{w}\|_2^2 \\ &= -\varepsilon^2 \lambda \left(\operatorname{Pr}^{-1} (\mathbf{v}_s \cdot \nabla \mathbf{w}, \mathbf{w}) + \operatorname{Pr}^{-1} (\mathbf{w} \cdot \nabla \mathbf{v}_s, \mathbf{w}) - \operatorname{Ra}(\theta, w^3) \right) + \varepsilon^2 (\mathbf{G}_\lambda, \mathbf{w}). \end{aligned} \quad (6.15)$$

Since $\lambda^2 = (\mu^2 - \varepsilon^{-2} \eta^2) + 2i\varepsilon^{-1} \mu \eta$ and $\operatorname{Re}(\mathbf{v}_s \cdot \nabla \mathbf{w}, \mathbf{w}) = 0$, the real and imaginary parts of (6.15) yield

$$\begin{aligned} & \frac{1}{\operatorname{Pr}} (\varepsilon^2 \mu^2 - \eta^2) \|\mathbf{w}\|_2^2 + \varepsilon^2 \mu \|\nabla \mathbf{w}\|_2^2 + \|\operatorname{div} \mathbf{w}\|_2^2 \\ &= -\varepsilon^2 \mu \operatorname{Re} \left(\operatorname{Pr}^{-1} (\mathbf{w} \cdot \nabla \mathbf{v}_s, \mathbf{w}) - \operatorname{Ra}(\theta, w^3) \right) \\ & \quad + \varepsilon \eta \operatorname{Im} \left(\operatorname{Pr}^{-1} (\mathbf{v}_s \cdot \nabla \mathbf{w}, \mathbf{w}) + \operatorname{Pr}^{-1} (\mathbf{w} \cdot \nabla \mathbf{v}_s, \mathbf{w}) - \operatorname{Ra}(\theta, w^3) \right) \\ & \quad + \varepsilon^2 \operatorname{Re} (\mathbf{G}_\lambda, \mathbf{w}) \end{aligned} \quad (6.16)$$

and

$$\begin{aligned} & \frac{2\varepsilon \mu \eta}{\operatorname{Pr}} \|\mathbf{w}\|_2^2 + \varepsilon \eta \|\nabla \mathbf{w}\|_2^2 \\ &= -\varepsilon^2 \mu \operatorname{Im} \left(\operatorname{Pr}^{-1} (\mathbf{v}_s \cdot \nabla \mathbf{w}, \mathbf{w}) + \operatorname{Pr}^{-1} (\mathbf{w} \cdot \nabla \mathbf{v}_s, \mathbf{w}) - \operatorname{Ra}(\theta, w^3) \right) \\ & \quad - \varepsilon \eta \operatorname{Re} \left(\operatorname{Pr}^{-1} (\mathbf{w} \cdot \nabla \mathbf{v}_s, \mathbf{w}) - \operatorname{Ra}(\theta, w^3) \right) \\ & \quad + \varepsilon^2 \operatorname{Im} (\mathbf{G}_\lambda, \mathbf{w}). \end{aligned} \quad (6.17)$$

Since $|\operatorname{Im} \lambda| = \frac{\eta}{\varepsilon}$, by Proposition 6.4, we have

$$|\operatorname{Ra}(\theta, w^3)| \leq \frac{\operatorname{Ra}\varepsilon}{\eta} \left(1 + \frac{2\|\mathbf{v}_s\|_\infty}{\beta}\right) \{(\|\nabla\theta_s\|_\infty + \operatorname{Ra})\|\mathbf{w}\|_2^2 + \|h\|_2\|\mathbf{w}\|_2\},$$

and hence, we see from (6.16) that

$$\begin{aligned} & \frac{1}{\operatorname{Pr}}(\eta^2 - \varepsilon^2\mu^2)\|\mathbf{w}\|_2^2 \\ & \leq \left(\varepsilon^2\mu + \frac{\eta}{\eta_*} + \varepsilon\eta\right)\|\nabla\mathbf{w}\|_2^2 + ((\varepsilon^2|\mu| + \varepsilon\eta)\operatorname{Pr}^{-1}\|\nabla\mathbf{v}_s\|_\infty + \varepsilon\eta\operatorname{Pr}^{-2}\|\mathbf{v}_s\|_\infty^2)\|\mathbf{w}\|_2^2 \\ & \quad + (\varepsilon^2|\mu| + \varepsilon\eta)\frac{\operatorname{Ra}\varepsilon}{\eta} \left(1 + \frac{2\|\mathbf{v}_s\|_\infty}{\beta}\right) \{(\|\nabla\theta_s\|_\infty + \operatorname{Ra})\|\mathbf{w}\|_2^2 + \|h\|_2\|\mathbf{w}\|_2\} \\ & \quad + \varepsilon^2\|\mathbf{G}_\lambda\|_2\|\mathbf{w}\|_2. \end{aligned} \tag{6.18}$$

Similarly, we see from (6.17) that

$$\begin{aligned} & \frac{2\mu\eta}{\operatorname{Pr}}\|\mathbf{w}\|_2^2 + \frac{3\eta}{4}\|\nabla\mathbf{w}\|_2^2 \\ & \leq -\eta\operatorname{Pr}^{-1}\operatorname{Re}(\mathbf{w} \cdot \nabla\mathbf{v}_s, \mathbf{w}) + \left(\frac{\varepsilon^2|\mu|^2\operatorname{Pr}^{-2}\|\mathbf{v}_s\|_\infty^2}{\eta} + \varepsilon|\mu|\operatorname{Pr}^{-1}\|\nabla\mathbf{v}_s\|_\infty\right)\|\mathbf{w}\|_2^2 \\ & \quad + (\varepsilon|\mu| + \eta)\frac{\operatorname{Ra}\varepsilon}{\eta} \left(1 + \frac{2\|\mathbf{v}_s\|_\infty}{\beta}\right) \{(\|\nabla\theta_s\|_\infty + \operatorname{Ra})\|\mathbf{w}\|_2^2 + \|h\|_2\|\mathbf{w}\|_2\} \\ & \quad + \varepsilon\|\mathbf{G}_\lambda\|_2\|\mathbf{w}\|_2. \end{aligned} \tag{6.19}$$

We set $M = \inf_{\mathbf{w} \in H_{0,per}^1, \mathbf{w} \neq \mathbf{0}} \frac{\operatorname{Re}(\mathbf{w} \cdot \nabla\mathbf{v}_s, \mathbf{w})}{\|\nabla\mathbf{w}\|_2^2}$. It then follows from $\frac{\eta_*}{4} \times (6.18) + (6.19)$ that

$$\begin{aligned} & \left(I_{\varepsilon,\mu,\eta} + \frac{\eta_*}{4\operatorname{Pr}}\eta^2\right)\|\mathbf{w}\|_2^2 + \left(\frac{\eta}{2} - \frac{\varepsilon^2\eta_*\mu}{4} - \frac{\varepsilon\eta_*\eta}{4}\right)\|\nabla\mathbf{w}\|_2^2 \\ & \leq -\eta\operatorname{Pr}^{-1}M\|\nabla\mathbf{w}\|_2^2 + C_{\varepsilon,\lambda}(\|F\|_{H^1 \times (L^2)^2})\|\mathbf{w}\|_2, \end{aligned} \tag{6.20}$$

where

$$\begin{aligned} I_{\varepsilon,\mu,\eta} &= \frac{2\mu\eta}{\operatorname{Pr}} - \frac{\varepsilon^2\mu^2}{\operatorname{Pr}} \left(\frac{\eta_*}{4} + \frac{\|\mathbf{v}_s\|_\infty^2}{\eta\operatorname{Pr}}\right) - \frac{\varepsilon|\mu|}{\operatorname{Pr}} \left(1 + \frac{\varepsilon\eta_*}{4}\right)\|\nabla\mathbf{v}_s\|_\infty \\ & \quad - \frac{\varepsilon\eta_*\eta}{4\operatorname{Pr}} \left(\|\nabla\mathbf{v}_s\|_\infty + \frac{\|\mathbf{v}_s\|_\infty^2}{\operatorname{Pr}}\right) \\ & \quad - \varepsilon\operatorname{Ra} \left(1 + \frac{\varepsilon\eta_*}{4}\right) \left(\frac{\varepsilon|\mu|}{\eta} + 1\right) \left(1 + \frac{2\|\mathbf{v}_s\|_\infty}{\beta}\right) (\|\nabla\mathbf{v}_s\|_\infty + \operatorname{Ra}), \\ C_{\varepsilon,\lambda}(\|F\|_{H^1 \times (L^2)^2}) &= \varepsilon \left(1 + \frac{\varepsilon\eta_*}{4}\right) \left[\operatorname{Ra} \left(\frac{\varepsilon|\mu|}{\eta} + 1\right) \left(1 + \frac{2\|\mathbf{v}_s\|_\infty}{\beta}\right) \|h\|_2 + \|\mathbf{G}_\lambda\|_2\right]. \end{aligned}$$

We take $\varepsilon > 0$ so small that $\varepsilon^2 \mu_* \leq \frac{1}{8}$ and $\varepsilon \eta_* \leq \frac{1}{8}$ in (6.20). Then by using the Poincaré inequality, we have

$$\left(I_{\varepsilon, \mu, \eta} + \frac{\eta_*}{4\text{Pr}} \eta^2 + \frac{\beta^2}{8} \eta \right) \|\mathbf{w}\|_2^2 + \frac{\eta}{16} \|\nabla \mathbf{w}\|_2^2 \leq -\eta \text{Pr}^{-1} M \|\nabla \mathbf{w}\|_2^2 + C_{\varepsilon, \lambda} (\|F\|_{H^1 \times (L^2)^2}) \|\mathbf{w}\|_2. \quad (6.21)$$

and hence, if

$$M \geq -\frac{\text{Pr}}{32},$$

we find from (6.21) that

$$\left(I_{\varepsilon, \mu, \eta} + \frac{\beta^2}{8} \eta \right) \|\mathbf{w}\|_2^2 + \frac{\eta}{32} \|\nabla \mathbf{w}\|_2^2 \leq C_{\varepsilon, \lambda} (\|F\|_{H^1 \times (L^2)^2}) \|\mathbf{w}\|_2. \quad (6.22)$$

Therefore, if we show $I_{\varepsilon, \mu, \eta} + \frac{\beta^2}{8} \eta \geq \frac{\beta^2}{16} \eta$ for $-c_2 \leq \mu \leq \mu_*$ and $\eta \geq \eta_*$ with some $c_2 > 0$ when $0 < \varepsilon \ll 1$, then the proof of Proposition 6.6 is complete.

Let us prove $I_{\varepsilon, \mu, \eta} + \frac{\beta^2}{8} \eta \geq \frac{\beta^2}{16} \eta$. When $0 \leq \mu \leq \mu_*$, we have

$$\begin{aligned} I_{\varepsilon, \mu, \eta} \geq & \eta \left\{ \mu \left(\frac{2}{\text{Pr}} - K_\varepsilon \right) - \frac{\varepsilon \eta_*}{4\text{Pr}} \left(\|\nabla \mathbf{v}_s\|_\infty + \frac{\|\mathbf{v}_s\|_\infty^2}{\text{Pr}} \right) \right. \\ & \left. - \frac{\varepsilon \text{Ra}}{\eta_*} \left(1 + \frac{\varepsilon \eta_*}{4} \right) \left(1 + \frac{2\|\mathbf{v}_s\|_\infty}{\beta} \right) (\|\nabla \mathbf{v}_s\|_\infty + \text{Ra}) \right\}, \end{aligned}$$

where

$$\begin{aligned} K_\varepsilon = & \frac{\varepsilon^2 \mu_*}{\text{Pr}} \left(\frac{1}{4} + \frac{\|\mathbf{v}_s\|_\infty^2}{\eta_*^2 \text{Pr}} \right) + \frac{\varepsilon}{\text{Pr}} \left(\frac{\varepsilon}{4} + \frac{1}{\eta_*} \right) \|\nabla \mathbf{v}_s\|_\infty \\ & + \frac{\varepsilon^2 \text{Ra}}{\eta_*^2} \left(1 + \frac{\varepsilon \eta_*}{4} \right) \left(1 + \frac{2\|\mathbf{v}_s\|_\infty}{\beta} \right) (\|\nabla \mathbf{v}_s\|_\infty + \text{Ra}). \end{aligned}$$

Therefore, if we take $\varepsilon > 0$ so that

$$\frac{2}{\text{Pr}} - K_\varepsilon \geq 0$$

and

$$\begin{aligned} \frac{\beta^2}{16} - \frac{\varepsilon \eta_*}{4\text{Pr}} \left(\|\nabla \mathbf{v}_s\|_\infty + \frac{\|\mathbf{v}_s\|_\infty^2}{\text{Pr}} \right) \\ - \frac{\varepsilon \text{Ra}}{\eta_*} \left(1 + \frac{\varepsilon \eta_*}{4} \right) \left(1 + \frac{2\|\mathbf{v}_s\|_\infty}{\beta} \right) (\|\nabla \mathbf{v}_s\|_\infty + \text{Ra}) \geq 0, \end{aligned}$$

then we have $I_{\varepsilon, \mu, \eta} + \frac{\beta^2}{8} \eta \geq \frac{\beta^2}{16} \eta$ for $0 \leq \mu \leq \mu_*$ and $\eta \geq \eta_*$, and hence,

$$\frac{\beta^2}{16} \eta \|\mathbf{w}\|_2^2 + \frac{\eta}{32} \|\nabla \mathbf{w}\|_2^2 \leq C_{\varepsilon, \lambda} (\|F\|_{H^1 \times (L^2)^2}) \|\mathbf{w}\|_2 \quad (6.23)$$

for $0 \leq \mu \leq \mu_*$ and $\eta \geq \eta_*$.

When $\mu < 0$, we assume that $|\mu| \leq \frac{\text{Pr}}{64} \beta^2$. We write $I_{\varepsilon, \mu, \eta}$ as $I_{\varepsilon, \mu, \eta} = \eta \tilde{I}_\varepsilon(\mu, \eta)$. Then we see that $\tilde{I}_\varepsilon(\mu, \eta) \geq \tilde{I}_\varepsilon(-\frac{\text{Pr}}{64} \beta^2, \eta_*) = -\frac{\beta^2}{32} + O(\varepsilon)$, and so we take $\varepsilon > 0$ in

such a way that $\tilde{I}_\varepsilon(-\frac{\text{Pr}}{64}\beta^2, \eta_*) + \frac{\beta^2}{16} \geq 0$. It then follows that $I_{\varepsilon, \mu, \eta} + \frac{\beta^2}{8}\eta \geq \frac{\beta^2}{16}\eta$, and hence, we have (6.23) for $-c_2 \leq \mu < 0$ and $\eta \geq \eta_*$ with $c_2 = \frac{\text{Pr}}{128}\beta^2$. Similarly to the proof of Proposition 6.1, we can derive the necessary estimate for $\|\partial_x p\|_2^2$ by combining estimate (6.23), Propositions 6.4 and 7.5. This completes the proof. $\square$

Theorem 3.3 now follows by taking $\eta_* = \frac{a_1}{2}$, $\mu_* = 2b_\varepsilon|_{\varepsilon=1, \delta=1}$ and $\varepsilon > 0$ sufficiently small.

7 Estimate for $\|\partial_x p\|_2$

In this section we give basic energy estimates which yield the estimate for $\|\partial_x p\|_2$ in the resolvent estimate. The following Propositions 7.1–7.4 can be obtained by the Matsumura-Nishida energy method [9]. We here give an outline of the proof.

Proposition 7.1. *Let $u = {}^\top(p, \mathbf{w}, \theta)$ be a solution of (6.2)–(6.4) satisfying boundary conditions (3.4) and (3.5). Then*

$$\begin{aligned} & (\text{Re } \lambda + \varepsilon^2 |\text{Im } \lambda|^2) \|\partial_{x'} u\|_2^2 + \frac{1}{4} D(\partial_{x'} \mathbf{w}, \partial_{x'} \theta) + \frac{1}{4} \|\varepsilon^2 \lambda \partial_{x'} p\|_2^2 \\ & \leq C(1 + \varepsilon^2) \{ \delta \varepsilon^2 \|\partial_{x'} p\|_2^2 + \delta^{-1} \varepsilon^2 \|\partial_{x'} f\|_2^2 + \|F\|_2^2 + M(\mathbf{w}, \theta) \} \end{aligned} \quad (7.1)$$

for any $\delta > 0$, where C is a positive constant independent of ε and δ .

Proposition 7.1 can be proved as in the proof of Proposition 6.1 by taking the inner product $\langle \partial_{x'}(6.1), \partial_{x'} u \rangle_\varepsilon$ and using the relation $\varepsilon^2 \lambda \partial_{x'} p = -\text{div } \partial_{x'} \mathbf{w} + \varepsilon^2 \partial_{x'} f$ which follows from (6.2).

We next compute the inner product $\langle (6.1), \lambda u \rangle_\varepsilon$ to obtain the following estimate.

Proposition 7.2. *Let $u = {}^\top(p, \mathbf{w}, \theta)$ be a solution of (6.2)–(6.4) satisfying boundary conditions (3.4) and (3.5). Then*

$$(\text{Re } \lambda + |\text{Im } \lambda|) D(\mathbf{w}, \theta) + \frac{1}{2} |\lambda|^2 \|u\|_2^2 \leq C \left\{ \frac{1}{\varepsilon^2} \|\text{div } \mathbf{w}\|_2^2 + M(\mathbf{w}, \theta) \right\}, \quad (7.2)$$

where C is a positive constant independent of ε .

As for the normal derivative of p , we have the following estimate.

Proposition 7.3. *Let $u = {}^\top(p, \mathbf{w}, \theta)$ be a solution of (6.2)–(6.4) satisfying boundary conditions (3.4) and (3.5). Then*

$$\begin{aligned} & (\text{Re } \lambda + \varepsilon^2 |\text{Im } \lambda|^2) \varepsilon^2 \|\partial_{x_3} p\|_2^2 + \frac{1}{4} \|\partial_{x_3} p\|_2^2 + \frac{1}{4} \|\varepsilon^2 \lambda \partial_{x_3} p\|_2^2 \\ & \leq C \{ \varepsilon^2 \|\partial_{x_3} f\|_2^2 + \|F\|_2^2 + M(\mathbf{w}, \theta) + |\lambda|^2 \|\mathbf{w}\|_2^2 + \|\nabla \partial_{x'} \mathbf{w}\|_2^2 \}, \end{aligned} \quad (7.3)$$

where C is a positive constant depending on Pr but independent of ε .

Outline of Proof. We denote $\mathbf{w}' = {}^\top(w^1, w^2)$, $\nabla' = {}^\top(\partial_{x_1}, \partial_{x_2})$ and $\Delta' = \partial_{x_1}^2 + \partial_{x_2}^2$. Applying ∂_{x_3} to (6.2), we have

$$\varepsilon^2 \lambda \partial_{x_3} p + \nabla' \cdot \partial_{x_3} \mathbf{w}' + \partial_{x_3}^2 w^3 = \varepsilon^2 \partial_{x_3} f. \quad (7.4)$$

The third equation of (6.3) is written as

$$\text{Pr}^{-1} \lambda w^3 - \Delta' w^3 - \partial_{x_3}^2 w^3 + \partial_{x_3} p = \tilde{g}^3, \quad (7.5)$$

where

$$\tilde{g}^3 = \text{Pr}^{-1} g^3 - \text{Pr}^{-1} (\mathbf{v}_s \cdot \nabla w^3 + \mathbf{w} \cdot \nabla v_s^3) + \text{Ra} \theta.$$

We compute (7.4)+(7.5) to obtain

$$\varepsilon^2 \lambda \partial_{x_3} p + \partial_{x_3} p = H, \quad (7.6)$$

where

$$H = \varepsilon^2 \partial_{x_3} f + \tilde{g}^3 - (\text{Pr}^{-1} \lambda w^3 - \Delta' w^3 + \nabla' \cdot \partial_{x_3} \mathbf{w}').$$

Taking the L^2 inner product of (7.6) with $\partial_{x_3} p$, we have

$$\begin{aligned} & (\text{Re } \lambda + \varepsilon^2 |\text{Im } \lambda|^2) \varepsilon^2 \|\partial_{x_3} p\|_2^2 + \frac{1}{4} \|\partial_{x_3} p\|_2^2 + \frac{1}{4} \|\varepsilon^2 \lambda \partial_{x_3} p\|_2^2 \\ & \leq C \{ \varepsilon^2 \|\partial_{x_3} f\|_2^2 + \|F\|_2^2 + M(\mathbf{w}, \theta) + |\lambda|^2 \|\mathbf{w}\|_2^2 + \|\nabla \partial_{x'} \mathbf{w}\|_2^2 \}. \end{aligned}$$

This, together with the relation $\varepsilon^2 \lambda \partial_{x_3} p = -\partial_{x_3} p + H$, gives the desired estimate. $\square$

The elliptic estimates yield the following estimate for the second order derivatives of $\mathbf{w}$ and θ and the first order derivatives of p .

Proposition 7.4. *Let $u = {}^\top(p, \mathbf{w}, \theta)$ be a solution of (6.2)–(6.4) satisfying boundary conditions (3.4) and (3.5). Then*

$$\begin{aligned} & \|\partial_x^2 \mathbf{w}\|_2^2 + \|\partial_x^2 \theta\|_2^2 + \|\partial_x p\|_2^2 \\ & \leq C \{ \varepsilon^4 \|f\|_{H^1}^2 + \|F\|_2^2 + \|\varepsilon^2 \lambda p\|_{H^1}^2 + |\lambda|^2 (\|\mathbf{w}\|_2^2 + \|\theta\|_2^2) + M(\mathbf{w}, \theta) \}, \end{aligned} \quad (7.7)$$

where C is a positive constant depending on Pr but independent of ε .

Outline of Proof. We rewrite (6.1) as

$$\begin{cases} \text{div } \mathbf{w} = \varepsilon^2 f - \varepsilon^2 \lambda p, \\ -\Delta \mathbf{w} + \nabla p = \text{Pr}^{-1} \mathbf{g} - (\text{Pr}^{-1} \lambda \mathbf{w} + \text{Pr}^{-1} (\mathbf{v}_s \cdot \nabla \mathbf{w} + \mathbf{w} \cdot \nabla \mathbf{v}_s) - \text{Ra} \theta \mathbf{e}_3), \\ -\Delta \theta = h - (\lambda \theta + \mathbf{v}_s \cdot \nabla \theta + \mathbf{w} \cdot \nabla \theta_s - \text{Ra} \mathbf{w} \cdot \mathbf{e}_3). \end{cases}$$

Applying the estimates for the Stokes system and the elliptic equation (see, e.g., [5, 11]), we obtain the desired estimate. $\square$

Combining the estimates in Propositions 7.1–7.4, we have the following estimate for $\|\partial_x p\|_2$ and $\|\partial_x^2 w\|_2$.

Proposition 7.5. *Let $0 < \varepsilon \leq 1$ and let $u = {}^\top(p, \mathbf{w}, \theta)$ be a solution of (6.2)–(6.4) satisfying boundary conditions (3.4) and (3.5). Then*

$$\begin{aligned} & (\operatorname{Re} \lambda + |\operatorname{Im} \lambda|) D(\mathbf{w}, \theta) + (\operatorname{Re} \lambda + \varepsilon^2 |\operatorname{Im} \lambda|^2) (|||\partial_{x'} u|||_2^2 + \varepsilon^2 \|\partial_{x_3} p\|_2^2) \\ & + b \left(\|\varepsilon^2 \lambda \partial_x p\|_{H^1}^2 + \frac{1}{2} |\lambda|^2 |||u|||_2^2 + \|\partial_x^2 \mathbf{w}\|_2^2 + \|\partial_x^2 \theta\|_2^2 + \|\partial_x p\|_2^2 \right) \\ & \leq C(1 + \varepsilon^2) \left\{ \varepsilon^2 \|\partial_x f\|_2^2 + \left(1 + \frac{1}{\varepsilon^2}\right) (\|\operatorname{div} \mathbf{w}\|_2^2 + M(\mathbf{w}, \theta) + |||F|||_2^2) \right\} \end{aligned} \quad (7.8)$$

Proof of Proposition 7.5. Let $0 \leq \varepsilon \leq 1$. We see from (7.1) + (7.2) + $d_2 \times$ (7.3) with suitably small $d_2 > 0$ that

$$\begin{aligned} & (\operatorname{Re} \lambda + |\operatorname{Im} \lambda|) D(\mathbf{w}, \theta) + (\operatorname{Re} \lambda + \varepsilon^2 |\operatorname{Im} \lambda|^2) (|||\partial_{x'} u|||_2^2 + \varepsilon^2 \|\partial_{x_3} p\|_2^2) \\ & + b \left(D(\partial_{x'} \mathbf{w}, \partial_{x'} \theta) + \|\partial_{x_3} p\|_2^2 + \|\varepsilon^2 \lambda \partial_x p\|_2 + \frac{1}{2} |\lambda|^2 |||u|||_2^2 \right) \\ & \leq C(1 + \varepsilon^2) \left\{ \delta \varepsilon^2 \|\partial_{x'} p\|_2^2 + \delta^{-1} \varepsilon^2 \|\partial_x f\|_2^2 \right. \\ & \quad \left. + \left(1 + \frac{1}{\varepsilon^2}\right) (\|\operatorname{div} \mathbf{w}\|_2^2 + M(\mathbf{w}, \theta) + |||F|||_2^2) \right\} \end{aligned} \quad (7.9)$$

for any $\delta > 0$ with positive constants b and C independent of δ and ε .

Adding $d_3 \times$ (7.3) to (7.9) and taking $d_3 > 0$ suitably small, we have

$$\begin{aligned} & (\operatorname{Re} \lambda + |\operatorname{Im} \lambda|) D(\mathbf{w}, \theta) + (\operatorname{Re} \lambda + \varepsilon^2 |\operatorname{Im} \lambda|^2) (|||\partial_{x'} u|||_2^2 + \varepsilon^2 \|\partial_{x_3} p\|_2^2) \\ & + b \left(\|\varepsilon^2 \lambda \partial_x p\|_2^2 + \frac{1}{2} |\lambda|^2 |||u|||_2^2 + \|\partial_x^2 \mathbf{w}\|_2^2 + \|\partial_x^2 \theta\|_2^2 + \|\partial_x p\|_2^2 \right) \\ & \leq C(1 + \varepsilon^2) \left\{ \delta \varepsilon^2 \|\partial_{x'} p\|_2^2 + \delta^{-1} \varepsilon^2 \|\partial_x f\|_2^2 \right. \\ & \quad \left. + \left(1 + \frac{1}{\varepsilon^2}\right) (\|\operatorname{div} \mathbf{w}\|_2^2 + M(\mathbf{w}, \theta) + |||F|||_2^2) \right\} \end{aligned} \quad (7.10)$$

for any $\delta > 0$ with positive constants b and C independent of δ and $0 < \varepsilon \leq 1$. Taking $\delta > 0$ suitably small, we arrive at

$$\begin{aligned} & (\operatorname{Re} \lambda + |\operatorname{Im} \lambda|) D(\mathbf{w}, \theta) + (\operatorname{Re} \lambda + \varepsilon^2 |\operatorname{Im} \lambda|^2) (|||\partial_{x'} u|||_2^2 + \varepsilon^2 \|\partial_{x_3} p\|_2^2) \\ & + b \left(\|\varepsilon^2 \lambda \partial_x p\|_{H^1}^2 + \frac{1}{2} |\lambda|^2 |||u|||_2^2 + \|\partial_x^2 \mathbf{w}\|_2^2 + \|\partial_x^2 \theta\|_2^2 + \|\partial_x p\|_2^2 \right) \\ & \leq C(1 + \varepsilon^2) \left\{ \varepsilon^2 \|\partial_x f\|_2^2 + \left(1 + \frac{1}{\varepsilon^2}\right) (\|\operatorname{div} \mathbf{w}\|_2^2 + M(\mathbf{w}, \theta) + |||F|||_2^2) \right\} \end{aligned} \quad (7.11)$$

This completes the proof. $\square$

Acknowledgements. Y. Kagei was partly supported by JSPS KAKENHI Grant Number 24340028, 15K13449, 24224003, 16H03947; T. Nishida was partly supported by JSPS KAKENHI Grant Number 26400163.

References

- [1] F. H. Busse, Non-linear properties of thermal convection, Rep. Prog. Phys., **41** (1978), pp. 1929–1967.

- [2] A. Chorin, The numerical solution of the Navier-Stokes equations for an incompressible fluid, *Bull. Amer. Math. Soc.*, **73** (1967), pp. 928–931.
- [3] A. Chorin, A numerical method for solving incompressible viscous flow problems, *J. Comput. Phys.*, **2** (1967), pp. 12–26.
- [4] A. Chorin, Numerical solution of the Navier-Stokes equations, *Math. Comp.*, **22** (1968), pp. 745–762.
- [5] G. P. Galdi, *An Introduction to the Mathematical Theory of the Navier-Stokes Equations*, Vol. 1, Springer-Verlag New York (1994).
- [6] V. I. Iudovich, On the origin of convection, *PMM. J. appl. Math. Mech.*, **30** (1966), pp. 1193–1199.
- [7] D. D. Joseph, *Stability of fluid motions, I, II*, Springer-Verlag, Berlin, Heidelberg, New York (1973).
- [8] T. Kato, *Perturbation Theory for Linear Operators*, Springer-Verlag, Berlin, Heidelberg, New York (1980).
- [9] A. Matsumura and T. Nishida, Initial boundary value problems for the equations of motion of compressible viscous and heat-conductive fluids. *Comm. Math. Phys.*, **89** (1983) , pp. 445–464.
- [10] P. H. Rabinowitz, Existence and nonuniqueness of rectangular solutions of the Bénard problem, *Arch. Rational Mech. Anal.*, **29** (1968), pp. 32–57.
- [11] H. Sohr, *The Navier-Stokes equations : an elementary functional analytic approach*, Birkhäuser, Basel (2001).

List of MI Preprint Series, Kyushu University

The Global COE Program

Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p-adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials
- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials

- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings
- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric τ -functions of the q-Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications

- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain
- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection
- MI2009-26 Manabu YOSHIDA
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance

- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA
Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI
Finite element computation for scattering problems of micro-hologram using DtN map
- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO
Abstract collision systems on groups

- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA
An algebraic approach to underdetermined experiments
- MI2010-10 Kei HIROSE & Sadanori KONISHI
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE
Lagrangian approach to weakly nonlinear stability of an elliptical flow
- MI2010-21 Hiroki MASUDA
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiro FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra

- MI2010-25 Toshimitsu TAKAESU
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI
Spin-spin correlation functions of the q -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and spike detection via Gaussian basis expansions
- MI2010-32 Nobutaka NAKAZONO
Hypergeometric τ functions of the q -Painlevé systems of type $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA
CAP representations of inner forms of $Sp(4)$ with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER
Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide FUKUMOTO & Alexander B. SAMOKHIN
Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI
Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI
Anderson-Darling test and the Malliavin calculus
- MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling via Compressed Sensing

- MI2011-5 Hiroshi INOUE
Implications in Compressed Sensing and the Restricted Isometry Property
- MI2011-6 Daeju KIM & Sadanori KONISHI
Predictive information criterion for nonlinear regression model based on basis expansion methods
- MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI
Group variable selection via relevance vector machine
- MI2011-8 Jan BREZINA & Yoshiyuki KAGEI
Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow
Group variable selection via relevance vector machine
- MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC
Recursive structures in the multispecies TASEP
- MI2011-10 Kazunori YASUTAKE
On projective space bundle with nef normalized tautological line bundle
- MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA
An explicit formula for the discrete power function associated with circle patterns of Schramm type
- MI2011-12 Yoshiyuki KAGEI
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow
- MI2011-13 Vladimír CHALUPECKÝ & Adrian MUNTEAN
Semi-discrete finite difference multiscale scheme for a concrete corrosion model: approximation estimates and convergence
- MI2011-14 Jun-ichi INOBUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves
- MI2011-15 Hiroshi INOUE
A generalization of restricted isometry property and applications to compressed sensing
- MI2011-16 Yu KAWAKAMI
A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space
- MI2011-17 Naoyuki KAMIYAMA
Matroid intersection with priority constraints
- MI2012-1 Kazufumi KIMOTO & Masato WAKAYAMA
Spectrum of non-commutative harmonic oscillators and residual modular forms
- MI2012-2 Hiroki MASUDA
Mighty convergence of the Gaussian quasi-likelihood random fields for ergodic Levy driven SDE observed at high frequency

- MI2012-3 Hiroshi INOUE
A Weak RIP of theory of compressed sensing and LASSO
- MI2012-4 Yasuhide FUKUMOTO & Youich MIE
Hamiltonian bifurcation theory for a rotating flow subject to elliptic straining field
- MI2012-5 Yu KAWAKAMI
On the maximal number of exceptional values of Gauss maps for various classes of surfaces
- MI2012-6 Marcio GAMEIRO, Yasuaki HIRAOKA, Shunsuke IZUMI, Miroslav KRAMAR, Konstantin MISCHAIKOW & Vidit NANDA
Topological Measurement of Protein Compressibility via Persistence Diagrams
- MI2012-7 Nobutaka NAKAZONO & Seiji NISHIOKA
Solutions to a q -analog of Painlevé III equation of type $D_7^{(1)}$
- MI2012-8 Naoyuki KAMIYAMA
A new approach to the Pareto stable matching problem
- MI2012-9 Jan BREZINA & Yoshiyuki KAGEI
Spectral properties of the linearized compressible Navier-Stokes equation around time-periodic parallel flow
- MI2012-10 Jan BREZINA
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a time-periodic parallel flow
- MI2012-11 Daeju KIM, Shuichi KAWANO & Yoshiyuki NINOMIYA
Adaptive basis expansion via the extended fused lasso
- MI2012-12 Masato WAKAYAMA
On simplicity of the lowest eigenvalue of non-commutative harmonic oscillators
- MI2012-13 Masatoshi OKITA
On the convergence rates for the compressible Navier- Stokes equations with potential force
- MI2013-1 Abuduwaili PAERHATI & Yasuhide FUKUMOTO
A Counter-example to Thomson-Tait-Chetayev's Theorem
- MI2013-2 Yasuhide FUKUMOTO & Hirofumi SAKUMA
A unified view of topological invariants of barotropic and baroclinic fluids and their application to formal stability analysis of three-dimensional ideal gas flows
- MI2013-3 Hiroki MASUDA
Asymptotics for functionals of self-normalized residuals of discretely observed stochastic processes
- MI2013-4 Naoyuki KAMIYAMA
On Counting Output Patterns of Logic Circuits
- MI2013-5 Hiroshi INOUE
RIPless Theory for Compressed Sensing

- MI2013-6 Hiroshi INOUE
Improved bounds on Restricted isometry for compressed sensing
- MI2013-7 Hidetoshi MATSUI
Variable and boundary selection for functional data via multiclass logistic regression modeling
- MI2013-8 Hidetoshi MATSUI
Variable selection for varying coefficient models with the sparse regularization
- MI2013-9 Naoyuki KAMIYAMA
Packing Arborescences in Acyclic Temporal Networks
- MI2013-10 Masato WAKAYAMA
Equivalence between the eigenvalue problem of non-commutative harmonic oscillators and existence of holomorphic solutions of Heun's differential equations, eigenstates degeneration, and Rabi's model
- MI2013-11 Masatoshi OKITA
Optimal decay rate for strong solutions in critical spaces to the compressible Navier-Stokes equations
- MI2013-12 Shuichi KAWANO, Ibuki HOSHINA, Kazuki MATSUDA & Sadanori KONISHI
Predictive model selection criteria for Bayesian lasso
- MI2013-13 Hayato CHIBA
The First Painleve Equation on the Weighted Projective Space
- MI2013-14 Hidetoshi MATSUI
Variable selection for functional linear models with functional predictors and a functional response
- MI2013-15 Naoyuki KAMIYAMA
The Fault-Tolerant Facility Location Problem with Submodular Penalties
- MI2013-16 Hidetoshi MATSUI
Selection of classification boundaries using the logistic regression
- MI2014-1 Naoyuki KAMIYAMA
Popular Matchings under Matroid Constraints
- MI2014-2 Yasuhide FUKUMOTO & Youichi MIE
Lagrangian approach to weakly nonlinear interaction of Kelvin waves and a symmetry-breaking bifurcation of a rotating flow
- MI2014-3 Reika AOYAMA
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Parallel flow in a cylindrical domain
- MI2014-4 Naoyuki KAMIYAMA
The Popular Condensation Problem under Matroid Constraints

- MI2014-5 Yoshiyuki KAGEI & Kazuyuki TSUDA
Existence and stability of time periodic solution to the compressible Navier-Stokes equation for time periodic external force with symmetry
- MI2014-6 This paper was withdrawn by the authors.
- MI2014-7 Masatoshi OKITA
On decay estimate of strong solutions in critical spaces for the compressible Navier-Stokes equations
- MI2014-8 Rong ZOU & Yasuhide FUKUMOTO
Local stability analysis of azimuthal magnetorotational instability of ideal MHD flows
- MI2014-9 Yoshiyuki KAGEI & Naoki MAKIO
Spectral properties of the linearized semigroup of the compressible Navier-Stokes equation on a periodic layer
- MI2014-10 Kazuyuki TSUDA
On the existence and stability of time periodic solution to the compressible Navier-Stokes equation on the whole space
- MI2014-11 Yoshiyuki KAGEI & Takaaki NISHIDA
Instability of plane Poiseuille flow in viscous compressible gas
- MI2014-12 Chien-Chung HUANG, Naonori KAKIMURA & Naoyuki KAMIYAMA
Exact and approximation algorithms for weighted matroid intersection
- MI2014-13 Yusuke SHIMIZU
Moment convergence of regularized least-squares estimator for linear regression model
- MI2015-1 Hidetoshi MATSUI & Yuta UMEZU
Sparse regularization for multivariate linear models for functional data
- MI2015-2 Reika AOYAMA & Yoshiyuki KAGEI
Spectral properties of the semigroup for the linearized compressible Navier-Stokes equation around a parallel flow in a cylindrical domain
- MI2015-3 Naoyuki KAMIYAMA
Stable Matchings with Ties, Master Preference Lists, and Matroid Constraints
- MI2015-4 Reika AOYAMA & Yoshiyuki KAGEI
Large time behavior of solutions to the compressible Navier-Stokes equations around a parallel flow in a cylindrical domain
- MI2015-5 Kazuyuki TSUDA
Existence and stability of time periodic solution to the compressible Navier-Stokes-Korteweg system on R^3
- MI2015-6 Naoyuki KAMIYAMA
Popular Matchings with Ties and Matroid Constraints

- MI2015-7 Shoichi EGUCHI & Hiroki MASUDA
Quasi-Bayesian model comparison for LAQ models
- MI2015-8 Yoshiyuki KAGEI & Ryouta OOMACHI
Stability of time periodic solution of the Navier-Stokes equation on the half-space under oscillatory moving boundary condition
- MI2016-1 Momonari KUDO
Analysis of an algorithm to compute the cohomology groups of coherent sheaves and its applications
- MI2016-2 Yoshiyuki KAGEI & Masatoshi OKITA
Asymptotic profiles for the compressible Navier-Stokes equations on the whole space
- MI2016-3 Shota ENOMOTO & Yoshiyuki KAGEI
Asymptotic behavior of the linearized semigroup at space-periodic stationary solution of the compressible Navier-Stokes equation
- MI2016-4 Hiroki MASUDA
Non-Gaussian quasi-likelihood estimation of locally stable SDE
- MI2016-5 Yoshiyuki KAGEI & Takaaki NISHIDA
On Chorin's method for stationary solutions of the Oberbeck-Boussinesq equation