Self-adjoint extensions of momentum operators: application of weak Weyl relations

Hiroshima, Fumio Faculty of Mathematics, Kyushu University

Kuribayashi, Sotaro Graduate School of Mathematics, Kyushu University

Sasaki, Itaru International Young Researchers Empowerment Center, Shinshu University

https://hdl.handle.net/2324/17009

出版情報:Journal of Math-for-Industry (JMI). 2 (A), pp.21-25, 2010-04-08. Faculty of Mathematics, Kyushu University バージョン: 権利関係:



Self-adjoint extensions of momentum operators: application of weak Weyl relations

Fumio Hiroshima, Sotaro Kuribayashi and Itaru Sasaki

Received on January 12, 2010

Abstract. By weak Weyl relations it is shown that momentum operators, $-i\partial_{x_j}$, defined on $C_0^{\infty}(\Omega)$ with some general open set $\Omega \subset \mathbb{R}^n$ are *not* essentially self-adjoint but have uncountably many self-adjoint extensions.

Keywords. canonical commutation relation, CCR, Weyl relation, weak Weyl relation, momentum operator

1. Preliminaries

In this paper we are concerned with uncountably many selfadjoint extensions of momentum operators defined on open sets in \mathbb{R}^n . Let us begin with considering a one dimensional momentum operator

$$p = -i\frac{d}{dx}.$$

It is well known that p is essentially self-adjoint on $C_0^{\infty}((-\infty,\infty))$, namely it has the unique self-adjoint extension. We can see however that p has no self-adjoint extension on $C_0^{\infty}((0,\infty))$ but has uncountably many self-adjoint extensions on $C_0^{\infty}((0,1))$. When the dimension $n \geq 2$, however, it is not trivial to see the essential or non-essential self-adjointness of momentum operators defined on $C_0^{\infty}(\Omega)$ with open set Ω . We are then interested in self-adjoint extensions of momentum operator $-i\partial_{x_j}$ on $L^2(\Omega)$ with a general open set $\Omega \subset \mathbb{R}^n$. In this paper we show a sufficient condition such that $-i\partial_{x_j}$ is not essentially self-adjoint but has uncountably many self-adjoint extensions by using the so-called weak Weyl relations.

Let \mathscr{H} be a *separable* Hilbert space. Let H be a selfadjoint operator and T a symmetric operator on \mathscr{H} . We say that $\{T, H\}$ satisfies weak Weyl relation if

(1)
$$e^{-itH} \operatorname{Dom}(T) \subset \operatorname{Dom}(T)$$

and

(2)
$$Te^{-itH}\psi = e^{-itH}(T+t)\psi$$

holds for all $\psi \in \text{Dom}(T)$ and $t \in \mathbb{R}$, where Dom(A)denotes the domain of operator A. A symmetric operator T satisfying (2) is called a time operator associated with H, which is, to our best knowledge, introduced by [Sch83A, Sch83B] and drastically investigated by [Ara05]. See also [Ara99-a, Ara99-b]. If T is self-adjoint instead of symmetric, and $\{T, H\}$ satisfies the Weyl relation

(3)
$$e^{-isT}e^{-itH} = e^{-ist}e^{-itH}e^{-isT},$$

then weak Weyl relation (2) can be derived from (3), but (2) does not necessarily imply (3) in general. The important facts on weak Weyl relations and Weyl relation are (1) and (2) below:

- (1) If T is a self-adjoint operator, then Weyl relation (3) can be derived from weak Weyl relation (2);
- (2) If {T, H} satisfies Weyl relation (3), then there exist Hilbert spaces *H_m* and unitary operators

$$U_m: \mathscr{H}_m \to L^2(\mathbb{R}), \quad 1 \le m \le M,$$

such that

- (1) $\mathscr{H} = \bigoplus_{m=1}^{M} \mathscr{H}_m,$
- (2) T and H are reduced to \mathscr{H}_m ,

(3)
$$UT|_{\mathscr{H}_m}U^{-1} = x$$
 and $UH|_{\mathscr{H}_m}U^{-1} = -i\frac{a}{dx}$

(2) is known as von Neumann's uniqueness theorem.

The key idea in this paper is that the momentum operator $-i\partial_{x_j}$ can be regarded as the time operator associated with multiplication operator $-x_j$. If $-i\partial_{x_j}$ is essentially self-adjoint on $C_0^{\infty}(\Omega)$ with an open set $\Omega \subset \mathbb{R}^n$, then the pair of self-adjoint operators

(4)
$$\left\{\overline{-i\partial_{x_j}\lceil_{C_0^{\infty}(\Omega)}}, -x_j\right\}$$

also satisfies Weyl relation. If the multiplication operator x_j on $L^2(\Omega)$ is bounded, then it contradicts von Neumann's uniqueness theorem. Thus we can conclude that $-i\partial_{x_j} \left[C_{0}^{\infty}(\Omega) \right]$ is not essentially self-adjoint if Ω is bounded. This kind of results may be already known, but our proof is new and simple. In this paper we discuss more general open sets Ω , which include unbounded sets.

2. Time operators and main results

In this section we introduce weak Weyl relations and time operators, and prove the main theorem.

Definition 1. (1) Let T_1, \dots, T_n and H_1, \dots, H_n be selfadjoint. We say that $\{T_j, H_j\}_{j=1}^n$ is Weyl relation if and only if

$$e^{-itT_j}e^{-isH_k} = e^{-i\delta_{jk}st}e^{-isH_k}e^{-itT_j}$$
$$e^{-itT_j}e^{-isT_k} = e^{-isT_k}e^{-itT_j}$$
$$e^{-itH_j}e^{-isH_k} = e^{-isH_k}e^{-itH_j}$$

hold for all $s, t \in \mathbb{R}$ and $j, k = 1, \cdots, n$.

(2) Let T_1, \dots, T_n be symmetric and H_1, \dots, H_n self-adjoint. We say that $\{T_j, H_j\}_{j=1}^n$ is weak Weyl relation if and only if

(5)
$$e^{-itH_k} \operatorname{Dom}(T_i) \subset \operatorname{Dom}(T_i)$$

and

(6)
$$T_j e^{-itH_k} \psi = e^{-itH_k} (T_j + \delta_{jk}t) \psi, \ \psi \in \text{Dom}(T_j)$$

hold for all $t \in \mathbb{R}$ and $j, k = 1, \cdots, n$.

We show some properties of Weyl relation and weak Weyl relation. Let \overline{T} denote the closure of T. Let $\{T_j, H_j\}_{j=1}^n$ be weak Weyl relation, then also is $\{\overline{T}_j, H_j\}_{j=1}^n$. As is easily seen that weak Weyl relation can be derived from Weyl relation. The converse is, however, not true. If $\{T, H\}$ is a weak Weyl relation, then the spectrum of H is purely absolutely continuous. In particular H has no eigenvalues. See [AM08-a, Gal02, Miy01]. Furthermore we can construct a time operator $T_{g(H)}$ associated with g(H), where g is some Borel measurable function, as

$$T_{g(H)} = \frac{1}{2} (\dot{g}(H)^{-1}T + T\dot{g}(H)^{-1}),$$

where $\dot{g} = dg/dx$. See [AM08-b, HKM09] for details.

An important relationship between weak Weyl relation and Weyl relation is as follows.

Proposition 1. Let $\{T_j, H_j\}_{j=1}^n$ be weak Weyl relation and T_1, \dots, T_n self-adjoint. Then $\{T_j, H_j\}_{j=1}^n$ is Weyl relation.

Proof. See [Ara05, Proposition 2.10].

Let us define operators $\hat{x}_1, \dots, \hat{x}_n$ and $\hat{p}_1, \dots, \hat{p}_n$ in $L^2(\mathbb{R}^n)$. Operator \hat{x}_j is the multiplication operator by x_j with the domain

$$\mathrm{Dom}(\hat{x}_j) = \left\{ f \in L^2(\mathbb{R}^n) \left| \int_{\mathbb{R}^n} x_j^2 |f(x)|^2 dx < \infty \right\},\right.$$

and

$$\hat{p}_j = -i\partial_{x_j}$$

with

$$\operatorname{Dom}(\hat{p}_j) = H^1(\mathbb{R}^n).$$

It is a fundamental fact that $\{\hat{x}_j, \hat{p}_j\}_{j=1}^n$ or $\{\hat{p}_j, -\hat{x}_j\}_{j=1}^n$ are Weyl relations. **Proposition 2.** (von Neumann's uniqueness theorem [Neu31]) Let $\{T_j, H_j\}_{j=1}^n$ be Weyl relation. Then there exist Hilbert spaces \mathscr{H}_m and unitary operators

$$U_m: \mathscr{H}_m \to L^2(\mathbb{R}^n), \quad 1 \le m \le M,$$

such that

- (1) $\mathscr{H} = \oplus_{m=1}^{M} \mathscr{H}_{m},$
- (2) for each j = 1, ..., n, T_j and H_j are reduced to \mathscr{H}_m ,
- (3) $U_m T_j \lceil_{\mathscr{H}_m} U_m^{-1} = \hat{x}_j \text{ and } U_m H_j \lceil_{\mathscr{H}_m} U_m^{-1} = \hat{p}_j.$

In particular $\operatorname{Spec}(T_j) = \operatorname{Spec}(H_j) = \mathbb{R}$.

The next lemma is a criterion to show non-essential selfadjointness of a time operator.

Lemma 1. Let $\{T_j, H_j\}_{j=1}^n$ be weak Weyl relation with $\operatorname{Spec}(H_j) \neq \mathbb{R}$ for some $j = 1, \dots, n$. Then T_j is not essentially self-adjoint.

Proof. Suppose that T_j is essentially self-adjoint. Then $\{\overline{T}_j, H_j\}$ is Weyl relation by Proposition 1. However this contradicts Proposition 2 since $\operatorname{Spec}(H_j) \neq \mathbb{R}$. Therefore T_j is not essentially self-adjoint.

Let Ω be an open subset in \mathbb{R}^n . Let x_j be the multiplication operator by x_j with

$$\operatorname{Dom}(x_j) = \left\{ f \in L^2(\Omega) \left| \int_{\Omega} x_j^2 |f(x)|^2 dx < \infty \right\},\right.$$

and $p_j = -i\partial_{x_j}$ with

$$\operatorname{Dom}(p_j) = C_0^{\infty}(\Omega)$$

Let $f, g \in C_0^{\infty}(\Omega)$. Since we know that

$$(f, p_j g)_{L^2(\Omega)} = (f, p_j g)_{L^2(\mathbb{R}^n)} = (p_j f, g)_{L^2(\mathbb{R}^n)} = (p_j f, g)_{L^2(\Omega)},$$

 p_j is symmetric and then closable.

Lemma 2. Let Ω be an open subset in \mathbb{R}^n . Then $\{\overline{p}_j, -x_j\}_{j=1}^n$ is weak Weyl relation on $L^2(\Omega)$.

Proof. Since e^{itx_j} leaves $C_0^{\infty}(\Omega)$ invariant and

$$p_k e^{itx_j} f = e^{itx_j} (p_k + \delta_{jk} t) f$$

for $f \in C_0^{\infty}(\Omega)$ by a direct calculation, $\{p_j, -x_j\}_{j=1}^n$ is weak Weyl relation on $L^2(\Omega)$. Hence $\{\overline{p}_j, -x_j\}_{j=1}^n$ is also weak Weyl relation.

We define the class \mathscr{O}_j of open subsets in \mathbb{R}^n . Let π_j : $\mathbb{R}^n \to \mathbb{R}$ be the projection defined by $\pi_j(x) = x_j$. We define \mathscr{O}_j by

$$\mathscr{O}_j = \left\{ \Omega \subset \mathbb{R}^n \, \middle| \, \Omega \text{ is an open subset and } \overline{\pi_j(\Omega)} \neq \mathbb{R} \right\}$$

for j = 1, ..., n.

Theorem 1. Let $\Omega \in \mathcal{O}_j$. Then p_j is not essentially selfadjoint on $C_0^{\infty}(\Omega)$. Proof. Note that $\operatorname{Spec}(x_j) = \overline{\pi_j(\Omega)} \neq \mathbb{R}$ for $\Omega \in \mathcal{O}_j$. Assume that p_j is essentially self-adjoint. Then $\{\overline{p}_j, -x_j\}$ is weak Weyl relation by Lemma 2. In particular, $\{\overline{p}_j, -x_j\}$ is also Weyl relation. It however contradicts Lemma 1 since $\operatorname{Spec}(x_j) \neq \mathbb{R}$. Hence p_j is not essentially self-adjoint. \Box Let \mathcal{O} be the set of open subsets Ω in \mathbb{R}^n such that $\overline{\Omega} \neq \mathbb{R}^n$.

Theorem 2. Let $\Omega \in \mathcal{O}$. Then at least one momentum operator is not essentially self-adjoint on $C_0^{\infty}(\Omega)$.

Proof. Assume that all the momentum operators p_j are essentially self-adjoint. Then $\{\overline{p}_j, -x_j\}_{j=1}^n$ is Weyl relation. Thus the joint distribution of $\{-x_1, ..., -x_n\}$ has to be \mathbb{R}^n by Theorem 1, but is indeed $\overline{\Omega}$. Then at least one momentum operator is not essentially self-adjoint. \Box The following corollary is immediate.

The following corollary is inillegrate.

Corollary 1. Let $\Omega \in \mathcal{O}$ be symmetric, i.e., $\{(x_{\pi(1)}, \dots, x_{\pi(n)}) | (x_1, \dots, x_n) \in \Omega\} \subset \Omega$ for all n-degree permutations π . Then, for each $j = 1, \dots, n$, p_j is not essentially self-adjoint on $C_0^{\infty}(\Omega)$.

Proof. By Theorem 2 at least one momentum operator is not essentially self-adjoint. Since all the momentum operators are unitarily equivalent by the symmetry, each p_j is not essentially self-adjoint on $C_0^{\infty}(\Omega)$.

3. Generalizations

We can extend Theorems 1 and 2 mentioned in the previous section. We say that an open set Ω is simple if and only if there is no open set $K \subset \mathbb{R}^{n-1}$ such that $\Omega = \mathbb{R} \times K$.

Theorem 3. Let Ω be simple. Then each momentum operator p_j , j = 1, ..., n, is not essentially self-adjoint on $C_0^{\infty}(\Omega)$.



Figure 1:

Proof. Let $1 \leq j \leq n$ be fixed. Then there exists a parallel transformation T on \mathbb{R}^n such that

(7)
$$O = (0, \cdots, 0) \notin T\Omega$$

and

(8)
$$X_j = (0, ..., t, ...0) \in T\Omega$$

with some t. Without loss of generality we may reset $T\Omega$ as Ω . The distance between Ω and the origin O is denoted by $\delta > 0$. Let $\delta > \epsilon > 0$ be such that

9)
$$B_{\epsilon}(X_j) \subset \Omega,$$

where $B_{\epsilon}(X_j)$ denotes the open ball centered at X_j with radius ϵ . (9) can be possible by (8) for sufficiently small ϵ .

Now we define the self-adjoint operator
$$D$$
 on $L^{2}(\Omega)$ by

(10)
$$D = \sum_{j=1}^{n} x_j^2.$$

Thus it follows that



Figure 2:

$$(f, Df)_{L^2(\Omega)} \ge \delta^2 \|f\|_{L^2(\Omega)}^2$$

for all $f \in C_0^{\infty}(\Omega)$. Let $\phi \in C_0^{\infty}(\mathbb{R}^n)$ be such that $\operatorname{supp} \phi \subset B_{\epsilon}(0)$. See Fig. 1. With $t \in \mathbb{R}$, the parallel transform of ϕ in the X_i -direction is denoted by ϕ_t ;

$$\phi_t(x) = \phi((x_1, \dots, x_j - t, \dots, x_n)), \quad x \in \Omega,$$

we see that $\phi_t \in L^2(\Omega)$ by (9). Suppose that p_j is essentially self-adjoint on $C_0^{\infty}(\Omega)$ and we denote the closure of $p_j \lceil_{C_0^{\infty}(\Omega)}$ by P_j . Thus it follows that

11)
$$(e^{isP_j}\phi_t, De^{isP_j}\phi_t)_{L^2(\Omega)} \ge \delta^2 \|\phi\|_{L^2(\mathbb{R}^n)}.$$

Now we compute the left hand side of (11). Since P_j is self-adjoint, $\{P_j, -x_j\}$ is not only weak Weyl relation but also Weyl relation. Then

$$e^{-isP_j}e^{itx_j} = e^{-ist}e^{itx_j}e^{-isP_j}.$$

From this the weak Weyl relation

$$x_j e^{-isP_j} = e^{-isP_j} (x_j + s)$$

(12)
$$x_j^2 e^{-isP_j} = e^{-isP_j} (x_j + s)^2$$

on $C_0^{\infty}(\Omega)$. Inserting -s into s in (12), we have

(13)
$$x_j^2 e^{isP_j} = e^{isP_j} (x_j - s)^2$$

and

$$e^{-isP_j}De^{isP_j} = x_1^2 + \dots + (x_j - s)^2 + \dots + x_n^2$$

on $C_0^{\infty}(\Omega)$. Inserting this into the left-hand side of (11) and setting t = s, we have

LHS (11) =
$$\int_{B_{\epsilon}(0)} |\phi(x)|^2 \left(\sum_{j=1}^n x_j^2\right) dx \le \epsilon^2 \|\phi\|_{L^2(\mathbb{R}^n)}^2.$$

Thus $\epsilon^2 \|\phi\|_{L^2(\mathbb{R}^n)}^2 \geq \delta^2 \|\phi\|_{L^2(\mathbb{R}^n)}^2$. See Fig. 2. This contradicts that $\epsilon < \delta$ and then p_j is not essentially self-adjoint.

Theorem 4. Let $\Omega = \mathbb{R}^m \times K \in \mathcal{O}$, where K is simple. Then $p_1, ..., p_m$ are essentially self-adjoint on $C_0^{\infty}(\Omega)$ but $p_{m+1}, ..., p_n$ not essentially self-adjoint.

Proof. Under the identification $L^2(\Omega) = L^2(\mathbb{R}^m) \otimes L^2(K)$, $p_j = p_j \otimes 1, \ j = 1, ..., m$, are essentially self-adjoint on $C_0^{\infty}(\Omega)$. While $p_j, \ m+1 \leq j \leq n$, are not essentially self adjoint by Theorem 3.

Now we investigate the existence of self-adjoint extensions of momentum operators defined in $L^2(\Omega)$. Let R_j : $\mathbb{R}^n \to \mathbb{R}^n$ be the reflection defined by replacing the *j*th coordinate x_j with $-x_j$. Let $\mathscr{O}_{\text{sym},j}$ be defined by

$$\mathscr{O}_{\mathrm{sym},j} = \{ \Omega \in \mathscr{O} | R_j \Omega = \Omega \}, \quad j = 1, ..., n.$$

Corollary 2. Let $\Omega \in \mathcal{O}_{\text{sym},j}$ and suppose that p_j is not essentially self-adjoint. Then uncountably many self-adjoint extensions of $p_j [_{C_0^{\infty}(\Omega)}$ exist.

Proof. Let $P_j = \overline{p_j |_{C_0^{\infty}(\Omega)}}$. Let $C_j : L^2(Q) \to L^2(Q)$ be the antilinear map defined by

(14)
$$(C_j f)(x) = f(R_j x),$$

where \overline{f} denotes the complex conjugate of f. We can see that $C_j : C_0^{\infty}(\Omega) \to C_0^{\infty}(\Omega)$ and $C_j p_j = p_j C_j$ on $C_0^{\infty}(\Omega)$, then a limiting argument yields that $C_j \text{Dom}(P_j) \subset$ $\text{Dom}(P_j)$ and $C_j P_j = P_j C_j$ on $\text{Dom}(P_j)$. From this it follows that P_j has equal deficiency indices by von Neumann's theorem [RS2, Theorem X.3]. Since P_j is not self-adjoint, the deficiency indices of P_j is (m, m) with $m \geq 1$. Hence P_j has uncountably many self-adjoint extensions. \Box

- **Example 1. (1)** Let $\Omega = \mathbb{R}^n \setminus \overline{B_r(0)}$. Then each p_j , j = 1, ..., n, is not essentially self-adjoint and has uncountably many self-adjoint extensions, since $\Omega \in \bigcap_{j=1}^n \mathscr{O}_{\text{sym},j}$. Refer to see [Hir00].
- (2) Let $\Omega = B_r(0)$. Then each p_j , j = 1, ..., n, is not essentially self-adjoint and has uncountably many self-adjoint extensions, since $\Omega \in \bigcap_{j=1}^n \mathscr{O}_{\text{sym},j}$.

(3) Let

$$\Omega_{+} = \{ (x, y) \in \mathbb{R}^{2} | -1 < xy < 1 \}, \Omega_{-} = \{ (x, y) \in \mathbb{R}^{2} | 1 < xy, xy < -1 \}.$$

Then p_j , j = 1, 2, are not essentially self-adjoint on $C_0^{\infty}(\Omega_{\pm})$ but have uncountably many self-adjoint extensions, since $\Omega_{\pm} \in \bigcap_{j=1,2} \mathscr{O}_{\text{sym},j}$.

Acknowledgments

We thank K. Schmüdgen for helpful advices and we also thank unknown referee for a lot of useful comments. FH acknowledges support of Grant-in-Aid for Science Research (B) 20340032 from JSPS.

References

- [Ara99-a] A. Arai, Mathematical structures of quantum mechanics I, Asakurashoten 1999.
- [Ara99-b] A. Arai, Mathematical structures of quantum mechanics II, Asakurashoten 1999.
- [Ara05] A. Arai, Generalized weak Weyl relation and decay of quantum dynamics, *Rev. Math. Phys.* 17 (2005), 1071–1109.
- [AM08-a] A. Arai and Y. Matsuzawa, Time operators of a Hamiltonian with purely discrete spectrum, *Rev. Math. Phys* **20** (2008), 951-978.
- [AM08-b] A. Arai and Y. Matsuzawa, Construction of a Weyl representation from a weak Weyl representation of the canonical commutation relation, *Lett. Math. Phys.* 83 (2008), 201-211.
- [Gal02] E. A. Galapon, Self-adjoint time operator is the rule for discrete semi-bounded Hamiltonians, Proc. R. Soc. Lond. A 458 (2002), 2671–2689.
- [Hir00] M. Hirokawa, Canonical quantization on a doubly connected space and the Aharonov-Bohm phase, J. Funct. Anal. 174 (2000), 322–363.
- [HKM09] F. Hiroshima, S. Kuribayashi, Y. Matsuzawa, Strong time operators associated with generalized Hamiltonian, *Lett. Math. Phys.* 87 (2009),115-123.
- [Miy01] M. Miyamoto, A generalised Weyl relation approach to the time operator and its connection to the survival probability, J. Math. Phys. 42 (2001), 1038–1052.
- [Neu31] J. von Neumann, Die Eindeutigkeit der Schrödingerschen Operatoren, Math. Ann. 104 (1931), 570-578.
- [RS2] M. Reed and B. Simon, Methods of Modern Mathematical Physics II, Academic Press, 1975.

- [Sch83A] K. Schmüdgen, On the Heisenberg Commutation Relation I, J. Func. Anal. 50 (1983), 8–49.
- [Sch83B] K. Schmüdgen, On the Heisenberg Commutation Relation II, Pub. RIMS Kyoto University 19 (1983), 601–671.

Fumio Hiroshima
Faculty of Mathematics, Kyushu University
819-0395, Fukuoka, Japan.
E-mail: hiroshima(at)math.kyushu-u.ac.jp
Sotaro Kuribayashi
Graduate School of Mathematics, Kyushu University
819-0395, Fukuoka, Japan.
E-mail: s-kuribayasi(at)math.kyushu-u.ac.jp
Itaru Sasaki
International Young Researchers Empowerment Center, Shinshu University,
390-8621, Matsumoto, Japan
E-mail: itasasa(at)gmail.com