

Composition, union and division of cellular automata on groups

Ito, Takahiro
九州大学大学院数理学府

Fujio, Mitsuhiro
Department of Systems Design and Informatics, Kyushu Institute of Technology

Inokuchi, Shuichi
Faculty of Mathematics, Kyushu University

Mizoguchi, Yoshihiro
Faculty of Mathematics, Kyushu University

<https://hdl.handle.net/2324/17105>

出版情報 : MI Preprint Series. 2010-23, 2010-05-19. Faculty of Mathematics, Kyushu University
バージョン :
権利関係 :



MI Preprint Series

Kyushu University
The Global COE Program
Math-for-Industry Education & Research Hub

Composition, union and division of cellular automata on groups

Takahiro Ito, Mitsuhiko Fujio,
Shuichi Inokuchi &
Yoshihiro Mizoguchi

MI 2010-23

(Received May 19, 2010)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

Composition, Union and Division of Cellular Automata on Groups

Takahiro Ito¹ and Mitsuhiro Fujio² and Shuichi Inokuchi³ and Yoshihiro Mizoguchi³

¹*Graduate School of Mathematics, Kyushu University, JAPAN*

²*Department of Systems Design and Informatics, Kyushu Institute of Technology, JAPAN*

³*Faculty of Mathematics, Kyushu University, JAPAN*

received 19th May 2010,

We introduce the notion of 'Composition', 'Union' and 'Division' of cellular automata on groups. A kind of notions of compositions was investigated by Sato (1994) and Manzini (1998) for linear cellular automata, we extend the notion to general cellular automata on groups and investigated their properties. We observe the all unions and compositions generated by one-dimensional 2-neighborhood cellular automata over \mathbf{Z}_2 including non-linear cellular automata. Next we prove that the composition is right-distributive over union, but is not left-distributive. Finally, we conclude by showing reformulation of our definition of cellular automata on group which admit more than three states. We also show our formulation contains the representation using formal power series for linear cellular automata in Manzini (1998).

Keywords: Cellular automata, Groups, Models of computation, Automata

1 Introduction

The study of cellular automata was initiated by von Neumann (1983) and have been developed by many researchers as a good computational model for physical systems simulation. Recently cellular automata have been investigated in various fields including computer science, biology, physics, since they provide simple and powerful models for parallel computation and natural phenomena.

In this paper, we investigate cellular automata on groups as a formal model of computation. To compose simple cellular automata into a complex cellular automaton, we introduce the notion of 'Composition' of cellular automata on groups. The notion of automata on groups was first treated as a special case for automata on graphs (Caley graphs) which represent groups in Róka (1994); Rémila (1998). Watanabe and Noguchi (1982) investigated the decomposition of finite automata from the view point of spatial networks using groups. Pries et al. (1986) investigated cellular automata as a tool for implementing hardware algorithms in VLSI. They considered configurations decided by a cellular automaton as a group and divided configurations into simple configurations using group properties. Sato (1994) introduced group structured linear cellular automata and the star operation of local transition rules. The star operation is a

kind of composition of cellular automata but the definition of it is different from ours. Manzini (1998) also investigated the linear cellular automata using the formal power series and their product to find inverse local transition functions. The product of formal power series are equal to our composition of cellular automata for linear cases. An abstract collision system in Ito et al. (2008) is considered as an extension of a cellular automaton, the notion of 'composition' for an abstract collision system on G -sets is investigated in Ito (2010).

This paper follow on from Fujio (2008). He introduced the composition of cellular automata on groups in order to reduce a complex behaved dynamics into simpler ones. We introduce a formal definition of cellular automata on group over \mathbf{Z}_2 . In our framework, operations on cellular automata 'Union', 'Division' and 'Composition' are introduced. Unions of all 2-neighborhood cellular automata are investigated. Compositions of all 2-neighborhood cellular automata are also investigated and determined the subset of 3-neighborhood cellular automata which are generated by composing two 2-neighborhood cellular automata. Next we prove that the composition is right-distributive over union, but is not left-distributive. Finally, we conclude by showing reformulation of our definition of cellular automata on group which admit more than three states. We also show our formulation contains the representation using formal power series for linear cellular automata in Manzini (1998).

2 Cellular Automata on Groups

Definition 1 Let G be a group. A cellular automaton on C is a triple $C = (G, V, V')$ of a group G , subsets $V \subset G$ and $V' \subset 2^V$. For V' , we define functions $l_{V'} : 2^V \rightarrow \{\phi, \{e\}\}$ by

$$l_{V'}(X) = \begin{cases} \phi & (X \notin V') \\ \{e\} & (X \in V'), \end{cases}$$

and $F_C : 2^G \rightarrow 2^G$ by $F_C(\mathbf{c}) = \bigcup_{g \in G} gl_{V'}(g^{-1}\mathbf{c} \cap V)$. We call the map l_V a local transition function and F_C a global transition function.

Proposition 2 Let $C_1 = (G, V_1, V'_1)$ and $C_2 = (G, V_2, V'_2)$ be cellular automata. If

$$e \in F_{C_1}(\mathbf{c}) \Leftrightarrow e \in F_{C_2}(\mathbf{c}) \text{ (for any } \mathbf{c} \in 2^G)$$

then $F_{C_1} = F_{C_2}$

Proof. Since $F_{C_1}(\mathbf{c}) = \{g \in G \mid l_{V'_1}(g^{-1}\mathbf{c} \cap V_1) = \{e\}\} = \{g \in G \mid g^{-1}\mathbf{c} \cap V_1 \in V'_1\}$, we have $g \in F_{C_1}(\mathbf{c}) \Leftrightarrow g^{-1}\mathbf{c} \cap V_1 \in V'_1 \Leftrightarrow e \in F_{C_1}(g^{-1}\mathbf{c}) \Leftrightarrow e \in F_{C_2}(g^{-1}\mathbf{c}) \Leftrightarrow g^{-1}\mathbf{c} \cap V_2 \in V'_2 \Leftrightarrow g \in F_{C_2}(\mathbf{c})$. \square

In the followings, we consider the set of all integers \mathbf{Z} as an additive group $\mathbf{Z} = (\mathbf{Z}, +, 0)$. So usual one dimensional cellular automata with 2-states are represented as cellular automata on the group \mathbf{Z} . We define 2-neighborhood and 3-neighborhood 2-states cellular automata in the next definition.

Definition 3 For $k \geq 1$ and $n \in \{0, 1, \dots, 2^{2^k} - 1\}$, we define cellular automata $CA(k, n)$ on \mathbf{Z} by $CA(k, n) = (\mathbf{Z}, V, V'_n)$ where $V = \{0, 1, \dots, k-1\}$, and V'_n is the subset of 2^V which satisfies $n = \sum_{X \in V'_n} 2^{\sum_{i \in X} 2^i}$.

We note $CA(1, 0) = (\mathbf{Z}, \{0\}, \phi)$ and $CA(1, 1) = (\mathbf{Z}, \{0\}, \{0\})$.

Example 4 Since $6 = 2 + 2^2 = 2^{2^0} + 2^{2^1}$, we have $CA(2, 6) = (\mathbf{Z}, \{0, 1\}, \{\{0\}, \{1\}\})$. Since $90 = 2 + 2^3 + 2^4 + 2^6 = 2^{2^0} + 2^{2^0+2^1} + 2^{2^2} + 2^{2^1+2^2}$, we have $CA(3, 90) = (\mathbf{Z}, \{0, 1, 2\}, \{\{0\}, \{2\}, \{0, 1\}, \{1, 2\}\})$. The elements X in V'_n represents the state of neighborhood which induce the next states '1'. For a rule number 90, we have the following table:

Neighborhood	111	110	101	100	011	010	001	000
$X \in V'_n$	$\{0, 1, 2\}$	$\{1, 2\}$	$\{0, 2\}$	$\{2\}$	$\{0, 1\}$	$\{1\}$	$\{0\}$	ϕ
$l'_V(X)$	ϕ	$\{e\}$	ϕ	$\{e\}$	$\{e\}$	ϕ	$\{e\}$	ϕ

The configuration $\mathbf{c} \in \mathbf{Z}$ represents places where the state is 1. Since $n \in F_C(\mathbf{c}) \Leftrightarrow l_{V'}(n^{-1}\mathbf{c} \cap V) = \{e\} \Leftrightarrow n^{-1}\mathbf{c} \cap V \in V' \Leftrightarrow \mathbf{c} \cap nV \in nV'$, the next state at n is 1 if $\mathbf{c} \cap nV \in nV'$. For 3-neighborhood case we are choosing $V = \{0, 1, 2\}$, the left-hand side of the state is changing. It seems to be better that we choose $V = \{-1, 0, 1\}$ but it is not convenient for even-neighborhood case. Our numbered 3-neighborhood cellular automata $CA(3, n)$ is a shifted version of usual numbered elementary cellular automata. Later, we define a cellular automaton SHIFT which represent a shift operation and a operator 'composition' (\diamond) of two cellular automata. After that the usual numbered elementary cellular automata is represented as $\text{SHIFT} \diamond CA(3, n)$.

Example 5 $\text{SHIFT} = (\mathbf{Z}, \{-1, 0\}, \{\{-1\}, \{-1, 0\}\})$ is a cellular automata on group \mathbf{Z} .

\mathbf{Z}^2 is also considered as a group, so it is easy to represent a multi-dimensional cellular automata such as The Game of Life (Berlekamp et al. (1982)) as a cellular automata on a group.

Example 6 $\text{LIFE} = (\mathbf{Z}^2, V_{\text{LIFE}}, V'_{\text{LIFE}})$ is a cellular automata on group \mathbf{Z}^2 , where

$$V_{\text{LIFE}} = \left\{ \begin{pmatrix} -1 \\ -1 \end{pmatrix}, \begin{pmatrix} 0 \\ -1 \end{pmatrix}, \begin{pmatrix} +1 \\ -1 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} +1 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ +1 \end{pmatrix}, \begin{pmatrix} 0 \\ +1 \end{pmatrix}, \begin{pmatrix} +1 \\ +1 \end{pmatrix} \right\}, \text{ and}$$

$$V'_{\text{LIFE}} = \{v \in 2^V \mid (\#v = 3) \vee (\#v = 4 \wedge \begin{pmatrix} 0 \\ 0 \end{pmatrix} \in v)\}.$$

We note that $\#v$ is the number of elements in a set v .

One dimensional cellular automaton on \mathbf{Z} is embedded into the two dimensional cellular automaton on \mathbf{Z}^2 . We define two natural embeddings EX and EY in the following.

Definition 7 For a cellular automata $C = (G, V, V')$, we define a cellular automata $EX(C)$ on \mathbf{Z}^2 by $EX(C) = (\mathbf{Z}^2, V_{EX(C)}, V'_{EX(C)})$ where

$$V_{EX(C)} = \left\{ \begin{pmatrix} x \\ 0 \end{pmatrix} \mid x \in V \right\}, \text{ and}$$

$$V'_{EX(C)} = \left\{ \left\{ \begin{pmatrix} x \\ 0 \end{pmatrix} \mid x \in X \right\} \mid X \in V' \right\}.$$

We also define a cellular automata $EY(C)$ on \mathbf{Z}^2 by $EY(C) = (\mathbf{Z}^2, V_{EY(C)}, V'_{EY(C)})$ where

$$V_{EY(C)} = \left\{ \begin{pmatrix} 0 \\ x \end{pmatrix} \mid x \in V \right\}, \text{ and}$$

$$V'_{EY(C)} = \left\{ \left\{ \begin{pmatrix} 0 \\ x \end{pmatrix} \mid x \in X \right\} \mid X \in V' \right\}.$$

$n \setminus m$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
0	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	1	1	3	3	5	5	7	7	9	9	11	11	13	13	15	15
2	2	3	2	3	6	7	6	7	10	11	10	11	14	15	14	15
3	3	3	3	3	7	7	7	7	11	11	11	11	15	15	15	15
4	4	5	6	7	4	5	6	7	12	13	14	15	12	13	14	15
5	5	5	7	7	5	5	7	7	13	13	15	15	13	13	15	15
6	6	7	6	7	6	7	6	7	14	15	14	15	14	15	14	15
7	7	7	7	7	7	7	7	7	15	15	15	15	15	15	15	15
8	8	9	10	11	12	13	14	15	8	9	10	11	12	13	14	15
9	9	9	11	11	13	13	15	15	9	9	11	11	13	13	15	15
10	10	11	10	11	14	15	14	15	10	11	10	11	14	15	14	15
11	11	11	11	11	15	15	15	15	11	11	11	11	15	15	15	15
12	12	13	14	15	12	13	14	15	12	13	14	15	12	13	14	15
13	13	13	15	15	13	13	15	15	13	13	15	15	13	13	15	15
14	14	15	14	15	14	15	14	15	14	15	14	15	14	15	14	15
15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15	15

Fig. 1: Table of unions : $CA(2, n) \cup CA(2, m)$

Definition 8 Let $1 \leq k < k'$, $0 \leq x \leq k' - k$ and $CA(k, n) = (\mathbf{Z}, V, V')$. $CA(k, n)_x^{k'}$ is defined by $CA(k, n)_x^{k'} = (\mathbf{Z}, \{0, 1, \dots, k' - 1\}, V'')$ where

$$\begin{aligned} V'' &= \{s_1 \cup v \cup s_2 \mid s_1 \in S_1, s_2 \in S_2, v \in V'\}, \\ S_1 &= \begin{cases} \{\emptyset\} & (x = 0) \\ 2^{\{0, \dots, x-1\}} & (x > 0) \end{cases}, \\ S_2 &= 2^{\{x+1, \dots, k\}} \end{aligned}$$

We note that $F_{CA(k, n)_0^{k'}} = F_{CA(k, n)}$ and $F_{CA(k, n)_1^{k'}} = SHIFT \diamond F_{CA(k, n)}$.

Definition 9 (Union) Let $C_1 = (G, V_1, V'_1)$ and $C_2 = (G, V_2, V'_2)$ be cellular automata on G . The union $C_1 \cup C_2$ of C_1 and C_2 is defined by $C_1 \cup C_2 = (G, V_1 \cup V_2, V'_1 \cup V'_2)$.

Definition 10 (Division) Let $C = (G, V, V')$ be a cellular automaton on G . If there exist $C_1 = (G, V_1, V'_1)$ and $C_2 = (G, V_2, V'_2)$ be cellular automata on G such that $V = V_1 \cup V_2$ and $V' = V'_1 \cup V'_2$, then we call C_1 and C_2 are division of C and C is dividable by C_1 and C_2 .

Example 11 The class of all 2-neighborhood cellular automata $\{CA(2, n) \mid n = 0, \dots, 15\}$ is generated by $\{CA(2, 0), CA(2, 1), CA(2, 2), CA(2, 4), CA(2, 8)\}$ using 'union' operations. For example, $CA(2, 13)$ is dividable by $CA(2, 1)$, $CA(2, 4)$, and $CA(2, 8)$. Fig 1 is the table of unions for $CA(2, n)$ ($n = 0, \dots, 15$).

Definition 12 (Composition) Let $C_1 = (G, V_1, V'_1)$ and $C_2 = (G, V_2, V'_2)$ be cellular automata on G . The composition $C_1 \diamond C_2$ of C_1 and C_2 is defined by $C_1 \diamond C_2 = (G, V_1 \cdot V_2, V'_1 \diamond V'_2)$ where

$$\begin{aligned} V_1 \cdot V_2 &= \{v_1 v_2 \in G \mid v_1 \in V_1, v_2 \in V_2\} \text{ and} \\ V'_1 \diamond V'_2 &= \{X \in 2^{V_1 \cdot V_2} \mid \{v \in V_1 \mid v^{-1} X \cap V_2 \in V'_2\} \in V'_1\}. \end{aligned}$$

Example 13 Let $V = \{0, 1\}$ and $V' = \{\{0\}, \{1\}\}$. We have $V \cdot V = \{0, 1, 2\}$. Since $1^{-1}\{0, 1\} \cap \{0, 1\} = \{0 - 1, 1 - 1\} \cap \{0, 1\} = \{0\}$ and $0^{-1}\{1, 2\} \cap \{0, 1\} = \{1 - 0, 2 - 0\} \cap \{0, 1\} = \{1\}$, we have $V' \diamond V' = \{\{0\}, \{2\}, \{0, 1\}, \{1, 2\}\}$. So we have $CA(2, 6) \diamond CA(2, 6) = CA(3, 90)$.

$n \setminus m$	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	255	236	209	192	139	136	129	128	55	36	17	0	3	0	1	0
2	0	16	34	48	68	68	66	64	8	24	34	48	12	12	2	0
3	255	252	243	240	207	204	195	192	63	60	51	48	15	12	3	0
4	0	2	12	12	48	34	24	8	64	66	68	68	48	34	16	0
5	255	238	221	204	187	170	153	136	119	102	85	68	51	34	17	0
6	0	18	46	60	116	102	90	72	72	90	102	116	60	46	18	0
7	255	254	255	252	255	238	219	200	127	126	119	116	63	46	19	0
8	0	1	0	3	0	17	36	55	128	129	136	139	192	209	236	255
9	255	237	209	195	139	153	165	183	183	165	153	139	195	209	237	255
10	0	17	34	51	68	85	102	119	136	153	170	187	204	221	238	255
11	255	253	243	243	207	221	231	247	191	189	187	187	207	221	239	255
12	0	3	12	15	48	51	60	63	192	195	204	207	240	243	252	255
13	255	239	221	207	187	187	189	191	247	231	221	207	243	243	253	255
14	0	19	46	63	116	119	126	127	200	219	238	255	252	255	254	255
15	255	255	255	255	255	255	255	255	255	255	255	255	255	255	255	255

Fig. 2: Table of compositions : $CA(2, n) \diamond CA(2, m)$

Example 14 The rule numbers of the 3-neighborhood cellular automata generated by composing 2-neighborhood cellular automata is $\{0, 1, 2, 3, 8, 12, 15, 16, 17, 18, 19, 24, 34, 36, 46, 48, 51, 55, 60, 63, 64, 66, 68, 72, 85, 90, 102, 116, 119, 126, 127, 128, 129, 136, 139, 153, 165, 170, 183, 187, 189, 191, 192, 195, 200, 204, 207, 209, 219, 221, 231, 236, 237, 238, 239, 240, 243, 247, 252, 253, 254, 255\}$. There are 62 kinds of 3-neighborhood cellular automata. Fig 2 is the table of compositions for $CA(2, n)$ ($n = 0, \dots, 15$).

Lemma 15 Let $C = (G, V, V')$ be a cellular automaton and $V_0 \subset G$. For any $\mathbf{c} \in 2^G$,

$$F_C(\mathbf{c}) \cap V_0 = F_C(\mathbf{c} \cap (V_0 \cdot V)) \cap V_0$$

Proof. We have $F_C(\mathbf{c}) \cap V_0 = \{v_0 \in V_0 \mid v_0^{-1} \mathbf{c} \cap V \in V'\} = \{v_0 \in V_0 \mid \mathbf{c} \cap v_0 V \in v_0 V'\} = \{v_0 \in V_0 \mid (\mathbf{c} \cap V_0 \cdot V) \cap v_0 V \in v_0 V'\} = F_C(\mathbf{c} \cap (V_0 \cdot V)) \cap V_0$. \square

The composition of cellular automata corresponds to find a cellular automaton which global transition function is the composition of global transition functions of original cellular automata.

Theorem 16 (Fujio (2008))

$$F_{C_1} \circ F_{C_2} = F_{C_1 \diamond C_2}$$

Proof. Since $F_{C_2}(\mathbf{c}) \cap V_1 = \{v \in V_1 \mid v^{-1} \mathbf{c} \cap V_2 \in V_2'\}$, we have

$$\begin{aligned} e \in F_{C_1}(F_{C_2}(\mathbf{c})) &\Leftrightarrow F_{C_2}(\mathbf{c}) \cap V_1 \in V_1' \\ &\Leftrightarrow F_{C_2}(\mathbf{c} \cap V_1 \cdot V_2) \cap V_1 \in V_1' \text{ (by Proposition. 2)} \\ &\Leftrightarrow \{v_1 \in V_1 \mid v_1^{-1} (\mathbf{c} \cap V_1 \cdot V_2) \cap V_2 \in V_2'\} \in V_1' \\ &\Leftrightarrow \mathbf{c} \cap V_1 \cdot V_2 \in V_1' \diamond V_2' \\ &\Leftrightarrow e \in F_{C_1 \diamond C_2}(\mathbf{c}) \end{aligned}$$

\square

Theorem 17 Let $C_1 = (G, V, V_1')$, $C_2 = (G, V, V_2')$ and $C_3 = (G, V_3, V_3')$ be cellular automata on a group G . Then,

$$(C_1 \cup C_2) \diamond C_3 = (C_1 \diamond C_3) \cup (C_2 \diamond C_3)$$

Proof. First, we note

$$\begin{aligned} (C_1 \cup C_2) \diamond C_3 &= (G, V \cdot V_3, (V'_1 \cup V'_2) \diamond V'_3), \text{ and} \\ (C_1 \diamond C_3) \cup (C_2 \diamond C_3) &= (G, V \cdot V_3, (V'_2 \diamond V'_1) \cup (V'_3 \diamond V'_1)). \end{aligned}$$

Next, we have

$$\begin{aligned} (V'_1 \cup V'_2) \diamond V'_3 &= \{X \in 2^{V \cdot V_3} \mid \{v \in V \mid v^{-1}X \cap V_3 \in V'_3\} \in V'_1 \cup V'_2\} \\ &= \{X \in 2^{V \cdot V_3} \mid \{v \in V \mid v^{-1}X \cap V_3 \in V'_3\} \in V'_1\} \\ &\quad \cup \{X \in 2^{V \cdot V_3} \mid \{v \in V \mid v^{-1}X \cap V_3 \in V'_3\} \in V'_2\} \\ &= (V'_1 \diamond V'_3) \cup (V'_2 \diamond V'_3) \end{aligned}$$

□

We note that $C_1 \diamond (C_2 \cup C_3) = (C_1 \diamond C_2) \cup (C_1 \diamond C_3)$ does not always holds for cellular automata C_1 , C_2 and C_3 . For example $CA(2, 6) \diamond (CA(2, 2) \cup CA(2, 4)) = CA(2, 6) \diamond CA(2, 6) = CA(3, 90)$, and $(CA(2, 6) \cup CA(2, 2)) \diamond (CA(2, 6) \cup CA(2, 4)) = CA(3, 46) \diamond CA(3, 116) = CA(3, 126)$.

Proposition 18 *Let $CA(1, n)_x^{k_1}$, $CA(k_2, n_2)$ and $CA(k_2, n_3)$ be cellular automata on \mathbf{Z} , where $0 \leq x < k_1$, and $n = 0, 1$. Then,*

$$CA(1, n)_x^{k_1} \diamond (CA(k_2, n_2) \cup CA(k_2, n_3)) = (CA(1, n)_x^{k_1} \diamond CA(k_2, n_2)) \cup (CA(1, n)_x^{k_1} \diamond CA(k_2, n_3)).$$

Proof. Let $V_1 = \{0, \dots, k_1 - 1\}$, $V'_1 = \{X \in 2^V \mid x \in X\}$, $\bar{V}'_1 = \{X \in 2^V \mid x \notin X\}$, $CA(k_2, n_2) = (\mathbf{Z}, V_2, V'_2)$, and $CA(k_2, n_3) = (\mathbf{Z}, V_2, V'_3)$. First, we note

$$\begin{aligned} CA(1, 0)_x^{k_1} &= (\mathbf{Z}, V_1, \bar{V}'_1), \\ CA(1, 1)_x^{k_1} &= (\mathbf{Z}, V_1, V'_1), \\ CA(1, 0)_x^{k_1} \diamond (CA(k_2, n_2) \cup CA(k_2, n_3)) &= (\mathbf{Z}, V_1 \cdot V_2, V'_1 \diamond (V'_2 \cup V'_3)), \text{ and} \\ (CA(1, n)_x^{k_1} \diamond CA(k_2, n_2)) \cup (CA(1, n)_x^{k_1} \diamond CA(k_2, n_3)) &= (\mathbf{Z}, V_1 \cdot V_2, (V'_1 \diamond V'_2) \cup (V'_1 \diamond V'_3)). \end{aligned}$$

Since

$$\begin{aligned} V'_1 \diamond (V'_2 \cup V'_3) &= \{X \in 2^{V_1 \cdot V_2} \mid \{v \in V \mid v^{-1}X \cap V_2 \in (V'_2 \cup V'_3)\} \in V'_1\} \\ &= \{X \in 2^{V_1 \cdot V_2} \mid x^{-1}X \cap V_2 \in (V'_2 \cup V'_3)\}, \text{ and} \\ (V'_1 \diamond V'_2) \cup (V'_1 \diamond V'_3) &= \{X \in 2^{V_1 \cdot V_2} \mid \{v \in V \mid v^{-1}X \cap V_2 \in V'_2\} \in V'_1\} \\ &\quad \cup \{X \in 2^{V_1 \cdot V_2} \mid \{v \in V \mid v^{-1}X \cap V_2 \in V'_3\} \in V'_1\} \\ &= \{X \in 2^{V_1 \cdot V_2} \mid x^{-1}X \cap V_2 \in V'_2\} \\ &\quad \cup \{X \in 2^{V_1 \cdot V_2} \mid x^{-1}X \cap V_2 \in V'_3\}, \end{aligned}$$

we have $V'_1 \diamond (V'_2 \cup V'_3) = (V'_1 \diamond V'_2) \cup (V'_1 \diamond V'_3)$, and

$$CA(1, 1)_x^{k_1} \diamond (CA(k_2, n_2) \cup CA(k_2, n_3)) = (CA(1, 1)_x^{k_1} \diamond CA(k_2, n_2)) \cup (CA(1, 1)_x^{k_1} \diamond CA(k_2, n_3)).$$

Similarly, we can prove

$$CA(1, 0)_x^{k_1} \diamond (CA(k_2, n_2) \cup CA(k_2, n_3)) = (CA(1, 0)_x^{k_1} \diamond CA(k_2, n_2)) \cup (CA(1, 0)_x^{k_1} \diamond CA(k_2, n_3)).$$

□

Example 19 We note $CA(3, 3) = (\mathbf{Z}, \{0, 1, 2\}, \{\phi, \{0\}\})$ and $CA(3, 102) = (\mathbf{Z}, \{0, 1, 2\}, \{\{0\}, \{1\}, \{0, 2\}, \{1, 2\}\})$. The composition $CA(3, 3) \diamond CA(3, 102) = (\{0, 1, 2, 3, 4\}, \{\{1\}, \{0, 1\}, \{1, 4\}, \{0, 1, 4\}, \{3\}, \{0, 3\}, \{3, 4\}, \{0, 3, 4\}\})$. Since $\{\{1\}, \{0, 1\}, \{1, 4\}, \{0, 1, 4\}, \{3\}, \{0, 3\}, \{3, 4\}, \{0, 3, 4\}\} = \bigcup\{\{0\} \cup s, s \cup \{4\}, \{0\} \cup s \cup \{4\} \mid s \in \{\{1\}, \{3\}\}\}$, we have $CA(3, 3) \diamond CA(3, 102) = CA(3, 18)_1^5$. (cf. Fig 3, Fig 4, Fig 5)




Fig. 3: An example of configurations of $CA(3, 3)$




Fig. 4: An example of configurations of $CA(3, 102)$




Fig. 5: An example of configurations of $CA(3, 18) = CA(3, 3) \diamond CA(3, 102)$

Example 20 A 2-neighborhood cellular automaton is considered as 3-neighborhood cellular automaton and 3-neighborhood cellular automaton is also considered as 5-neighborhood cellular automaton. The followings is an observation of the embeddings and compositions.

$$\begin{aligned} CA(2, 1) &= (\mathbf{Z}, \{0, 1\}, \{\phi\}) \\ CA(2, 1)_0^3 &= (\mathbf{Z}, \{0, 1, 2\}, \{\phi, \{2\}\}) \\ &= CA(3, 17) \\ CA(2, 1) \diamond CA(2, 1) &= (\mathbf{Z}, \{0, 1, 2\}, \{\{0, 1\}, \{0, 2\}, \{1, 2\}, \{1\}\}) \\ &= CA(3, 236) \\ CA(3, 17) \diamond CA(3, 17) &= (\mathbf{Z}, \{0, 1, 2, 3, 4, 5\}, V') \\ &= CA(5, 3974950124) = CA(3, 236)_0^5 \\ V' &= \bigcup\{\{s, s \cup \{3\}, s \cup \{4\}, s \cup \{3, 4\}\} \mid s \in CA(3, 236)\} \end{aligned}$$

3 Generalization

A subset V of G is considered as a characteristic function $V : G \rightarrow 2$ where $2 = \{0, 1\}$. That is V is a function which values are

$$V(g) = \begin{cases} 0 & (g \notin V) \\ 1 & (g \in V). \end{cases}$$

Sometimes V is represented as an injection $i_V : V \rightarrow G$ where $i_V(g) = g$.

Extending our 2-states cellular automata on groups to many-states cellular automata on groups, we replace the set $2 = \{0, 1\}$ to a finite set S .

Definition 21 Let G be a group, S a finite set. A generalized cellular automaton on G is a four-tuple $C = (G, S, i_V, V')$ of the group G , an injection $i_V : V \rightarrow G$, and a function $V' : S^V \rightarrow S$ where S^V is the set of all functions from V to S . A configuration $\mathbf{c} : G \rightarrow S$ is a function. The global transition function $F_C : S^G \rightarrow S^G$ is defined by $F_C(\mathbf{c})(g) = V'(\mathbf{c} \circ g \circ i_V)$.

Proposition 22 Let G be a group, and $S = 2 = \{0, 1\}$. The global function $F_C : 2^G \rightarrow 2^G$ is the same as defined in Definition 1. That is $F_C(\mathbf{c}) = \{g \in G \mid V'(\mathbf{c} \circ g \circ i_V) = 1\} = \bigcup_{g \in G} g \cdot l_{V'}(g^{-1} \cdot \mathbf{c} \cap V)$.

Proof. For $g \in G$, we have

$$\begin{aligned} g \in \bigcup_{g \in G} g \cdot l_{V'}(g^{-1} \cdot \mathbf{c} \cap V) &\Leftrightarrow l_{V'}(g^{-1} \cdot \mathbf{c} \cap V) = \{e\} \\ &\Leftrightarrow g^{-1} \cdot \mathbf{c} \cap V \in V' \\ &\Leftrightarrow g^{-1}\{x \mid \mathbf{c}(x) = 1\} \cap V \in V' \\ &\Leftrightarrow \{g^{-1}x \mid \mathbf{c}(x) = 1\} \cap V \in V' \\ &\Leftrightarrow \{v \mid \mathbf{c}(gv) = 1\} \cap V \in V' \text{ (cf. } (x = gv)) \\ &\Leftrightarrow \{v \mid \mathbf{c}(gv) = 1, v \in V\} \in V' \\ &\Leftrightarrow \{v \mid \mathbf{c} \circ g \circ i_V(v) = 1\} \in V' \\ &\Leftrightarrow V'(\mathbf{c} \circ g \circ i_V) = 1 \\ &\Leftrightarrow g \in F_C(\mathbf{c}). \end{aligned}$$

□

Example 23 Let $G = \mathbf{Z}$, $S = \mathbf{Z}_m$, and $V = \{-r, -r+1, \dots, 0, \dots, +r\}$. For a polynomial $f(X) = \sum_{i=-r}^{+r} a_i X^i$, ($a_i \in \mathbf{Z}_m$), we define the function $V'_{f(X)} : \mathbf{Z}_m^V \rightarrow \mathbf{Z}_m$ by $V'(x_{-r}, x_{-r+1}, \dots, x_0, \dots, x_{+r}) = \sum_{i=-r}^{+r} a_{-i} x_i$, ($(x_{-r}, x_{-r+1}, \dots, x_0, \dots, x_{+r}) \in \mathbf{Z}_m^V$). A configuration $\mathbf{c} \in \mathbf{Z}_m^{\mathbf{Z}}$ is represented as a formal power series $\sum c_i X^i$ where $c_i = \mathbf{c}(i)$ (cf. Sato (1994); Manzini (1998)). Since $\mathbf{c} \circ j \circ i_V(i) =$

$\mathbf{c}(j+i) = c_{j+i}$, and $\mathbf{c} \circ j \circ i_V = (c_{j-r}, c_{j-r+1}, \dots, c_j, \dots, c_{j+r})$, we have

$$\begin{aligned}
(\sum \mathbf{c}(i)X^i)f(X) &= (\sum c_i X^i)f(X) \\
&= (\sum c_i X^i)(\sum_{i'=-r}^{+r} a_{i'} X^{i'}) \\
&= (\sum c_i X^i)(\sum_{i'=-r}^{+r} a_{-i'} X^{-i'}) \\
&= \sum_{i'=-r}^{+r} (\sum c_i a_{-i'} X^{i-i'}) \\
&= \sum_{i'=-r}^{+r} ((\sum a_{-i'} c_{j+i'}) X^j) \text{ (cf. } j = i - i') \\
&= \sum (V'(c_{j-r}, c_{j-r+1}, \dots, c_j, \dots, c_{j+r}) X^j) \\
&= \sum (V'(\mathbf{c} \circ j \circ i_V) X^j). \\
&= \sum (F_C(\mathbf{c})(j) X^j).
\end{aligned}$$

The transition of the cellular automaton $C = (\mathbf{Z}, \mathbf{Z}_m, i_V, V'_{f(X)})$ is corresponding to the product of polynomials (the formal power series).

Acknowledgements

The authors thank Professor Yasuo Kawahara for his valuable suggestions and discussions. This work has been partially supported by Kyushu University Global COE Program “Education-and-Research Hub for Mathematics-for-Industry” and Regional Innovation Cluster Program (Global Type 2nd Stage) “Fukuoka Cluster for Advanced System LSI Technology Development”.

References

- E. Berlekamp, J. Conway, and R. Guy. *Winning Ways for Your Mathematical Plays, 2*. Academic Press, 1982.
- M. Fujio. XOR² = 90 - graded algebra structure of the boolean algebra of local transition rules -. In *Papers Res. Inst. Math. Sci., Kyoto Univ.*, volume 1599, pages 97–102, 2008.
- T. Ito. Abstract collision systems on g -sets. *J. of Math for Industry*, 2(A):57–73, 2010.
- T. Ito, S. Inokuchi, and Y. Mizoguchi. An abstract collision system. In *Automata-2008 Theory and Applications of Cellular Automata*, pages 339–355. Luniver Press, 2008.
- G. Manzini. Invertible linear cellular automata over \mathbf{Z}_m . *J. Comput. Syst. Sci.*, 56:60–67, 1998.

- W. Pries, A. Thanailakis, and H. Card. Group properties of cellular automata and VLSI applications. *IEEE Trans. on Computers.*, C-35(12):1013–1024, 1986.
- E. Rémila. An introduction to automata on graphs. In *Cellular Automata*, pages 345–352. Kluwer Academic Publishers, 1998.
- Z. Róka. *Automates cellulaires sur les graphes de Caley*. PhD thesis, Université Lyon I et Ecole Normale Supérieure de Lyon, 1994.
- T. Sato. Group structured linear cellular automata over \mathbf{Z}_m . *J. Comput. Syst. Sci.*, 49:18–23, 1994.
- J. von Neumann. *Theory of self-reproducing automata*. Univ. of Illinois Press, 1983.
- T. Watanabe and S. Noguchi. On the uniform decomposition of automata and spatial networks. *IEICE Trans. Inf. and Syst.*, 11(2):1–9, 1982.

List of MI Preprint Series, Kyushu University

The Global COE Program
Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p -adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials

- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials
- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings

- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric q -functions of the q -Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications
- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain

- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection

- MI2009-26 Manabu YOSHIDA
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance
- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA
Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI
Finite element computation for scattering problems of micro-hologram using DtN map

- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO
Abstract collision systems on groups
- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA
An algebraic approach to underdetermined experiments

- MI2010-10 Kei HIROSE & Sadanori KONISHI
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE
Lagrangian approach to weakly nonlinear stability of an elliptical flow

MI2010-21 Hiroki MASUDA

Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test

MI2010-22 Toshimitsu TAKAESU

A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs

MI2010-23 Takahiro ITO, Mitsuhiko FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Composition, union and division of cellular automata on groups