

Synthetic Photochemistry. XL. Photochemical Cycloaddition of p-Tropoquinone Bisacetal to Olefins. Formations of Cycloadducts with C=O and C=C Groups

Mori, Akira

Department of Molecular Science and Technology, Interdisciplinary Graduate School of
Engineering Sciences, Kyushu University

Kasai, Seishi

Department of Molecular Science and Technology, Interdisciplinary Graduate School of
Engineering Sciences, Kyushu University

Isayama, Yasutoshi

Department of Molecular Science and Technology, Interdisciplinary Graduate School of
Engineering Sciences, Kyushu University

Takeshita, Hitoshi

Department of Molecular Science and Technology, Interdisciplinary Graduate School of
Engineering Sciences, Kyushu University

<https://doi.org/10.15017/17694>

出版情報：九州大学大学院総合理工学報告. 9 (2), pp.197-203, 1988-01-01. 九州大学大学院総合理工学研究科

バージョン：

権利関係：

Synthetic Photochemistry. XL¹⁾. Photochemical Cycloaddition of *p*-Tropoquinone Bisacetal to Olefins. Formations of Cycloadducts with C=O and C=C Groups

Akira MORI[†], Seishi KASAI^{††}, Yasutoshi ISAYAMA^{††}

and Hitoshi TAKESHITA^{*†}

(Received September 8, 1987)

The photochemical cycloaddition of *p*-tropoquinone bisacetal to electron-rich olefins mainly gave oxetanes; an alkylidenecycloheptadienone derivative from cyclohexene and a hemiacetal from cyclopentene were explained by subsequent fragmentation of the oxetane. The reaction with 2,3-trimethoxycyclohept-6-ene-1,5-dione and 2,3-dimethyl-2-butene gave *cis*- and *trans*-fused cyclobutane derivatives and an ene product. These results were different from those of the photochemical reactions of *p*-tropoquinone and its monoacetal.

Recently, we have found practical methods for the preparation of *p*-tropoquinone acetals (**1**) and (**2**) from 2-alkoxy-5-hydroxytropones by means of thallium (III) nitrate (TTN) - or cerium (IV) ammonium nitrate (CAN) - oxidations^{2,3)} and the anodic oxidation⁴⁾ and have verified their usefulness as the masked forms of *p*-tropoquinone (**3**) for synthesizing functional seven-membered ring derivatives.

For the photochemistry of **3** and its derivatives, only a few examples are known: Itô et al. have observed a formation of the dioxene derivative (**4**) from **3** and styrene.⁵⁾ On the other hand, we ourselves have found the photoreaction of **3** in methanol to a ring contracted product, methyl 2,5-dioxo-3-cyclopentenyl acetate (**5**)⁶⁾ and 2,10-dihydroxydicyclohepta-*[b, d]* furan-3, 9-dione (**6**)^{6,7)} while the attempted reaction in benzene gave no identifiable product. The formation of **5** is explained as the hemiacetal formation from **3** prior to photolysis; this was supported since *p*-tropoquinone acetal (**7**) in either benzene or methanol afforded alkyl 2,5-dioxo-3-cyclopentenyl acetate (**8**)⁶⁾. The formation of **6** is probably due to a hydrogen-abstraction of the excited **3** followed by dimerization.

In this paper we will show the result of the photochemical cycloaddition of acetals to several electron-rich olefins to afford oxetanes and cyclobutane derivatives without a contraction of seven-membered ring.


When a benzene solution of **2** and 2,3-dimethyl-2-butene was irradiated by means of high pressure mercury lamp through a Pyrex-glass filter, the product (**9**), having no carbonyl and hydroxyl absorptions in the IR spectrum, was obtained in 39% yield and it was immediately identified to be an oxetane; its ¹H NMR spectrum⁸⁾ showed four olefinic protons signals remained and three singlets newly appeared at $\delta = 1.15$ (6H), 1.34 (3H), and 1.42 (3H).

The reactions with unsymmetrical olefins, 2-methyl-2-butene, butyl vinyl ether, isobutyl

[†]Department of Molecular Science and Technology

^{††}Department of Molecular Science and Technology, Graduate Student


vinyl ether, and isopropenyl methyl ether were also investigated; similar results, formations of oxetanes (**10**–**13**) were recognized as summarized in Table 1.



[Scheme 1]

Table 1 Photochemical Cycloadditions of **2** with Various Olefins.


Olefins	Products	Yields (ratio)
	9 R ₁ =R ₂ =R ₃ =R ₄ =Me	39%
	10 R ₁ =R ₂ =R ₃ =Me, R ₄ =H	29% (2 : 1)
	11 R ₁ : R ₂ =On-Bu : H, R ₃ =R ₄ =H	5%
	12 R ₁ : R ₂ =Oi-Bu : H, R ₃ =R ₄ =H	10%
	13 R ₁ : R ₂ =Me : OMe, R ₃ =R ₄ =H	32% (2 : 1)
	15 R ₁ =Me, R ₂ =R ₃ =-(CH ₂) ₄ -, R ₄ =H	5%
	16 R ₁ =R ₃ =-(CH ₂) ₅ -, R ₂ =R ₄ =H	8%
	17 R ₁ =R ₃ =H, R ₂ =R ₄ =-(CH ₂) ₅ -	7%



Their structures were elucidated by the spectral analyses. Namely, the ^1H NMR spectrum of **10** showed an isomeric mixture of oxetanes in 2:1 ratio. Regiochemistry of **10** was determined by the considerations of the chemical shifts⁸⁾ of the methine protons on the carbon bearing the oxygen atom, which appeared at 4.48 (q, $J=6.5$ Hz) for the major isomer and at 4.64 (q, $J=6.5$ Hz) for the minor isomer, but the stereochemistry was not further investigated.

The reactions of **2** with butyl and isobutyl vinyl ethers yielded the corresponding single oxetanes (**11**) and (**12**); their regiochemistry was determined similarly on the basis of the chemical shift analysis (4.2–4.8) of three protons on the oxetane ring. With isopropenyl methyl ether, two isomeric oxetanes (**13**) were isolated and the structures were deduced as depicted.

Next, the irradiation of a mixture of cyclohexene and **2** yielded an aldehyde (**14**) in 20% yield. Its ^1H NMR spectrum exhibited a formyl proton signal at 9.72 ppm as a triplet ($J=1$ Hz) and a newly-formed olefinic proton signal at 5.79 ppm as a triplet ($J=7.5$ Hz). Therefore, the structure of **14** is the bisacetal of alkylidenecycloheptadienone. Since the formyl proton signal was already recognized in the NMR spectrum of the crude reaction mixture immediately after the irradiation, **14** might be formed during the irradiation through a secondary photochemical cleavage⁹⁾ of the oxetane (**A**). With 1-methylcyclohexene, an oxetane (**15**) was obtained. From cycloheptene, two isomeric oxetanes (**16** and **17**) formed. The stereochemistry of the junction in **16** and **17** was assigned to be trans from the coupling constant ($J=10$ Hz)¹⁰⁾ of a methine proton at 4.56. Furthermore, the reaction with cyclopentene gave a diastereomeric mixture of ene products (**18**) and a hemiacetal (**19**). The latter **19** might be derived from the oxetane (**B**). In **B**, the oxetane ring cleavage led to an alkylidenecycloheptadienone derivative (**C**), which followed the hydration process to give **19**.




[Scheme 2]

Other than these, the reactions of styrene, diphenylacetylene and tetracyanoethylene with **2** gave no product.

Subsequently, the photoreaction of 2,2,3-trimethoxycyclohept-6-ene-1,5-dione,²⁾ a Michael-type methanol adduct of *p*-tropoquinone monoacetal (**20**), with 2,3-dimethyl-2-butene was carried out; a mixture (3:2) of the cyclobutane derivatives (**21**) and (**22**) in 21% yield and an ene product (**23**) in 6% yield were isolated. The cyclobutane products showed a carbonyl absorption at 1740 cm^{-1} in IR spectrum and no olefinic protons in the ^1H NMR spectrum. Since protons on the juncture of **22** appeared as a pair of *AB* system with a large spin-spin coupling constant ($J=12.5$ Hz)¹⁰⁾ at 2.84 and 3.23, it is a trans-fused

cyclobutane. On the other hand, the structure of **21** was suggested to be a cis-fused one in terms of the chemical transformation to the bicyclic hemiacetal (**24**) by the treatment with perchloric acid in aqueous acetone. The isolation of trans-ring-fused product, **22**, is interesting. The ^1H NMR spectrum of **23** showed the signals due to 1,1,2-trimethylpropenyl group, but the position of the alkenyl group in **23** could not be determined from the spectral data. The photoadditions of **20** with isopropenyl methyl ether and 2-methoxyfuran were resulted the unseparable mixture.

The oxetane formation for **2** was explained from the facts that the electrophilic oxygen atom of the n, π^* triplet state of the carbonyl group attacked the electron-rich olefins. From the reaction of **20** and electron-rich olefin, 2,3-dimethyl-2-butene, cis- and trans-fused cyclobutanes afforded with the competitive formation of the ene product. Presumably, the 1,4-diradical (**D**) cyclized to give cyclobutanes and abstracted a hydrogen to give ene product.



In conclusion, by acetalization of α -dicarbonyl groups in *p*-tropoquinone, the photochemical behavior showed a marked difference from that of *p*-tropoquinone itself; bisacetal **2** gave oxetane derivatives with electron-rich olefins without contraction of seven-membered ring.

Experimental

The elemental analyses were carried out by Miss S. Hirashima, of Institute of Advanced Material Study, Kyushu University. The NMR spectra were measured by a JEOL FX 100 Spectrometer in CDCl_3 solution unless otherwise specified, and the chemical shifts expressed were in δ unit. The mass spectra were measured with a JEOL OISG-2 Spectrometer. The IR spectra were taken as a liquid film inserted between NaCl plates using a Jasco IR-A Spectrometer. The UV spectra were measured by a Hitachi U-3200 Spectrophotometer.

Photoreaction of 2 and 2,3-Dimethyl-2-butene. A benzene solution (1.5 cm^3) of **2** (37 mg) and 2,3-dimethyl-2-butene (150 mg) was irradiated for 2 h with 400 W high pressure mercury lamp under N_2 atmosphere at $5-10^\circ\text{C}$. After the evaporation of the volatile materials, the residue was purified with a silica-gel chromatography (EtOAc:hexane=1:3) to give **9** [a colorless oil, 20 mg; 39%. Found: m/z , 310.1808 (M^+). Calcd for $\text{C}_{17}\text{H}_{26}\text{O}_5$: 310.1779. ^1H NMR $\delta = 1.15$ (6H, s), 1.34 (3H, s), 1.42 (3H, s), 3.23 (6H, s), 3.3–3.6 (2H,

m), 3.9–4.1 (2H, m), 5.48 (1H, d, $J=13$ Hz), 5.54 (1H, d, $J=13$ Hz), 6.12 (1H, dd, $J=13, 2$ Hz), and 6.19 (1H, dd, $J=13, 2$ Hz). ^{13}C NMR $\delta = 20.7, 22.1, 27.0$ (2C), 47.6, 49.4, 49.9, 58.4, 58.7, 84.8, 85.2, 95.1, 97.5, 127.1, 129.2, 134.9, and 138.3. IR ν : 2950, 1455, 1410, 1390, 1370, and 1270 cm^{-1} . UV $\lambda_{\text{max}}^{\text{MeOH}}$: 218 nm ($\epsilon = 1700$)].

Photoreaction of 2 and 2-Methyl-2-butene. A benzene solution (2 cm^3) of **2** (35 mg) and 2-methyl-2-butene (130 mg) was irradiated for 2 h as same as above procedure. After the evaporation of the volatile materials, the residue was purified with a silica-gel chromatography (EtOAc:hexane = 1:5) to give an isomeric mixture (2:1 ratio) of oxetanes, **10** [a colorless oil, 13.5 mg; 29%. Found: m/z , 296.1621 (M^+). Calcd for $\text{C}_{16}\text{H}_{24}\text{O}_5$: 296.1621. ^1H NMR $\delta = 1.10$ (3H, s), 1.18 (3H, s), 1.31 (3H, d, $J=6.5$ Hz), 3.23 (6H, s), 3.4–3.6 (2H, m), 3.9–4.1 (2H, m), 4.48 (1H, q, $J=6.5$ Hz), 5.47 (1H, d, $J=12.5$ Hz), 5.58 (1H, d, $J=12.5$ Hz), 6.05 (1H, dd, $J=12.5, 2$ Hz), and 6.19 (1H, dd, $J=12.5, 2$ Hz) for a major isomer; 1.08 (3H, s), 1.12 (3H, s), 1.22 (3H, d, $J=6.5$ Hz), 3.23 (6H, s), 3.4–3.6 (2H, m), 3.9–4.1 (2H, m), 4.64 (1H, q, $J=6.5$ Hz), 5.52 (1H, d, $J=12.5$ Hz), 5.53 (1H, d, $J=12.5$ Hz), 5.97 (1H, dd, $J=12.5, 2$ Hz), and 6.25 (1H, dd, $J=12.5, 2$ Hz) for a minor isomer. IR ν : 2950, 1455, 1410, 1390, and 1270 cm^{-1} . UV $\lambda_{\text{max}}^{\text{MeOH}}$: 208 nm ($\epsilon = 6550$)].

Photoreaction of 2 and Butyl Vinyl Ether. A benzene solution (4 cm^3) of **2** (70 mg) and butyl vinyl ether (390 mg) was irradiated for 4 h. After the purification with a silica-gel chromatography (AcOEt:hexane = 1:5–1:2), **11** [a pale yellow oil, 5 mg; 5%. Found: m/z , 326.1727 (M^+) Calcd for $\text{C}_{17}\text{H}_{26}\text{O}_6$: 326.1727. ^1H NMR $\delta = 0.90$ (3H, t, $J=8$ Hz), 1.1–1.6 (4H, m), 3.20 (3H, s), 3.23 (3H, s), 3.4–3.7 (4H, m), 3.9–4.1 (2H, m), 4.19 (1H, dd, $J=7, 6.5$ Hz), 4.41 (1H, dd, $J=7, 6.5$ Hz), 4.67 (1H, t, $J=7$ Hz), 5.47 (1H, d, $J=13$ Hz), 5.63 (1H, d, $J=13$ Hz), 6.10 (1H, dd, $J=13, 2$ Hz), and 6.33 (1H, dd, $J=13, 2$ Hz). ^{13}C NMR $\delta = 13.9, 19.4, 31.8, 49.2, 49.6, 58.5, 58.7, 68.6, 69.2, 73.3, 80.6, 88.9, 96.4, 128.8$ (2C), 131.1, and 135.7. IR ν : 2950, 1460, 1150, 1100, and 1000 cm^{-1}] was obtained.

Photoreaction of 2 with Isobutyl Vinyl Ether. A benzene solution (4 cm^3) of **2** (70 mg) and isobutyl vinyl ether (390 mg) was irradiated for 4 h. After the purification with a silica-gel chromatography (AcOEt:hexane = 1:5–1:2), **12** [a pale yellow oil, 10 mg; 10%. Found: m/z , 326.1730 (M^+). Calcd for $\text{C}_{17}\text{H}_{26}\text{O}_6$: 326.1727. ^1H NMR $\delta = 0.88$ (6H, d, $J=6.5$ Hz), 1.7–2.2 (1H, m), 3.20 (3H, s), 3.24 (3H, s), 3.3–3.6 (4H, m), 3.9–4.1 (2H, m), 4.18 (1H, dd, $J=7, 6$ Hz), 4.42 (1H, dd, $J=7, 6$ Hz), 4.77 (1H, t, $J=7$ Hz), 5.48 (1H, d, $J=13$ Hz), 5.64 (1H, d, $J=13$ Hz), 6.08 (1H, dd, $J=13, 2$ Hz), and 6.34 (1H, dd, $J=13, 2$ Hz). ^{13}C NMR $\delta = 19.4, 28.5, 39.5, 49.2, 49.5, 54.8, 58.5, 58.7, 73.2, 80.7, 88.9, 96.5, 97.4, 128.6, 128.7, 131.2, \text{ and } 135.6$. IR ν : 2950, 1460, 1150, 1095, 1060, and 1000 cm^{-1}] was obtained.

Photoreaction of 2 with Isopropenyl Methyl Ether. A benzene solution (2 cm^3) of **2** (50 mg) and isopropenyl methyl ether (190 mg) was irradiated for 2 h. After the same work up, an isomeric mixture (2:1) of oxetanes, **13** [a colorless oil, 21 mg; 32%. Found: m/z , 298.1411 (M^+). Calcd for $\text{C}_{15}\text{H}_{22}\text{O}_6$: 298.1414. ^1H NMR $\delta = 1.48$ (3H, d, $J=1$ Hz), 3.21 (3H, s), 3.24 (6H, s), 3.4–3.6 (2H, m), 3.9–4.1 (2H, m), 4.25 (1H, d, $J=7$ Hz), 4.50 (1H, dd, $J=7, 1$ Hz), 5.57 (1H, d, $J=13$ Hz), 5.61 (1H, d, $J=13$ Hz), 5.96 (1H, dd, $J=13, 2$ Hz), and 6.42 (1H, dd, $J=13, 2$ Hz) for a major isomer; 1.40 (3H, s), 3.18 (3H, s), 3.21 (3H, s), 3.24 (3H, s), 3.4–3.6 (2H, m), 3.9–4.1 (2H, m), 4.30 (1H, d, $J=7$ Hz), 4.55 (1H, d, $J=7$ Hz), 5.55 (1H, d, $J=12.5$ Hz), 5.56 (1H, d, $J=12.5$ Hz), 5.96 (1H, dd, $J=12.5, 2$ Hz), and

6.16 (1H, dd, $J=12.5$, 2 Hz) for a minor isomer. IR ν : 2950, 1450, 1410, 1380, 1320, 1150, 1090, 1070, and 1000 cm^{-1}] was obtained.

Photoreaction of 2 with Cyclohexene. A benzene solution (2 cm^3) of **2** (30 mg) and cyclohexene (130 mg) was irradiated for 1.5 h. By a silica-gel chromatography (AcOEt:hexane=1:2), **14** [a pale yellow oil, 8 mg; 20%. Found: m/z , 308.1614 (M^+). Calcd for $\text{C}_{17}\text{H}_{24}\text{H}_5$: 308.1622. ^1H NMR $\delta = 1.4-1.8$ (4H, m), 2.2–2.5 (4H, m), 3.19 (3H, s), 3.21 (3H, s), 3.50 (2H, d, $J=7.5$ Hz), 4.06 (2H, d, $J=7.5$ Hz), 5.39 (1H, dd, $J=13$, 1.5 Hz), 5.58 (1H, dd, $J=12.5$, 1.5 Hz), 5.79 (1H, t, $J=7.5$ Hz), 6.08 (1H, dd, $J=13$, 1.5 Hz), 6.46 (1H, ddd, $J=12.5$, 1.5, 1 Hz), and 9.72 (1H, t, $J=1$ Hz). IR ν : 2950, 1725, 1680, 1450, 1290, and 1095 cm^{-1} . UV $\lambda_{\text{max}}^{\text{MeOH}}$: 208 nm ($\epsilon = 7200$), 218 (5800), and 255 (3250)] was obtained.

Photoreaction of 2 with 1-Methylcyclohexene. A benzene solution (3 cm^3) of **2** (70 mg) and 1-methylcyclohexene (360 mg) was irradiated for 2 h. After a silica-gel chromatography of the reaction mixture (AcOEt:hexane=1:5), **15** [a pale yellow oil, 5 mg; 5%. Found: m/z , 322.1771 (M^+). Calcd for $\text{C}_{18}\text{H}_{26}\text{O}_5$: 322.1777. ^1H NMR $\delta = 1.0-2.1$ (8H, m), 1.16 (3H, s), 3.21 (3H, s), 3.23 (3H, s), 3.3–3.6 (2H, m), 3.9–4.15 (2H, m), 4.45 (1H, t, $J=5$ Hz), 5.52 (1H, d, $J=13$ Hz), 5.61 (1H, d, $J=13$ Hz), 6.18 (1H, dd, $J=13$, 2 Hz), and 6.27 (1H, dd, $J=13$, 2 Hz). ^{13}C NMR $\delta = 19.0$, 20.6, 24.7, 28.8, 29.6, 46.9, 49.5, 49.9, 58.6 (2C), 82.7, 83.3, 95.2, 97.7, 128.7, 129.7, 133.7, and 136.9. IR ν : 2950, 1450, 1405, 1380, 1150, 1090, 1060, and 1000 cm^{-1}] was obtained.

Photoreaction of 2 with Cycloheptene. A benzene solution (5 cm^3) of **2** (80 mg) and cycloheptene (400 mg) was irradiated for 4 h. After a silica-gel chromatography (AcOEt:hexane=1:5) of the reaction mixture, two isomeric products, **16** [a pale yellow oil, 8.6 mg; 8%. Found: m/z , 322.1781 (M^+). Calcd for $\text{C}_{18}\text{H}_{26}\text{O}_5$: 322.1781. ^1H NMR $\delta = 1.3-1.8$ (10H, m), 2.7–2.9 (1H, m), 3.18 (3H, s), 3.22 (3H, s), 3.4–3.6 (2H, m), 3.9–4.1 (2H, m), 4.56 (1H, td, $J=10$, 4 Hz), 5.43 (1H, d, $J=13$ Hz), 5.65 (1H, d, $J=13$ Hz), 5.85 (1H, dd, $J=13$, 2 Hz), and 6.32 (1H, dd, $J=13$, 2 Hz). ^{13}C NMR $\delta = 23.6$, 24.7, 25.6, 27.9, 35.4, 49.4, 49.5, 55.0, 58.6, 58.7, 82.3, 84.5, 96.1, 97.7, 128.8, 130.7, 133.4, and 136.5. IR ν : 2950, 1450, 1260, 1210, 1150, 1090, 1070, and 1000 cm^{-1}] and **17** [a pale yellow oil, 7.3 mg, 7%. Found: m/z , 322.1776 (M^+). ^1H NMR $\delta = 1.2-1.9$ (10H, m), 2.8–3.1 (1H, m), 3.15 (3H, s), 3.26 (3H, s), 3.4–3.6 (2H, m), 3.9–4.1 (2H, m), 4.56 (1H, td, $J=10$, 4 Hz), 5.38 (1H, d, $J=12.5$ Hz), 5.59 (1H, d, $J=13$ Hz), 6.09 (1H, dd, $J=12.5$, 2 Hz), and 6.33 (1H, dd, $J=13$, 2 Hz). ^{13}C NMR $\delta = 23.4$, 24.9, 26.5, 28.1, 35.0, 48.9, 49.5, 53.4, 58.5, 58.7, 82.2, 85.2, 95.7, 97.8, 127.2, 128.4, 131.0, and 139.5. IR ν : 2950, 1450, 1405, 1260, 1150, 1090, 1060, and 1000 cm^{-1}] were obtained.

Photoreaction of 2 with Cyclopentene. A benzene solution (5 cm^3) of **2** (100 mg) and cyclopentene (360 mg) was irradiated for 4 h. The reaction mixture was purified by a silica-gel chromatography (AcOEt:hexane=1:5) to give **18** [a pale yellow oil, 13.4 mg; 10%. Found: m/z , 294.1461 (M^+). Calcd for $\text{C}_{16}\text{H}_{22}\text{O}_5$: 294.1464. ^1H NMR $\delta = 1.4-2.4$ (4H, m), 2.9–3.1 (1H, m), 3.23 (3H, s), 3.27 (3H, s), 3.3–4.2 (4H, m), and 5.3–6.0 (6H, m) for a-isomer; 1.4–2.4 (4H, m), 2.9–3.1 (1H, m), 3.24 (3H, s), 3.28 (3H, s), 3.4–4.1 (4H, m), and 5.3–6.0 (6H, m) for b-isomer. ^{13}C NMR $\delta = 24.0$, 32.5, 49.2, 49.3, 56.8, 57.1, 58.7, 74.5, 96.5, 97.1, 127.7, 130.2, 130.7, 134.6, 136.3, and 137.9 for a-isomer; 24.0, 32.5, 49.2, 49.3, 56.8, 57.1, 58.7, 74.5, 96.5, 97.1, 128.5, 129.3, 130.2, 134.0, 134.5, and 136.2 for b-isomer. IR ν :

3600, 2950, 1610, 1450, 1415, 1265, 1150, 1090, 1070, and 1000 cm^{-1}] and **19** [a pale yellow oil, 2 mg; 2%. Found: m/z , 294.1461 ($M^+ - 18$). Calcd for $C_{16}H_{22}O_5$: 294.1464. $^1\text{H NMR}$ δ = 1.2–2.2 (6H, m), 2.90 (1H, dd, $J=8, 5$ Hz), 3.19 (3H, s), 3.25 (3H, s), 3.4–3.6 (2H, m), 3.9–4.2 (2H, m), 5.20 (1H, dd, $J=5, 4$ Hz), 5.43 (1H, d, $J=12.5$ Hz), 5.46 (1H, d, $J=13$ Hz), 5.95 (1H, dd, $J=13, 2$ Hz), and 6.50 (1H, dd, $J=12.5, 2$ Hz). IR ν : 3450, 1450, 1400, 1145, 1090, 1065, and 1000 cm^{-1}].

Photoreaction of 20 with 2, 3-Dimethyl-2-butene. A benzene solution (6 cm^3) of **20** (130 mg) and 2, 3-dimethyl-2-butene (1.03 g) was irradiated for 3 h. After a silica-gel chromatography (AcOEt:hexane = 1:5) of the reaction mixture, two isomeric products (**21** and **22**) [a pale yellow oil, 38.2 mg; 21%. Found: m/z , 298.1776 (M^+). Calcd for $C_{16}H_{26}O_5$: 298.1778. $^1\text{H NMR}$ δ = 0.98 (3H, s), 1.02 (3H, s), 1.03 (3H, s), 1.19 (3H, s), 2.70 (1H, dd, $J=16, 5$ Hz), 2.91 (1H, dd, $J=16, 3$ Hz), 3.0–3.4 (2H, underneath methoxyl signals), 3.25 (3H, s), 3.33 (3H, s), 3.36 (3H, s), and 3.77 (1H, dd, $J=5, 3$ Hz) for **21**; 0.98 (3H, s), 1.02 (3H, s), 1.04 (3H, s), 1.32 (3H, s), 2.83 (1H, dd, $J=17, 6.5$ Hz), 2.84 (1H, d, $J=12.5$ Hz), 3.09 (1H, dd, $J=17, 1.5$ Hz), 3.18 (3H, s), 3.23 (1H, d, $J=12.5$ Hz), 3.35 (6H, s), and 3.77 (1H, dd, $J=6.5, 1.5$ Hz) for **22**. $^{13}\text{C NMR}$ δ = 19.8, 20.2, 23.3, 23.6, 40.2, 42.4, 42.9, 48.4, 49.5, 50.4, 54.2, 58.7, 76.4, 102.5, 202.0, and 209.6. IR ν : 1740, 1710, 1110, and 755 cm^{-1}] and **23** [a pale yellow oil, 11.1 mg, 6%. Found: m/z , 298.1761 (M^+). $^1\text{H NMR}$ δ = 1.14 (6H, s), 1.72 (3H, d, $J=1.5$ Hz), 2.34 (1H, dd, $J=15.5, 4$ Hz), 2.72 (1H, d, $J=12$ Hz), 2.8–2.95 (3H, m), 3.29 (3H, s), 3.38 (3H, s), 3.39 (3H, s), 3.61 (1H, dd, $J=6, 5$ Hz), 4.76 (1H, s), and 4.82 (1H, q, $J=1.5$ Hz). $^{13}\text{C NMR}$ δ = 19.4, 22.5, 25.2, 38.8, 41.8, 45.8, 50.8, 51.0, 52.8, 58.4, 76–78.5 (1C, underneath CDCl_3), 103.5, 112.1, 150.2, 204.3, and 209.0. IR ν : 1720, 1640, 1100, and 900 cm^{-1}] were obtained.

Hydrolysis of a Mixture of 21 and 22. Formation of 24. A mixture of an acetone solution (2 cm^3) of **21** and **22** (**21:22** = 3:2, 20 mg) was stirred with 2 M perchloric acid (2 cm^3) at room temperature for 3 d. After a silica-gel chromatography (AcOEt:hexane = 1:5) of the reaction mixture, **24** [a pale yellow oil, 3 mg; 26%. Found: m/z , 284.1622 (M^+). Calcd for $C_{15}H_{24}O_5$: 284.1622. $^1\text{H NMR}$ δ = 0.99 (3H, s), 1.02 (3H, s), 1.11 (3H, s), 1.20 (3H, s), 2.4–3.1 (4H, m), 3.31 (3H, s), 3.35 (3H, s), and 4.1–4.3 (1H, m). IR ν : 3450, 1710, 1380, and 1050 cm^{-1}] and unreacted **22** [6 mg].

References

- 1) Part XXXIX. A. Mori, H. Suizu, and H. Takeshita, *Bull. Chem. Soc. Jpn.*, **60**, 3817 (1987).
- 2) H. Takeshita, A. Mori, and Y. Isayama, *Bull. Chem. Soc. Jpn.*, **58**, 1678 (1985).
- 3) A. Mori, Y. Isayama, and H. Takeshita, *Bull. Chem. Soc. Jpn.*, **59**, 511 (1986).
- 4) A. Mori, T. Kubota, S. Kasai, and H. Takeshita, *Chem. Lett.*, **1987**, 1931.
- 5) M. Hirama, Y. Koyama, Y. Shoji, and S. Itô, *Tetrahedron Lett.*, **1978**, 2289.
- 6) A. Mori and H. Takeshita, *Kyushu Daigaku Sogo Rikogaku Kenkyuka Hokoku*, **3**, 125 (1981).
- 7) H. Takeshita, T. Kusaba, and A. Mori, *Chem. Lett.*, **1982**, 701.
- 8) K. Shima and H. Sakurai, *Bull. Chem. Soc. Jpn.*, **42**, 849 (1969).
- 9) G. Jones, II, "Organic Photochemistry", Marcel Dekker Inc., 1981, Vol. 5, pp. 1-122.
- 10) The coupling constants between the protons on the ring junction were reported to be $J_{\text{cis}} = 8$ Hz and $J_{\text{trans}} = 10$ Hz in cis- and trans-fused bicyclo[5.2.0]nonane systems; See E. Dunkelblum, H. Hart, and M. Suzuki, *J. Am. Chem. Soc.*, **99**, 5074 (1977).