Chloroplastic Glucose 6-Phosphate Dehydrogenase from *Chlorella vulgaris* Alleviates Freezing and Menadione-Induced Oxidative Stresses in *Saccharomyces cerevisiae*

Machida, Takeshi

Laboratory of Food Hygienic Chemistry, Division of Food Biotechnology, Department of Bioscience and Biotechnology, Faculty of Agriculture, Graduate School, Kyushu University

Ohashi, Naoto

Laboratory of Food Hygienic Chemistry, Division of Food Biotechnology, Department of Bioscience and Biotechnology, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

Mimura, Ayano

Laboratory of Food Hygienic Chemistry, Division of Food Biotechnology, Department of Bioscience and Biotechnology, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

Honjo, Ken-ichi

Laboratory of Food Hygienic Chemistry, Division of Food Biotechnology, Department of Bioscience and Biotechnology, Faculty of Agriculture, Graduate School, Kyushu University

他

<https://doi.org/10.5109/17797>

出版情報 : 九州大学大学院農学研究院紀要. 55 (1), pp.29-38, 2010-02-26. Faculty of Agriculture, Kyushu University

バージョン :

権利関係 :

Chloroplastic Glucose 6-Phosphate Dehydrogenase from *Chlorella vulgaris* Alleviates Freezing and Menadione-Induced Oxidative Stresses in *Saccharomyces cerevisiae*

Takeshi MACHIDA, Naoto OHASHI¹, Ayano MIMURA¹, Ken-ichi HONJOH*, Masayoshi IIO and Takahisa MIYAMOTO

Laboratory of Food Hygienic Chemistry, Division of Food Biotechnology,
Department of Bioscience and Biotechnology, Faculty of Agriculture,
Graduate School, Kyushu University, Fukuoka 812–8581, Japan
(Received October 29, 2009 and accepted November 19, 2009)

Enhanced glucose 6-phosphate dehydrogenase (E.C.1.1.1.49, G6PDH) activity has been identified as a hardening-induced intracellular change of *Chlorella vulgaris*, which acquires freezing tolerance during hardening. In the present study, a full-length cDNA clone corresponding to a gene encoding a chloroplastic isoform of G6PDH, designated *CvchG6PDH*, was isolated from *C. vulgaris* C-27. By comparing the deduced amino acid sequence of *CvchG6PDH* with the N-terminal amino acid sequence of mature G6PDH₂ protein isolated previously, a DNA region encoding mature *CvchG6PDH* was determined and designated *mCvchG6PDH*. The deduced amino acid sequence of *CvchG6PDH* showed higher homology to those of plant plastidic G6PDH genes than those of cytosolic ones. A recombinant mCvchG6PDH protein expressed in *Escherichia coli* showed similar enzymatic properties to previously-isolated *Chlorella* G6PDH₂, suggesting that the gene encoded plastidic G6PDH₂ protein. Expression of *CvchG6PDH* was induced transcriptionally throughout 24-h hardening, while the translation was induced up to 9-h hardening and then decreased, and the change did not reflect the enhanced G6PDH activity during hardening. Furthermore, the mCvchG6PDH alleviated both freezing and menadione-induced oxidative stresses in yeast. We showed the contribution of *CvchG6PDH* in menadione stress tolerance as one of its functions in the acquisition of freezing tolerance of *Chlorella*.

INTRODUCTION

Freezing injury is one of the most severe constraints limiting plant growth and it has been a major preoccupation in northern latitude (Levitt, 1980). It could be said that to investigate the developing mechanisms of plants against freezing is very important for the purpose of stable food production and establishment of the method for long-term food storage. Some plants acquire freezing tolerance when exposed to non-freezing low temperatures, and their physiological strategies used to tolerate freezing stress vary (Sadakane *et al.*, 1980; Sarhan and Perras, 1987; Thomashow, 1990; Uemura *et al.*, 1997).

We have studied the acquisition of freezing tolerance of plants, using *Chlorella vulgaris* C-27 as a convenient eukaryotic model. Hardened cells of *C. vulgaris* C-27 can survive after freezing treatment even at –196 °C (Hatano *et al.*, 1976). In our previous studies, we have found an increase in the activity of glucose 6-phosphate dehydrogenase (G6PDH) as an intracellular change of *Chlorella* during hardening (Sadakane *et al.*, 1980). The enzyme, G6PDH (EC 1.1.1.49), catalyses the rate-limiting step of the oxidative pentose phosphate pathway (OPPP), an important catabolic route for the provision of NADPH and sugar phosphates (Wendt *et al.*, 2000). In plants, G6PDH isoenzymes reside in two com-

partments, cytosol and plastids (Schnarrenberger *et al.*, 1973). The plastidic G6PDH is known to be reductively inactivated, achieved via reversible dithiol–disulphide interchange of two highly conserved regulatory cysteine residues specific to the plastidic one (Wenderoth *et al.*, 1997). The main role of G6PDH in plastids is the supply of reducing equivalents in the form of NADPH required for various reductive biosyntheses (e.g. amino acids, fatty acids) and to provide sugar-phosphate intermediates that serve as precursors of nucleotides and secondary plant products. In plants, substantial evidences show that G6PDH might be involved in responses to stresses, such as oxidative stress (Hauschild and von Schaewen, 2003; Debnam *et al.*, 2004) and salt stress (Nemoto and Sasakuma, 2000).

A number of factors have appeared to be involved in freezing injury of plants, including water deficit due to ice formation, lipid phase separation, and inactivation of ATPase in plasma membrane (Tao *et al.*, 1998). In addition, production of reactive oxygen species (ROS) during freezing–thawing may be involved. In research on coniferous plant, increases in superoxide dismutase activity and ascorbate content have been found when suffering photooxidative damage in fall and winter (Tao *et al.*, 1990; 1992). Many researchers have studied the relationship between ROS-scavengers and freezing tolerance, and clarified the importance and the role in the acquisition of freezing tolerance (Seppanen and Fagerstedt, 2000; Kocsy *et al.*, 2001; Yang *et al.*, 2007). Those reports suggest that protection system against ROS-mediating oxidative damage plays an important role in improvement of freezing tolerance of plants.

¹ Laboratory of Food Hygienic Chemistry, Division of Food Biotechnology, Department of Bioscience and Biotechnology, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

* Corresponding author (E-mail: honjoh@agr.kyushu-u.ac.jp)

We have previously purified and characterized two isoforms of G6PDHs from *Chlorella*. The two G6PDHs appeared to be the cytosolic and the chloroplastic enzymes based on their sensitivities to DTT (Honjoh *et al.*, 2003). Furthermore, we have isolated and characterized a cDNA encoding cytosolic G6PDH, and investigated the involvement in the acquisition of freezing tolerance using transformed yeast (Honjoh *et al.*, 2007). Although the gene encoding cytosolic G6PDH, designated *CvcG6PDH*, was induced at a transcriptional level during hardening, leading to the change in the activity, it did not reflect the increase in the activity of G6PDH during hardening. The previous study also showed that the freeze-thaw stress tolerance of the transformed yeast expressing *CvcG6PDH* was not improved sufficiently in spite of the 8.7-fold high activity of G6PDH compared to that of the control strain. Thus, involvement of other isoform in freezing tolerance should be investigated.

In the present study, we have isolated and characterized a cDNA for chloroplastic *G6PDH* gene from *C. vulgaris* C-27. The effect of low temperatures, responsible for the freezing tolerance of *Chlorella*, on the expression of the gene was investigated at transcriptional and post-transcriptional levels. Furthermore, the involvements of two isoforms of *Chlorella* G6PDH in the acquisition of stress tolerance were investigated using transformed yeast expressing the gene.

MATERIALS AND METHODS

Strains and conditions

Chlorella vulgaris Beijerinck IAM C-27 was obtained from the Algal Culture Collection of the Institute of Applied Microbiology, Tokyo University, Japan. Cells of *C. vulgaris* C-27 were cultured at 25 °C synchronously and hardened at 3 °C as described previously (Honjoh *et al.*, 2007).

Escherichia coli BL21(DE3)pLysS (Novagen, Madison, WI, USA) was used for production of a large amount of recombinant mCvchG6PDH protein in *E. coli*.

Saccharomyces cerevisiae INVSc2 (*Mat* α , *his3* Δ 200, *ura3*-167; Invitrogen, Carlsbad, CA, USA) was used for expression of *mCvchG6PDH* in yeast.

Isolation of a full-length cDNA clone encoding CvchG6PDH and sequence analysis

Hardened cells of *Chlorella* were harvested, washed twice with sterilized water, and suspended in an extraction buffer, which contained 0.1 M Tris-HCl (pH 7.0), 0.1 M NaCl, 10 mM EDTA, and 0.1% SDS, dissolved in diethylpyrocarbonate-treated water. Purification of poly(A) + RNA and cDNA synthesis were performed based on the method of Machida *et al.* (2009). A partial *CvchG6PDH* cDNA fragment was amplified using two degenerated primers, 5'- GAT GCT TTT A(C/T)C ATT C(C/G/T)G GTC AGT A(C/T)A A -3' and 5'- CTC (C/T) TT (A/G)CC CA(A/G) GTA (A/G)TG GTC (A/G)AT -3', designed based on conserved regions of G6PDHs. DNA regions that extended 5'- and 3'-ends of the full-length *CvchG6PDH* cDNA were amplified using rapid ampli-

cation of cDNA ends (RACE) method. Full-length *CvchG6PDH* cDNA was amplified with two primers, 5'- GGC CCT TTT CGT TGT TGA GCC TG -3' and 5'- CCA GGC AAC AGC AAT TAC AGG GGA -3'. Amplified fragments were subcloned into a pGEM-T easy vector (Promega, Madison, WI, USA), and nucleotide sequences of the fragments were determined based on the method of Sanger *et al.* (1977).

Homologous genes to the *CvchG6PDH* were searched for using the tBlastX program (Altschul *et al.*, 1990) on the National Center of Biotechnology Information web site (<http://www.ncbi.nlm.nih.gov/>), and the identity (%) was calculated using the clustal W program (<http://align.genome.jp/>). Phylogenetic tree was constructed with the Neighbor-Joining method based on the method of Wendt *et al.* (1999). Subcellular localization of the protein encoded by the cloned cDNA was predicted using the WoLF PSORT program (Houton *et al.*, 2006).

Northern blot analysis

Total RNAs from unhardened, 3, 9, and 24-h hardened cells were isolated as described above. Ten micrograms of total RNA were separated on a 1.0% agarose gel containing 1.8% formaldehyde, and blotted onto a Hybond-N⁺ nylon membrane (GE healthcare, Heidelberg, Germany). The membrane was subjected to Northern blot analysis using a DIG High Prime DNA Labeling and Detection Starter Kit II (Roche, Basel, Switzerland). The full-length *CvchG6PDH* cDNA was labeled by the random-labeling method with DIG-dUTP, and then used as a probe. The hybridized bands were detected using a chemiluminescent substrate, [3-(5-Chloro-4'-methoxyspiro[adamantane-2,3'-[1,2]dioxetan]-4'-yl)phenyl] phosphate phosphoric acid 3-[4'-methoxy-5-chlorospiro [tricyclo[3.3.1.1^{3,7}]decane-2,3'-[1,2]dioxetane]-4'-yl]phenyl ester, supplied in the kit. The loading amount of total RNA was standardized by comparison with the electrophoretic band intensity of rRNA.

Expression of mCvchG6PDH in *E. coli*

A gene encoding a mature form of CvchG6PDH, designated *mCvchG6PDH*, was amplified by PCR using the full-length cDNA as a template. To amplify *mCvchG6PDH*, two primers with the recognition sites of *NheI* and *XhoI* were designed (5'- TAT GCT AGC GGG TTG CAA GAG GAG AAC TGG -3' and 5'- GGC CTC GAG TCA CTC GTC GTC AGC G -3'; recognition sites of *NheI* and *XhoI* are underlined). The amplified DNA fragment was digested with the corresponding enzymes and ligated into the *NheI*-*XhoI* site of a pET-17xb vector (Novagen). *E. coli* BL21(DE3)pLysS was transformed with the plasmid by the electroporation method according to the manufacturer's instructions (BTX Division of Genetronics, Inc., Holliston, MA, USA). The transformants were selected by cultivation on LB/Amp/Cam agar containing ampicillin (100 μ g/ml) and chloramphenicol (34 μ g/ml) at 37 °C for 16 h.

The *mCvchG6PDH* was separately amplified by PCR to be ligated into a pET-32a(+) vector (Novagen) with two primers, 5'- TAT GGA TCC GGG TTG CAA GAG GAG

AAC TGG -3' and 5'- CGG CGT CGA CTC ACT CGT CGT CAG CG -3' containing *Bam*HI and *Sal*I recognition sites (underlined). The amplified DNA fragment was digested and incorporated into the *Bam*HI-*Sal*I site of the vector. The plasmid generated was introduced into *E. coli* BL21(DE3)pLysS by the electroporation method. The transformants were selected by cultivation on LB/Amp/Cam agar plates at 37 °C for 16 h.

Expression of *mCvchG6PDH* in *E. coli* and purification of His-tagged *mCvchG6PDH* protein

The *E. coli* transformants were cultured in LB/Amp/Cam medium at 37 °C with shaking. When the OD₆₀₀ reached 0.5, isopropyl- β -D-thiogalactopyranoside (IPTG) was added to the culture at a final concentration of 1 mM. After 6 h of cultivation at 25 °C, the cells were collected by centrifugation, then resuspended in 50 mM potassium-phosphate buffer (pH 7.5; K-Pi) containing 1 mM phenylmethylsulfonyl fluoride (PMSF), and disrupted at 4 °C by sonication using a Tomy Ultrasonic Disrupter UP-201 (Tomy Seiko, Tokyo, Japan) for 10 min at 48 W with 0.5-s pulses at 0.5-s intervals. The suspension was centrifuged at 20,000 $\times g$ for 10 min at 4 °C and the resulting supernatant was used for further experiments. His-tagged *mCvchG6PDH* was purified by Ni²⁺-affinity chromatography using a His-Bind resin (Novagen) according to the manufacturer's instructions. Expression and purification of the proteins were checked by SDS-PAGE on a 10% polyacrylamide gel by the method of Laemmli (1970).

Preparative SDS-PAGE and N-terminal amino acid sequencing

The partially-purified His-tagged *mCvchG6PDH* protein was treated with enterokinase at 23 °C for 16 h or thrombin (Novagen) at 25 °C for 16 h to remove tagging peptides. And then, the digested proteins were purified additionally by preparative electrophoresis. The purified *mCvchG6PDH* was subjected to SDS-PAGE and electroblotted onto a polyvinylidene difluoride membrane (Towbin *et al.*, 1970). The N-terminal sequence of the detected protein was analyzed with a gas-phase protein sequence analyzer (model PPSQ-21; Shimadzu, Kyoto, Japan). Rabbit anti-*mCvchG6PDH* polyclonal antibodies were raised using the purified *mCvchG6PDH* as an antigen.

Enzyme assays

G6PDH activity was assayed in a double-beam mode using a Shimadzu UV-visible recording spectrophotometer (Shimadzu) as described previously (Honjoh *et al.*, 2003). The assay mixture contained 55 mM Tris-HCl (pH 7.8), 3.3 mM MgCl₂, 0.2 mM NADP⁺, 3.3 mM glucose 6-phosphate, and protein extract. One unit of activity was defined as the amount of enzyme forming 1.0 μ mol of NADPH per minute at 25 °C.

Dithiothreitol (DTT)-sensitivity test was performed as follows. Protein samples were incubated on ice for 3 h in the presence of 100 mM DTT. Residual activity (%) was expressed as the percentage of specific activity of

treated sample against that of untreated sample.

Protein extraction from *Chlorella*

Chlorella cells were harvested, washed twice, and suspended in 50 mM K-Pi containing 1 mM PMSF. The suspension was frozen in liquid nitrogen and stored at -80 °C until use. The frozen suspension of cells was thawed at 4 °C and homogenized with 0.5 mm diameter glass beads on a reciprocal shaker (Vibrogen-Zellmühle, Edmund Bühler Co., Tübingen, Germany), operated at 4,500 rpm at 4 °C for 20 min. The homogenate was freed from the beads by passage through a sintered-glass funnel and then centrifuged at 20,000 $\times g$ for 30 min at 4 °C. The upper phase was recovered and centrifuged again at 20,000 $\times g$ for 60 min at 4 °C. The resulting upper phase was used in further experiments as a *Chlorella* cell extract.

Western blot analysis

Protein extracts of unhardened, 3, 9, and 24-h hardened *Chlorella* cells were separated by SDS-PAGE, and then electroblotted onto a nitrocellulose membrane. The membrane was blocked with 3% skimmed milk in TBS containing 0.1% Tween-20 (TTBS) for 1 h, incubated with antibodies for *CvchG6PDH* at a 1:500 (v/v) dilution in TTBS containing 3% skimmed milk for 1 h, washed four times with TTBS, and then incubated with anti-rabbit IgG (whole molecule)-peroxidase (Sigma-aldrich, St. Louis, MO, USA) at a 1:500 (v/v) dilution in TTBS containing 3% skimmed milk for 1 h. After washing the membrane four times with TTBS, bands were detected with 3,3'-diaminobenzidine tetrahydrochloride and hydrogen peroxide. All procedures in this section were performed at room temperature.

Quantification of band intensity

The bands shown in the results of Northern blot and Western blot were quantified using the Yab's Gel Image software (<http://homepage.mac.com/yabyab/rb/gelimage-j.html>).

Plasmid construction and transformation of yeast

To express the *mCvchG6PDH* genes in *S. cerevisiae*, the *mCvchG6PDH* was amplified by PCR using Phusion DNA polymerase (Finnzymes, Keilaranta, Finland). To amplify *mCvchG6PDH*, two oligonucleotides with the recognition sites of *Bgl*II and *Eco*RI were designed (5'-TAT AGA TCT ATG GGG TTG CAA GAG GAG AAC -3' and 5'-CGG CGA ATT CTC ACT CGT CGT CAG CGG -3'; recognition sites of *Bgl*II and *Eco*RI are underlined). The amplified DNA fragment was digested with the corresponding enzymes and introduced into the *Bgl*II-*Eco*RI site of a pTG887 vector (Transgene S. A., France), downstream of the *PGK* promoter to allow a high level of expression of introduced genes. The plasmid generated was designated pTG887/*mCvchG6PDH*.

S. cerevisiae INVSc2 was transformed with the plasmids by the lithium acetate method (Ito *et al.*, 1983). Transformants were screened onto an uracil-deficient synthetic raffinose (SR-ura) agar plate, which contained

0.67% yeast nitrogen base, 2% raffinose, 0.2% complete supplement mixture without uracil (CSM-URA, BIO 101, Vista, CA, USA), and 2% agar, at 30 °C for 48 h.

Expression of the introduced genes in yeast

The transformed yeast cells were cultured overnight in SR-ura medium with shaking at 30 °C. The overnight cultures were inoculated into SR-ura containing 2% glucose to express the introduced genes. After 24 h of cultivation at 30 °C, the yeast cells were harvested by centrifugation at $1,000 \times g$ for 10 min, and suspended in 10 mM Tris-HCl (pH 7.5) containing 1 mM PMSF. The cell suspension was mixed with an equal volume of 0.5 mm diameter glass beads and homogenized with a vortex mixer. The homogenates were centrifuged at $20,000 \times g$ at 4 °C for 10 min. G6PDH activity in yeast was determined using the soluble lysate by the method described above.

Freezing tolerance test

Yeast cells expressing *mCvchG6PDH* were pre-cultured in 2 ml of SR-ura medium with shaking at 30 °C for 48 h. Five microliters of pre-cultured cells were inoculated into 5 ml of SR-ura medium containing 2% glucose. When the OD₆₀₀ reached 1.0, the yeast cells were harvested by centrifugation at $560 \times g$ for 5 min at room temperature, and resuspended in a 0.9% NaCl solution at a cell concentration of $2-3 \times 10^6$ cells/ml (OD₆₀₀ = 0.1). One hundred microliters of the cell suspensions were transferred to 1.5-ml microcentrifuge tubes and pre-chilled in a deep-freezer at -80 °C for 2 min for ice formation. Then, the samples were subjected to freezing treatment in a program freezer at -20 °C for 24 h. After freezing treatment, the samples were thawed at 25 °C, serially diluted with 0.9% NaCl, and immediately plated onto YPD agar plates, containing 1% yeast extract, 2% polypeptone, 2% glucose, and 1.5% agar. After 2 d of incubation at 30 °C, colonies were counted. Viability (%) was calculated by dividing the counts of frozen sample by those of unfrozen one. Yeast cells carrying pTG887 and pTG887/*CvcG6PDH* (Honjoh *et al.*, 2007) were also tested.

Oxidative stress tolerance test

The pre-cultured transformants were inoculated into 5 ml of SR-ura medium containing 2% glucose, and cultured at 30 °C with shaking until OD₆₀₀ reached 0.5. Then, the cells were harvested by centrifugation at $560 \times g$ for 5 min at room temperature, washed twice with 100 mM sodium-phosphate buffer (pH 7.0, Na-Pi), and resuspended in Na-Pi at a cell concentration of $2-3 \times 10^6$ cells/ml (OD₆₀₀ = 0.1). One milliliter of the cell suspensions was mixed with menadione, which induces production of superoxide anion inside the cells, at a final concentration of 200 μ M and then incubated at 30 °C for 1 h with shaking. After stress treatment, the samples were serially diluted with Na-Pi and immediately plated onto YPD agar plates. After 2 d of incubation at 30 °C, the colonies were counted. Viability (%) was calculated by dividing the counts of frozen sample by those of unfro-

zen one. Yeast cells carrying pTG887 and pTG887/*CvcG6PDH* were also tested.

Statistical analysis

Statistical analysis was performed using Excel 2004 (Microsoft, USA) with the add-in software Statcel 2 (Yanai, 2002). The Tukey-Kramer test was used to evaluate significance among the groups by one-way ANOVA.

RESULTS

Isolation of full-length cDNA encoding *CvchG6PDH*

A 309 bp *CvchG6PDH* DNA fragment was amplified by PCR using degenerated primers. Nucleotide sequence of a full-length cDNA clone of *CvchG6PDH* gene was determined using 5'- and 3'-RACE procedures based on the sequence of the 309-bp fragment. The full-length cDNA fragment was amplified by PCR, subcloned into a pGEM-T easy vector, and sequenced. The length of *CvchG6PDH* was 2,397 bp in length and contained an open reading frame encoding 598 amino acid residues. A putative active region (-RIDHYLGKE-) and two coenzyme-binding sites (-GXXGXXG/A-) (Cosgrove *et al.*, 1998) were found in the deduced amino acid sequence of *CvchG6PDH* (Fig. 1).

A computer search of the databases using the Clustal W program revealed that the deduced amino acid sequence of the obtained cDNA exhibited 58–59% identity to those of the plastidic G6PDHs from higher plants and 41–44% identity to those of the cytosolic G6PDHs (Table 1). Phylogenetic analysis was performed to estimate the subcellular localization of *CvchG6PDH* based on the deduced amino acid sequence (Fig. 2). Moreover, the result of analysis using the WoLF PSORT program also indicated that the encoded protein is targeted or localized in the chloroplast. The N-terminal (NH₂-GLQEENWEKAALSIV-) and internal (-LVIRIQPNEGIIYLKVNKNKVP-) amino acid sequences of chloroplastic G6PDH protein (G6PDH_c, Honjoh *et al.*, 2003) were found in the deduced amino acid sequence

Table 1. Sequence homology between the deduced amino acid sequence of *CvchG6PDH* and those of other cytosolic and plastidic G6PDH genes as given by Clustal W program

Origin	Identity (%)	Accession number
Plastidic G6PDH		
<i>Solanum tuberosum</i>	58.9	AJ010712
<i>Dunaliella bioculata</i>	58.6	AJ132346
<i>Arabidopsis thaliana</i>	58.5	AJ001359
<i>Nicotiana tabacum</i>	58.5	AJ001772
<i>Spinacia oleacea</i>	57.6	AJ000182
<i>Petroselinum crispum</i>	56.5	AF012861
Cytosolic G6PDH		
<i>A. thaliana</i>	43.9	AJ010971
<i>S. tuberosum</i>	43.4	X74421
<i>N. tabacum</i>	43.4	AJ001770
<i>Oryza sativa</i>	42.8	AY078072
<i>Chlorella vulgaris</i>	40.9	AB085846

1 GGCCCTTTTCGTTGTTGAGCCTGCTTTTCATTCTTGCCTTTGTCAACCCAGCCACCAGGGGTGTTGCTACTATCAGCGGGCGCACGTTTC

91 TTGAGCAGCGATCGGGGCGCGCAGCAGGCGGACCCGTGATGCCCGCTCTGCACAGCATGAGGCTCGTTGCACAAGTTCTGACCCTCAGCG
M P A L H S M R L V A Q V L T L S

181 GCAAGGTGGCGCAACGCCATCGCGCTGACCATCAACCCAGCGAGCTGTCTGCGCAGCCTGCCGACAGCACCAGGCTGCCGACAC
G K V A Q R H A A D H Q P Q R A V C R Q P A R Q H H R L R A

271 TGACAGCACAAAGCAGTGCGCCATCGGCATCAACCACTGGCAGCGGCGCAGCGACCACTCAGAGTCCAGAGATTGAAGCAGCGGAGGTGG
L T A Q S S A P S A S T T G S G A A T T Q S P E I E A A E V

361 AGGCAGCAGCAGCAGCAGCAGCTCAAAATGGCTCGCCTTCTTTCTCAAAATATGAAGGTGTATGGGTGCAAGAGGAGAACTGGGAGAAGG
E A A A A D A R Q N G S P S F S N M K V Y **G L Q E E N W E K**

451 CTGCACTGAGCATCGTGGTGGGCGCGTCTGGTGACCTGGCCAAGAAGAAATCTTCCCTCCCTCTTTGCGCTGTACTATGAAAAACA
A A L S I V V G A S G D L A K K K I F P S L F A L Y Y E N

541 TGCTTCCGAGAAGTTCAAGGTGTATGGCTACGCCCGCAGCAAGATGAACGACGAGGAGTTCGAGACCTGATTGCCGCGCAGCTGACCT
M L P Q N F K V Y G Y A R S K M N D E E F R D L I A G S L T

631 GCAGGCTGACGACGCTGGTGACTGCGGCAAGAGATGGACGAGTTCTTGGAGCGCTGCTTACCAGGCGGGCAGTACGATCTGAGG
C R L N D A G D C G K K M D E F L E R C F Y Q A G Y A S E

721 ACGACTTTCGCGCGCTGGACAAGCGCATGGCTGACGGGGAGAGCGAGCAGTCTGCGCAGACAGAATGTTCTACCTCTCGATCCCGCCAA
D D F A A L D K R M A D G E S E Q S C A D R M F Y L S I P P

811 ACATCTTCAACCGGTGGCAGCTGCGCCTCCAAGGCAGCATCCAGCAAGTGTGGCTGGACGCGCATGATTGTTGAGAAGCCGTTTGCA
N I F T T V A A C A S K A A S S K C G W T R M I V E K P L G

901 AGGACAGCAGTTCGTTCCAGGAGTTGAGCGCTGCGCTGTACGAGCAGCTTCGAGAGGACCAGATATACCGCATTGACCACTACCTGGGCA
K D S E S F Q E L S A A L Y E H L R E D Q I Y R I D H Y L G

991 AGGAGCTGATCGAGAAGCTGACGCTGCTGCGCTTTCAAACCTGGTGTGTGAGCGCTGTGGAGCAGGCAGTACATCAGAAACCTGCGAGG
K E L I E N L T V L R F A N L V F E P L W S R Q Y I R N V Q

1081 TCATCTTCAGCGAGAACTTTGGGACAGAGGGCCGAGGGGGCTACTTTGACCAAGTACGGCATTGTGAGGAGCGTCACAAACCCACCTGTC
V I F S E N F G T E G R G G Y F D Q Y G I V R D V I Q N H L

1171 TCCAAATACTGGCCCTCTTCGCCATGGAGCAGCGGCCAGCCTGGATGCCGAGGACATTTCGTAACGAGAAGGTCAAGGTGCTGAAGAGCA
L Q I L A L F A M E Q P A S L D A E D I R N E K V K V L K S

1261 TGGGCCAGTTCGCGCTGGAGGACATGGTGTGGGCGAGTACCGCTCAGCACCACACGCGGGACACGCTGCCAGGATACCTGGGACGAGC
M G Q V R L E D M V L G Q Y R S R T T R G T T L P G Y L D D

1351 ACACGGTACCGCCAAACAGCATCACCCACCTTTGCCGCTGCTCCGTGTTTCATCAACAATGCCAGGTGGGACGCGCTGCCCTTCCTGC
D T V P P N S I T P T F A A C S V F I N N A R W D G V P F L

1441 TCAAGCGCGGCAAGGCGCTGGCCAACAAGCGCGGAGATCCGGGTGCAAGTTCGGCATGTGCCGGCAACCTGTACCGCAACAACTGG
L K A G K A L A N K A A E I R V Q F R H V P G N L Y R N K L

1531 GGCTGGACCTGGACAAGGCCCAACGAGCTGGTCATCCGCGTCCAGCCCAACGAGGGCATCTACCTGAAAGTGAACAACAGGTGCCGG
G L D L D K A T N K **L V I R V Q P N E G I Y L K V N N K V P**

1621 GGCTGGGCTGCGCATAGACACAACGCGCTCGACCTACCTACAAGTCCAAGTACCAGGCAACACTGCCAGATGCATATGAGCGGCTGA
G L G L R I D T T R L D L T Y K S K Y Q A T L P D A Y E R L

1711 TCCTGGAGTGCATCAACGGCGACCGCGCTGTTTCATCCGCAACGACGAGCTGGAGGTGGCCTGGGAGAAAGTTCACTCCGCTGCTCAAGG
I L D C I N G D R R L F I R N D E L E V A W E K F T P V L K

1801 AGATTGAGGATCGGGGTGTCAGCCGAGCTGTACCCCTACGGCAGCGCGGCCCGCTTGGCGCGCACTACCTGGCAGCCAAGCAGCGCG
E I E D R G V Q P E L Y P Y G S R G P V G A H Y L A A K H G

1891 TGCGGTGGGGGACCTGGCCGCTGACGACGAGTGAGCCGCCCGCTGGCTGGCAACACAAGTGCCTGCCAGTCCATTGTCCACAAACCC
V R W G D L A A D D E *

1981 CGTCCCTCTCTCCAAACACTTGACCTCTACGCTCCGGCTGGCATGCCACGCAACTTCATTTCGACGCCGCCCTTTGACGACGCCCC

2071 ATCCAAGGCATAAACTGTACGCGCTATCGCCACACTGCTGGCCGATGCCCCGTACGCGCCCGGCTGCCGACGCGGTTGGCACACCCACG

2161 TCCTGTACACCACACTACACATACCTCCTCTGCCTTCCTTACCCCTGTTTAGCTGCTGTTTAGGTGCTGCTGCTGCACTTTTGGTTGCC

2251 CCACTGCGCATGGTGCAGCGGTCACACGACAGGCCACACACCTGTGACCTTGTACCGCTGTTCTGTCCCTCTCTCGCCCTTAGC

2341 TGACCTTAGCTGACACCATGTACCAACCCATGTCCCTGTAATTGCTGTTGCTGCTG

Fig. 1. Nucleotide and deduced amino acid sequences of the cDNA of *CvchG6PDH* are shown. The putative active site (–RIDHYLGKE–) of G6PDH is double-underlined and two NADP-binding sites (–GXXGXXG/A–) are boxed. The two regions that were identical to the internal amino acid sequences of purified G6PDH₂ (Honjoh *et al.*, 2003) are shown in bold letters. Two conserved cysteine residues found in common plastidic G6PDH are circled.

Fig. 2. Phylogenetic tree of G6PDHs was constructed with the Neighbor-Joining method using the clustal W program. Accession numbers for the sequences are shown in parentheses as follows: *Duna P-G6PDH*: *Dunaliella biocata* plastidic G6PDH (AJ132346), *Petro P-G6PDH*: *Petroselinum crispum* plastidic G6PDH (AF012861), *Sola P-G6PDH*: *Solanum tuberosum* plastidic G6PDH (AJ010712), *Arabi P-G6PDH*: *Arabidopsis thaliana* plastidic G6PDH (AJ001770), *Nico C-G6PDH*: *Nicotiana tabacum* cytosolic G6PDH (AJ001770), *Sola C-G6PDH*: *S. tuberosum* cytosolic G6PDH (X74421), *Arabi C-G6PDH*: *Arabidopsis thaliana* cytosolic G6PDH (AJ010971), *CvcG6PDH*: *Chlorella vulgaris* C-27 cytosolic G6PDH (AB085846).

of *CvchG6PDH* cDNA, and thus the DNA region encoding mature *CvchG6PDH* (*mCvchG6PDH*) was determined. The length of *mCvchG6PDH* was 1,500 bp in length and it encoded 500 amino acid residues with a predicted molecular mass of about 56.6 kDa and an isoelectric point of 5.35. The region containing 98 amino acids from N-terminus of *CvchG6PDH* was determined as a transit peptide region. Based on the results, *CvchG6PDH* was suggested to encode the chloroplastic isoform of the enzyme. The nucleotide sequence of the cloned *CvchG6PDH* cDNA was deposited at DDBJ/EMBL/GenBank under accession number AB331729.

Expression of *mCvchG6PDH* in *E. coli* and G6PDH assays of the expressed protein

To confirm that the protein encoded by *mCvchG6PDH* is functionally active, the gene was introduced into *E. coli* using a pET-17xb vector. According to the result of SDS-PAGE, the *mCvchG6PDH* (56.9 kDa) protein was expressed as a soluble protein when treated with IPTG at 25 °C (Fig. 3A). G6PDH activity and DTT-sensitivity of the *mCvchG6PDH* was examined. An extract of *E. coli* expressing *CvcG6PDH* (Honjoh *et al.*, 2007) was also prepared to compare DTT-sensitivity of *CvcG6PDH* with that of *mCvchG6PDH*. The *mCvch*-

G6PDH protein showed remarkably high G6PDH activity (Table 2), and its activity was inhibited in the presence of DTT, while the activity of *CvcG6PDH* was not inhibited with DTT (Fig. 3B).

Expression and purification of His-tagged *mCvchG6PDH*

To produce anti-*mCvchG6PDH* antibodies, *mCvchG6PDH* gene was expressed in *E. coli* as a His-tagged protein. *E. coli* cells expressing the gene were cultured at 25 °C for 6 h after addition of 1 mM IPTG. Expression, purification, and removal of His-tag region of His-tagged *mCvchG6PDH* protein were confirmed by SDS-PAGE

Table 2. Specific activity of enzyme extracts of *E. coli* transformed with pET-17xb/*CvcG6PDH* and pET-17xb/*mCvchG6PDH*

Plasmid	Specific activity (U/mg protein)
pET-17xb	Not detected
pET-17xb/ <i>CvcG6PDH</i>	19.9 ± 2.7
pET-17xb/ <i>mCvchG6PDH</i>	51.0 ± 2.5

Values are means ± SD obtained from three independent experiments.

Fig. 3. (A) The *mCvchG6PDH* was expressed in *E. coli* with a pET-17xb vector. Lane 1, insoluble protein; lane 2, soluble protein from the transformed *E. coli*. (B) DTT sensitivity of *mCvchG6PDH* was examined using *E. coli* crude extract. Closed symbols: incubation without DTT; (●) *CvcG6PDH*; (■) *mCvchG6PDH*. Open symbols: incubation with 100 mM DTT; (○) *CvcG6PDH*; (□) *mCvchG6PDH*. All values are means ± SD obtained from three independent experiments. (C) The *mCvchG6PDH* was expressed in *E. coli* as a His-tagged protein with a pET-32a(+) vector to purify the expressed protein. Lane 1, purified His-*mCvchG6PDH* by Ni²⁺-affinity chromatography; lane 2, His-*mCvchG6PDH* digested with thrombin; His-*mCvchG6PDH* digested with enterokinase. The arrowheads indicate the bands of the target proteins in each figure.

(Fig. 3C). The N-terminal sequence of the purified protein was determined as NH₂-AMADIGSGLQEENWEK (tagging peptide region was underlined), and the sequence without an underline was identical to the N-terminal sequence of G6PDH₂ identified previously (Honjoh *et al.*, 2003). Then, antibodies against mCvch-G6PDH were raised using the purified protein as an antigen.

Change in the expression pattern of *CvchG6PDH* during hardening

Northern blot analysis of *CvchG6PDH* gene was performed using total RNAs isolated from 0, 3, 9, and 24-h hardened *Chlorella* cells. The transcription of *CvchG6PDH* increased up to 3-h hardening and the high level of transcription continued thereafter (Fig. 4A).

Western blot analysis was performed using protein extracts of cells hardened for 0, 3, 9, and 24 h to investigate change in the CvchG6PDH protein amount in cells during hardening. CvchG6PDH protein was detected with anti-mCvchG6PDH antibodies. The protein amount of CvchG6PDH decreased at the early phase of hardening and then increased, but it did not show dramatic change in the amount (Fig. 4B).

Fig. 4. Hardening-induced change in the expression level of *CvchG6PDH* in *Chlorella* cells was investigated at (A) the transcriptional and (B) the translational levels.

Expression of the *mCvchG6PDH* in yeast

The *mCvchG6PDH* was expressed in yeast with a pTG887 vector. Expression of the genes was confirmed by enzyme assays (Table 3). Specific activity of the extract of INVSc2/pTG887/*mCvchG6PDH* (4.58 ± 0.37 U/mg protein) was remarkably higher than those of control yeast (INVSc2/pTG887; 0.28 ± 0.01 U/mg protein) and INVSc2/pTG887/*CvcG6PDH* (1.01 ± 0.09 U/mg protein).

Table 3. Specific activity of enzyme extracts of yeast transformants expressing *CvcG6PDH* and *mCvchG6PDH*

Plasmid	Specific activity (U/mg protein)
pTG887 (control)	0.28 ± 0.01
pTG887/ <i>CvcG6PDH</i>	$1.01 \pm 0.09^*$
pTG887/ <i>mCvchG6PDH</i>	$4.58 \pm 0.37^{**}$

Values are means $\pm$ SD obtained from three independent experiments. Significance against the value of control yeast is indicated as: $^*p < 0.05$; $^{**}p < 0.01$.

Fig. 5. (A) Freezing and (B) menadione-induced oxidative stress tolerances of yeast expressing *Chlorella* G6PDH genes. Values are means $\pm$ SD obtained from three independent experiments. Significance against the value of control yeast is indicated as: $^*p < 0.01$.

Stress tolerances of the transformed yeast

Effects of mCvchG6PDH on freezing and oxidative stress tolerances were investigated using *S. cerevisiae* INVSc2 as a model eukaryote. As shown in Fig. 5A, the viability of INVSc2/pTG887/mCvchG6PDH ($26.7 \pm 2.2\%$) after freezing and thawing was significantly higher than those of control yeast ($12.7 \pm 1.9\%$), while it was slightly lower than that of INVSc2/pTG887/CvcG6PDH ($29.5 \pm 3.6\%$).

Oxidative stress treatment was performed using menadione as an intracellular ROS generator. As shown in Fig. 5B, the viability of INVSc2/pTG887/mCvchG6PDH ($37.17 \pm 3.33\%$) after 1-h incubation with $200 \mu\text{M}$ menadione was significantly higher than those of control yeast ($4.64 \pm 2.30\%$) and INVSc2/pTG887/CvcG6PDH ($9.47 \pm 1.35\%$).

DISCUSSION

G6PDH is a key-enzyme of the pentose phosphate pathway (Wendt *et al.*, 2000). *C. vulgaris* C-27 cells acquire freezing tolerance after cold-hardening, at which the G6PDH activity increased approximately 1.5-fold (Sadakane *et al.*, 1980). Two types of G6PDH isozymes, which were predicted as cytosolic and chloroplastic G6PDHs, were identified in *C. vulgaris* C-27 and characterized previously (Honjoh *et al.*, 2003). In the present study, we isolated a full-length cDNA clone encoding the chloroplastic isoform, and investigated its involvement and function in freezing tolerance.

A characteristic feature of chloroplastic G6PDH is considered as inactivation by covalent redox modification mediated by the ferredoxin-thioredoxin system (Scheibe, 1990; Buchanan, 1991) or DTT (Honjoh *et al.*, 2003), and this inactivation is due to the reduction of two cysteine residues specific to chloroplastic G6PDH (Wendt *et al.*, 1999). The cysteine residues were also observed in the deduced amino acid sequence of CvcG6PDH as shown in Fig. 1. G6PDH activity in crude extract of *E. coli* expressing mCvchG6PDH was suppressed when the extract was treated with DTT, while that of *E. coli* expressing CvcG6PDH did not show DTT-sensitivity (Fig. 3B). Our results suggested that the CvcG6PDH cDNA isolated in the present study encoded chloroplastic G6PDH in *C. vulgaris* C-27.

The CvchG6PDH protein amount in *C. vulgaris* C-27 did not show similar change to the up-regulation of the total G6PDH activity during hardening (Fig. 4B), although the transcript level of CvcG6PDH increased after 3-h hardening and was kept at a high level throughout 24-h hardening (Fig. 4A). Our previous study showed the change in CvcG6PDH activity during hardening did not reflect the hardening-induced increase in the total G6PDH activity (Honjoh *et al.*, 2007). Those results on expression changes in CvcG6PDH and CvchG6PDH were not sufficient to explain the increase in G6PDH activity of *C. vulgaris* C-27 during hardening. Several higher plants have several genes encoding G6PDH, for example, *Arabidopsis* has two genes encoding cytosolic G6PDH and four genes encoding plastidic G6PDH (Wakao and

Benning, 2005), and tobacco has two genes encoding cytosolic G6PDH and two genes encoding plastidic G6PDH (Wendt *et al.*, 1999). To clarify the involvement of G6PDH in the acquisition of freezing tolerance of *Chlorella*, further isotypes of G6PDH should be searched for and isolated.

The mCvchG6PDH improved freezing tolerance of yeast significantly, and the improvement of freezing tolerance was also observed in yeast expressing CvcG6PDH (Fig. 5A). Since there was no significant difference between freezing tolerances of CvcG6PDH- and mCvchG6PDH-expressing yeasts, the observed effect was considered as common effect of G6PDH isozymes. One of the roles of G6PDH in the acquisition of freezing tolerance is perhaps due to the production of a reducing equivalent NADPH, which serves as an activator of several antioxidant enzymes, such as glutathione reductase, thioredoxin reductase (Carmel-Harel and Storz, 2000), monodehydroascorbate reductase (Foyer *et al.*, 1994), and catalase (Salvemini *et al.*, 1999). Those reports suggest that G6PDH is involved in the acquisition of freezing tolerance by activating antioxidant enzymes that can play a role in the acquisition of freezing tolerance. Freezing-thawing process can generate oxidative stress such as production of reactive oxygen species (O_2^- , H_2O_2 , $\text{OH}\cdot$), reactive nitrogen species (NO , ONOO^-), and lipid peroxides (Carmel-Harel and Storz, 2000; Xin and Browse, 2000; Chatterjee and Gagnon, 2001; Mittler, 2002; Neil *et al.*, 2002). Since the target peroxides of the antioxidant enzymes vary, the activities of such enzymes and the resulting products should be investigated to clarify the function of G6PDH in the acquisition of freezing tolerance of *Chlorella*.

The mCvchG6PDH also improved menadione-induced oxidative stress tolerance of yeast, while the improvement was not significant in CvcG6PDH-expressing yeast (Fig. 5B). McAims *et al.* (2003) have demonstrated that menadione inactivated G6PDH activity when tested using bovine endothelial cells, causing decrease in cellular antioxidant capacity. Thus, the intrinsic G6PDH activity of yeast might be suppressed due to such function of menadione in our studies, and the introduced *Chlorella* G6PDHs may enhance G6PDH activity in yeast cells to maintain cellular reactions under such unfavorable condition in which G6PDH activity was inhibited. Based on the consideration, the difference between the viabilities of the CvcG6PDH- and mCvchG6PDH-expressing strains (Fig. 5B) is perhaps due to the difference between their G6PDH activities (Table 3). Although we could not find any specific function of CvchG6PDH under stress condition, the contribution of enhanced G6PDH activity to alleviation of cellular damages induced by menadione was first identified in the present study. The effect would be important to survive under freezing-thawing stress since the stress is accompanied by several types of oxidative stress as mentioned above. Our data suggest that the one of the functions of G6PDH in the acquisition of freezing tolerance of *Chlorella* is to alleviate oxidative stress generated under low-temperature and freezing-thawing processes.

ACKNOWLEDGMENTS

The authors are grateful to Dr. M. Nakao of Kyushu University for amino acid sequencing. We also thank to Transgene S. A. for use of the pTG887 plasmid and to Mr. Hsu-Ming Wen for preparation of the manuscript.

REFERENCES

- Altschul, S. F., T. L. Madden, A. A. Schäffer, J. Zhang, Z. Zhang, W. Miller and D. J. Lipman 1997 Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.*, **25**: 3389–3402
- Buchanan, B. B. 1991 Regulation of CO₂ assimilation in oxygenic photosynthesis: the ferredoxin/thioredoxin system. *Arch. Biochem. Biophys.*, **288**: 1–9
- Carmel-Harel, O. and G. Storz 2000 Roles of the glutathione- and thioredoxin-dependent reduction systems in the *Escherichia coli* and *Saccharomyces cerevisiae* responses to oxidative stress. *Annu. Rev. Microbiol.*, **54**: 439–461
- Chatterjee, S. and C. Gagnon 2001 Production of reactive oxygen species by spermatozoa undergoing cooling, freezing, and thawing. *Mol. Reprod. Dev.*, **59**: 451–458
- Cosgrove, M. S., C. Naylor, S. Paludan, M. J. Adams and H. R. Levy 1998 On the mechanism of the reaction catalyzed by glucose 6-phosphate dehydrogenase. *Biochemistry*, **37**: 2759–2767
- Debnam, P. M., A. R. Fernie, A. Leisse, A. Golding, C. G. Bowsher, C. Grimshaw, J. S. Knight and M. J. Emes 2004 Altered activity of the P2 isoform of plastidic glucose 6-phosphate dehydrogenase in tobacco (*Nicotiana tabacum* cv. Samsun) causes changes in carbohydrate metabolism and response to oxidative stress in leaves. *Plant J.*, **38**: 49–59
- Esposito, S., E. Carillo and S. Carfagna 1998 Ammonium metabolism stimulation of glucose-6P dehydrogenase and phosphoenolpyruvate carboxylase in young barley roots. *J. Plant Physiol.*, **153**: 61–66
- Foyer, C. H., M. Lelandais and K. J. Kunert 1994 Photooxidative stress in plants. *Physiol. Plant.*, **92**: 696–717
- Hatano, S., H. Sadakane, M. Tutumi and T. Watanabe 1976 Studies on frost hardiness in *Chlorella ellipsoidea* II. Effects of inhibitors of RNA and protein synthesis and surfactants on the process of hardening. *Plant Cell Physiol.*, **17**: 643–651
- Hauschild, R. and A. von Schaewen 2003 Differential regulation of glucose-6-phosphate dehydrogenase isoenzyme activities in potato. *Plant Physiol.*, **133**: 47–62
- Honjoh, K., A. Mimura, E. Kuroiwa, T. Hagiwara, K. Suga, H. Shimizu, R. S. Dubey, T. Miyamoto, S. Hatano and M. Iio 2003 Purification and characterization of two isoforms of glucose 6-phosphate dehydrogenase (G6PDH) from *Chlorella vulgaris* C-27. *Biosci. Biotechnol. Biochem.*, **67**: 1888–1896
- Honjoh, K., T. Machida, T. Hagiwara, K. Suga, M. Yonekura, H. Shimizu, N. Ohashi, T. Miyamoto, S. Hatano and M. Iio 2007 Molecular cloning and characterization of a cDNA for low-temperature-inducible cytosolic glucose 6-phosphate dehydrogenase gene from *Chlorella vulgaris* and expression of the gene in *Saccharomyces cerevisiae*. *Plant Sci.*, **172**: 649–658
- Houton, P., K.-J. Park, T. Obayashi and K. Nakai 2006 “Protein subcellular localization prediction with WoLF PSORT”. *Proceedings of Asian Pacific Bioinformatics Conference, APBC06*: 39–48
- Ito, H., Y. Fukuda, K. Murata and A. Kimura 1983 Transformation of intact yeast cells treated with alkali cations. *J. Bacteriol.*, **153**: 163–168
- Kocsy, G., G. Galiba and C. Brunold 2001 Role of glutathione in adaptation and signaling during chilling and cold acclimation in plants. *Physiol. Plant.*, **113**: 158–164
- Laemmli, U. K. 1970 Cleavage of structural proteins during the assembly of the head of bacteriophage T₄. *Nature*, **227**: 680–685
- Levitt, J. 1980 Chilling, freezing and high temperature stresses, In “Responses of Plants to Environmental Stresses” Vol. 1, 2nd ed., Academic Press, New York, pp. 27–64
- Machida, T., E. Kato, A. Ishibashi, J. Sato, S. Kawasaki, Y. Niimura, K. Honjoh and T. Miyamoto 2009 Expression pattern of a chloroplast NADPH-dependent thioredoxin reductase in *Chlorella vulgaris* during hardening and its interaction with 2-Cys peroxiredoxin. *Biosci. Biotechnol. Biochem.*, **73**: 695–701
- McAims, W. C., R. C. Schaeffer Jr, J. W. Baynes and M. B. Wolf 2003 Menadione causes endothelial barrier failure by a direct effect on intracellular thiols, independent of reactive oxydant production. *Biochim. Biophys. Acta*, **1641**: 43–53
- Mittler, R. 2002 Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.*, **7**: 405–410
- Neil, S., R. Desikan and J. Hancock 2002 Hydrogen peroxide signaling. *Curr. Opin. Plant Biol.*, **5**: 388–395
- Nemoto, Y. and T. Sasakuma 2000 Specific expression of glucose-6-phosphate dehydrogenase (G6PDH) gene by salt stress in wheat (*Triticum aestivum* L.). *Plant Sci.*, **158**: 53–60
- Sadakane, H., K. Kabata, K. Ishibashi, T. Watanabe, and S. Hatano 1980 Studies on frost hardiness in *Chlorella ellipsoidea*. V. The role of glucose and related compounds. *Environ. Exp. Bot.*, **20**: 297–305
- Salvemini, F., A. Franze, A. Iervolino, S. Filosa, S. Salzano and M. V. Ursini 1999 Enhanced glutathione levels and oxidoreductase mediated by increased glucose-6-phosphate dehydrogenase expression. *J. Biol. Chem.*, **274**: 2750–2757
- Sanger, F., S. Nicklen and A. R. Coulson 1977 DNA sequencing with chain-terminating inhibitions. *Proc. Natl. Acad. Sci. USA*, **74**: 403–410
- Sarhan, F. and M. Perras 1987 Accumulation of high molecular weight protein during cold hardening of wheat (*Triticum aestivum* L.). *Plant Cell Physiol.*, **28**: 1173–1179
- Scheibe, R. 1990 Light/dark modulation: regulation of chloroplast metabolism in a new light. *Bot. Acta.*, **103**: 327–334
- Schnarrenberger, C., A. Oeser and N. E. Tolbert 1973 Two isoenzymes each of glucose-6-phosphate dehydrogenase in spinach leaves. *Arch. Biochem. Biophys.*, **154**: 438–448
- Seppanen, M. M. and K. Fagerstedt 2000 The role of superoxide dismutase activity in response to cold acclimation in potato. *Physiol. Plant.*, **108**: 279–285
- Tao, D. L., Y. H. Jin and Y. J. Du 1990 PSII electron transport and superoxide dismutase of Korean pine during winter. *Sci. Silvae. Sin.*, **26**: 289–293
- Tao, D. L., Y. H. Jin, Y. J. Du, Y. Y. Liu, G. Z. Liu and R. Y. Zhan 1992 Responses of organic free-radicals production of over-wintering conifer needles to low temperature and light. *Chin. J. Appl. Ecol.*, **3**: 120–124
- Tao, D. L., D. Oquist and D. Wingsle 1998 Active oxygen scavengers during cold acclimation of scots pine seedlings in relation to freezing tolerance. *Cryobiology*, **37**: 38–45
- Thomashow, M. F. 1998 Molecular genetics of cold acclimation of higher plants. *Adv. Genet.*, **28**: 99–131
- Towbin, H., T. Staehelin and J. Gordon 1979 Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure and some applications. *Proc. Natl. Acad. Sci. USA*, **76**: 4350–4354
- Uemura, M., R. A. Joseph and P. L. Steponkus 1995 Cold acclimation of *Arabidopsis thaliana*. *Plant Physiol.*, **109**: 15–30
- Wakao, S. and C. Benning 2005 Genome-wide analysis of glucose-6-phosphate dehydrogenases in *Arabidopsis*. *Plant J.*, **41**: 243–256
- Wenderoth, I., R. Scheibe and A. von Schaewen 1997 Identification of the cysteine residues involved in redox modification of plant plastidic glucose-6-phosphate dehydrogenase. *J. Biol. Chem.*, **272**: 26985–26990
- Wendt, U. K., R. Hauschild, C. Lange, M. Pietersma, I. Wenderoth and A. von Schaewen 1999 Evidence for functional convergence of redox regulation in G6PDH isoforms of cyanobacteria and higher plants. *Plant Mol. Biol.*, **40**: 487–494

- Wendt, U. K., I. Wenderoth, A. Tegeler and A. von Schaewen 2000 Molecular characterization of a novel glucose-6-phosphate dehydrogenase from potato (*Solanum tuberosum* L.). *Plant J.*, **23**: 723–733
- Xin, Z. and J. Browse 2000 Cold comfort farm: the acclimation of plants to freezing temperatures. *Plant Cell Environ.*, **23**: 893–902
- Yanai, H. 2002 Statcel– The useful add-in software forms on Excel (2nd ed), OMS, Tokyo, Japan
- Yang, S. H., L. J. Wang and S. H. Li 2007 Ultraviolet–B irradiation–induced freezing tolerance in relation to antioxidant system in winter wheat (*Triticum aestivum* L.) leaves. *Environ. Exp. Bot.*, **60**: 300–307