Comparative Analysis of Paralytic Shellfish Toxin Content and Profile Produced by Dinoflagellate *Gymnodinium Catenatum* Isolated from Inokushi Bay, Japan

Oh, Seok Jin

Korea Inter-University Institute of Ocean Science, Pukyong National University

Matsuyama, Yukihiro

National Research Institute of Fisheries and Environment of Inland Sea, Fisheries Research Agency

Yoon, Yang Ho

College of Fisheries and Ocean Science, Chonnam National University

Miyamura, Kazuyoshi

Fisheries Research Institute Oita Prefectural Agriculture, Forestry and Fisheries Research Center

他

<https://doi.org/10.5109/17799>

出版情報：九州大学大学院農学研究院紀要. 55 (1), pp.47-54, 2010-02-26. Faculty of Agriculture, Kyushu University

バージョン：

権利関係：

Comparative Analysis of Paralytic Shellfish Toxin Content and Profile Produced by Dinoflagellate *Gymnodinium Catenatum* Isolated from Inokushi Bay, Japan

Seok Jin OH¹, Yukihiro MATSUYAMA², Yang Ho YOON³, Kazuyoshi MIYAMURA⁴,
Chang Geun CHOI¹, Han-Soeb YANG¹ and Ik Joon KANG*

Aquatic Biomonitoring and Environmental Laboratory, Department of Bioscience and
Biotechnology, Faculty of Agriculture, Kyushu University,
Fukuoka 812–8581, Japan

(Received November 2, 2009 and accepted November 19, 2009)

Paralytic shellfish toxin contents and profiles were investigated using shellfish toxin-producing dinoflagellate *Gymnodinium catenatum* strains isolated from Inokushi Bay during the March and July 2002 outbreak. The March strains, which were 72 strains isolated from bloom water, exhibited average growth rates of 0.25 divisions day⁻¹ due to sub-optimal temperature. The average toxin content was 321 f mole cell⁻¹. The principal toxin were C1+2 (79.8 mole%) and GTX 5+6 (18.0 mole%), and minor components were characterized as GTX 2+3, dcGTX2+3, and neoSTX+STX. Interestingly, 7 strains were clearly distinguished from those of other strains by the complete presence of GTX1+4. In the July strains, which were 33 strains isolated from bloom water, the average growth rate and toxin content was 0.34 divisions day⁻¹ and 160 f mole cell⁻¹, respectively. C1+2 (83.3 mole%) and GTX 5+6 (10.6 mole%) were the principal toxins, as in the March strains. However, carbamoyl toxins in the July strains were increasing with decreasing N-sulfocarbamoyl toxins in the March strains. We also observed major changes in toxin profile, evident by the increase of C1+2 and the decrease of GTX 5+6 in response to increasing temperature, using a single March strain. Minor components of July strains were characterized by GTX2+3, neoSTX, STX and dcGTX2+3. Thus, if we were to consider the toxin profile percentage based on temperature variation, *G. catenatum* outbreaks in March and July may have originated from one and the same population.

Keywords: *Gymnodinium catenatum*; paralytic shellfish toxin; toxin content; toxin profile; Inokushi Bay

INTRODUCTION

Gymnodinium catenatum Graham is a toxic dinoflagellate that causes paralytic shellfish poisoning (PSP) in coastal waters (Hallegraeff *et al.*, 1989). The first *G. catenatum* bloom associated with PSP was reported from Spain in 1977 (Estrada *et al.*, 1984), followed by cases of human fatalities from the Gulf of California in 1979 (Morey–Gaines, 1982). Reports that this species had spread across many parts of the world began to appear (Hallegraeff *et al.*, 1989). In Japan, *G. catenatum* was first suspected to be involved with PSP outbreaks in Senzaki Bay of western Japan, 1986. Since then, contaminations have occurred almost every year in western Japan (Ikeda *et al.*, 1989; Nishioka *et al.*, 1993; Baba *et al.*, 2001). The introduction (e.g. through ballast water) or migration of *G. catenatum* to central and northern Japan is possible where the maximum surface water temperature of 20 °C is favorable for this species

(Matsuoka and Fukuyo, 1994).

It has been proven that the capability to produce certain toxin profile is inherited (Anderson, 1990; Oshima *et al.*, 1993a, b). Given that differences in toxin profiles have a genetic basis, specific toxin profile might be used as biochemical markers for a number of PSP-producing dinoflagellate species. However, toxin profiles can vary due to differing environmental conditions (Ogata *et al.*, 1987; Maranda *et al.*, 1985; Cembella *et al.*, 1987; Boyer *et al.*, 1987; Boczar *et al.*, 1988; Anderson, 1990; Ishida *et al.*, 1993; Taroncher–Oldenburg *et al.*, 1997). Given this fact, a number of studies investigated species population dynamics based on toxin profiles. For example, Chang *et al.* (1997) reported on the toxin profile of *Alexandrium minutum* isolated from coastal areas of New Zealand, showing that is was quite different from that of species isolated from other parts of the world. It is quite possible that *A. minutum* may have been introduced to New Zealand via ballast water given its isolated location and high maritime traffic from all around the world. In Australia, *G. catenatum* is regarded as an exotic species since only being noticed in coastal waters from the 1980's, and supported by the fact that its cysts were not observed in sediment prior to 1980s. Hallegraeff and Bolch (1992) suggested that cysts of *G. catenatum* were brought to Tasmania by ballast water in ships from Japan and Korea. However, toxic profiles of the Tasmanian strains are clearly distinguishable from those isolated from Japan (Oshima *et al.*, 1993a). This might indicate whether they are autochthonal species or intro-

¹ Korea Inter–University Institute of Ocean Science, Pukyong National University, 599–1 Daeyeon-dong, Nam-gu, Busan 608–737 Korea

² National Research Institute of Fisheries and Environment of Inland Sea, Fisheries Research Agency 2–17–5 Maruishi, Hatsukaichi, Hiroshima 739–0452, Japan

³ College of Fisheries and Ocean Science, Chonnam National University San 96–1, Dundok-dong, Yeosu 550–749, Korea

⁴ Fisheries Research Institute Oita Prefectural Agriculture, Forestry and Fisheries Research Center, Kamiura, Saiki, Oita, 879–2602, Japan

* Corresponding author (E-mail: kangnew@agr.kyushu-u.ac.jp)

duced from countries other than Japan and Korea. Recently, Park *et al.* (2004) observed strains produced C3 and C4 toxins around Korean coastal waters, similar to the Tasmanian strains.

A major causative organism of PSP in Inokushi Bay, which has the highest production of cultured short-necked clam in Japan, was reported to be *G. catenatum* (Miyamura *et al.*, 2007). This species has been observed in the bay throughout the year, which is of interest as water temperatures between winter and summer are significantly different. Moreover, *G. catenatum* attributed PSP events in summer did not occur in conjunction with *G. catenatum* outbreak in the bay, in spite of increasing cell density. This may have been due to specific physiological conditions related to environmental parameters between summer and winter. The aim of this study was to compare the population dynamics of *G. catenatum* between the May and July 2002 blooms, based on toxin profiles in the bay.

MATERIALS AND METHODS

Culture isolates of *Gymnodinium catenatum*

G. catenatum was isolated from Inokushi Bay, Oita Prefecture, Japan (Fig. 1) in March and July 2002 on a light microscope using a micropipette (ca. ϕ 50–100 μ m), coinciding with high cell numbers in the water column (Fig. 2). Temperature and salinity were observed using a conductivity–temperature–depth system (Model 58, YSI/Nanotech Inc., Tokyo, Japan). The isolated cells were rinsed at least four times by filtered seawater (0.22 μ m pore size; GSWP, Millipore, MA, USA), and then placed in culture tubes with 5 ml modified f/2 medium (Guillard and Ryther, 1962) sterilized at 121 $^{\circ}$ C for 1 hr. From sea-

Fig. 1. Study area in Inokushi Bay, Japan (●: sampling station).

Fig. 2. Temporal changes of temperature, salinity and cell density of *Gymnodinium catenatum* in Inokushi Bay, 2002.

water sampled in March and July, 72 and 33 clonal strains of *G. catenatum* were isolated, respectively (hereafter referred to as the March and July strains). Isolated cells were cultured at 30 psu, with an illumination of 100 μ mol $m^{-2} s^{-1}$ using cool-white fluorescent lamps (12 h light: 12 h dark cycle). Incubation temperature was set at the *in situ* temperature (18 $^{\circ}$ C for March strains and 25 $^{\circ}$ C for July strains). Culturable cells (i.e. viable strains after isolation) were 55 of the 73 March strains (76.4%) and 28 of the 33 July strains (84.8%). To remove substances such as ammonia and metals, all equipment and glassware were washed with approximately 30% v/v HCl and thoroughly rinsed with distilled water before being autoclaved (202 kPa, 20 min).

Growth rate (division day^{-1}) estimates of each strain were calculated using a least square regression on log-transformed cell density data during exponential growth phase:

$$\mu = \frac{1}{\Delta t} \ln \frac{N_t}{N_0} ,$$

where

N_0 , N_t : initial and final *in vivo* fluorescence of the exponential period (cells ml^{-1}),

Δt : period of exponential growth phase (day).

Cell harvest and toxin analysis

Cultured cells were harvested by a plankton net (10 μ m pore size) in late exponential phase at 6:00 p.m. Pellets were extracted with 0.03 N acetic acid by sonication, and cell debris removed by centrifugation (10,000 $\times$ 10 min). Analysis of PSP toxins using high-performance

liquid chromatography with fluorescence detection (Model-700, Hitachi, Tokyo, Japan) was performed following the post-column oxidative fluorescence method of Oshima (1995). Toxins were resolved by reverse-phase chromatography using a silica-base column (Inertsil C8; 0.46×15 cm, GL Science, Tokyo, Japan). Three separate isocratic elutions were employed to separate the spectrum of PSP-toxins at a flow of 0.8 ml min⁻¹. The gonyautoxins (GTX1–4) and the N-sulfocarbamoyl toxin were resolved using 10 mM aqueous ammonium phosphate at pH 7.1, with 2 mM heptane sulfonate (Na salt) serving as the ion-pairing reagent. To separate neosaxitoxin (neoSTX) and saxitoxin (STX), 2 mM heptane sulfonate (Na salt) was substituted as the ion-pairing reagent in 30 mM aqueous ammonium phosphate buffer (pH 7.1) with 4% acetonitrile. The N-sulfocarbamoyl (C-) toxins were resolved using 2 mM tetrabutyl ammonium phosphate adjusted to pH 6 as the mobile phase. The eluate from the column was continuously mixed with 7 mM periodic acid in 50 mM sodium phosphate buffer (pH 9.0) at 0.4 ml min⁻¹, heated at 65 °C by passing through Teflon tubing and then mixed with 0.5 N acetic acid at 0.4 ml min⁻¹ just before entering the monitor. The fluorometer was set at an excitation wavelength of 330 nm and emission of 390 nm.

Gonyautoxins 5 and 6 (GTX 5+6) were determined in accordance with the increased yield of neosaxitoxin (neoSTX) and saxitoxin (STX) formed after heating (1 hr, 100 °C) with addition of 0.1 N HCl (Nagashima *et al.*, 1987), as the standard toxins for GTX 5+6 were not available. Meanwhile, STX, neoSTX, gonyautoxins 1–4 (GTX1–4), decarbamoylgonyautoxins 2+3 (dcGTX2+3) and C1–4 were estimated from standard toxins obtained

from Japan Fisheries Resource Conservation Association, and/or from Prof. Yasukatsu Oshima of Tohoku University, Japan. The results were expressed in relative amounts of each toxin on a molar basis (mol%) and toxic content (f mole cell⁻¹).

Toxin contents and profiles of *Gymnodinium catenatum* in response to changing temperature

To examine the toxin contents and profiles of *G. catenatum* in response to changing temperatures, we used a single strain (GCINO2002M14) isolated from Inokushi Bay in March 2002. The effect of temperature was determined at 18 °C and 25 °C, which corresponded to the in situ temperature of each sampling period. Salinity and irradiance was constant during incubation period (30 psu and 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$). To avoid culture shock due to change in incubated temperature, temperatures of cultures were raised about 1 °C a day, and were finally maintained at 25 °C. The acclimated cells were inoculated into five replicate flasks with fresh modified f/2 medium at 18 °C and 25 °C. Toxin content was estimated using cells in late exponential phase as mentioned above.

RESULTS

Relationship between hydrographic condition and cell density of *Gymnodinium catenatum*

Temperature and salinity during the sampling period ranged from 16.0–28.9 °C and 31.15–34.72 psu, respectively (Fig. 2). *G. catenatum* exhibited increasing cell density from January to March (16–21 °C and 34.0–35.0 psu), and from July to August (25–27 °C and 31–32 psu). Maximum cell density was obtained at

Fig. 3. Growth rates of *Gymnodinium catenatum* isolated from Inokushi Bay in March and July, 2002. Experimental conditions; 18 °C for strains isolated from the March strain and 22 °C for the July strains. Salinity 30, pH 8.0, 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (cool-white fluorescent lamps, 12:12 h LD cycle).

Fig. 4. Toxin contents of *Gymnodinium catenatum* isolated from Inokushi Bay in March and July, 2002. Cultured cells were harvested by plankton net (20 μ m pore size) in late exponential phase (See Fig. 3 for incubation conditions).

16.1 °C and 34.72 psu (1,808 cells L^{-1}). However, cell densities from July to August remained at 265 cells L^{-1} , which was lower than that of February (temperature 26.3 °C and salinity 32.09 psu).

Toxin contents and profiles of *Gymnodinium catenatum* strains isolated from Inokushi Bay

Growth rates of individual strains are shown in Fig. 3. The average growth rates were significantly different at 0.25 division day^{-1} for the March strains and 0.34 division day^{-1} for the July strains (one-way ANOVA; $P < 0.001$). The average toxin content was 321 ± 160 f mole $cell^{-1}$ in the March strains and 160 ± 92.8 f mole $cell^{-1}$ in July strains. Non-toxin strains were not observed. Toxin

content of the strains varied greatly between the minimum (56.6 f mole $cell^{-1}$) and maximum values (740 f mole $cell^{-1}$; Fig. 4). Moreover, a one-way ANOVA indicated that variability of toxin contents were significant among individual strains ($p < 0.001$). When toxin contents were plotted against individual growth rates, the relationship showed an inverse linear trend, despite the low correlation coefficient ($r = -0.38$; $P < 0.001$; Fig. 5).

In the toxin profile of the March strains, C1+2 toxin was the dominant toxin (ca. 79.8 ± 10.0 mole% among whole PSP toxin; Fig. 6). However, C3+4 toxin was completely absent in all strains. GTX5+6 toxin was occupied by proportions of 18.0 ± 8.42 mole%. Minor components of the March strains were characterized by GTX2+3 (0.21 ± 0.37 mole%), dcGTX2+3 (0.32 ± 0.25 mole%) and neoSTX+STX (1.65 ± 3.11 mole%) (Fig. 6). In the July strains, proportions of C1+2 and GTX 5+6 toxins were 83.3 ± 6.63 mole% and 10.6 ± 5.78 mole%, respectively. In contrast, minor components were characterized by GTX2+3 (2.26 ± 2.78 mole%), dcGTX2+3 (0.72 ± 0.79 mole%) and neoSTX+STX (3.14 ± 3.37 mole%; Fig. 7).

Toxin content and profiles of *Gymnodinium catenatum* under different water temperatures

Table 1 shows the toxin content of *G. catenatum* strain (GCINO2002M14) incubated at 18 °C and 22 °C with in situ temperature. The toxin content of strain incubated at 18 °C and 25 °C were 136 ± 21.6 f mole $cell^{-1}$ and 58.0 ± 20.3 f mole $cell^{-1}$, respectively. The toxin contents in 18 °C were ca. 2 times higher than those of 25 °C. This corresponded well with results which showed that average toxin contents of the March strains were 2 times higher. The toxin profile also varied with that of C1+2

Fig. 5. Correlation between growth rate and toxin contents of *Gymnodinium catenatum* isolated from Inokushi Bay in March and July, 2002.

Fig. 6. (a) Toxin profiles in *Gymnodinium catenatum* isolated from Inokushi Bay in March, 2002. (b) Toxin profiles of *G. catenatum* strains, with the exception of C1+2 (See Fig. 3 for incubation conditions).

Fig. 7. Toxin profiles in *Gymnodinium catenatum* isolated from Inokushi Bay in July, 2002. (b) Toxin profiles of *G. catenatum* strains, with the exception of C1+2 (See Fig. 3 for incubation conditions).

increasing from 50.6 ± 1.30 mole% to 76.9 ± 1.68 mole%, while GTX 5+6 decreased from 48.5 ± 1.31 mole% to 22.0 ± 1.71 mole% with increasing temperature, although GTX2+3 did not vary between the two temperature ranges (neoSTX+STX was not detected; Fig. 8).

DISCUSSION

G. catenatum blooms occur from January to March and from July to August in Inokushi Bay. Water temperature during these outbreaks ranged from 16 °C to 21 °C

Fig. 8. Changes in toxin profile of *Gymnodinium catenatum* with increasing temperature from 18 °C to 25 °C.

and from 25 °C to 27 °C, respectively. Bravo and Anderson (1994) reported that the optimum temperature of the Spanish strain was between 22–28 °C. In tropical regions, such as Palau, Venezuela and Philippines, optimum growth temperature is between 23–29 °C (Hallegraeff and Fraga, 1998). In laboratory experiments using strains isolated from Hiroshima Bay of Japan and Yosue Bay of Korea, optimum temperatures ranged from 20–30 °C (Yamamoto *et al.*, 2002; Oh and Yoon, 2004). These temperatures are consistent with temperatures observed in this study from July to August. However, the Tasmanian strain was able to better grow in lower temperatures (Hallegraeff *et al.*, 1989). Blooms in Tasmanian waters occur from December to June when water temperature ranges from 12 °C to 18 °C, which were replicated in laboratory experiments by Blackburn *et al.* (1989). These temperatures are consistent with temperature from January to March of this study.

G. catenatum outbreaks in Inokushi Bay during different temperature ranges indicate the presence of various populations with different physiological characteristics. Sakamoto and Kotani (1998) reported that *Alexandrium tamarense* isolated from Hiroshima Bay have plural populations due to wide variation of toxin profiles among strains isolated from the same seawater. Ichimi *et al.*, (2002) also suggested the presence of various planktonic populations around the same area. Based on these informations, we cannot rule out the possibility that the same population is in fact present during both sampling periods, compared to the notion that two different populations were present.

Average of growth rate was 0.25 division day⁻¹ in the March strains and 0.34 division day⁻¹ in the July strains (Fig. 3). Different studies have reported maximum growth rates of *G. catenatum* at 0.4 divisions day⁻¹ from the Denmark strain (Ellegaard *et al.*, 1993), 0.53 divisions day⁻¹ from the Spanish strain (Bravo and Anderson 1994) and 0.44 divisions day⁻¹ (0.31 day⁻¹ in the original paper) from the Japan and Korea strains (Yamamoto *et al.*, 2002; Oh *et al.*, 2004). These maximum growth rates were similar to the growth rate of the July strains in this study, which was incubated near its optimum temperature. On the other hand, growth rates of the March

strains were similar to that of laboratory experiments conducted at the same temperature (18 °C) and salinity (30 psu) (i.e. ca. 0.22 divisions day⁻¹; Yamamoto *et al.*, 2002). However, we can not explain the contrasting growth rates despite being isolated from the same season and area. Further studies are required to investigate if the difference is related to hereditary characteristics or other physiologically related factors.

Average toxin content of the March strains was 2 times higher than that of the July strains. This discrepancy was similar to that of the laboratory experiment. Causative factors may be related to the production of intracellular amino acid, which is a precursor in toxin biosynthesis. In amino acid synthesis nitrate should be reduced to ammonium by the sequential action of nitrate reductase (Lomas and Glibert, 1999). Enzyme activity increases with decreased temperature (Kristiansen, 1983; Gao *et al.*, 1993; 2000). Thus, increasing of toxin content might have been caused by increasing of intercellular amino acid at relatively low temperature. Anderson *et al.* (1990) also proposed the hypothesis that the effect of low temperature was to reduce protein synthesis, and surplus of arginine within the cell could be used for toxin synthesis. The other reason for increase of toxin content with decreasing temperature may be related to growth rate. High growth rate with optimum temperature conditions and no nutrient limitation may decrease toxin content (Fig. 5). This may be because of transfer of toxic content to daughter cells with faster division (Anderson *et al.* 1990). These results may be a possible explanation for the *G. catenatum* related PSP event in Inokushi Bay in summer.

Our study confirmed that the major toxins in the March and July strains were C1+2 and GTX5+6. In previous studies, major *G. catenatum* toxins isolated from Australia, Japan, Philippine, Portugal and Spain were shown to be C1–4 and GTX 5+6 (Franca *et al.*, 1993; Fukuyo *et al.*, 1993; Ikeda *et al.*, 1989; Oshima *et al.*, 1993a, b; Takatani *et al.*, 1998). However, Japan strains have been reported to be devoid of C3+4 (Oshima *et al.*, 1993a, b). Singapore strains were also clearly distinguished from those of other countries by the complete absence of C3+4 (Holmes *et al.*, 2002).

One-way ANOVA indicated that the C1+2 and GTX5+6 proportions of the March strains were significantly different from those of the July strains ($P < 0.001$), attributed to carbamoyl toxins in the July strains increasing with decreasing N-sulfocarbamoyl toxins in the March strains. Oshima *et al.* (1993a) and Flynn *et al.* (1996) reported that toxin profile in *G. catenatum* is not significantly affected by temperature, light or nutrient status. If toxin profile is stable as reported by Oshima *et al.* (1993a) and Flynn *et al.* (1996), the March population might be different to the July population.

However, we observed major changes in toxin profile with increased temperature (Fig. 9). Some authors have reported changes in the composition of PSP affected by physical and chemical environmental parameters (Anderson, 1990; Sakamoto and Kotani, 1998; Hwang and Lu, 2000; Ichimi *et al.*, 2002). Thus, by considering

the percentage of toxin profile based on temperature variation, the March and July strains could probably be the same population. However, some strains are significantly different from those of other strains. Our study observed GTX1+4 from 7 strains (GCINO2002M17, 27, 29, 56, 57 and 67) among the March strains. This toxin has not been previously reported, except from Singapore strains (Holmes *et al.*, 2002). Furthermore, in Neo-STX+STX, 3 strains (GCINO2002M17, 25 and 56) from the March strains reached more than 10 mol%. These results suggested that some strains could be distinguished from populations of *G. catenatum* species in Inokushi Bay, raising the probability that these strains were introduced from other regions (Ichimi *et al.*, 2002). Based on toxicological analysis, we showed that *G. catenatum* blooms in March and July may have originated from one and the same population. Future research initiatives need to investigate the population dynamics of strains based on molecular biological methods, such as gene analysis, in order to validate the results of this study.

ACKNOWLEDGEMENTS

We are grateful to Prof. Y. Oshima of Tohoku University, as well as the Fisheries Agency of Japan for providing the PSP toxin standards.

REFERENCES

- Anderson, D. M. 1990 Toxin variability in *Alexandrium* species. In "Toxic Marine Phytoplankton", ed. by E. Graneli, B. Sundström, L. Edler and D. M. Anderson, Elsevier, New York, pp. 41–51
- Anderson, D. M., J. J. Sullivan and B. Reguera 1989 Paralytic shellfish poisoning in Northwest Spain: the toxicity of the dinoflagellate *Gymnodinium catenatum*. *Toxicon*, **27**: 665–674
- Baba, T., S. Hiyama and T. Tainaka 2001 Vertical migration of the toxic dinoflagellate *Gymnodinium catenatum* toxicity of cultured oyster in Senzaki Bay, Yamaguchi Prefecture. *Bull. Plankton Soc. Japan*, **48**: 95–99 (In Japanese with English abstract)
- Blackburn, S. I., G. M. Hallegraeff and C. J. Bolch 1989 Vegetative reproduction and sexual life cycle of the toxic dinoflagellate *Gymnodinium catenatum* from Tasmania, Australia. *J. Phycol.*, **25**: 577–590
- Boczar, B. A., M. K. Beitler, J. Liston, J. J. Sullivan, and R. A. Cattolico 1988 Paralytic shellfish toxins in *Protogonyaulax tamarens* and *Protogonyaulax catenella* in axenic culture. *Plant. Physiol.*, **88**: 1285–1290
- Boyer, G. L., J. J. Sullivan, R. J. Anderson, P. J. Harrison and F. J. R. Taylor 1987 Effects of nutrient limitation on toxin production and composition in the marine dinoflagellate *Protogonyaulax tamarens*. *Mar. Biol.*, **96**: 123–128
- Bravo, I. and D. M. Anderson 1994 The effects of temperature, growth medium and darkness on excystment and growth of the toxic dinoflagellate *Gymnodinium catenatum* from north-west Spain. *J. Plankton Res.*, **16**: 513–525
- Cembella, A. E., J. J. Sullivan, G. L. Boyer, F. J. R. Taylor and R. J. Anderson 1987 Variation in paralytic shellfish toxin composition within the *Protogonyaulax tamarens*/*catenella* species complex: red tide dinoflagellates. *Biochem. Syst. Ecol.*, **15**: 171–186
- Chang, F. H., D. M. Anderson, D. M. Kulis and D. Till 1997 Toxin production of *Alexandrium minutum* (Dinophyceae) from the Bay of Plenty, New Zealand. *Toxicon*, **35**: 393–409
- Ellegaard, M., N. F. Christensen and Ø. Moestrup 1993 Temperature and salinity effects on growth of a non-chain-forming strain of *Gymnodinium catenatum* (Dinophyceae) established from a cyst from recent sediments in the sound (Øresund), Denmark. *J. Phycol.*, **29**: 418–426
- Estrada, M., F. J. Sanchez and S. Fraga 1984 *Gymnodinium catenatum* (Graham) en las rías gallegas (NO de Espana). *Invest. Pesq.*, **48**: 31–40
- Franca, S., P. Alvito, I. Sousa and V. Mascarenhas 1993 The dinoflagellate *Gymnodinium catenatum* isolated from the coast of Portugal: Observations on development, toxicity and ultrastructure. In "Toxic Phytoplankton Blooms in the Sea", ed. by T. J. Smayda and Y. Shimizu, Elsevier, Amsterdam, pp. 869–874
- Flynn, K. J., K. Flynn, E. H. John, B. Reguera, M. I. Reyero and J. M. Franco 1996 Changes in toxins, intracellular and dissolved free amino acids of the toxic dinoflagellate *Gymnodinium catenatum* in response to changes in inorganic nutrients and salinity. *J. Plankton Res.*, **18**: 2093–2111
- Fukuyo, Y., M. Kodama and T. Ogata 1993 Occurrence of *Gymnodinium catenatum* in Manila Bay, the Philippines. In "Toxic Phytoplankton Blooms in the Sea", ed. by T. J. Smayda and Y. Shimizu, Elsevier, Amsterdam, pp. 875–880
- Gao, Y., G. J. Smith and R. S. Alberte 1993 Nitrate Reductase from the Marine Diatom *Skeletonema costatum*. *Plant Physiol.*, **103**: 1437–1445
- Gao, Y., G. J. Smith and R. S. Alberte 2000 Temperature dependence of nitrate reductase activity in marine phytoplankton: biochemical analysis and ecological implications. *J. Phycol.*, **36**: 304–313
- Guillard, R. R. L. and D. Ryther 1962 Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt and *Detonula confervacea* (Cleve) Gran. Can. *J. Microbiol.*, **8**: 229–239
- Hada, Y. 1967 Protozoan plankton of the Inland Sea, Setonaikai I. *The mastigophora. Bull. Suzugamine Woman's Coll., Nat. Sci.*, **13**: 1–26
- Hallegraeff, G. M. and C. J. Bolch 1992 Transport of diatom and dinoflagellate resting spores in ships' ballast water: implications for plankton biogeography and aquaculture. *J. Plankton Res.*, **14**: 1067–1084
- Hallegraeff, G. M. and S. Fraga 1998 Bloom dynamics of the toxic dinoflagellate *Gymnodinium catenatum*, with emphasis on Tasmanian and Spanish coastal waters. In "Physiological Ecology of Harmful Algal Blooms", ed. by D. M. Anderson, A. D. Cembella and G. M. Hallegraeff, Springer, Berlin, pp. 59–80
- Hallegraeff, G. M., S. O. Stanley, C. J. Bolch and S. I. Blackburn 1989 *Gymnodinium catenatum* blooms and shellfish toxicity in southern Tasmania, Australia. In "Red Tides: Biology, Environmental Science and Toxicology", ed. by T. Okaichi, D. M. Anderson and T. Nemoto, Elsevier, New York, pp. 77–80
- Hang, D. F. and Y. H. Lu 2000 Influence of environmental and nutritional factors of growth, toxicity, and toxin profile of dinoflagellate *Alexandrium minutum*. *Toxicon*, **38**: 1491–1503
- Holmes, M. J., C. J. Bolch, D. H. Green, A. D. Cembella, S. Lay and M. Teo 2002 Singapore isolates of the dinoflagellate *Gymnodinium catenatum* (Dinophyceae) produce a unique profile of paralytic shellfish poisoning toxins. *J. Phycol.*, **38**: 96–106
- Ichimi, K., T. Suzuki and A. Ito 2002 Variety of PSP toxin profiles in various culture strains of *Alexandrium tamarens* and change of toxin profile in natural *A. tamarens* population. *J. Exp. Mar. Biol. Ecol.*, **273**: 51–60
- Ishida, Y., C. H. Kim, Y. Sako, N. Hirooka and A. Uchida 1993 PSP toxin production is chromosome dependent in *Alexandrium* spp. In "Toxic Phytoplankton Blooms in the Sea", ed. by T. J. Smayda and U. Shimizu, Elsevier, Amsterdam, pp. 881–887
- Ikeda, T., S. Matsuno, S. Sato, T. Ogata, M. Kodama, Y. Fukuy and H. Takayama 1989 First report on paralytic shellfish poisoning caused by *Gymnodinium catenatum* (Dinophyceae) in Japan. In "Red Tides", ed. by T. Okaichi, D. M. Anderson and T. Nemoto, Elsevier, New York, pp. 411–414
- Kristiansen, S. 1983 The temperature optimum of the nitrate

- reductase assay for marine phytoplankton. *Limnol. Oceanogr.*, **28**: 776–780
- Lomas, M. W. and P. M. Glibert 1999 Temperature regulation of nitrate uptake: A novel hypothesis about nitrate uptake and reduction in cool-water diatom. *Limnol. Oceanogr.*, **44**: 556–572
- Maranda, L., D. M. Anderson and Y. Shimizu, 1985 Comparison of toxicity between populations of *Gonyaulax tamarensis* of eastern North American waters. *East. Coast. Shelf. Sci.*, **21**: 401–410
- Matsuoka, K. and Y. Fukuyo 1994 Geographical distribution of the toxic dinoflagellate *Gymnodinium catenatum* Graham in Japanese coastal waters. *Botanica Marina*, **37**: 495–503
- Miyamura, K., Y. Matsuyama and S. J. Oh 2007 Occurrence of the toxic dinoflagellate *Gymnodinium catenatum*, toxin contents of suspended matter and prediction of PSP outbreak in the scallop *Chlamys nobilis* in Inokushi Bay, Oita Prefecture. *Nippon Suisan Gakkaishi*, **73**: 32–42 (in Japanese with English abstract)
- Morey-Gaines, G. 1982 *Gymnodinium catenatum* (Dinophyceae): morphology and affinities with armored forms. *Phycologia*, **21**: 154–163
- Nishioka J., Y. Wada and Y. Imanishi 1993 On the occurrence of *Gymnodinium catenatum* (Dinophyceae) in Kumihama Bay. *Bull. Kyoto Inst. Ocean. Fish. Sci.*, **16**: 43–49 (in Japanese with English abstract)
- Ogata T., T. Ishimaru and M. Kodama 1987 Effect of water temperature and light intensity on growth rate and toxicity change in *Protogonyaulax tamarensis*. *Mar. Biol.*, **95**: 217–220
- Oh, S. J. and Y. H. Yoon 2004 Effects of water temperature, salinity and irradiance on the growth of the toxic dinoflagellate, *Gymnodinium catenatum* (Graham) isolated from Yeosuhae Bay, Korea. *Algae*, **19**: 293–302 (in Korean with English abstract)
- Oshima, Y. 1995 Post-column derivatization HPLC methods for paralytic shellfish poisons. In “Manual on Harmful Marine Microalgae”, ed. by G. M. Hallegraeff, D.M. Anderson and A. D. Cembella, UNESCO, Paris, pp. 81–94
- Oshima, Y., S. I. Blackburn and G. M. Hallegraeff 1993a Comparative study on paralytic shellfish toxin profiles of the dinoflagellate *Gymnodinium catenatum* from three different countries. *Mar. Biol.*, **116**: 471–476
- Oshima, Y., H. Itakura, K. Lee, T. Yasumoto, S. Blackburn and G. Hallegraeff 1993b Toxin production by the dinoflagellate *Gymnodinium catenatum*. In “Toxic phytoplankton blooms in the sea”, ed. by T. J. Smayda and Y. Shimizu, Elsevier, New York, pp. 907–912
- Oshima, Y., M. Hasegawa, T. Yasumoto, G. Hallegraeff and S. Blackburn 1987 Dinoflagellate *Gymnodinium catenatum* as the source of paralytic shellfish toxins in Tasmanian shellfish. *Toxicon*, **25**: 1105–1111
- Park, T. G., C. H. Kim and Y. Oshima 2004 Paralytic shellfish toxin profiles of different geographic populations of *Gymnodinium catenatum* (Dinophyceae) in Korean coastal waters. *Phycol. Res.*, **52**: 300–305
- Sakamoto, S. and Y. Kotani 1998 Variation of paralytic shellfish contents and composition in *Alexandrium tamarensis* collected at Kure Bay, Hiroshima prefecture. *Bull. Nansei. Natl. Fish. Res. Inst.*, **31**: 45–52 (in Japanese with English abstract)
- Takatani, T., T. Morita, A. Anami, H. Akaeda, Y. Kamijo, K. Tsutsumi, T. Noguchi 1998 Appearance of *Gymnodinium catenatum* in association with the toxification of bivalves in Kamae, Oita Prefecture, Japan. *J. Food Hyg. Soc. Japan.*, **39**: 275–280
- Taroncher–Oldenburg, G., D. M. Kulis and D. M. Anderson 1997 Toxin variability during the cell cycle of the dinoflagellate *Alexandrium fundyense*. *Limnol. Oceanogr.*, **42**: 1178–1188
- Yamamoto, T., S. J. Oh and Y. Kataoka 2002 Effect of temperature, salinity and irradiance on the growth of the toxic dinoflagellate *Gymnodinium catenatum* (Dinophyceae) isolated from Hiroshima Bay, Japan. *Fisheries Sci.*, **68**: 356–363