Net Fine Root Carbon Production in *Pinus densiflora*, *Pinus koraiensis*, *Larix leptolepis* and *Quercus acutissima* Stands, Gongju area, Chungnam Province, Korea

Park, Gwan-Soo

Division of Environmental Science and Forest Resources, College of Agriculture & Life Sciences, Chungnam National University

Kang, Gil-Nam

Chungnam Forest Environment Research institute

Lee, Sang-Jin

Division of Environmental Science and Forest Resources, College of Agriculture & Life Sciences, Chungnam National University

Lee, Hang-Goo

Lee_Hang-Goo]

他

<https://doi.org/10.5109/17803>

出版情報：九州大学大学院農学研究院紀要. 55 (1), pp.73-76, 2010-02-26. Faculty of Agriculture, Kyushu University

バージョン：

権利関係：

Net Fine Root Carbon Production in *Pinus densiflora*, *Pinus koraiensis*, *Larix leptolepis* and *Quercus acutissima* Stands, Gongju area, Chungnam Province, Korea

Gwan-Soo PARK¹, Gil-Nam KANG², Sang-Jin LEE¹, Hang-Goo LEE¹,
Si-Young LEE³, Hee-Mun CHAE⁴ and Shoji OHGA*

Laboratory of Forest Resources Management, Division of Forest Ecosystem Management,
Department of Forest Products Science, Kyushu University,
Sasaguri, Fukuoka 811-2415, Japan

(Received November 17, 2009 and accepted November 19, 2009)

This study was carried out to estimate annual carbon production by fine root in *Pinus densiflora*, *Pinus koraiensis*, *Larix leptolepis* and *Quercus acutissima* stands of Gongju region, Korea. Soil samples were taken at 0–30 cm, 30–60 cm, and 60–90 cm soil depths from April to November (monthly increment method) in 2007 using soil sampler. Fine root carbon biomass was higher at 0–30 cm soil depth than in the other soil depths. Total fine root carbon biomass at 0–90 cm soil depth was highest in *Larix leptolepis* stand among the four study stands. Net fine root carbon production ($\text{kg ha}^{-1} \text{yr}^{-1}$) in *Pinus densiflora*, *Pinus koraiensis*, *Larix leptolepis* and *Quercus acutissima* stands were 889 kg, 2,266 kg, 851 kg, and 1,121 kg at 0–30 cm soil depth, respectively. Total net fine root carbon production ($\text{kg ha}^{-1} \text{yr}^{-1}$) was highest in *Pinus koraiensis* stand among the four study stands. Fine root turnover rates in the four stands were ranged from 0.59 to 2.63 at 0–30 cm, 30–60 cm, and 60–90 cm soil depth.

INTRODUCTION

After the earth's average air temperature reached an all-time high recently, many scientist expressed concern about the potential for significant global warming as a result of the increased carbon dioxide (CO_2) and other greenhouse gases (CFC_s , CH_4 , and N_2O) within the coming century (Schlesinger, 1991; Hansen, 1993). Even small changes in temperature can have dramatic impacts on the earth complex atmosphere, ocean, land, and life systems (Hair and Sampson, 1992). Increasing population and economic activity will increase the concentration of CO_2 in the atmosphere and may accelerate changes in global climate which may have important consequences for the earth's ecology.

Forests have received considerable attention because they are a major sink for C cycle. The carbon sinks of plants include roots, shoots, and when plants are exposed to large seasonal climatic changes, storage reserves for maintenance during and after the period of dormancy. Not like a shoots, fine root carbon dynamics are seldom measured directly because these studies are labor intensive (Burke and Raynal, 1994) and also sampling methods are not standardized (Laurenroth *et al.*, 1986; Nadelhoffer and Raich, 1992; Singh *et al.*, 1984; Vogt *et al.*, 1986). However, it is true that a large part of the carbon in

deciduous forests is allocated in fine root, yet this process is seldom measured (Burke and Raynal, 1994).

The objective of this study was to estimate the amount of annual C production by fine root in *Pinus densiflora*, *Pinus koraiensis*, *Larix leptolepis*, and *Quercus acutissima* stands of Gongju region, Chungnam Province, Korea.

MATERIALS AND METHODS

This study was conducted in 2007 in *Pinus densiflora*, *Pinus koraiensis*, *Larix leptolepis* and *Quercus acutissima* stands at the Gongju city of Chungnam Province, Korea. Annual mean air temperature is 11.9°C and total annual precipitation averages 1,534 mm. The elevation range of the study site was 94–214 m. The average age of the *Pinus densiflora* stand was 36 years and the number of trees per ha are 1,331. The average age of the *Pinus koraiensis* stand was 39 years and the number of trees per ha was 1,206. The average age of the *Larix leptolepis* stand was 36 years and the number of trees per ha was 588. The average age of the *Quercus acutissima* stand was 42 years and the number of trees per ha was 363. The study sites was dominated by *Smilax china*, *Oplismenus undulatifolius*, *Corylus heterophylla*, *Rhus trichocarpa*, *Lindera obtusiloba* and *platycarya strobilacea*. The soil is brown forest soil, a representative soil of Korea.

For this study, three 10 m×10 m permanent plots, which was established on a relatively similar slope, aspect and soil, were randomly selected in each stand. Fine root biomass was established by monthly soil coring from April to November at five random locations in each plot (Hwang 1996; Makkonen and Helmisaari, 1999). Sample points for coring were randomly selected from the permanent grid. A stainless steel core (7.4 cm diameter and

¹ Division of Environmental Science and Forest Resources, College of Agriculture & Life Sciences, Chungnam National University, Daejeon, Korea

² Chungnam Forest Environment Research institute, Gongju, Korea

³ Division of the Professional School of Disasters Prevention Technology, Kangwon National University, Samcheok-Si, Korea

⁴ Korea Climate Change Countermeasure Research center, Chuncheon-Si, Korea

* Corresponding author (E-mail: ohga@forest.kyushu-u.ac.jp)

[illegible]

Table 4. Fine root carbon biomass (kg ha⁻¹) in *Larix leptolepis* stand at 0–30 cm, 30–60 cm, and 60–90 cm soil depths. The number in parenthesis is one standard error of the mean (n=12 at 0–30 cm and 30–60 cm, n=6 at 60–90 cm)

Soil depth	22 Apr.	20 May.	20 Jun.	22 Jul.	22 Aug.	21 Sep.	20 Oct.	23 Nov.	Mean
0–30 cm	1326.0 (195)	1340.0 (181)	1833.1 (146)	1542.3 (263)	1770.3 (202)	1765.7 (343)	1279.5 (222)	663.0 (110)	1440.0
30–60 cm	519.7 (107)	274.5 (54)	497.8 (122)	483.9 (173)	593.2 (123)	339.6 (71)	481.5 (82)	428.0 (114)	452.3
60–90 cm	160.9 (67)	437.3 (120)	214.0 (68)	239.6 (81)	244.3 (110)	181.5 (47)	323.4 (57)	274.5 (71)	259.4
0–90 cm									2151.7

Table 5. Fine root carbon biomass (kg ha⁻¹) in *Quercus acutissima* stand at 0–30 cm, 30–60 cm, and 60–90 cm soil depths. The number in parenthesis is one standard error of the mean (n=12 at 0–30 cm and 30–60 cm, n=6 at 60–90 cm)

Soil depth	22 Apr.	20 May.	20 Jun.	22 Jul.	22 Aug.	21 Sep.	20 Oct.	23 Nov.	Mean
0–30 cm	1493.5 (172)	1677.3 (244)	1565.6 (127)	1107.3 (144)	1777.3 (316)	1267.8 (187)	895.6 (120)	665.3 (138)	1306.2
30–60 cm	469.9 (85)	467.6 (95)	465.3 (72)	362.9 (84)	586.2 (165)	325.7 (57)	456.0 (151)	402.5 (67)	442.0
60–90 cm	323.4 (93)	430.4 (117)	307.1 (85)	172.1 (56)	367.6 (134)	162.8 (409)	172.1 (52)	69.8 (56)	250.7
0–90 cm									1998.9

root biomass in Northeast hardwood stands of U.S.A. was peak in May. Also, McLaugherty *et al.* (1982) reported highest fine root biomass in spring season.

Net fine root carbon production (kg ha⁻¹ yr⁻¹) in *Pinus densiflora*, *Pinus koraiensis*, *Larix leptolepis* and *Quercus acutissima* stands were 889 kg, 2,266 kg, 851 kg, and 1,121 kg at 0–30 cm soil depth, respectively (Table 6, 7, 8, and 9). The net fine root carbon production was more than in *Pinus rigida* plantation forests of Korea (671 kg at 0–20 cm; Hwang, 1997) and more or lower than in Northeast hardwood stands of U.S.A. (1,100 kg ha⁻¹; Burke and Raynal, 1994). The net fine root carbon production in the four study stands was also higher at 0–30 cm soil depth than in other soil depths.

As shown in Tables 6, 7, 8 and 9, the net fine root carbon production 0–30 cm soil depth was highest in *Pinus koraiensis* stand among the study stand.

Total fine root NPP at 0–90 cm were 1,731 kg, 3,092 kg, 2,249 kg and 2,068 kg in *Pinus densiflora*, *Pinus koraiensis*, *Larix leptolepis* and *Quercus acutissima* stands, respectively (Tables 6, 7, 8 and 9). These were very similar to those for Northeast hardwood stands of U.S.A. (2,000–2,500 kg at 0–100 cm soil depths; Raynal and Burke, 1994). Park (2001) reported that net primary carbon production by leaves was 1,100 kg, 1,200 kg, and 1,200 kg in *Quercus mongolica* stands in Gongju, Pohang, and Yanyang area of Korea. This result indicates the important role of fine root to carbon cycle in for-

Table 6. Net fine root carbon production (kg ha⁻¹ yr⁻¹) in *Pinus densiflora* stand of Gongju area at 0–30 cm, 30–60 cm, and 60–90 cm soil depths

Soil depth (cm)	22 Apr. ~20 May.	20 May. ~20 Jun.	20 Jun. ~22 Jul.	22 Jul. ~22 Aug.	22 Aug. ~21 Sep.	21 Sep. ~20 Oct.	20 Oct. ~23 Nov.	Total
0–30	0	212	330	0	0	0	347	889
30–60	0	91	105	0	0	0	435	630
60–90	0	0	95	0	0	0	116	212
0–90								1731

Table 7. Net fine root carbon production (kg ha⁻¹ yr⁻¹) in *Pinus koraiensis* stand of Gongju area at 0–30 cm, 30–60 cm, and 60–90 cm soil depths

Soil depth (cm)	22 Apr. ~20 May.	20 May. ~20 Jun.	20 Jun. ~22 Jul.	22 Jul. ~22 Aug.	22 Aug. ~21 Sep.	21 Sep. ~20 Oct.	20 Oct. ~23 Nov.	Total
0–30	0	309	0	1333	0	130	493	2266
30–60	0	130	0	167	0	0	142	440
60–90	140	14	14	219	0	0	0	387
0–90								3092

Table 8. Net fine root carbon production ($\text{kg ha}^{-1} \text{ yr}^{-1}$) in *Larix leptolepis* stand of Gongju area at 0–30 cm, 30–60 cm, and 60–90 cm soil depths

Soil depth (cm)	22 Apr. ~20 May.	20 May. ~20 Jun.	20 Jun. ~22 Jul.	22 Jul. ~22 Aug.	22 Aug. ~21 Sep.	21 Sep. ~20 Oct.	20 Oct. ~23 Nov.	Total
0–30	14	439	0	344	0	0	0	851
30–60	0	244	0	330	0	142	0	717
60–90	293	0	26	65	0	298	0	681
0–90								2249

Table 9. Net fine root carbon production ($\text{kg ha}^{-1} \text{ yr}^{-1}$) in *Quercus acutissima* stand of Gongju area at 0–30 cm, 30–60 cm, and 60–90 cm soil depths

Soil depth (cm)	22 Apr. ~20 May.	20 May. ~20 Jun.	20 Jun. ~22 Jul.	22 Jul. ~22 Aug.	22 Aug. ~21 Sep.	21 Sep. ~20 Oct.	20 Oct. ~23 Nov.	Total
0–30	249	0	0	872	0	0	0	1121
30–60	0	0	0	344	0	130	0	475
60–90	153	0	0	319	0	0	0	472
0–90								2068

Table 10. Fine root turnover rate (%/yr) in *Pinus densiflora*, *Larix leptolepis*, *Pinus koraiensis*, and *Quercus acutissima* Stand in Gongju area at 0–30cm, 30–60cm, and 60–90cm soil depths.

Soil depth (cm)	<i>Pinus densiflora</i>	<i>Pinus koraiensis</i>	<i>Larix leptolepis</i>	<i>Quercus acutissima</i>
0–30	0.83	1.68	0.59	0.86
30–60	1.63	1.46	1.58	1.07
60–90	0.85	2.24	2.63	1.88

est ecosystems like other reports (Hansen, 1993; Cooper, 1983; Burke and Raynal, 1994). As shown in Tables 6, 7, 8 and 9, the total net fine root carbon production in 0–90 cm soil depth was highest in *Pinus koraiensis* stand among the study stand.

Fine root turnover rate (yr) was ranged from 0.59 to 2.63 (Table 10). This rate was in the range of the rate reported by Vogt and Bloomfield (1991). In this study, the fine root turnover rates were lower at 0–30 cm soil depth than the other soil depths except *Pinus koraiensis* stand. It was reported that low rates of fine root turnover was on relatively poor site (Mooney and Gulman, 1982). However, fine root turnover may vary with site quality and species composition (Shaver and Billings, 1975).

REFERENCES

- Burke, M. K. and D. J. Raynal 1994 Fine root growth phenology, production, and turnover in a northern hardwood forest ecosystem. *Plant and Soil* **162**: 135–146
- Cooper, C. F. 1983 Carbon storage in managed forests. *Can. J. For. Res.* **13**: 155–166
- Fairly, R. I. and I. J. Alexander 1985 Methods of calculating fine root production in forests. In: Ecological interactions in the soil. special publication no. 4 of the British Ecol. Soc. pp. 37–41. London
- Hair, D. and R. N. Sampson 1992 Climate change—history. Prospects, and possible impacts. pp. 1–10 In: Sampson, R. N. and Hair, D. (eds.) Forests and Global Change. Vol. I: Opportunities for Increasing Forest Cover. Am. For. Washington, D. C.
- Hansen, E. A. 1992 Soil carbon sequestration beneath hybrid poplar plantations. Nor. Cen. For. Exp. Sta. USDA For. Serv. For. Sci. Lab. Grand Rapids, MN, 55744: p. 13
- Hwang, J.-H. 1996 Effects of fertilization on fine root dynamics in pitch pine and Japanese Larch Plantations. Master Thesis, Korea University
- Laurenroth, W. K., H. W. Hunt, D. M. Swift and J. S. Singh 1998 Reply to Vogt *et al.* *Ecology* **67**: 580–582
- Makkonen, K. and H. S. Helmisaari 1999 Assessing fine root biomass and production in a Scots pine stand—comparison of soil core and root in growth core methods. *Plant and Soil* **210**: 43–50
- McClagherty, C. A., J. D. Aberand and J. M. Melillo 1982 Decomposition dynamics of fine roots in forested ecosystems. *Oikos* **42**: 378–386
- Nadelhoffer, K. J. and J. W. Raich 1992 Fine root production estimates and belowground carbon allocation in forest ecosystems. *Ecology* **73**: 1139–1147
- Park, G. S. and S. W. Lee 2001 Biomass and net primary production of *Quercus variabilis* natural forest ecosystems in Gongju, Pohang, and Yangtang of Korea. *J. Korean For. Soc.* **90**: 692–698
- Schlesinger, W. H. 1991 Biogeochemistry; an Analysis of Global change. Academic press, N. Y. p. 443
- Shaver, G. R. and W. D. Billings 1975 Root production and root turnover in a wet Tundra ecosystem, Barrow, Alaska. *Ecology* **56**: 401–409
- Singh, J. S., W. K. Laurenroth, H. W. Hunt and D. M. Swift 1984 Bias and random errors in estimators of net root production: a simulation approach. *Ecology* **65**: 1760–1764
- Vogt, K. A., C. C. Grier, S. T. Bower, D. G. Sprugel, and D. J. Vogt 1986 Overestimation of net root production: a real or imaginary problem? *Ecology* **67**: 577–579